

Global topology transport from monotonicity
and singularity of complex manifold.
Component of manifold in differential and integral
from quantum level of deprivate equation.

Masaaki Yamaguchi

Global assemble manifold defined with infinity of differential fields to determine of definition create from Euler product and Gamma function for Beta function being potten from integral manifold system. Therefore, this defined from global assemble manifold from being Gamma of global equation in topological expalanations.

$$\begin{aligned}
\int f(x)dx &= \int \Gamma(\gamma)' dx_m \\
&= 2(\cos(ix \log x) - i \sin(ix \log x)) \\
\left(\frac{\int f(x)dx}{\log x} \right) &= \lim_{\theta \rightarrow \infty} \left(\frac{\int f(x)dx}{\theta} \right) = 0, 1 \\
e^{i\theta} &= \cos \theta + i \sin \theta \\
\left(\int f(x)dx \right)' &= 2(i \sin(ix \log x) - \cos(ix \log x)) \\
&= 2(-\cos(ix \log x) + i \sin(ix \log x)) \\
&= (\cos(ix \log x) - i \sin(ix \log x))' \\
&= \frac{d}{de^{i\theta}} ((\cos, -\sin) \cdot (\sin, \cos)) \\
\int \Gamma(\gamma)'' dx_m &= \left(\int \Gamma(\gamma)' dx_m \right)^{\nabla L} = \left(\int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma \right)^{\nabla L} \leq \left(\int \Gamma dx_m + \frac{d}{d\gamma} \Gamma \right)^{\nabla L} \leq (e^f - e^{-f} \leq e^{-f} + e^f)' \\
&= 0, 1
\end{aligned}$$

Then, the defined of global integral and differential manifold from being assembled world lines and partial equation of deprivate formula in Homology manifold and gamma function of global topology system, this system defined with Shwaltsschild cicle of norm from black hole of entropy exsented with result of monotonicity for constant of equations. Therefore, this calicurate function resore with result of seed of two in three manifold of dimension.

川島隆太先生が、私みたいな人が、頭の中で偏微分や多重積分、部分積分、共変微分や、置換積分や、複素微分や複素積分を考えていると思っていたと言っていたが、それ以上の大域的微分や大域的 2 重微分や、大域的多重積分や大域的偏微分や大域的部分積分、微分幾何の量子化、単体積分、単体微分などの、大学院生でも考えない、思いもしない、実際の式を書くと誰でも、その関数と多様体が存在すると思うしかできない、私の記述式を見ると納得する理論式でもあると、私も自慢できる理論であると自負できる。

$$\log x|_{g_{ij}}^{\nabla L} = f^{f'}, F^f|_{g_{ij}}^{\nabla L} : x \rightarrow y, x^p \rightarrow y$$

$$f(x) = \log x = p \log x, f(y) = p \log x$$

It's defined with global partial deprivate formula and assemble manifold.

$$\frac{d^2}{df^2} F = F^{f'} \cdot f^{f''},$$

$$\int \int F dx_m = F^f \cdot F^{(f)'}$$

$$\frac{d}{df dg}(f, g) = (f \cdot g)^{f' + g'}$$

And, global parcial integral equation also established with global topology computations. therefore, this circutation exclude for all of topology extension ideas.

$$(F^f \cdot G^g) = \int (f \cdot g)^{f' + g'}$$

$$\int \int F \cdot G dx_m = [F^f \cdot G^g] - \int (f \cdot g)^{f' + g'}$$

$$\int \frac{d}{df dg} FG = \int F^{f'} G + \int FG^{g'}$$

$$\int F^{f'} G dx_m = [F^f G^g] - \int FG^{g'} dx_m$$

Global quato algebra computations also,

$$\left(\frac{F(x)}{G(x)} \right)^{(fg)'} = \frac{F^{f'} G - FG^{g'}}{G^g}$$

Global partial manifold explain with reduction of operator for being represented from partial equation of references.

$$\frac{d}{df dg} FG = \frac{d}{df_m} FG$$

$$\frac{\partial}{\partial f_m} FG = F^{f^{\mu\nu}} \cdot G + F \cdot G^{g^{\mu\nu}}, \int F dx_m = F^f$$

Entorpy computation from integrate for being manifold of global deprivate and integrate formulas. Gamma function sented for Beta function of inspiration with monitonicity of component with differential and integral being definitions.

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

This area defined with different and integrate of component with global topology.

$$t = \Gamma(x)$$

$$T = \int \Gamma(x) dx$$

for this system used for being is combine with beta function for component of deprive equation.

$$\beta(p, q) = - \int \frac{1}{t^2} dt$$

This point be for global differential variable exchanged,

$$T_m = \int \Gamma(x) dx_m$$

This also point be for being retried from ordinary differential computation for component of integral manifold.

$$T' = \frac{t'}{\log t} dt + C (C \text{ は積分定数})$$

This exceed of proof being for being defined with deprive variable of global differential formula, this succeeded for true is,

$$dx_m = \frac{1}{\log x} dx, dx_m = (\log x^{-1})'$$

Therefore, this exchanged from monotonicity deprivation from global partial integral manifold is able to,

$$T' = \int \Gamma(\gamma)' dx_m$$

After all, these exchanged of monotonicity deprivation succeeded from being catastrophe of summative of partial and assemble of deprivations for differential geometry of quantum level to global integral and differential manifold.

$$\bigoplus T^\nabla = \int T dx_m, \delta(t) = t dx_m$$

this excluded of being conclusion are which beta function evaluate with mononicity from ordinary differential equation be resulted from component of deprivation and integral explanations. This circulation be resembled to define with global topology of extension external of deprive and integral of manifolds estourced with quantum level of differential geometry be proofed with all of equation anbraced from Euler product.

虚数の虚数倍した値が超越数の x 倍と同じとすると、超越数の $2\pi m$ 倍が $n = 1$ で i となると定義すると、次の式たちが導かれる。 Imaginary pole circulate with twigled of pole in step function from Naipia number of assembled from equalation of defined are escorted to be defined with next equations.

These defined equation are climbate with idea of equation from Caltan of imaginary number of circulation.

$$i^i = (\sqrt{i})^{\sqrt{i}} = e^{x \log x}$$

$$e^x = i, e^{2\pi m} = i^n, \frac{d}{dx} e^{2\pi m} = i^n$$

$$e^{i\theta} = \cos \theta + i \sin \theta, e^{2\pi m} = i^n$$

$$2\pi m = n, e = i(e = i \text{ と } i \text{ となる } e^{i\theta} = \cos \theta + i \sin \theta \text{ の範囲で } 2\pi m \text{ がある。})$$

$$\frac{d}{di} i = i^i = e^{x \log x}, m = \frac{1}{2\pi}, l = 2\pi r, \pi = \frac{l}{2r}$$

$$||ds^2|| = e^{-2\pi T||\psi||} [\eta_{\mu\nu} + \bar{h}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

$$e^x = i, e^{2\pi m} = i^n, e = i$$

$$[\eta_{\mu\nu} + \bar{h}(x)] dx^\mu dx^\nu / i = H\Psi = i\hbar\psi, H\Psi = \frac{1}{i}[H, \Psi]$$

Hisenbelg equation are repressed from AdS5 manifold to particle level equalation of quantum levl of differential geometry entranced.

$$\bigoplus (i\hbar\nabla)^{\oplus L} = e^{ix \log x^{e^{ix \log x}'}}$$

$$\Gamma = \int e^{-x} x^{1-t} dx$$

$$\gamma = \int e^{-x} x^{1-t} \log x dx$$

$$= \int \Gamma(\gamma)' dx_m$$

$$= \frac{d}{d\gamma} \Gamma$$

Eight differential geometry are each intersect with own level of concept from expalanation of Euler product system. This component of three manifold sergeried with geometry of destroy and desect with time element of Stokes equation. This manifold gut theory from described with zeta function to catastrophe for non tree of routs result on sergery of space system. 各微分構造においての時間発展での熱エネルギーの変化は $E = mc^2, E = m^2c^2$ の three manifold が微分幾何構造の変化にそれぞれ対応する。Each geometry point interacte with exchange of three manifold in Seifert structure from special relativity of equation for heat energy fluentations on time developed from this extermaination.

$$\nabla(i\hbar\nabla)^{\oplus L}, \bigoplus (i\hbar\nabla)^{\oplus L}, \square(i\hbar\nabla)^{\oplus L}$$

These operator of equation on summatative manifolds from emerged with element of particle conclution, Euler product is resulted from this operator expalanation of locality insectations. Moreover, this eight geometry of differential operator are constructed with four pattern of Jones manifold from summatative formula and this system extate with special relativity references. This decieved of elemet on summatative equation routed to internext in real and imaginary pole on complex dimension, and this dimension explation with Stokes theorem defined too. And this defined circutation of Yacobi matrix is repressed with Knot theory with anstate with Jones manifold. その上に、8種類の微分幾何は、Jones 多項式の4パターンでの構成される差分と加群方程式から、特殊相対性理論をも思わせる、この差分方程式

$$\frac{d}{d\gamma} \Gamma = e^f + e^{-f} \geq e^f - e^f$$

から、8種類の代数幾何が、ストークスの定理と同じく、この Jones 多項式の固有周期が、すべての時間の流れの基軸をも内包していることが、わかる。この Jones 多項式は、結び目の理論をも、この周期が言い表してもいる。

$$\begin{aligned}\frac{d}{d\gamma}\Gamma + \int C dx_m &= 2(\cos(ix \log x) - i \sin(ix \log x)) \\ e^{-\theta} &= \cos(ix \log x) - i \sin(ix \log x) \\ \int \Gamma(\gamma)' dx_m &= 2(\cos(ix \log x) - i \sin(ix \log x))\end{aligned}$$

カルーツァ・クライン空間の方程式は、

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \psi^2(x)(dx^2 + \kappa^2 A_\mu(x)dx^\nu)^2$$

この式は、

$$\begin{aligned}ds^2 &= -N(r)^2 dt^2 + \psi^2(x)(dr^2 + r^2 d\theta^2) \\ ds^2 &= -dt^2 + r^{-8\pi G m}(dr^2 + r^2 d\theta^2)\end{aligned}$$

とまとめ、

$$\begin{aligned}dx^2 &= g_{\mu\nu}(x)(g_{\mu\nu}(x)dx^2 - dx g_{\mu\nu}(x)) \\ dx &= (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x)dx g_{\mu\nu}(x))^{\frac{1}{2}}\end{aligned}$$

と反重力と正規部分群の経路和となり、

$$\pi(\chi, x) = i\pi(\chi, x)f(x) - f(x)\pi(\chi, x)$$

と基本群にまとまる。シュワルツシルト半径は、

$$ds^2 = -(1 - \frac{r_s}{r})c^2 dt^2 + \frac{1}{1 - \frac{r_s}{r}} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\psi^2$$

これは、カルーツァ・クライン空間と同型となる。一般相対性理論は、

$$R^{\mu\nu} + \frac{1}{2}g_{ij}\Lambda = \kappa T^{\mu\nu}$$

この多様体積分は、

$$\int \kappa T^{\mu\nu} \mathrm{dvol} = \int \left(R^{\mu\nu} + \frac{1}{2}g_{ij}\Lambda \right) \mathrm{dvol}$$

ガンマ関数のおける大域的微分多様体は、これと同型により、

$$\begin{aligned}\int \Gamma \cdot \frac{d}{d\gamma} \Gamma dx_m &\leq \int \Gamma dx_m + \frac{d}{d\gamma} \Gamma \\ &= \int \Gamma(\gamma)' dx_m\end{aligned}$$

オイラーの定数の多様体積分は、

$$\int \left(\int \frac{1}{x^s} dx - \log x \right) \mathrm{dvol}$$

この解は、シュワルツシルト半径と同型より、

$$= e^f - e^{-f} \leq e^f + e^{-f}$$

次元の単位は多様体より、大域的微分多様体とオイラーの定数の多様体積分の加群分解は、オイラーの公式の三角関数の虚数の度解より、

$$\frac{d}{df}F + \int C dx_m = 2(\cos(ix \log x) - i \sin(ix \log x))$$

すべては、大域的微分多様体の重力と反重力方程式に行き着き、

$$\frac{d}{df}F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

と求まる。

8種類の微分幾何での、それぞれの時間の固有値の動静は、ストークスの流体力学での、熱エネルギーの各流れの仕方で表されている。種数1の2種が種数0の球体に両辺で交わっていると、閉3次元多様体は、種数は2となり、固有値とホモロジー多様体、特異点が同時に計算できる結果にもいけることがわかる。

一般相対性理論における重力理論は大域的微分多様体と積分多様体についての単体量を求めるための空間の対数関数における不変性を記述するプランクスケールと異次元の宇宙におけるスケール、ウィークスケールと言われている anti-D-brane として、この2種類の計量をガンマ関数とベータ関数としてオイラーの定数とそれによる連分数が微分幾何の量子化と数式の値を求めると同じという予知と推測値から、広中平祐定理の4重帰納法のオイラーの公式による多様体積分と同類値として、ペレルマン多様体がサーストン空間に成り立つと同じく、広中平祐定理がヒルベルト空間で成り立つとしての、これら2種の多様体の場の理論が、ゼータ関数が AdS5 多様体で成り立つことと、不変量として、ゼータ関数がこの3種の多様体の場の理論のバランスをとる理論として言えることである。

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

This norm is component of fields in Hilbert manifold of space theorem rebuilt with Mebius space in Gamma Function on Beta function fill into power of rout fundamental group.

Gravity of general relativity theory describe with Cutting of space in being discatastrophed from Global differential and integral manifold of scaled levetivity in plank scal and the other vector of universe scale, these two scale inspectivity of Gamma and Beta function escourted with these manifold experted in result of Differentail geometry of quantum level manifold equal with Euler product and continue parameter, moreover this product is relativity of Hironaka theorem in Four assembled integral of Euler equation.

重力場理論の式は、Gravity equation is

$$\square = -\frac{16\pi G}{c^4} T^{\mu\nu}$$

This equation quated with being logment of formula, and this formula divided with universe of number in prime zone, therefore, this dicided with varint equation is monotonicity of being composited with Weil's theorem united for Gamma function. この式は、対数を宇宙における数により求める素数分布論として、この大域的積分分断多様体がガンマ関数をヴェイユ予想を根幹とする単体量として決まることに起因する商代数として導かれる。

$$\not\square = \frac{8\pi G}{c^4} T^{\mu\nu} / \log x$$

$$\frac{d}{d\gamma}\Gamma$$

$$= [i\pi(\chi, x), f(x)]$$
$$\frac{d}{d\gamma}\Gamma = \int C dx_m = \int (\int \frac{1}{x^s} dx - \log x) d\text{vol}$$
$$\bigoplus (i\hbar^\nabla)^{\oplus L}$$

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$= C$$
$$C = b_0 + \frac{c_1}{b_1 + \frac{c_2}{b_2 + \frac{c_3}{b_3 + \frac{c_4}{b_4 + \dots}}}}$$
$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$(2.71828)^{2.828} = a^{a+b^b+c^c+d^d\ldots}$$

$$= \int e^f \cdot x^{1-t} dx$$

This represent is Gamma function in Euler product. Therefore this product is zeta function of global differential equation.

$$\begin{aligned} \frac{d}{df} F(v_{ij}, h) &= \int e^{-f} dV [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) \left(\frac{v}{2} - h \right)] \\ &= \frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm \end{aligned}$$

これらの方程式は 8 種類の微分幾何の次元多様体として、そして、これらの多様体は曲平面による双対性をも生成している。そして、このガウスの曲平面は、大域的微分多様体と微分幾何の量子化から素因数を形成している。These equation are eight differential geometry of dimension calyement. And these calyement equation excluded into pair of dimension surface. This surface of Gauss function are global differential manifold, and differential geometry of quantum level.

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

微分幾何の量子化はオイラーの定数とガンマ関数が指数による連分数としての不変性として素因数を形成していて、このガウスの曲平面による量子力学における重力場理論は、ダランベルシアン of 切断多様体がこの大域的切断多様体を付加している。Differential geometry of quantum level constructed with Euler product and Gamma function being discatastrophed from continued fraction style. Gravity equation lend with varint of monotonicity of level expressed from gravity of letter varient formula. これらの方程式は基本群と大域的微分多様体をエスコートしている。ベータ関数とガンマ関数がこれらのフォームラの方程式を放出している。結果、これらの方程式は広中平祐定理の複素多様体とグリーシャ教授によるペレルマン多様体からサポートされている。この 2 名の教授は、一つは抽象理論をもう一方は具象理論を説明としている。These equation escorted into Global differential manifold and fundemental equation. Weil's theorem is imported from this equation in gravity of letters. Beta function and Gamma function are excluded with these formulas. These equation comontius from Heisuke Hironaga of complex manifold and Gresha professor of Perelman manifold. These two professors are one of abstract theorem and the other of visual manifold theorem.

1 Hilbert manifold in Mebius space

this element of Zeta function on integrate of fields

Hilbert manifold equals with Von Neumann manifold, and this fields is concluded with Glassman manifold, the manifold is duality of twister into created of surface built. These fields is used for relativity theorem and quantum physics.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

This norm is component of fields in Hilbert manifold of space theorem rebuilt with Mebius space in Gamma Function on Beta function fill into power of rout fundamental group. And this result with AdS_5 space time in Quantum caos of Minkowsky of manifold in abel manifold. Gravity of metric in non-commutative equation is Global differential equation conbult with Kaluza-Klein space. Therefore this mechanism is $T^{\mu\nu}$ tensor is equal with $R^{\mu\nu}$ tensor. And This moreover inspect with laplace operator in stimulate with sign of differential operator. Minus of zone in Add position of manifold is Volume of laplace equation rebuilt with Gamma function equal with summative of manifold in Global differential equation, this result with setminus of zone of add summative of manifold, and construct with locality theorem straight with fundamental group in world line of surface, this power is boson and fermion of cone in hyper function.

$$V(\tau) = [f(x), g(x)] \times [f^{-1}(x), h(x)]$$

$$\Gamma(p, q) = \int e^{-x} x^{1-t} dx$$

$$= \beta(p, q)$$

$$= \pi(f(\chi, x), x)$$

$$||ds^2|| = \mathcal{O}(x)[(f(x) \circ g(x))^{\mu\nu}] dx^\mu dx^\nu$$

$$= \lim_{x \rightarrow \infty} \sum_{k=0}^{\infty} a_k f^k$$

$$G^{\mu\nu} = \frac{\partial}{\partial f} \int [f(x)^{\mu\nu} \circ G(x)^{\mu\nu} dx^\mu dx^\nu]^{\mu\nu} dm$$

$$= g_{\mu\nu}(x) dx^\mu dx^\nu - f(x)^{\mu\nu} dx^\mu dx^\nu$$

$$[i\pi(\chi, x), f(x)] = i\pi f(x) - f(x)\pi(\chi, x)$$

$$T^{\mu\nu} = (\lim_{x \rightarrow \infty} \sum_{k=0}^{\infty} \int \int [V(\tau) \circ S^{\mu\nu}(\chi, x)] dm)^{\mu\nu} dx^\mu dx^\nu$$

$$G^{\mu\nu} = R^{\mu\nu} T^{\mu\nu}$$

$$\begin{vmatrix} D^m & dx \\ dx & \sigma^m \end{vmatrix} \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix} \begin{vmatrix} x \\ y \end{vmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\sigma^m \begin{bmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{bmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

$$V(M) = \frac{\partial}{\partial f} ({}^N \int [f \setminus M]^{\oplus N})^{\mu\nu} dx^\mu dx^\nu$$

$$V(M) = \pi(2 \int \sin^2 dx) \oplus \frac{d}{df} F^M dx_m$$

$$\lim_{x \rightarrow \infty} \sum_{k=0}^{\infty} a_k f^k = \int (F(V) dx_m)^{\mu\nu} dx^\mu dx^\nu$$

$$\begin{aligned}
\bigoplus_{k=0}^{\infty} [f \searrow g] &= \vee(M \wedge N) \\
\pi_1(M) &= e^{-f^2 \int \sin^2 x dm} + O(N^{-1}) \\
&= [i\pi(\chi, x), f(x)] \\
M \circ f(x) &= e^{-f \int \sin x \cos x dx_m} + \log(O(N^{-1}))
\end{aligned}$$

Non-Symmetry space time.

$$\begin{aligned}
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
\epsilon S(\nu) &= \square_v \cdot \frac{\partial}{\partial \chi} (\sqrt[5]{\sqrt{\wedge g^2}}) d\chi
\end{aligned}$$

Differential Volume in AdS_5 graviton of fundamental rout of group.

$$\wedge(F_t^m)'' = \frac{1}{12} g_{ij}^2$$

Quarks of other dimension.

$$\pi(V_\tau) = e^{-(\sqrt{\frac{\pi}{16}} \log x)^\delta} \times \frac{1}{(x \log x)}$$

Universe of rout, Volume in expanding space time.

$$\begin{aligned}
\frac{d}{dt} (g_{ij})^2 &= \frac{1}{24} (F_t^m)^2 \\
m^2 &= 2\pi T \left(\frac{26 - D_n}{24} \right)
\end{aligned}$$

This quarks of mass in relativity theorem, and fourth of universe in three manifold of one dimension surface, and also this integrate of dimension in conbult of quarks.

$$g_{ij} \wedge \pi(\nu_\tau) = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\psi^2$$

Out of rout in AdS_5 space time.

$$||ds^2|| = g_{ij} \wedge \pi(\nu_v)$$

AdS_5 norm is fourth of universe of power in three manifold out of rout.

Sphere orbital cube is on the right of hartshorne conjecture by the right equation, this integrate with fourth of piece on universe series, and this estimate with one of three manifold. Also this is blackhole on category of symmetry of particles. These equation conbult with planck volume and out of universe is phologram field, this field construct with electric field and magnitic field stand with weak electric theorem. This theorem is equals with abel manifold and AdS_5 space time. Moreover this field is anti-brane and brane emerge with gravity and antigravity equation restructed with. Zeta function is existed with Re pole of $\frac{1}{2}$ constance reluctances. This pole of constance is also existed with singularity of complex fields. Von Neumann manifold of field is also this means.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

$$\begin{aligned}
e^{i\theta} &= \cos \theta + i \sin \theta \\
e^{x \log x} &= x^{\frac{1}{2}+iy}, x \log x = \log(\cos \theta + i \sin \theta) \\
&= \log \cos \theta + i \log \sin \theta
\end{aligned}$$

This equation is developed with frobenius theorem activate with logment equation. This theorem used to

$$\begin{aligned}
\log(\sin \theta + i \cos \theta) &= \log(\sin \theta - i \cos \theta) \\
\log\left(\frac{\sin \theta}{i \cos \theta}\right) &= -2R_{ij}, \frac{d}{dt}g_{ij}(t) = -2R_{ij}
\end{aligned}$$

$$\mathcal{O}(x) = \frac{\zeta(s)}{\sum_{k=0}^{\infty} a_k f^k}$$

$$\text{Im} f = \ker f, \chi(x) = \frac{\ker f}{\text{Im} f}$$

$$H(3) = 2, \nabla H(x) = 2, \pi(x) = 0$$

This equation is world line, and this equation is non-integrate with relativity theorem of rout constance.

$$[f(x)] = \infty, ||ds^2|| = \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

$$T^2 d^2\psi = [f(x)], T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

These equation is equals with M theorem. For this formula with zeta function into one of universe in four dimensions. Sphere orbital cube is on the surface into AdS_5 space time, and this space time is Re pole and Im pole of constance reluctances.

Gauss function is equals with Abel manifold and Seifert manifold. Moreover this function is infinite time, and this energy is cover with finite of Abel manifold and Other dimension of seifert manifold on the surface.

Dilaton and fifth dimension of seifert manifold emerge with quarks and Maxwell equation, this power is from dimension to flow into energy of boson. This boson equals with Abel manifold and AdS_5 space time, and this space is created with Gauss function.

Relativity theorem is composited with infinite on D-brane and finite on anti-D-brane, This space time restructured with element of zeta function. Between finite and infinite of dimension belong to space time system, estrald of space element have with infinite oneselves. Anti-D-brane have infinite themselves, and this comontend with fifth dimension of AdS_5 have for seifert manifold, this asperal manifold is non-move element of anti-D-brane and move element of D-brane satisfid with fifth dimension of seifert algebra liner. Gauss function is own of this element in infinite element. And also this function is abel manifold. Infinite is coverd with finite dimension in fifth dimension of AdS_5 space time. Relativity theorem is this system of circumstance nature equation. AdS_5 space time is out of time system and this system is belong to infinite mercy. This hiercyent is endrol of memolite with geniue of element. Genie have live of

telomea endore in gravity accesorlity result. AdS5 space time out of over this element begin with infinite assentance. Every element acknowlege is imaginary equation before mass and spiritual envy.

$$\begin{aligned}
T^2 d^2 \psi &= \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k \\
\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k &= [T^2 d^2 \psi] \\
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M ({}^5\sqrt{x^2}) d\Lambda + \frac{d}{df} \int \int_M {}^N({}^3\sqrt{x})^{\oplus N} d\Lambda \\
{}^M(\vee(\wedge f \circ g)^N)^{\frac{1}{2}} &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
||ds^2|| &= \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2 \psi \\
\mathcal{O}(x) &= e^{-2\pi T|\psi|} \\
G^{\mu\nu} &= R_{\mu\nu} T^{\mu\nu} \\
&= -\frac{1}{2} \Lambda g_{ij}(x) + T^{\mu\nu}
\end{aligned}$$

Fifth dimension of eight differential structure is integrate with one geometry element, Four of universe also integrate to one universe, and this universe represented with symmetry formula. Fifth universe also is represented with seifert manifold peices. All after universe is constructed with six element of circuatance. This aquire maniculate with quarks of being esperaled belong to.

2 Imaginary equation in AdS5 space time create with dimension of symmetry

D-brane and anti-D-brane is composited with all of series universe emerged for one geometry of dimension, this gravity of power from D-brane and anti-D-brane emelite with ancestor. Seifert manifold is on the ground of blackhole in whitehole of power pond of senseivility. Six of element of quarks and universe of pieces is supersymmetry of mechanism resolved with

hyper symmetry of quarks constructed to emerge with darkmatter. This darkmatter emerged with big-ban of heircyent in circumstance of phenomena.

D-brane and anti-D-brane equations is

$$\begin{aligned}
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
C &= \int \int \frac{1}{(x \log x)^2} dx_m
\end{aligned}$$

Euler constance is quantum group theorem rebuild with projective space involved with.

虚数方程式は、反重力に起因するフーリエ級数の励起を生成する。それは、人工知能を生み出す、5次元時空にも、この虚数方程式は使われる。AdS5 の次元空間は、反ド・シッター時空の D-brane と anti-D-brane

の conformal 場を生み出す。ホログラフィー時空は、この量子起因によるものである。2次元曲面によるブラックホールは、ガンマ線バーストによる5次元時空の構造から観測される。空間の最小単位によるプランク定数は、宇宙の大域的微分多様体から導き出される、AdS5の次元空間の準同型写像を形成している。これは、最小単位から宇宙の大きさを導いている。最大最小の方程式は、相加相乗平均を形成している。時間と空間は、宇宙が生成したときから、宇宙の始まりと終わりを既に生み出している。宇宙と異次元から、ブラックホールとホワイトホールの力がわかり、反重力を見つけられる。オイラーの定数は、この量子定数からわかる。虚数の仕組みはこの量子スピンの産物である。オイラーの定数は、この虚時空の斥力の現存である。それは、非対称性理論から導かれる。不確定性原理は、AdS5のブラックホールとホワイトホールを閉3次元多様体に統合する5次元時空から求められる。位置と運動エネルギーが、空間の最小単位であるプランク定数を宇宙全体にする微細構造定数からわかり、面積確定から、アーベル多様体を母関数に極限值として、ゼータ関数をこの母関数に不変式として、2種類ずつにまとめる4種類の宇宙を形成する8種類のサーストンの幾何化予想から導き出される。この閉3次元多様体は、ミラー対称性を軸として、6種類の次元空間を一種類の異次元宇宙と同質ともしている。複素多様体による特異点解消理論は、この原理から求められる。この特異点解消理論は、2次元曲面を3次元多様体に展開していく、時空から生成される重力の密度を反重力と等しくしていく時間空間の4次元多様体と虚時空から求められる。ヒルベルト空間は、フォン・ノイマン多様体とグラスマン多様体をこのサーストンの幾何化予想を場の理論既定値として形成される。この空間は、ミンコフスキー時空とアーベル多様体全体を表している。そして、この空間は、球対称性を複素多様体を起点として、大域的トポロジーから、偏微分を作用素微分として時空間をカオスからずらすと5次元多様体として成立している。これらより、3次元多様体に2次元射影空間が異次元空間として、AdS5空間を形成される。偏微分、全微分、線形微分、常微分、多重微分、部分積分、置換微分、大域的微分、単体分割、双対分割、同調、ホモロジー単体、コホモロジー単体、群論、基本群、複体、マイヤー・ビートリス完全系列、ファン・カンペンの定理、層の理論、コホモルティズム単体、CW複体、ハウスドロフ空間、線形空間、位相空間、微分幾何構造、モースの定理、カタストロフの空間、ゼータ関数系列、球対称性理論、スピン幾何、ツイスター理論、双対被覆、多重連結空間、プランク定数、フォン・ノイマン多様体、グラスマン多様体、ヒルベルト空間、一般相対性理論、反ド・シッター時空、ラムダ項、D-brane, anti-D-brane, コンフォーマル場、ホログラム空間、ストリング理論、収率による商代数、ニュートンポテンシャルエネルギー、剛体力学、統計力学、熱力学、量子スピン、半導体、超伝導、ホイーストン・ブリッチ回路、非可換確率論、Connes理論、これら、演算子代数を形成している、微分・積分作用素が、ヒルベルト空間に存在している多様体の特質を全面に押し出して、いろいろな多様体と関数そして、群論を形作っている。

$$\begin{aligned} \begin{vmatrix} D^m & dx \\ dx & \sigma^m \end{vmatrix} \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix} \begin{vmatrix} x \\ y \end{vmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}} \\ \sigma^m \begin{bmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{bmatrix}^{\frac{1}{2}} &= \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \\ (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) & \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ = \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \end{aligned}$$

$$l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\left(\frac{\partial}{\partial \tau} f(x,y,z)\right)^{3'} = A^{\mu\nu}$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.