

$$\not{\Box}=\frac{8\pi G}{c^4}T^{\mu\nu}/\log x$$

$$\begin{array}{l} {}^t\overline{\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits}\mathrm{cohomD}_\chi[\mathrm{I}_\mathrm{m}]\\ =\oint (px^n+qx+r)^{\nabla l}\end{array}$$

$$\frac{d}{dl}L(x,y)=2\int ||\sin 2x||^2d\tau$$

$$\frac{d}{d\gamma}\Gamma$$

$$||ds^2||=\lim_{x\rightarrow\infty}[\delta(x)\int\int\int\pi\left(\sum_{k=0}^{\infty}\frac{{}^n\sqrt{p},x}{n}\right)^{\frac{1}{2}}d\tau]^{\mu\nu}$$

$$\oint\!\!\!\oint\cong ||ds^2||=\lim_{x\rightarrow\infty}[\delta(x)\int\int\int\pi\left(\sum_{k=0}^{\infty}\frac{{}^n\sqrt{p},x}{n}\right)^{\frac{1}{2}}d\tau]^{\mu\nu}$$

$$e^{-2\pi T||\psi||}[\eta+\bar{h}]dx^\mu dx^\nu+T^2d^2\psi$$

$$\frac{-16\pi G}{c^4}T^{\mu\nu}/\log x=\not{\Box}$$

$$\begin{array}{l} \frac{-16\pi G}{c^4}T^{\mu\nu}/e^{-2\pi T||\psi||}\\ =4\pi G\rho\end{array}$$

$$\frac{\partial}{\partial x\partial y}=\nabla_i\nabla_j$$

$$\Box\int\!\!\!\int\!\!\!\int={^t\overline{\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits\mathop{\mathfrak{f}}\nolimits}}$$

$$\frac{\partial}{\partial x}\int\!\!\!\int\!\!\!\int=\nabla_i\nabla_j\int\nabla f(x)d\eta$$

$$\frac{\partial}{\partial f}\int\!\!\!\int\!\!\!\int=\Box\int\!\!\!\int\!\!\!\int$$