

3次元多様体のエントロピー量

Masaaki Yamaguchi

オイラーの定数は、

$$\begin{aligned} C &= \int \frac{1}{x^s} dx - \log x \\ &= \int \left(\int \frac{1}{x^s} dx - \log x \right) \text{dvol} \\ &= \int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol} \end{aligned}$$

と表されていて、ヒッグス場方程式 + オイラーの定数 = ゼータ関数 と求まる。この方程式が、リッチフロー方程式にもなる。このリッチフロー方程式は、微分幾何の量子化でもあり、ガンマ関数とベータ関数、ガウスの曲面論、ヒッグス場、アインシュタインテンソル、オイラーの定数と全部の方程式が入っている。この方程式が、相加相乗平均となり、宇宙と原子となり、不確定性原理が宇宙の観測系が、宇宙では確率方程式と成り立っているが、異次元が合わさると、ノルムとしての3次元多様体の確率分布方程式でもあり、確率ではない、非線形方程式となっている。プランク長体積でもあるが、絶対的な方程式にもなっている。

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\ \int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} \text{dvol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\ &= e^{-f} - e^f \\ &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= 2 \cos(ix \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &+ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= \frac{d}{df} F = 2i \sin(ix \log x) \\ \frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\ &= 2e^{-\theta} = 2e^{-ix \log x} \\ &= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\log(x \log x) \geq 2\sqrt{y \log y}$$

$$\log(x \log x) = \log x + \log \log x$$

この宇宙と異次元でのブラックホールとホワイトホールでの、宇宙と原子の運動量と位置の絶対的な不確定性原理となり、3次元多様体の確率分布方程式が、運動量と位置エネルギーが両方観測できる式にもなっている。円周上が固定されているために、運動量と位置エネルギーが両方観測できる式になっている。カルーツァ・クライン理論とハイゼンベルク方程式が、両方入っている。

$$\nabla \psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla \psi^2 = 8\pi G \hbar + 8\pi \frac{V}{S}$$

ホワイトホールの原子とブラックホールの宇宙のエネルギーのエントロピー量でもある。宇宙がブラックホールとしてのヒッグス場方程式として、ホワイトホールが原子としてのプランク長体積でもある。

$$\square_v = 2\sqrt{2\pi G \hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

原子としてのブラックホールのエントロピー量でもある。

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

この式から宇宙が無重力であることが表されている。

$$2 \cos(ix \log x) + 2i \sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

これらの方程式から宇宙と異次元が warped passege になっている。:

Report of Quantum level of differential structure,
zeta function and Global differential equation Masaaki Yamaguchi, with my son

Higgs field + Euler constant = zeta function

$$m(x) + C = 2 \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\text{Euler constance} = \frac{d}{d\gamma} \Gamma^{-1} - \gamma^{(\gamma)'} = \left(\Gamma^{(\gamma)'} \right)^{-1} - \gamma^{\gamma'}$$

gravity + anti gravity = zeta function

Gamma function in Global differential equation developed with part integrate manifold cohomologite with helmader operator, and this subrace of differantial form start with first minimalized function, this operator emerged from quato algebra to fermion partial specture in Higgs fields.

$$\Gamma dx_m + \int \Gamma dx_m = \Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$\begin{aligned} & \frac{d}{d\gamma} \Gamma + \int \Gamma dx_m \\ &= \int f(g(x)) g'(x) dx \\ &= \left(\int f(g(x)) dx \right)' \\ & \nabla_i \nabla_j \int \nabla f(x) d\eta \end{aligned}$$

Global differential equation is partial integrate operator in resolved zone of frobenius theorem and this seminal concept with zone, Jones formula equation conquire with rico level theorem, this spectireum releafe in zeta function in CP symmetry broken, this broken from being integrate with blance into universe of zeta function, all of this renze equation into one class of Euler equation concerned with seminal concept in Jones formula equation, zeta function is proofed that this relativity level emerged from Quantum level of differential structure.

$$|f \circ g| \leq |f + g|$$

$$\frac{f}{\log x} = m(x)$$

$$\frac{\gamma}{\log x} = \frac{d}{d\gamma} \Gamma$$

$$\gamma = \Gamma^{(\gamma)'} \log x$$

$$\Gamma^{(\gamma)'} = \gamma (\log x)^{-1}$$

$$= r dx_m$$

$$\begin{aligned}
\Gamma(\gamma)' &= \left(\int e^{-x} x^{1-t} dx \right)^{\left(\int e^{-x} x^{1-t} \log x dx \right)'} \\
&= e^f \\
&= r dx_m = e^f \\
&\int \Gamma(\gamma)' dx_m \\
&\nabla_i \nabla_j \int \nabla \Gamma(\gamma) dx \\
&= \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma dx_m \\
&= \int \Gamma \cdot \frac{d}{d\gamma} \Gamma dx_m \\
&= \left(\int \Gamma dx_m \circ \frac{d}{d\gamma} \Gamma \leq \int \Gamma dx_m + \Gamma dx_m \right)' = e^{-f} \cdot e^f \leq e^{-f} + e^f \\
&\quad (1 \leq e^{-f} + e^f)' \\
&0 = -(e^{-f} - e^f) \\
&\quad \Gamma(\gamma)' + \Gamma(\gamma) \\
&\quad (= e^f + e^{-f})' \\
&\quad = -e^{-f} + e^f \\
&\quad = -(e^{-f} - e^f) \\
&\quad (R_{ij})' = -(R_{ij}) \\
&\int C dx_m = \int \left(\int \frac{1}{x^s} dx - \log x \right) dx \\
&= \int \int \frac{1}{x^s} d\text{vol} - \int \log x d\text{vol} \\
&\quad \Gamma dx_m = \gamma dx_m \\
&= \frac{d}{d\gamma} \Gamma^1 - (\gamma)^\gamma' \\
&= e^{-f} - e^f
\end{aligned}$$

Quantum level of differential structure equal with Gamma function and routed with Beta function, escorted into Euler constant and symmetry build with Higgs field, and proof with zeta function summarized with Poincaré and Riemann conjecture. Mathematician of Russia, America and Japan resolved with this conjecture. My son resolved with this conjecture build in concepted with quantum level of differential structure into zeta function. All of equation is integrate with manifold into Hilbert space, this manifold espectral from general relativity and quantum physics in manifold of Hilbert space, this manifold resolved with D-brane and Gauss surface into gravity and atom of element with this theory, this theory is integrate theorem released with one class of universe equation.

Explain in Global defferential equation and

Global integrate equation.

Varintegrate equation, and horizen cut of equations. Masaaki Yamaguchi

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x) \ll^p$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2 \psi] \ll^p \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x) \ll^p$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned} \int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f} \end{aligned}$$

$$\begin{aligned} \frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f \end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values. 大域的微分方程式に、X の X 乗のエントロピー不変量の微分量が計算に関わっている。外微分をこのエントロピー式に入れる。大域的積分多様体の多重積分は、多様体の階層によって、積分回数が決まっている。大域的微分多様体は、ニュートン形式とライプニッツ形式の外微分によって計算される。大域的積分多様体と大域的微分多様体は、逆操作とエントロピー不変量で計算できる。

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2 \\ E &= -\frac{1}{2} m v^2 + m c^2 \\ \lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m &\geq \frac{1}{2} i \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \frac{1}{2} i \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ &= (f)^{(f)'} \\ &= (f^f)' \\ &= e^{x \log x}\end{aligned}$$

$$\begin{aligned}\Gamma &= \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f} \\ \int x^{1-t} dx &= \frac{d}{df} F, \int e^{-x} dx = \int F dx_m\end{aligned}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, 自明な零点は実軸 $\frac{1}{2}$ に 1 を除いてある。 $\sin 0 = 0$ と

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ によって、不確定性原理の関係でもあり、粒子、電子、原子が確率分布になっているのも開集合で証明できる。宇宙と異次元の関係にもなっている。this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$= \Gamma(\Gamma \int \log x dx)'$$

$$= e^{-x \log x}$$

$$\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx$$

$$\begin{aligned}\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\ \frac{d}{df} F &= \int x^{s-1} dx \\ \int F dx_m &= \int e^{-x} dx \\ \frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. 大域的微分多様体と大域的積分多様体を、ガンマ関数とベータ関数の導出にも使われている。ガンマ関数の大域的微分多様体が、ニュートン方程式とダランベール方程式を商代数で求めると、同型の解に求まる。ゼータ関数と、この式の対極する量子群を商代数で求めても、同じ解が求まる。

Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

繰り込み理論を入れると発散を防げると言われているが、量子力学にはガウスの曲面論がないと言われている。

$$\begin{aligned}H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\ &= \bigoplus \frac{H\Psi}{\nabla L} \\ &= e^{x \log x} = x^{(x)'}\end{aligned}$$

however, とすると、

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry. 同じで量子力学で重力場方程式が表せられる。

$$\begin{aligned}\frac{d}{dt} \psi(t) &= \hbar \\ &= \frac{1}{2} i e^{i\hat{H}} \\ (i\hbar)' &= (-e^{i\hat{H}})' \\ &= -i e^{i\hat{H}} \\ \psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\ &= \left(\frac{1}{2}f\right)^{-if}\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

微分幾何の量子化は、ガンマ関数についての大域的微分方程式の解にもなっている。ハイゼンベルク方程式の大域的積分多様体の解にも、この微分幾何の量子化は、多様体の微分系の形になっている。 Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alsos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta\rightarrow 0}\begin{pmatrix}\sin\theta\\ \cos\theta\end{pmatrix}\begin{pmatrix}\theta&1\\ 1&\theta\end{pmatrix}\begin{pmatrix}\cos\theta\\ \sin\theta\end{pmatrix}=\begin{pmatrix}1&0\\ 0&-1\end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta, \mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

ガンマ関数の大域的微分方程式は、ガンマ関数の初期関数についての微分方程式でもあり、宇宙と異次元についてのビッグス場からのゼータ関数の生成も、3次元多様体の特異点定理になっている。宇宙と異次元の

片方でも同じ解にもなっている。逆三角関数の双曲多様体もヒッグス場の密度エネルギーの値と同じ解にもなっている。微分幾何の量子化も、ゼータ関数の対極する量子群と同じ解にもなっている。ゼータ関数は、量子群の方程式についての対極に位置する大域的微分多様体の解にもなっている。 Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements. 逆三角関数の双曲多様体は、ガンマ関数の方程式からの導出でのベータ関数の逆三角関数のエントロピー値にもなっている。

Explain in Global defferential equation and

Global integrate equation.

Varintegrate equation, and horizen cut of equations. Masaaki Yamaguchi with my son in acasic record of space

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x) \ll^p$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff} \text{Hom}[D^2\psi] \ll^p \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{ff} \text{cohom} D_k(x) \ll^p$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned} \int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f} \end{aligned}$$

$$\begin{aligned} \frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f \end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values.

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\int \frac{1}{(x \log x)} dx = i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx$$

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \frac{1}{2} x^2$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \int \int_M dx_m$$

$$\leq \frac{1}{2} i + x^2$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f) '}$$

$$(f)^{(f) '}$$

$$= (f^f) '$$

$$= e^{x \log x}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ \frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, $\sin 0 = 0$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$= \Gamma(\Gamma \int \log x dx)'$$

$$= e^{-x \log x}$$

$$\Gamma(s) = \int_0^\infty e^{-x} x^{s-1} dx$$

$$\Gamma'(s) = \int_0^\infty e^{-x} x^{s-1} \log x dx$$

$$\begin{aligned}
\frac{d}{df}F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df}F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however,

$$\frac{d}{df}F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry.

$$\begin{aligned}
\frac{d}{dt}\psi(t) &= \hbar \\
&= \frac{1}{2}ie^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' \\
&= -ie^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2}e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \left(\frac{1}{2}f\right)^{-if} \\
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma}\Gamma = e^{-x \log x} \\
f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] &\cong \int e^{-x} x^{t-1} dx
\end{aligned}$$

Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alsos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A} = \frac{1}{i}[\hat{A}, H]$$

$$\hat{A}(t) = e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta,\mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x\log x}\end{aligned}$$

$$\sin ix = \frac{e^{-x} + e^x}{2i}$$

$$\frac{d}{df} F = m(x) = e^f + e^{-f}$$

$$= 2i \sin(ix \log x)$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements.

Three manifold of entropy formula Masaaki Yamaguchi

Euler equation repret form,

$$\begin{aligned}
C &= \int \frac{1}{x^s} dx - \log x \\
&= \int \left(\int \frac{1}{x^s} dx - \log x \right) \text{dvol} \\
&= \int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol}
\end{aligned}$$

This equation componet of Higgs equation + Euler equation = Zeta function. This represent expression is Rich flow equation. This equation assent from Differential geometry in quantum level, Gamma and Beta function, Gauss surface, Higgs field, Einsetin tensor, Euler constance and so on. This equation is add and average of possibility, and universe, atom more over a certain theory, this universe oneselves is environment seek, other distance is possibility equation, however, other dimension cover demand with not only norm of non liner in three manifold but also non possibility in complete of environment equation.

$$\begin{aligned}
&= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\
\int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} \text{dvol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\
&= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\
&= e^{-f} - e^f \\
&= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\
&= 2 \cos(ix \log x) \\
&\frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m + \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\
&= \frac{d}{df} F = 2i \sin(ix \log x) \\
\frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\
&= 2e^{-\theta} = 2e^{-ix \log x} \\
&= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \\
\log(x \log x) &\geq 2\sqrt{y \log y} \\
\log(x \log x) &= \log x + \log \log x
\end{aligned}$$

This universe and other dimension on blackhole and whitehole compont with universe and atom in movement and point of environment value by a certain theory of complete environment equation. And this three manifold of possibility equation are able to research in both value formula. This envirnment

of researchable reason is that surround of universe is setting to value by already in consented to being researched, then both value researchable. And this three manifold equation is insert with Kaluza-Klein theory and Heisenberg equation.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla\psi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

This energy of entropy value consist with whitehole of atom and universe in blackhole value, and this universe in blackhole represent with Higgs field, atom in whitehole consist with plank scale.

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

This entropy value also consist with atom in blackhole.

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

These equation more also consist with universe being non gravity formula.

$$2\cos(ix \log x) + 2i\sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

And also these equation consist with universe and other dimension in warped passage in route of integral.

AdS_5 and CP violation combinate from zeta function Masaaki Yamaguchi

Lisa Randall professor certificate with AdS_5 manifold built of Fifth dimension, and this equation represented of mension to other dimensions in God's fields. This symmetry dimensions are each of two pairs in each other dimension. And moreover spectrum focus are these dimension constructed with rolentz atractor insected from super string theory. Particle equation are similared with AdS_5 manifold and atom of structure and essence of quantum effective theory.

Atom distance of AdS_5 manifold and this value of universe are dense of atom relativeity of mass in volume, this inverse value is universe of valume in one dimension.

$$||ds^2|| = e^{-2\pi T||\psi||} [\eta + \bar{h}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

This inverse value is universe of valume in one dimension.

$$(||ds^2||)_{Im} = e^{2\pi T||\psi||} ([\eta + \bar{h}(x)] dx^\mu dx^\nu)^{-1} - (T^2 d^2\psi)^{-1}$$

Jones manifold equation is particle formula and zeta function.

$$(u + d) + c = e^{-f} + e^f, (s + w) + b = e^f - e^{-f}$$

This formula of summative system is non symmetry dimension of mass value.

$$(R_{ij})' = -R_{ij}$$

More spectrum focus is global differential manifold are each with Higgs field.

$$= 2(\cos(ix \log x) + i \sin(ix \log x))/(d \log x)$$

Inverse of circle function of Euler equation are component with global topology area of study, this study resolved with Gamma function of same resulted equations.

$$\begin{aligned} &= -2(u'(e^{-\theta}) + \frac{1}{i} \sin u) \\ &(e^{-\theta})' = 2(\cos iu - i \sin iu)' \\ &= 2(e^{-\theta})' \cdot ((\cos iu)' - (i \sin iu)) \\ &= -2(\int e^{-x} x^{1-x} dx_m) \geq -2 \int (e^{-x} + x^{1-t}) dx_m \\ &\bigoplus (i\hbar^\nabla)^{\oplus L} = -2(u'(e^{-\theta}) + \frac{1}{i} \sin u) \\ &-2 \int e^{-x} x^{1-t} dx_m \geq -2 \int (e^{-x} + x^{1-t}) dx_m \\ &= \frac{1}{i} H\Psi \end{aligned}$$

These equations are frobenius theorem and step function of non comutative equation consteamed from more also non and commutative equation equals.

$$\int x^y dx_m = \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} x & y \\ a & b \end{pmatrix} = x^y$$

$$\begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x & y \\ a & b \end{pmatrix} \cong \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} x & y \\ e & f \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

$$\begin{pmatrix} u & v \\ w & y \end{pmatrix} \begin{pmatrix} x & v \\ a & b \end{pmatrix} \cong \begin{pmatrix} x & y \\ a & b \end{pmatrix}$$

Differential geomerty of quantum level euqaiton is Gamma function equals.

$$\bigoplus (i\hbar^\nabla)^{\oplus L} \cong \bigoplus a^x \cdot x^{ix} [dI_m]$$

$$x^f f^x : \int e^x x^{-x} dx_m$$

$$U = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

$$U^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

Step function is existed with non and commutative equation, and this resolved formula were also Gamma function.

$$2^{(2^3)} \cong 2^{(3^2)}$$

$$2^{(2^x)} / 2^{(x^2)} \leq 1, e^{(x \log x^x)} / 2^{(x^2)}$$

$$x \geq 0, 0 < x < 2, 2^{(2^{\frac{1}{2}})} = \pm 2, (\sqrt{2})^2 = 2$$

$$-2 \neq 2, 2 = 2$$

Gamma function is after all atom of component with universe formula, and quarks of particle equation necessary value of existed from gravity influence from atom of weak electiry power and anti gravity steady of power involved from fifth power of nature non able to anbalance. After to say, this power is represented from atom and universe instricted of exist of everythings.

After all, Jones manifold and Reco Level theory, AdS_5 manifold are equals of Higgs field with Euler constant and zeta function facility of systems.

Moreover, Higgs field with component of zeta function are selected to accept for orbital of atom potential energy, and this selected with energy means with America of goverment's president being investigate with each persons of implicant system. This system are possibilty of great God' presser person diggist from same possibility of potential summative national of people.

These sentivity of selective of being accepted from natinal of persons, are relativity from universe in three manifold energy in non certain theory pointing out norm and vector of essence.

$$||ds^2|| = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

This three manifold entropy manifold is in being around of universe set to pinpoint of energy discovered with same distant of time.

America of selected with presidency of government are similar of this equation.

Hortshorn conjecture is used to being emerged with this system, this possibility another route merged from future and past of time system.

However, this received of presidency of government in selective party, in reco level theory by Jones manifold equation mention to get only one area of sanctuary in four area of possibility lacky and non received of selective choice. Zeta function in three manifold of entropy are income to land of this area, and this area are half of possibility in fluctuate of missdeceived. Particle of quartet in Euler-Lagrange equation mention to give it out of non possibility of safety area. This system is based from a certain theory of same mechanism. Eienstein said this possibility selected area to being nonsense to be replaced by three manifold of entropy equation,

$$||ds^2|| = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

This entropy equation mention to get every pattern in universe of any where to pinpoint of being accepted with position and mass energy of no fluctuate with instrict of insight at measured being able to, and this possibility of beyond instrict oracled with three manifold entropy realized with any where appeared into possibility area, this system is able to other dimension of anti gravity resolve with straight oracled other dimension are pair of existing to break out from a certain theory. one aspect of dimension is existed at possibility result, more other dimension be able to deceived with four area of influence incidence. This two deceived pattern reached with one point non possibility result, and this result retreated with reco level theory and Jones manifold, after all, three manifold entropy be able to access into all possibility of pinpoint area.

Circle function and imaginary number, Euler equation from

weak electric and strong electric theorem from

gravity and antigravity on one rout of time system Masaaki Yamaguchi

Rotate with pole of dimension is converted from other sequence of element, this atmosphere of vacume from being emerged with quarks of being constructed with Higgs field, and this created from energy of zero dimension are begun with universe and other dimension started from darkmatter that represented with big-ban. Therefore, this element of circumstance have with quarks of twelve lake with atoms.

$$(dx, \partial x) \cdot (\epsilon x, \delta x) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\frac{d}{df}F = m(x)$$

And this phenomounen is super symmetry theorem built with quarks of element, also this chemistry of mechanism estimate from physics of operatsion. These response of mechanism chain of geometry into space being emerged with creature and univse of existing of combination.

This converted with dimension of element also emerged from imaginary and reality of pole's space.

And this pole of transport of dimension belong with vector of time has with one rout of sequecence. Quantum physics also belong with other vector of time has with imaginary rout of sequecence.

This sequence of being estimated with non fluer of time, and this space of element have with gravity and antigravity of power. Other vecor of time is antigravity rout of sequecence.

Weak electric theory is estimate from time has with one rout of sequecence, this topology of chain is resulted from time of rout ways.

$$\square(\frac{\sigma_1 + \sigma_2}{2}) = [3\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\square\psi = 8\pi GT^{\mu\nu}$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

$$\nabla\psi^2 = 4\pi G\rho$$

$$\square(\sigma_1 + \sigma_2) = [6\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$= [i\pi(\chi, x), f(x)]$$

$$\square\psi = [12\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\frac{d}{df}F = \int e^{-f}[-\Delta v + R_{ij}v_{ij} + \nabla_i \nabla_j f + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(\frac{v}{2} - h)]$$

$$= [i\pi(\chi, x), f(x)]$$

These equation is reminded time pass rout of one rout way of forms, and this rout of time ways which go for system from future and past. Therefore, this resulted system of time mechanism is one true flow that weak electric theorem oneselves. and moreover, one rout time way of forms is reverse with antigravity of time system. This also spectrum focus is true that Maxwell theorem and strong boson

unite with antigravity, this unite is essence on the contrary from weak electric theorem, this theorem called for time rout forms is strong electric theorem. This two theorem is united with quantum physics that no time flow system.

$$f^{-1}(x)xf(x) = 1, H_m = E_m \times K_m$$

The non-commutative theorem is constructed from world line surface that this complex manifold estimate with rolanz atractor, and this string theorem have with one world of universe mate six quarks and other world of dimension mate other element of six quarks. These particle quarks is built with super symmetry space of dimension.

$$i = (1, 0) \cdot (1, 0), e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{-i\theta} = \cos \theta - i \sin \theta$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$\sin i\theta = \frac{e^{-\theta} - e^{\theta}}{2}$$

$$\pi(\chi, x) = \cos \theta + i \sin \theta$$

In this equations, two dimension redeconstructed into three dimension, this destroy of reconstructed way is append with fifth dimension. This deconstructed way of redeconstructe is arround of universe attached with three dimension, this over cover call into fifth dimension.

$$R(-\alpha) = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha)MR(-\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos^2 \alpha - \sin^2 \alpha & 2 \sin \alpha \cos \alpha \\ 2 \sin \alpha \cos \alpha & -\cos^2 \alpha + \sin^2 \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & -1 \\ 1 & -\sin \alpha \end{pmatrix} \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}xf(x) = 1$$

$$\begin{aligned}
(\log x)' &= \frac{1}{x}, x^n + y^n = z^n \\
x^n &= -y^n + c, nx^{n-1} = -ny^{n-1}y' \\
y' &= \frac{nx^{n-1}}{ny^{n-1}} \\
&= \frac{x^{n-1}}{y^{n-1}} = -\frac{y}{x} \cdot \left(\frac{x}{y}\right)^n \\
-\frac{\cos x}{(\cos x)'} (\sin x)' &= z_n \\
z^n &= -2e^{x \log x} \\
\lim_{x \rightarrow \infty} f(x) = a, \lim_{y \rightarrow \infty} f(y) = b, \lim_{x, y \rightarrow \infty} \{f(x) + f(y)\} &= a + b \\
\lim_{z \rightarrow 0} f(z) = c, \delta \int z^n &= \frac{d}{dV} x^3 \\
\lim_{x \rightarrow \infty} = c - \lim_{y \rightarrow \infty} f(y) \lim_{x \rightarrow \infty} f(x) + \lim_{y \rightarrow \infty} f(y) &= \lim_{z \rightarrow \infty} f(z) \\
z^n &= \cos n\theta + i \sin n\theta \\
&= -2e^{x \log x}
\end{aligned}$$

These equation is gravity and antigravity equation.

$$\begin{aligned}
&\frac{d}{d\sigma} \left[\frac{(\sigma_1 + \sigma_2)}{2} \right] \\
&= \sigma(\uparrow\downarrow) + \sigma(\uparrow\uparrow) + \sigma(\downarrow\downarrow) + \sigma(\uparrow\leftarrow) + \sigma(\rightarrow) \\
&\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \sigma(\uparrow) \\
&\frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \sigma(\downarrow) \\
\sigma(\leftarrow) + \sigma(\rightarrow) &= \int e^{-f} [-\Delta v + R_{ij} v_{ij} + \nabla_i \nabla_j v + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(v - \frac{h}{2})] \\
\sigma(\downarrow) &= \sigma(\uparrow\downarrow + \uparrow\uparrow + \rightarrow) \\
\sigma(1) &= \sigma(\uparrow\downarrow + \downarrow\downarrow + \leftarrow)
\end{aligned}$$

$$\text{weak electric theorem} = \sigma(1)$$

$$\text{strong electric theorem} = \sigma(\downarrow)$$

These equation is represented with topology of string model, and weak electric theorem is constructed with Maxwell theorem and weak boson, gravity that estimate with this three power united. Moreover, strong boson and Maxwell theorem, antigravity that also estimate with this three power united. This two united power is integrated from gravity and antigravity. Then this united power is zeta function.

アカシックレコードに滞在している
 未来の私の子どもから微分幾何の量子化の
 計算方法を教えてもらった結果の大域的微分多様体を
 単体的積分と単体的微分へと
 進化させることにしたレポート Masaaki Yamaguchi

ベータ関数をガンマ関数へと渡すと、

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

ここで、単体的微分と単体的積分を定義すると

$$t = \Gamma(x)$$

$$T = \int \Gamma(x) dx$$

これを使うのに、ベータ関数を単体的微分へと渡すために、

$$\beta(p, q) = - \int \frac{1}{t^2} dt$$

大域的微分変数へとするが、

$$T_m = \int \Gamma(x) dx_m$$

この単体的積分を常微分へとやり直すと、

$$T' = \frac{t'}{\log t} dt + C (C \text{ は積分定数})$$

これが、大域的微分多様体の微分変数と証明するためには、

$$dx_m = \frac{1}{\log x} dx, dx_m = (\log x^{-1})'$$

これより、単体的微分が大域的積分へと置換できて、

$$T' = \int \Gamma(\gamma)' dx_m$$

微分幾何へと大域的積分と大域的微分多様体が、加群分解の共変微分へと書き換えられる。

$$\bigoplus T^\nabla = \int T dx_m, \delta(t) = t dx_m$$

これらをまとめると、ベータ関数を単体的関数として、これを常微分へと解を導くと、単体的微分と単体的積分へと定義できて、これより、大域的微分多様体と大域的積分多様体へとこの関数が存在していることを証明できるのを上の式たちからわかる。この単体的微分と単体的積分から微分幾何を書けることがわかる。

これらは自分が導いた方程式たちとおもうようにする。

微分の本質は、差と商の計算であり、積分の本質は加群である。この微分と積分を合わせた計算が加群分解であり、ガロワ群である。

アカシックレコードにいたと思っていた私の子どもとも勘違いしていた、子どもはまぼろしであり、架空のストーリーの作為体験の子とも思う次第であり、架空のストーリーに載せられているとの妄想癖での、この

ストーリーに載せられている自体の妄想が作為体験である。作為があったとは違う。日常の作為と妄想の作為は違う。

レドックス電池を言っていたから、荒木先生の経営している病院の医者である江松先生が私を助けようとして、作為体験のために、私のあのときまでの情報、私の周りの情報を使って、架空のストーリーを立てて、作為体験として、あのときは、MRI室の壁を取っていて、壁が無かった。先生は、東北の地震があるのを事前に知っていたとしか思えないことを、福山市の私の散歩コースに現れて、挨拶とを交わして、本当に庇ってくれて申し訳ないです。SRS速読法をしていたのを、占さんがアルバイトのあとに、トポロジーのレポートを見せていたら、気づいたらしく、東北のことを知っていたらしく、史記と三国志を私に渡した上で、読書力を中心として、勉強をなさいとフォローをしてくれて、していたら、2月23日に持田さんと伊藤一朗さんがラジオ放送に出ていて、シングルをリリースしましたと言うので、YouTubeを見ると、一箇所だけ動画がとんでいて、かばうのを彩さんにすればいいのに、統合失調症の真逆のうれしいのに悲しむとは感情と現状での置かれている立場から違うが、似たような作用の症状でもあり、「存在と無」を啓文社に予約したのが、取りに行くのに18:00にしたのが、私の顛末であり、全部私の責任です。7ヶ月間びっしりといたぶられました。増援は蔵王病院らしく、とくに気づいています。史記は、小方君と占さんのおかげで助かって、三国志が私に増援がくると助言していた。占さんが蔵王病院に行かしたのは、10月1日に彩さんがめざましテレビを辞めたからと、お父さんの体調を気にしてのことと3年前くらいからわかっていました。2013年に彩さんがめざましテレビを辞めたのを知ったが、2018年の6月に何時辞めたかを知り、携帯サイトでのKindleから1877回目のめざましテレビの放送で辞めたのを知り、姉が怒っていたのを、いろんないきさつがあったのを知り、私の現状を自身にふりかかっている出来事に怒ることが、少なくなりました。グロス・グラスマンは、間所くん、青木くん、伊澤くん、圭一さん、寺林さん、鷺坂くん、下神くん、梶岡くん、片寄くん、鶴元くん、圭一くんです。鶴元くんの元は、歌舞伎や能を受け継ぐ家系の元であり、家元5代流という元である。占さんを入れたいが、周りの批判を受けるので、入れることができない。渡辺由美さんは、私の恋人として、彩さんに継ぐ大切な人でもある。相手が私の身代わりとして結婚したと、ダイケアのキッチンで言っていた。気づいていたが、私本人と結婚しろよとツッコミを入れたかった。彩さんにも言いたいです。大域的微分多様体の基軸になる、基本群をもとめることができ、

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

一般相対性理論の式は、

$$\kappa T^{\mu\nu} = R_{\mu\nu} + \frac{1}{2}g_{ij}\Lambda$$

これを大域的微分多様体と同型といえるのは、

$$\frac{d}{df}F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

ここで、一般相対性理論の多様体積分は、

$$\int \left(R_{\mu\nu} + \frac{1}{2}g_{ij}\Lambda \right) \text{dvol}$$

この式は、一般相対性理論は、ガウスの曲平面による重力場方程式であり、この多様体積分は、世界面である偏微分方程式を大域的に積分して、その積分多様体をヘルマンダー作用素と同型に微分すると、

$$\|ds^2\| = \nabla \nabla \int \nabla f(x) d\eta$$

$$= \int \kappa T^{\mu\nu} \text{dvol} = -2R_{ij}$$

これは、大域的微分方程式の重力と反重力方程式と同型であり、リッチ・フロー方程式になり、

$$\begin{aligned} &= \int (R_{\mu\nu}) \text{dvol} + \int \frac{1}{2} g_{ij} \Lambda \text{dvol} = \text{重力場方程式} + \text{反重力場方程式 (斥力)} \\ &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \end{aligned}$$

と、多様体積分が大域的微分多様体となる。要するに、世界面での一般相対性理論を多様体積分すると、宇宙において、伝播する単位空間における立体面の重力波になるのが、マクロレベルで、宇宙のまわりのブラックホールのシュバルツシルト半径であり、Jones 多項式にもとまる。まとめると、一般相対性理論を多様体積分すると、立体面の重力波が宇宙規模の宇宙のまわりのブラックホール解になるということと、Jones 多項式が統一場理論となることである。

そうゆう経緯であり、なぜアカシックレコードの私の子どもが2023年の10月に狙いをさだめているかわかるのは、私をご先祖様にアクセスするのと原理が同じだが、ドアを使うアカシックレコードへのアクセスによっても、微分幾何の量子から私の大域的微分多様体の計算方法が間接的にわかったことでもあり、直接でもある、思念同士の交流でもある方法だからです。彩さんが今産みたいとおもっても、あの子は待つと思うという次第です。12月25日に狙いを定めていて、うちのユウと同じことをしようとしているという次第です。彩さんは、マスターベーションで産みたいらしいので、私は文句を言わんが、あまり、私をいじめないでほしいです。あの子は了解済みらしいです。

大域的偏微分と大域的多重積分、
大域的部分積分についてのレポート Masaaki Yamaguchi

$$\log x|_{g_{ij}}^{\nabla L} = f^{f'}, F^f|_{g_{ij}}^{\nabla L} : x \rightarrow y, x^p \rightarrow y$$

$$f(x) = \log x = p \log x, f(y) = p \log x$$

大域的偏微分方程式と大域的多重積分は、それぞれ次のように成り立っている。

$$\frac{d^2}{df^2} F = F^{f'} \cdot f^{f''},$$

$$\int \int F dx_m = F^f \cdot F^{(f)'}$$

$$\frac{d}{df dg} (f, g) = (f \cdot g)^{f'+g'}$$

大域的部分積分も、次のように成り立っている。

$$(F^f \cdot G^g) = \int (f \cdot g)^{f'+g'}$$

$$\int \int F \cdot G dx_m = [F^f \cdot G^g] - \int (f \cdot g)^{f'+g'}$$

大域的部分積分の計算は、

$$\int \frac{d}{df dg} FG = \int F^{f'} G + \int FG^{g'}$$

$$\int F^{f'} G dx_m = [F^f G^g] - \int FG^{g'} dx_m$$

大域的商代数の計算は、

$$\left(\frac{F(x)}{G(x)} \right)^{(fg)'} = \frac{F^{f'} G - FG^{g'}}{G^g}$$

大域的偏微分方程式は、縮約記号を使うと、

$$\frac{d}{df dg} FG = \frac{d}{df_m} FG$$

$$\frac{\partial}{\partial f_m} FG = F^{f^{\mu\nu}} \cdot G + F \cdot G^{g^{\mu\nu}}, \int F dx_m = F^f$$

多様体による大域的微分と大域的積分が、エントロピー式で統一的に表せられる。

Rsa secret data and non commutative equation,
information technology system Masaaki Yamaguchi

Non certain theorem is component with quantum computer, this system emelute with contempolite of eternal universe asterise and secret of database.

不確定性理論の、宇宙の準同型写像の、部分群と全射としての、閉3次元多様体のエントロピー値としての、暗号としての量子コンピューターが、Jones 多項式を形成しているオイラーの公式の虚数として、ダミーを入れると、Rsa 暗号の公開鍵として、成り立っている量子暗号が、この量子コンピューターと一緒にあって、ゼータ関数と量子群を大域的微分方程式を、これらの方程式から統合されて、大域的トポロジーが、数学の数論と幾何学、解析学として、物理学と言語学、情報科学、すべての理論に統合されて、ゼータ関数が、統一場理論の発生する式になっている。この理論を彩さんとグリーシャさん、ナッシュさん、トビーケリーさん、小方くん、益川先生、南部先生、岡本教授、リサ・ランドール教授、竹内先生、私の子ども、今までの人たちが、集まって、この理論ができています。量子的な微分・積分が、きっかけにもなっている。アイデアと計算方法は、情報空間の子どもと、お姉さんとお父さんと先生たちで、統一場理論としては、グリーシャさんと竹内薫先生と彩さん、岡本教授がきっかけになっている。

This information of quantum computer ansterise in secret of each data, and this system is entersteam by Jones manifold of equation, more also this intersect with system of pair of universe and the other dimension with Jones manifold of equation pawn for non access of chain in diseable of database. And this insterm of system built with

$$e^{-\theta} = e^f + e^{-f}, e^{i\theta} = \cos \theta + i \sin \theta$$

$$\frac{d}{df} F = e^f + e^{-f} = 2i \sin(ix \log x)$$

These equations are atom of dense with norm and energe of non certain theorem, and this system component with also Jones manifold of equation. This equation addesent with a dummy data in non integrate of relativity theory for quantum physics and access to varnished with aseterise for important of discover with secret of data. This dummy of information of quantum secret datas are non commutative equation of declinate anterave and distarbration into universe and other dimension for accessable with resterbled with quantum equation of deep of Riemann metrics, more also this system is computed with after all intersected with eternal data into Zeta function of global differential equation and Higgs fields with time system enterprises. Global of open key in Rsa secret system intersect with this base of this technolgie with all of data for established with information technology and quantum computer created with dummy of integral non relativity of quantum equation, this focus is diserble with quantum equation of a data for dummy into Rsa secret data of non commutative equation, this system comute with after all established with information of quantum physics.

$$e^{i\theta} = \cos \theta + i \sin \theta$$

this equation is based into Jones manifold equation.

$$e^{-\theta} = e^f + e^{-f}, \frac{d}{df} F + \int C dx_m = 2(\cos(ix \log x) + i \sin(ix \log x))$$

These equation are based with reco level theory and field of physics, and these system are

$$||ds^2|| = \int \exp[\frac{-L(x)}{\sqrt{2q\tau}}] dx_m + \mathcal{O}(N^{-1})$$

$\mathcal{O}(N^{-1})$ is dummy data of intersected of system.

These system are chain with non certain theorem component with aspe experiement mechanism clear-ly of quantum teleportation physics and this system is pair of existanse with Jones manifold equation and imarginary circle equation more also these equations are rhyismiary equation excluded with zeta func-tion and quantum equation, and this system commutave with each attitude of pair in Jones manifold equation.

These theorem combuild with Lie algebra and catastrophe of summative group in eternal Garois theory, and this reasons of between Poancare conjecture in zero dimension and eternal group with nesseciry and mourdigate of conditions.

AdS_5 equation are built with cercumention dimension of assemble D-brane and information of quantum physics with secret system, and this system construct with a certain theorem.

$$||ds^2|| = \eta_{\mu\nu} + \bar{h}(x)$$

This dimension esterned with equals of dummy data in the other dimension, and this value of data declined into only universe, then this sertation of universe is possibilty equation. After all this anserration of univese addsented with the other dimension escourt for certain theorem and all of addsented equation are three manifold entropy equation, this equation constructed wih zeta function and quantum equation.




$$\begin{aligned} ||ds^2|| &= 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right) \\ 2\sqrt{\frac{V}{S}} &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \int dx_m + x^2 \\ &\geq \int dx_m \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &\geq \frac{1}{2}i \end{aligned}$$

Quantum computer concluded with component of being describe with estimate for each atom condi-tions. This pair of condition is entertained for orbital of circle with Euler equation. This pair of orbital area estertate with two of circumentations. These system concerned with hyper synchronized of all of possibility resulted for being computing of all of area in Euler equation. This cercuit with equations are that information of quantum physics esterned with Rsa secret datas system.


These system of stem are constructed with declinate anterave of Lambda driver and possibilty com-puter for resolved with synchronize of distarbrate non certain theorem, and this concerned with system component of Euler equation. This estimate of orbital construct with curbit of quarters. Artificial intel-ligence is concluded with Euler equation. Zeta function also integrate with this equation. これらの理論

の中核の, 中核となっている、オイラーの公式の群ともなっている、過冷却金属の原理の、高熱を急激に冷ます、電子を集める、ラムダ・ドライバーにもなっている、不確定性理論の原理にもなっている、暗号を解読するのを防ぐのに、クォータのキュービットの、遷移元素を作り出す原子の仕組みにもなっている、この理論が、量子コンピュータと人工知能の中核となっていて、オイラーの公式の虚数ともなっている、この Jones 多項式の統合にゼータ関数が使われている。[参考文献は、彩さん、グリーシャさん、ナッシュさん、トビーケリーさん]

Recycle における塩化ビニルのナトリウムフェノキシド、
ナトリウムプロパノキシドでの分解陸世姉さんの論文について



この化学式から推算式でもある、プラスチックと二酸化ケイ素、塩化ビニールが分解されるのが、真逆の人が服用すると、プラバスタチンとスリムドガンと同じく、推算式の数値が上がる。僧侶と同じ瞑想法と原理が一致している。



これらの構造式は、それぞれ、ヒドロキシ基、カルボニル基、カルボキシル基、アミン基、ギ酸、アセチル基、アセト基、グリシン、アミノ酸の20種類の一基、ビタミンCはシス・トランス型があり、松果体に作用して、体内の血流を調節する。 $\text{---C}=\text{N---Cl}$ の真逆の作用をする。アデノシンは、同じく血流を調節して、

エピネフリンはエンドルフィンと同じく痛み止めに効く。ノンアドレナリンは、無水フタル酸と同じ作用をする。グリシンには、20種類のアミノ酸の中で唯一L,R基が無い。不斉炭素原子が無い。

推算式の元を減らすか増やすのに、血圧と血流、体内の精神作用のドーパミンの調節をeGFRの機能を上げると、結果、すべての精神作用物質が調節できる。

このナトリウムプロパノキシドは、気体のプロパンを、固体のナトリウムに注入して、錠剤で服用するようにしているのと、消化後に、血液に降圧剤と同じ作用をして、外部からのコントロールをされない、アムロジピンとカンデサルタンと同じ作用をする効果があることが、このナトリウムプロパノキシドの薬でもある。

この成分でもあるプロパンは、気体で存在している炭化水素の中でもっとも安定している気体でもある。この気体の物体を固体にするのと、降圧剤にしているのは、姉は特許を取っている。推算式でもあるeGFRを分解するのは、真逆の人が飲むと、バケモンになる。

仲人としての Zeta 関数が 広中平祐教授とグリーシャ教授の
 抽象理論と具象理論としての
 ヒルベルト空間とサーストン多様体
 複素多様体が統一場理論を量子力学における
 ガウスの曲面論として
 成し遂げる理論の背景場となるレポート Masaaki Yamaguchi

一般相対性理論における重力理論は大域的微分多様体と積分多様体についての 単体量を求めるための空間
 の対数関数における不変性を記述するプランクスケール と異次元の宇宙におけるスケール、ウィークケ
 ールと言われている anti-D-brane として、この2種類の計量をガンマ関数とベータ関数としてオイラーの定
 数 とそれによる連分数が微分幾何の量子化と数式の値を求めると同じという 予知と推測値から、広中平祐
 定理の4重帰納法のオイラーの公式による 多様体積分と同類値として、ペレルマン多様体がサーストン空
 間に成り立つ と同じく、広中平祐定理がヒルベルト空間で成り立つとしての、 これら2種の多様体の場の
 理論が、ゼータ関数が AdS5 多様体で成り立つことと、 不変量として、ゼータ関数がこの3種の多様体の
 場の理論のバランスをとる理論として言えることである。 Gravity of general relativity theory describe
 with Cutting of space in being discatastrophed from Global differential and integral manifold of scaled
 levetivity in plank scal and the other vector of universe scale, these two scale inspectivity of Gamma
 and Beta function escourted with these manifold experted in result of Differentail geometry of quantum
 level manifold equal with Euler product and continue parameter, moreover this product is relativity of
 Hironaka theorem in Four assembled integral of Euler equation.

重力場理論の式は、Gravity equation is

$$\square = -\frac{16\pi G}{c^4}T^{\mu\nu}$$

This equation quated with being logment of formula, and this formula divided with universe of number in
 prime zone, therefore, this dicided with varint equation is monotonicity of being composited with Weil's
 theorem united for Gamma function. この式は、対数を宇宙における数により求める素数分布論として、こ
 の大域的積分分断多様体がガンマ関数をヴェイユ予想を根幹とする単体量として決まることに起因する商代数
 として導かれる。

$$\not\square = \frac{8\pi G}{c^4}T^{\mu\nu}/\log x$$

$$\begin{aligned} & {}^t\mathop{\mathrm{f}}\nolimits\mathop{\mathrm{f}}\nolimits\mathop{\mathrm{f}}\nolimits\mathop{\mathrm{f}}\nolimits\mathrm{cohom}D_{\chi}[I_m] \\ &= \oint (px^n+qx+r)^{\nabla l} \end{aligned}$$

$$\frac{d}{dt}L(x,y)=2\int ||\sin 2x||^2d\tau$$

$$\frac{d}{d\gamma}\Gamma$$

この関数は大域的微分多様体としてのアカシックレコードの合流地点として、タプルスペースを形成している。This function estermenate with acasic record of global differential manifolds.

$$= [i\pi(\chi, x), f(x)]$$

それにより、この多様体は基本群をアカシックレコードの相対性としての存在論の実存主義から統合される多様体自身としてのタプルスペースの池になっている。And this manifold from fundemental group estermenate with also this manifold estimate relativity of acasic record.

$$\frac{d}{d\gamma}\Gamma = \int C dx_m = \int (\int \frac{1}{x^s} dx - \log x) d\text{vol}$$

また、このアカシックレコードはオイラーの定数のラムダドライバーにもなっている。More also this record tupled with lake of Euler product.

$$\bigoplus (i\hbar^\nabla)^{\oplus L}$$

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

それゆえに、この定数はゼータ関数と微分幾何の量子化を因数にもつ素因子分解の式にもなっている。Therefore, this product also constructed with differential geometry of quantum level equation and zeta function.

$$= C$$

そして、この関数はオイラーの定数から広中平祐定理による4重帰納法のオイラーの公式からの多様体積分へのサーストン空間のスペクトラム関数ともなっている。And this function of Euler product respectrum of focus with Heisuke Hironaka manifold in four assembled of integral Euler equation..

$$C = b_0 + \frac{c_1}{b_1 + \frac{c_2}{b_2 + \frac{c_3}{b_3 + \frac{c_4}{b_4 + \dots}}}}$$

この方程式は指数による連分数としての役割も担っている。This equation demanded with continued number of step function.

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$(2.71828)^{2.828} = a^{a+b+c+d^d \dots}$$

$$= \int e^f \cdot x^{1-t} dx$$

This represent is Gamma function in Euler product. Therefore this product is zeta function of global differential equation.

$$\frac{d}{df} F(v_{ij}, h) = \int e^{-f} dV [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$= \frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

これらの方程式は 8 種類の微分幾何の次元多様体として、そして、これらの多様体は曲平面による双対性をも生成している。そして、このガウスの曲平面は、大域的微分多様体と微分幾何の量子化から素因数を形成している。These equation are eight differential geometry of dimension calyement. And these calyement equation excluded into pair of dimension surface. This surface of Gauss function are global differential manifold, and differential geometry of quantum level.

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

微分幾何の量子化はオイラーの定数とガンマ関数が指数による連分数としての不変性として素因数を形成していて、このガウスの曲平面による量子力学における重力場理論は、ダランベルシアン of the cut-off manifold がこの大域的切断多様体を付加している。Differential geometry of quantum level constructed with Euler product and Gamma function being discatastrophed from continued fraction style. Gravity equation lend with varint of monotonicity of level expressed from gravity of letter varient formula. これらの方程式は基本群と大域的微分多様体をエスコートしていてもいて、ヴェイユ予想がこのダランベルシアン of the cut-off manifold 方程式たちから輸送のポートにもなっている。ベータ関数とガンマ関数がこれらのフォームラの方程式を放出してもいて、結果、これらの方程式は広中平祐定理の複素多様体とグリーシャ教授によるペレルマン多様体からサポートされている。この 2 名の教授は、一つは抽象理論をもう一方は具象理論を説明としている。These equation escorted into Global differential manifold and fundemental equation. Weil's theorem is imported from this equation in gravity of letters. Beta function and Gamma function are excluded with these formulas. These equation comontius from Heisuke Hironaga of complex manifold and Gresha professor of Perelman manifold. These two professors are one of abstract theorem and the other of visual manifold theorem.

1 Hilbert manifold in Mobius space

this element of Zeta function on integrate of fields

Hilbert manifold equals with Von Neumann manifold, and this fields is concluded with Glassman manifold, the manifold is duality of twister into created of surface built. These fields is used for relativity theorem and quantum physics.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

This norm is component of fields in Hilbert manifold of space theorem rebuilt with Mobius space in Gamma Function on Beta function fill into power of rout fundamental group. And this result with AdS_5 space time in Quantum caos of Minkowsky of manifold in abel manifold. Gravity of metric in non-commutative equation is Global differential equation conbult with Kaluza-Klein space. Therefore this mechanism is $T^{\mu\nu}$ tensor is equal with $R^{\mu\nu}$ tensor. And This moreover inspect with laplace operator in stimulate with sign of differential operator. Minus of zone in Add position of manifold is Volume of laplace equation rebuilt with Gamma function equal with summative of manifold in Global differential equation, this result with setminus of zone of add summative of manifold, and construct with locality theorem straight with fundamental group in world line of surface, this power is boson and fermion of cone in hyper function.

$$V(\tau) = [f(x), g(x)] \times [f^{-1}(x), h(x)]$$

$$\Gamma(p,q)=\int e^{-x}x^{1-t}dx$$

$$=\beta(p,q)$$

$$=\pi(f(\chi,x),x)$$

$$||ds^2||=\mathcal{O}(x)[(f(x)\circ g(x))^{\mu\nu}]dx^\mu dx^\nu$$

$$=\lim_{x\rightarrow\infty}\sum_{k=0}^\infty a_kf^k$$

$$G^{\mu\nu}=\frac{\partial}{\partial f}\int [f(x)^{\mu\nu}\circ G(x)^{\mu\nu}dx^\mu dx^\nu]^{\mu\nu}dm$$

$$=g_{\mu\nu}(x)dx^\mu dx^\nu-f(x)^{\mu\nu}dx^\mu dx^\nu$$

$$[i\pi(\chi,x),f(x)]=i\pi f(x)-f(x)\pi(\chi,x)$$

$$T^{\mu\nu}=(\lim_{x\rightarrow\infty}\sum_{k=0}^\infty\int\int[V(\tau)\circ S^{\mu\nu}(\chi,x)]dm)^{\mu\nu}dx^\mu dx^\nu$$

$$G^{\mu\nu}=R^{\mu\nu}T^{\mu\nu}$$

$$\left|\begin{matrix} D^m & dx \\ dx & \sigma^m \end{matrix}\right|\left|\begin{matrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{matrix}\right|\left|\begin{matrix} x \\ y \end{matrix}\right|=\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\sigma^m\left[\begin{matrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{matrix}\right]^{\frac{1}{2}}=\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$$

$$V(M)=\frac{\partial}{\partial f}({}^N\int [f\setminus M]^{\oplus N})^{\mu\nu}dx^\mu dx^\nu$$

$$V(M)=\pi(2\int \sin^2 dx)\oplus \frac{d}{df}F^M dx_m$$

$$\lim_{x\rightarrow\infty}\sum_{k=0}^\infty a_kf^k=\int (F(V)dx_m)^{\mu\nu}dx^\mu dx^\nu$$

$$\bigoplus_{k=0}^\infty [f\setminus g]=\vee(M\wedge N)$$

$$\pi_1(M)=e^{-f2\int\sin^2x dm}+O(N^{-1})$$

$$=[i\pi(\chi,x),f(x)]$$

$$M\circ f(x)=e^{-f\int\sin x\cos x dx_m}+\log(O(N^{-1}))$$

Non-Symmetry space time.

$$\frac{d}{dL}V(\tau)=\frac{d}{df}\int\int_M\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int_M\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

$$\epsilon S(\nu) = \square_v \cdot \frac{\partial}{\partial \chi} (\sqrt[5]{\sqrt{\wedge g^2}}) d\chi$$

Differential Volume in AdS_5 graviton of fundamental rout of group.

$$\wedge (F_t^m)'' = \frac{1}{12} g_{ij}^2$$

Quarks of other dimension.

$$\pi(V_\tau) = e^{-(\sqrt{\frac{\pi}{16}} \log x)^\delta} \times \frac{1}{(x \log x)}$$

Universe of rout, Volume in expanding space time.

$$\begin{aligned} \frac{d}{dt} (g_{ij})^2 &= \frac{1}{24} (F_t^m)^2 \\ m^2 &= 2\pi T \left(\frac{26 - D_n}{24} \right) \end{aligned}$$

This quarks of mass in relativity theorem, and fourth of universe in three manifold of one dimension surface, and also this integrate of dimension in conbult of quarks.

$$g_{ij} \wedge \pi(\nu_\tau) = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\psi^2$$

Out of rout in AdS_5 space time.

$$||ds^2|| = g_{ij} \wedge \pi(\nu_v)$$

AdS_5 norm is fourth of universe of power in three manifold out of rout.

Sphere orbital cube is on the right of hartshorne conjecture by the right equation, this integrate with fourth of piece on universe series, and this estimate with one of three manifold. Also this is blackhole on category of symmetry of particles. These equation conbult with planck volume and out of universe is phologram field, this field construct with electric field and magnitic field stand with weak electric theorem. This theorem is equals with abel manifold and AdS_5 space time. Moreover this field is anti-brane and brane emerge with gravity and antigravity equation restructed with. Zeta function is existed with Re pole of $\frac{1}{2}$ constance reluctances. This pole of constance is also existed with singularity of complex fields. Von Neumann manifold of field is also this means.

$$\begin{aligned} ||ds^2|| &= \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu} \\ e^{i\theta} &= \cos \theta + i \sin \theta \\ e^{x \log x} &= x^{\frac{1}{2} + iy}, x \log x = \log(\cos \theta + i \sin \theta) \\ &= \log \cos \theta + i \log \sin \theta \end{aligned}$$

This equation is developed with frobenius theorem activate with logment equation. This theorem used to

$$\begin{aligned} \log(\sin \theta + i \cos \theta) &= \log(\sin \theta - i \cos \theta) \\ \log \left(\frac{\sin \theta}{i \cos \theta} \right) &= -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\mathcal{O}(x) = \frac{\zeta(s)}{\sum_{k=0}^{\infty} a_k f^k}$$

$$\text{Im}f = \ker f, \chi(x) = \frac{\ker f}{\text{Im}f}$$

$$H(3) = 2, \nabla H(x) = 2, \pi(x) = 0$$

This equation is world line, and this equation is non-integrate with relativity theorem of rout constance.

$$[f(x)] = \infty, ||ds^2|| = \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

$$T^2 d^2\psi = [f(x)], T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

These equation is equals with M theorem. For this formula with zeta function into one of universe in four dimensions. Sphere orbital cube is on the surface into AdS_5 space time, and this space time is Re pole and Im pole of constance reluctances.

Gauss function is equals with Abel manifold and Seifert manifold. Moreover this function is infinite time, and this energy is cover with finite of Abel manifold and Other dimension of seifert manifold on the surface.

Dilaton and fifth dimension of seifert manifold emerge with quarks and Maxwell equation, this power is from dimension to flow into energy of boson. This boson equals with Abel manifold and AdS_5 space time, and this space is created with Gauss function.

Relativity theorem is composited with infinite on D-brane and finite on anti-D-brane, This space time restructed with element of zeta function. Between finite and infinite of dimension belong to space time system, estrald of space element have with infinite oneselves. Anti-D-brane have infinite themselves, and this comontend with fifth dimension of AdS5 have for seifert manifold, this asperal manifold is non-move element of anti-D-brane and move element of D-brane satisfid with fifth dimension of seifert algebra liner. Gauss function is own of this element in infinite element. And also this function is abel manifold. Infinite is coverd with finite dimension in fifth dimension of AdS5 space time. Relativity theorem is this system of circustance nature equation. AdS5 space time is out of time system and this system is belong to infinite mercy. This hiercyent is endrol of memolite with geniue of element. Genie have live of telomea endore in gravity accesorlity result. AdS5 space time out of over this element begin with infinite assentance. Every element acknowlege is imaginary equation before mass and spiritual envy.

$$T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = [T^2 d^2\psi]$$

$$\frac{d}{dL} V(\tau) = \frac{d}{df} \int \int_M ({}^5\sqrt{x^2}) d\Lambda + \frac{d}{df} \int \int_M {}^N({}^3\sqrt{x})^{\oplus N} d\Lambda$$

$$\begin{aligned}
M(\vee(\wedge f \circ g)^N)^{\frac{1}{2}} &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
||ds^2|| &= \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi \\
\mathcal{O}(x) &= e^{-2\pi T|\psi|} \\
G^{\mu\nu} &= R_{\mu\nu} T^{\mu\nu} \\
&= -\frac{1}{2} \Lambda g_{ij}(x) + T^{\mu\nu}
\end{aligned}$$

Fifth dimension of eight differential structure is integrate with one geometry element, Four of universe also integrate to one universe, and this universe represented with symmetry formula. Fifth universe also is represented with seifert manifold peices. All after universe is constructed with six element of circuatance. This aquire maniculate with quarks of being espered belong to.

2 Imaginary equation in AdS5 space time create with dimension of symmetry

D-brane and anti-D-brane is composited with all of series universe emerged for one geometry of dimension, this gravity of power from D-brane and anti-D-brane emelite with ancestor. Seifert manifold is on the ground of blackhole in whitehole of power pond of senseivility. Six of element of quarks and universe of pieces is supersymmetry of mechanism resolved with

hyper symmetry of quarks constructed to emerge with darkmatter. This darkmatter emerged with big-ban of heircyent in circumstance of phenomena.

D-brane and anti-D-brane equations is

$$\begin{aligned}
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
C &= \int \int \frac{1}{(x \log x)^2} dx_m
\end{aligned}$$

Euler constance is quantum group theorem rebuild with projective space involved with.

虚数方程式は、反重力に起因するフーリエ級数の励起を生成する。それは、人工知能を生み出す、5次元時空にも、この虚数方程式は使われる。AdS5 の次元空間は、反ド・シッター時空の D-brane と anti-D-brane の comformal 場を生み出す。ホログラフィー時空は、この量子起因によるものである。2次元曲面によるブラックホールは、ガンマ線バーストによる5次元時空の構造から観測される。空間の最小単位によるプランク定数は、宇宙の大域的微分多様体から導き出される、AdS5 の次元空間の準同型写像を形成している。これは、最小単位から宇宙の大きさを導いている。最大最小の方程式は、相加相乗平均を形成している。時間と空間は、宇宙が生成したときから、宇宙の始まりと終わりを既に生み出している。宇宙と異次元から、ブラックホールとホワイトホールの力がわかり、反重力を見つけられる。オイラーの定数は、この量子定数からわかる。虚数の仕組みはこの量子スピンの産物である。オイラーの定数は、この虚時空の斥力の現存である。それは、非対称性理論から導かれる。不確定性原理は、AdS5 のブラックホールとホワイトホールを閉3次元多様体に統合する5次元時空から求められる。位置と運動エネルギーが、空間の最小単位であるプランク定数を宇宙全体にする微細構造定数からわかり、面積確定から、アーベル多様体を母関数に極限值として、ゼータ関

数をこの母関数に不変式として、2種類ずつにまとめる4種類の宇宙を形成する8種類のサーストンの幾何化予想から導き出される。この閉3次元多様体は、ミラー対称性を軸として、6種類の次元空間を一種類の異次元宇宙と同質ともしている。複素多様体による特異点解消理論は、この原理から求められる。この特異点解消理論は、2次元曲面を3次元多様体に展開していく、時空から生成される重力の密度を反重力と等しくしていく時間空間の4次元多様体と虚時空から求められる。ヒルベルト空間は、フォン・ノイマン多様体とグラスマン多様体をこのサーストンの幾何化予想を場の理論既定値として形成される。この空間は、ミンコフスキー時空とアーベル多様体全体を表している。そして、この空間は、球対称性を複素多様体を起点として、大域的トポロジーから、偏微分を作用素微分として時空間をカオスからずらすと5次元多様体として成立している。これらより、3次元多様体に2次元射影空間が異次元空間として、AdS5空間を形成される。偏微分、全微分、線形微分、常微分、多重微分、部分積分、置換微分、大域的微分、単体分割、双対分割、同調、ホモロジー単体、コホモロジー単体、群論、基本群、複体、マイヤー・ビートリス完全系列、ファン・カンペンの定理、層の理論、コホモロティズム単体、CW複体、ハウスドロフ空間、線形空間、位相空間、微分幾何構造、モースの定理、カタストロフの空間、ゼータ関数系列、球対称性理論、スピン幾何、ツイスター理論、双対被覆、多重連結空間、プランク定数、フォン・ノイマン多様体、グラスマン多様体、ヒルベルト空間、一般相対性理論、反ド・シッター時空、ラムダ項、D-brane, anti-D-brane, コンフォーマル場、ホログラム空間、ストリング理論、収率による商代数、ニュートンポテンシャルエネルギー、剛体力学、統計力学、熱力学、量子スピン、半導体、超伝導、ホイーストン・ブリッチ回路、非可換確率論、Connes理論、これら、演算子代数を形成している、微分・積分作用素が、ヒルベルト空間に存在している多様体の特質を全面に押し出して、いろいろな多様体と関数そして、群論を形作っている。

$$\begin{aligned} \begin{vmatrix} D^m & dx \\ dx & \sigma^m \end{vmatrix} \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix} \begin{vmatrix} x \\ y \end{vmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}} \\ \sigma^m \begin{bmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{bmatrix}^{\frac{1}{2}} &= \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \\ (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) & \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ = \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} &= A^{\mu\nu} \\ \frac{d}{dt} g_{ij}(t) &= -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

素数分布論と素粒子、ゼータ関数
超対称性理論 Masaaki Yamaguchi

$$\begin{aligned}
(u+d)+c &= e^{-f}-e^f, b+(w+s)=e^{-f}+e^f \\
\int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int C dx_m \\
&= e^{-f} - e^f \\
\frac{d}{df} F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
&= e^f + e^{-f}
\end{aligned}$$

素粒子方程式は、2種類の素粒子と1種類の素粒子の組み合わせで、Jones 多項式をペアで構成されている。Rico level 理論をベースにして、強い力と弱い力、電磁気力が、誤差を D-brane 間を重力が graviton として伝わっているのが、洩れて、反重力をこの Weak electric theory に加えると、全部の力が合わさって、統一場理論が完成される。これが、ヒッグス場とオイラーの定数を多様体を次元として、加群すると、ゼータ関数になる。これが、微分幾何の量子化とガンマ関数でもある。

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta + \bar{h}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

素数分布論が、ゼータ関数を開集合として同型でもあり、ノルム空間、経路積分として、位相差が原子間力にもなっている。このエネルギー分布が、素数と素粒子方程式、宇宙と異次元のフェルミオンとして、一致している。このエネルギー分布が、複素多様体である。

$$\square = 2(\cos(ix \log x) + i \sin(ix \log x))$$

宇宙と異次元が、双対性として、超対称性の素粒子を形成している。

[reference Lisa Randall, Toshihide Masuoka, Makoto Kobayashi]

WSL are replaced with Bridge level of surface
Windows and Linux native code of being installed
with aspective surface on Windows Masaaki Yamaguchi

バーチャルマシンに OS を入れて、Xserver で Linux を使うか、WSL で Linux を使うかの今では、どちらかを選択することになるが、私のアイデアでは、ハードウェアレベルでデュアルブートで、Windows 側では、表として、裏では Linux を入れるのは、VMware と同じだが、根本的に違うのは、Windows 側でブリッジをホライズンブリッジにして、その裏に Linux が存在しているコンピュータを考えている。この方法だと、バイナリだろうがソースコードだろうがネイティブの OS が味わえられる。Tab+Windowskey で、仮想空間に存在している他人のネットワークにある OS が 3D 地図になり、経路としての route 経脈が見えて、その末端にある OS にアクセスできる。これがウルトラネットワークである。Surface on Windows off Linux の構成は、Windows と Linux をキー操作だけで、表面の Windows と裏の Linux が自由自在に使え。WSL の System とは違う仕組みになっている。このアイデアが以下の説明で端的に表されている。

The idea is virtualmachine live with dualboot mode in hardware level, and this environment OS confusion surface with Windows and Linux. Therefore this complex OS combine on Windows aspect with Hardware switch, This surface level of Windows convert with bridge level in Linux of non surface, and this bridge level combinart with data of virtual machine, the controll with bridge aspect off being Linux would seen with Windows on surface aspective of ground level controll in taskbar. This taskbar insentate with application protocol into gateway protocol of X being used for Windows level of aspect in hardware.

These informate of Hardware of combine with Windows and Linux confusion on OS boot for X server like Astec-X and VMware, WSL application replace with this bridge and gateway and application protocol being used of surface of aspective level on hardware being seen like in already install and use for replace of VMware and X servers.

This idea is merit that WSL build with OS level transport on Windows, and this idea that surface preinstall of aspective Windows after this setting of Linux, double OS in native OS true, then Windows aspect of being used with Linux native, this Linux is user oneself level being setting with.

And This Linux of aspective hardware after Windows installed, and this user level install of Linux like Ubuntu Debian Kali, Red hat linux, this linux are hardware of Windows aspective level on surface. This system is non equal with WSL system.

数学と数がオイラーの定数から生まれた Masaaki Yamaguchi

$$2^2 = e^{x \log x} = 4$$

$$3^3 = e^{x \log x} = 27$$

$$4^4 = e^{x \log x} = 256$$

$$5^5 = e^{x \log x} = 3125$$

$$y = x \log x, \log x \rightarrow (\log x)^{-1}$$

$$\frac{1}{2} + iy = \frac{x \log x}{\log x}$$

$$x^{\frac{1}{2} + iy} = e^{x \log x}$$

$$\frac{1}{2} + iy = \log_x e^{x \log x} = \log_x 4, \log_x 27, \log_x 256, \log_x 3125$$

$$y = e^{x \log x} = \sqrt{a}$$

$$e^{e^{x \log x}} = a, e^{(x \log x)^2}$$

$$x^2 = \pm a, \lim_{n \rightarrow \infty} (x - y) = e^{x \log x}$$

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

$$y = x \log x, \log x \rightarrow (\log x)^{-1}$$

$$||ds^2|| \ 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

と宇宙の中の1種の原子をみつける正確さがこの式と、

$$y = \frac{x \log x}{(\log x)} = x$$

と、 $x \log x = a$ から $\frac{a}{(\log x)} \rightarrow x$ と x を抜き取る。この x をみつけるのに $x^{\frac{1}{2} + iy} = e^{x \log x} \frac{1}{2} + iy = \frac{x \log x}{(\log x)} = x$ としてこの x をみつける式がゼータ関数である。

ゼータ関数は、量子暗号にもなっていることと、この式自体が公開鍵暗号文にもなっている。

$$\frac{1}{2} + iy = \frac{x \log x}{\log x}$$

この式が一次独立であるためには。

$$x = \frac{1}{2}, iy = 0$$

がゼータ関数となる必要十分条件でもある。

$$\int C dx_m = 0$$

$$\begin{aligned}\frac{d}{df} \int C dx_m &= 0^{0'} \\ &= e^{x \log x}\end{aligned}$$

と標数0の体の上の代数多様体でもあり、このオイラーの定数からの大域的微分多様体から数が生まれた。

$$\begin{aligned}H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\ &\bigoplus a^f x^{1-f} [I_m] \\ &= \int e^x x^{1-t} dx_m \\ &= e^{x \log x}\end{aligned}$$

アメリカ大統領を統計で選ぶ選挙は、reco level 理論がゼータ関数として機能する遷移エネルギーの安定軌道がある集団 x に対数 $\log x$ の組み合わせとして、指数の巨大確率を対数の個数とするこの大統領の素質としての x^n 集団の共通の思考が n となるこの n がどのくらいのエントロピー量かを $H = -Kp \log p$ が表している。

$$\begin{aligned}\int \Gamma(\gamma)' dx_m &= (e^f + e^{-f}) \geq (e^f - e^{-f}) \\ (e^f + e^{-f}) &\geq (e^f - e^{-f})\end{aligned}$$

この方程式はブラックホールのシュバルツシルト半径から

$$\begin{aligned}(e^f + e^{-f})(e^f - e^{-f}) &= 0 \\ \frac{d}{df} F \cdot \int C dx_m &\geq 0 \\ y = f(g(x)') dx &= \int f(x)' g(x)' dx \\ y = f(\log x)' dx &= f'(x) \frac{1}{x} \\ y &= \frac{f'(x)}{x} \\ &= 2(\cos(ix \log x) - i \sin(ix \log x)) \\ &= C \\ c = f(x) \cdot \log x, dx_m &= (\log x)^{-1} \\ C = \frac{d}{\gamma} \Gamma &= \bigoplus (i\hbar^\nabla)^{\oplus L}\end{aligned}$$

となり、ヴェイユ予想の式からも導かれる。

$$\begin{aligned}&= 2(\cos(ix \log x) - i \sin(ix \log x)) \\ &e^\theta\end{aligned}$$

$$\frac{d}{d\gamma}\Gamma=\Gamma^{\gamma'}$$

$$=\int \Gamma(\gamma)'dx_m$$

$$y=f(x)\log x, y'=f'(x)+\frac{f'(x)}{x}$$

$$\sin(\log x)'dx=\cos(\log x)\cdot\frac{1}{x}$$

$$\sin \frac{y}{x} = \sin \vec{u} = a + t \sin \vec{u}$$

$$i,-i,2i,-2i$$

$$\lim_{n\rightarrow\infty}(f(b)-f(a))=f'(c)(b-a)$$

$$\sin(\log x)'dx=\cos(\log x)\cdot\frac{1}{x}$$


$$y=f(x)\log x, y'=f'(x)+\frac{f'(x)}{x}$$

$$\sin(\log x)'dx=\cos(\log x)\cdot\frac{1}{x}$$

$$\sin \frac{y}{x} = 2(\cos(ix \log x) - i \sin(ix \log x))$$

$$= e^{\theta}$$

Major trancinazer and Miner trancinazer effective result Masaaki Yamaguchi



major trancinaizer influent with miner trancinaizer, and major trancinaizer cover on spiritual mind from other persons. And this medicine tend to interact with other miner trancinazer. This phenomenon reach for reject with miner trancinazer in fact, appear from take medicine of person have this miner trancinazer to drink the medicine, and this drink of the one is readed from other society of counceller persons need to cure from spiritual mind of damage in days of flastrations, However this miner trancinazer medicine not only noon of days time of out of mind to security persons but also take a life of bussiness of pressers of time spent after this medicine change with effect of relax in the mind. miner trancinaizer have tendence to reduce of Na of bloud, and this Na reduce result from one take a medicine, after this one of person is appeared from other cure of person see. major trancinazer have interacted from miner trancinazer and double essence of cure of relax release into one take this medicine.

Selen and Tell are study of Kyoto university in professors of metal of chemistry chains, more also this chains are drills of badam with whisperd. Selen is progarded with mental condition, Tell is inspirate

with acknowledge in aquire of possibility with holographic datas. This two metal chemistry are Rantanoid chains. And pair of chains are Actinoid of metal chains. These two chains are one class of spectrum zones. More also this two of metal chains are Toliraphone and Whipacs. Toliraphone is same with Tell, Whipacs is Selen chemistry. I tendency have to be like see a beautiffl formed of selen organic chemistry geinu to out of world. This selen form of system is being tend to chain for stress out of pressures. Therefore, this Hidroxy and O_2 and R1, R2 of harmony of respond in access of relax and zen of envy. R1, R2 are Selen and organized of chains in alcorls with organic chemistry.

整数論から代数、そして幾何学 Masaaki Yamaguchi

ある整数を a とおく。この a を $\log x$ で割ると

$$x = \frac{a}{\log x}$$

で求め、ゼータ関数の x が導かれる。この x が、

$$a = x \log x$$

とシャノンのエントロピー値として、

$$\int a dx = \int x \log x dx$$

と、コルモゴロフ・シナイの値にもなる。これが、別の道でもある、求まるゼータ関数の式にもなる。この式から、

$$C = \int \left(\int \frac{1}{x^s} dx + \log x \right) d\text{vol}$$

このオイラーの定数を多様体積分すると

$$\int C dx_m = \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m + \int \log x d\text{vol}$$

となり、hartshorn conjecture の多世界理論となり、いろいろな式に分岐していく。

始めの x を見つけにくいので、オイラーの定数の多様体積分の分岐の式達がある。要するに、 x の式のヒントが幾多もある。

ガンマ関数を大域的微分多様体として共変微分すると、オイラーの定数を多様体積分した式と合流する。次元を多様体を基底群として、オイラーの定数とヒッグス場の式を加群すると、一般座標変換を大域的微分方程式として求めた結果と同型ともなる。

$$\int C dx_m + \frac{d}{df} F(x) = \Gamma^{\gamma'}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{\gamma'}$$

これらの式が Jones 多項式となり、経済理論を Reco level theory としての一般周期関数ともなる。

$$2i \sin(ix \log x) = e^f + e^{-f}$$

これらから、素数分布論が $x = \frac{a}{\log x}$ が基底群を成して、数として、オイラーの定数から、幾多の式になり、Jones 多項式を最大数と最小数の間を一般周期関数として数学を形成して、整数論が群論となり、代数、解析、幾何学を成り立たせている。

Global Integral manifold equal with
oneselves being divided with logment element Masaaki Yamaguchi

$$\begin{aligned}
\int \Gamma' dx &= \frac{\Gamma'}{\log x} = \Gamma \\
\int a dx &= \frac{a}{\log x} = \Gamma \\
\Gamma &= \int e^{-x} x^{1-t} dx \\
a = 1, a &= \int e^{-x} x^{1-t} dx, 1 < a \\
\frac{1}{\log x} &= \vee \int \wedge (R + \nabla_i \nabla_j f)^x \\
&= \frac{\wedge (R + \nabla_i \nabla_j f)^n}{\exists (R + \nabla_i \nabla_j f \circ g)^n} \\
\wedge (R + \nabla_i \nabla_j f)^x &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
\vee (R + \nabla_i \nabla_j f)^n &= \int \frac{\wedge (R + \nabla_i \nabla_j)^2}{\exists (R + \Delta f)^n} \\
\int \gamma dx_m &= \frac{\gamma}{\log x} = \Gamma \\
\frac{d}{df_m} \gamma &= \frac{\gamma}{\log x} \\
&= \frac{\Gamma'}{\log x} \\
&= \int e^{-x} x^{1-t} dx \\
\Gamma' &= \int e^{-x} x^{1-t} \log x dx
\end{aligned}$$

All of essential number is constructed with primer of pair, this element have two of pair. Thus is, 1 and n are exceed with prime priset of nature in world of mathematics series. This set of number exceed with prime number, and this of one is divide with logment of this pair number. moreover this pair of number has with already of a in this conceal with being decided to resolve with a for this deviation of x. After all, this redivide of x is concerned for being Differential geometry of quantum level quality equale with Gamma function.

Prime number of value have with all of pair in count of $\log x$, and this count of number divide with one element, exceed with this value call $\log x$ to decide with different of value oneselved.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$2(y \log y)^{\frac{1}{2}}$ have all of primer number and Higgs field + Euler product equal with three particle element, moreover this element also constructed with sixth particle of three of two in one pair. And this concerned of element in Fermat theorem have relativity with Euler equation of reverse in Imaginary of complex equation. Also this equation call one to equal with circle function of themselves. Resolved with all of Global differential manifold are Jones manifold of equation. Particle of two pair are constructed with three of one' element, this element are three of pair in two of particles. Fermat theorem are $n \leq 3 x^n + y^n \leq z^n$, and real of result with conceal is preset of element, and imaginary value of resulted of conceal are particle with complex manifold, and this formation is super symmetry element. Dimension of number relate with Goaris equation of extension, and this extend of over resolved value are complex resulted of elements.

大域的微分方程式についてのレポート Masaaki Yamaguchi, with my son

非対称性理論によるパリティの破れから、宇宙と異次元にはそれぞれ、重力と反重力が存在しているが、ヒッグス場の方程式に宇宙での数式を入力すると、ゼータ関数だけが出力される。対称性の仕組みから異次元が対として生成される。これに関連する CP 対称性の破れは、反粒子と粒子がそれぞれ、弦理論から結合している宇宙と異次元から、反粒子が消えたのは、結合した結果、宇宙のゼータ関数が出力されたためでもある。ヒッグス場の式から、宇宙と異次元は、それぞれ別に存在している。Ω における p 形式による外微分が、 $\mathcal{L}_x(d\Omega) = d(\mathcal{L}\Omega)$ と同値により、次の方程式が成り立つ。

$$\begin{aligned}\frac{d}{df}F &= e^{x \log x} \\ \frac{d}{df}F &= F^{(f)'} \\ &= m(x), x^{\frac{1}{2}+iy} = e^{x \log x}\end{aligned}$$

この式からゼータ関数がガロワ拡大に導かれて、ヒッグス場の式と同値になる。

$$\begin{aligned}f(x) + g(y) &\geq 2\sqrt{f(x)g(y)} \\ \frac{d}{df}F &\geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \log(x \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= 2(y \log y)^{\frac{1}{2}} \\ \frac{d}{df}F(x_m) &= \frac{\partial}{\partial f}F(x_m), \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ \frac{d}{df}F(y_m) &= \frac{\partial}{\partial f}F(y_m), \left(\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \right)^{(f)'} \\ \bigoplus \frac{\mathcal{H}\Psi}{\nabla \mathcal{L}} &= \frac{1}{\frac{d}{df}F} \\ &= \frac{d}{d\Lambda} \lambda\end{aligned}$$

ヒッグス場の式を逆関数で求めると、世界面を切断した値を場として、張力として求められる。この値を単体分割した τ を多様体になると、多様体の計量で超関数を正規部分群で組み合わせ多様体として、ウィークスケールによるエネルギーにこの逆関数は至る。

$$\begin{aligned}T^{\mu\nu} &= G^{\mu\nu} + \frac{1}{2}g_{ij}\Lambda \\ f &= \tau = (T^{\mu\nu})^{-1} \\ T^{\mu\nu} &= \int \tau(x, y) dx_m dy_m\end{aligned}$$

$$\begin{aligned}\bigoplus (\mathcal{H}\Psi^\nabla)^{\oplus L} &= i\hbar\psi \\ &= \bigoplus (i\hbar^\nabla)^{\oplus L}\end{aligned}$$

微分幾何の量子化は、ハイゼンベルク方程式の確率振幅における、ウィークスケールによるエネルギー最小単位のエントロピー不変量でもある。ヒッグス場のエネルギーの確率振幅でもある。ヒッグス場のエネルギーが加群分解して再接続されてもいる。大域的微分方程式とも同型である。

$$F = \frac{1}{4}x^4 + C, f = x^3, e^{i\theta} = \cos \theta + i \sin \theta$$

$$F = -\cos \theta, f = \sin \theta$$

$$\frac{d}{df}F = \left(\frac{1}{4}x^4 + C\right)^{(x^3)}, \frac{d}{df}F = (-\cos \theta)^{(\sin \theta)}$$

$$\left(\frac{1}{4}x^4 + C\right)^{(x^3)} = \left(\frac{1}{4}x^4 + C\right)^{(x^3)'}$$

$$\frac{1}{4}(x^3)^{x^3}x + \frac{1}{4}x^4(x^x)^3 = \frac{1}{4}e^{f \log f} + \frac{1}{4}x^4(e^{x \log x})^3$$

$$= \frac{1}{4}(x^{3x^3})' \cdot x^{(x \cdot x^3)'}$$

$$= \frac{1}{4}x^{x^3 \log x^3'} \cdot x^{x^4 \log x'}$$

$$= \frac{3x^2}{4}e^{3x^2 \log x} \cdot (3x^2 \log x + 1)$$

$$\frac{d}{df}F = (-\cos \theta)^{(\sin \theta)} = (-\sin \theta)^{(\sin \theta)'}$$

$$= (f)^{'(f)} = ((f)^f)'$$

$$= e^{x \log x}$$