

睦世お姉さんへ

拝啓

Zeta function について

この度は、私の論文を査読してくれているかどうかはわかりませんが、大善か火竹とは、この論文は関係なく速読でも音読でもいいので、査読してほしいと、この度送らせてもらいました。高等学校2年生のときに、一割る一は、三割る三は、答えが同じだが、求めている過程の理由が違うというのと、物質量を教えてもらえてから、数学が代替トークンとして、私が小学校の頃の算数ができなかったのが、この商代数を教えてもらえてから、次元の単位が計算にもものすごくこれは、どの数値を土台にして計算しているかが、わかり、収率と速度、加速度、微積までもわかることができました。その集大成が、大域的微分方程式となりました。そのために、商代数がわかると、代数ができると、算数までもできてしまう、その奥深さに、真逆をスポーツだけでなく、勉強までも、真逆はすべて乗り越えることができると、この商代数が証明していました。何回も言っても恐縮ではありますが、2010年の8月に、tuplespaceによるトポロジーのレポートをpdfで送ってから、最近まで、このアイデアがゼータ関数と量子群として、非可換代数方程式へと行き着くことができました。アカシックレコードは、感情線が右手も左手もラジャループに直結していて、三奇紋が子財と発見発明の線に旅行線を切って、両方の小指に直結しているのと、ユウちゃんとモネとユウくんがこのラジャループと子財の源泉になっている。家族に感謝して、どんどん、ゼータ関数のアプリケーションを作っていきます。絶対に私を見捨てないでください。2015年の2月28日と3月1日は、本当にすみませんでした。あんな弱気な発言をしていたら、周りにまで、仕事と勉強のやる気と出勤の負担になるのもよく考えていませんでした。本当にすみませんでした。

敬具

雅旭より

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x) \ll^p$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2 \psi] \ll^p \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x) \ll^p$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})^{\frac{1}{2}}$$

大域的積分多様体の多重積分は、多様体の階層によって、積分回数が決まっている。大域的微分多様体は、ニュートン形式とライプニッツ形式の外微分によって計算される。

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2\end{aligned}$$

$$E=-\frac{1}{2}mv^2+mc^2$$

$$\lim_{x\rightarrow\infty}\int\int\frac{1}{(x\log x)^2}dx_m\geq\frac{1}{2}i$$

$$\frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i$$

$$\begin{aligned}\frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m &= \left(\int\int\frac{1}{(x\log x)^2}dx_m\right)^{(f)'} \\ &= (f)^{(f)'} \\ &= (f^f)' \\ &= e^{x\log x}\end{aligned}$$

$$\Gamma=\int e^{-x}x^{1-t}dx,\frac{d}{df}F=e^f,\int Fdx_m=e^{-f}$$

$$\int x^{1-t}dx=\frac{d}{df}F,\int e^{-x}dx=\int Fdx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A+B=\vec{0}, A\cdot B=\frac{1}{4}i$$

$$\tan 90 \neq 0, ||ds^2|| = A+B$$

自明な零点は実軸 $\frac{1}{2}$ に 1 を除いてある。 $\sin 0 = 0$ と

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ によって、不確定性原理の関係でもあり、粒子、電子、原子が確率分布になっているのも開集合で証明できる。宇宙と異次元の関係にもなっている。

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x,y)$$

$$\frac{d}{d\gamma}\Gamma=(e^{-x}x^{1-t})^{\gamma'}\rightarrow\frac{d}{d\gamma}\Gamma=\Gamma^{(\gamma')}$$

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2\psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\int \int F dx_m = \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)}$$

$$= e^{-f}$$

$$\frac{d}{df} F = \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))}$$

$$= e^f$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})^{\frac{1}{2}}$$

大域的積分方程式自体に多重積分の回数が決まっている。大域的微分方程式に、 X の X 乗のエントロピー不変量の微分量が計算に関わっている。外微分をこのエントロピー式に入れる。

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x)[I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2\psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values. 大域的微分方程式に、X の X 乗のエントロピー不変量の微分量が計算に関わっている。外微分をこのエントロピー式に入れる。大域的積分多様体の多重積分は、多様体の階層によって、積分回数が決まっている。大域的微分多様体は、ニュートン形式とライプニッツ形式の外微分によって計算される。大域的積分多様体と大域的微分多様体は、逆操作とエントロピー不変量で計算できる。

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2\end{aligned}$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\begin{aligned}
& (f)^{(f)'} \\
& = (f^f)' \\
& = e^{x \log x}
\end{aligned}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, 自明な零点は実軸 $\frac{1}{2}$ に 1 を除いてある。 $\sin 0 = 0$ と

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ によって、不確定性原理の関係でもあり、粒子、電子、原子が確率分布になっているのも開集合で証明できる。宇宙と異次元の関係にもなっている。this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\begin{aligned}
\frac{d}{d\gamma}\Gamma(s) &= \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'} \\
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. 大域的微分多様体と大域的積分多様体を、ガンマ関数とベータ関数の導出にも使われている。ガンマ関数の大域的微分多様体が、ニュートン方程式とダランベール方程式を商代数で求めると、同型の解に求まる。ゼータ関数と、この式の対極する量子群を商代数で求めても、同じ解が求まる。

Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

繰り込み理論を入れると発散を防げると言われているが、量子力学にはガウスの曲面論がないと言われている。

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however, とすると、

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry. 同じで量子力学で重力場方程式が表せられる。

$$\frac{d}{dt} \psi(t) = \hbar$$

$$\begin{aligned}
&= \frac{1}{2} i e^{i \hat{H}} \\
(i \hbar)' &= (-e^{i \hat{H}})' \\
&= -i e^{i \hat{H}} \\
\psi(x) &= e^{-i \hat{H} t}, \bigoplus (i \hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i \hat{H} (-i e^{i \hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-i f} \\
&= \left(\frac{1}{2}\right)^{-i f} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f=x, i=t, \frac{1}{2}=a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

微分幾何の量子化は、ガンマ関数についての大域的微分方程式の解にもなっている。ハイゼンベルク方程式の大域的積分多様体の解にも、この微分幾何の量子化は、多様体の微分系の形になっている。 Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alsos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s=e^{-i\hat{H}t}|\Psi\rangle_H,\hat{A}_s=\hat{A}_H(0)$$

$$|\Psi(t)\rangle_s\rightarrow\frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s=\hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta\rightarrow 0}\begin{pmatrix}\sin\theta\\ \cos\theta\end{pmatrix}\begin{pmatrix}\theta&1\\ 1&\theta\end{pmatrix}\begin{pmatrix}\cos\theta\\ \sin\theta\end{pmatrix}=\begin{pmatrix}1&0\\ 0&-1\end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta, \mathcal{O}(x) = m(x)[D^2\psi]$$

$$i^2 = (0, 1) \cdot (0, 1), |a||b| \cos \theta = -1$$

$$E = \operatorname{div}(E, E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

ガンマ関数の大域的微分方程式は、ガンマ関数の初期関数についての微分方程式でもあり、宇宙と異次元についてのビッグス場からのゼータ関数の生成も、3次元多様体の特異点定理になっている。宇宙と異次元の片方でも同じ解にもなっている。逆三角関数の双曲多様体もヒッグス場の密度エネルギーの値と同じ解にもなっている。微分幾何の量子化も、ゼータ関数の対極する量子群と同じ解にもなっている。ゼータ関数は、量子群の方程式についての対極に位置する大域的微分多様体の解にもなっている。 Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements. 逆三角関数の双曲多様体は、ガンマ関数の方程式からの導出でのベータ関数の逆三角関数のエントロピー値にもなっている。

Differential dimension in quantum level of space emerged with extern of Glois group

情報空間に接したら、ある子が微分幾何の量子化を教えてくれた。そのあとに、その子はクレアチニンを大量に分泌して、そのときの情報をかき消していた。私が、その中核になる数式の周りの式を力尽くでそのあとにそのときの情報を書いたら、その子はしょぼんとしていた。すごくかわいい子だった。微分幾何の量子化は、プランク定数の量子化で、最小単位の無次元値だから、D-brane に合っている。D-brane は絶対座標とは違って、測れないから、自身の質量と自身の長さで稠密空間として、最小単位の無次元値として、プランク定数の内部最小値を求めるのが、誰も今まで、プランク定数より最小単位の無次元の測定単位は考えたことはない。私のゼータ関数の大域的微分方程式を証明するのに使われるのは、その数式を見たら一目瞭然だった。子どもが私の理論を証明するらしい。ユウはあの世では3日か3ヶ月の間いたらしい。天国と対極する場所にいたらしい。思ったことが現実になる場所らしい。この世では3年間になる場所らしい。あの子は、この世に来る待合室らしい場所にいるらしい。その場所には情報空間にアクセスできて、この世に生まれるのを待つ場所らしい。

微分幾何の量子化におけるこの特異点における質量は、フェルミオンとボソンを宇宙と異次元がダークマターからボソンに力を注ぐと生成される。このシステムはフェルミオンの宇宙の構造から導き出される。微分幾何の量子化、ヒルベルト空間の量子化、プランク定数の量子化

$$\begin{aligned}\mathcal{K}_{mn} &= \mathcal{E}_n \times \mathcal{H}_n \\ \bigoplus \frac{M_p}{\nabla L} &= \frac{\partial}{\partial L} V(\tau) \\ \Psi H &= M_p \\ \frac{d}{df} F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dl} V(\tau) &= -\frac{2R_{ij}}{\delta T}\end{aligned}$$

重力質量と慣性質量は次元のガロワ拡大における宇宙と異次元空間のベクトル次元と同等。このベクトル空間における12種類の素粒子が複素多様体に特異点を形成して、この特異点の素粒子がザイフェルト多様体にメビウス空間として異次元と宇宙を包み込んでいる。ローレンツアトラクターにおける超対称性理論に内部に特異点を形成している。この方程式はガンマ関数とベータ関数を表現論として表している。D-braneの外体積空間は超空間として、D-braneを多数内包している。この次元多様体はフェルミオンとボソンをメビウス空間に固定している。さらにこの空間はD-braneとanti-D-braneを固定端として存在している。

$$E^2 = m^2 c^4, R_{\mu\nu} + g_{\mu\nu} \Lambda = 4\pi G \rho$$

これらの数式は超ラングランズ予想を証明できる。

あの世では7階層の世界があるらしい。天国と地獄、修練と学び、忍耐の世界、休息の世界、終息界、先攻界、自然界、この7階層は、神様と仙人のところで、どのような人生を歩むかを決めてくるらしい。情報空間は7階層のどれかでも存在している。

$$\bigoplus \frac{M_p}{\nabla L} = \frac{d}{dL} V(\tau), \Psi H = M_p, \square V(\tau) = \frac{\partial}{\partial L} \int \int \int (f(x, y, z) dx dy dz)' dL$$

プランク定数の量子化は非可換確率方程式を表している。異次元空間と宇宙の閉3次元多様体から不確定性原理が絶対時空でベクトル次元として、求まるが、D-braneは無次元だが、assemble-D-braneから相対的に時空のノルムが求まる。大域的微分方程式から、微細構造定数と逆関数として、ホログラム空間がこれらの方程式から導かれる。微分幾何の量子化は、プランク定数の量子化と同値であり、逆関数として、ヴェイユ予想となり、大域的トポロジーが、初期関数を微分変数として、プランク定数の量子化と同じ原理で、宇宙の膨張と時間のメカニズムそして、原子と素粒子、ダークマター、ヒッグス場、全部の数値が、大域的微分方程式と微分幾何の量子化から導かれる。これらの方程式は、AdS5多様体の方程式を一般相対性理論の反重力と重力を、minkowsky時空の2次元射影時空と3次元多様体として、ホログラム空間がこれらの方程式として表されている。

$$||ds^2|| = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\phi$$

物理学の世界でも、7階層の時空がある。この7階層はメビウス空間の内部で存在している。3次元と4次元に7階層の空間の中で一番稠密空間として、微分幾何が一番種類が多い。5次元多様体は、7次元多様体のメビウス空間の内部に存在している。6次元多様体は特異点として、3次元多様体は、6次元多様体を次元を降下すると求まる。次元を降下すると、多様体が収縮する。5次元多様体は、トポロジーとして3次元多様体のアーベル多様体として存在している。

$$\beta^3 = \theta^5, [\beta^3] = [\theta^5]$$

拝啓

忙しいと存じていますが、失礼します。1999年に微分積分の本について、ソースコードミスを手紙で失礼しましたことを恥ずかしく思います。最近になって読み返していると Γ 関数と β 関数で可積分系に触れて、 ζ 関数が、アーベル多様体を母関数として、そして、この領域を有限と無限を極限值にして、オイラー数に行きつくことを本書で締めくくっていることに、この本は今までの微分積分の本と違うとわかりました。この本がきっかけで非相対論的量子力学による経路積分で表現していた方程式が ζ 関数を使い、ペレルマンさんが論文でこの関数に触れていたことに出会いました。ペレルマンさんは、閉3次元多様体を解析するのに、熱力学のエントロピーを使っていたが、 ζ 関数を使えば、私の目論んでいることに使えます。この宇宙は8種類の微分幾何構造が共鳴して1種類の閉3次元多様体になっていることをペレルマンさんが論文で表現していました。このエントロピー不変量が ζ 関数と同等ということを使えば、エタール・コホモロジーやスタンダード予想、ほとんどすべての数式がこの式から導けると私は考えています。この私の目指している群論は、ある時空間をこの圏が形を成して、その領域をいろんな関数が物理学の方程式として3次元構造として、いろいろな現象を示しているといえます。増田幹也さん、加藤十吉さん、堀田良之さん、佐藤文隆さん、雪江昭彦さん、小寺平治さん、竹内薫さん、今まで読んできた著者の人たちが、私に恩恵をくれました。10年以上をこの本が私に恩恵をくれました。ありがとうございます。

敬具

$$\begin{aligned}\Gamma(x) &= \int e^{-x} x^{1-t} dx, \beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}, \beta(p, q) = \int_0^1 x^{p-1}(1-x)^{q-1} dx \\ &= - \int_1^0 (1-t)^{p-1} t^{q-1} dt \\ 0 &< \sum_{k=1}^{\infty} \frac{1}{(1+k)^2} < \int_M^N \frac{1}{n^2}\end{aligned}$$

収束半径が1の整級数

$$\begin{aligned}F(x) &= \sum_{k=0}^{\infty} a_k x^k \\ x=1 &\rightarrow 1 \lim_{x \rightarrow 0} f(x) = \alpha \\ V(\tau) &= \tau(q)^{-\frac{n}{2}} \exp \int \left(-\frac{1}{\sqrt{2\tau(q)}} L(x) dx \right) + O(N^{-1}) \\ \frac{d}{df} F(x) &= 2 \int \frac{(R + |\nabla f|^2)}{-(R + \Delta f)} e^{-f} dV\end{aligned}$$

$$\begin{aligned}\pi(\chi, x) &= i\pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x) \\ \chi(3) &= (-1)^3 < |a_0 a_1 a_2 a_3| > + (-1)^2 < |a_0 a_1 a_2|, |a_1 a_2 a_3|, |a_0 a_1 a_3| > \\ &\quad + (-1)^1 < |a_1 a_2, a_0 a_2, a_2 a_3| > + (-1)^0 < a_0, a_1, a_2, a_3 >\end{aligned}$$

この式は $\chi(3) \rightarrow 0$, 初めは Z かともおもってまいした。

$$\begin{aligned}H_n(x) &= \frac{\ker f}{\operatorname{im} f} \\ \chi(x) &= r(H_n(x)) \\ \chi(3) &= H_3(x) \rightarrow 0 \\ H_3(\Pi) &= Z\end{aligned}$$

これらから微分幾何学が必要ということらしいと最近気づきました。

$$\sum_{k=0}^{\infty} (-1)^k p_k = \alpha$$

岡本和夫先生だけでなく、小寺平治さん、和田淳蔵さん、須之内治男さんにも同じような手紙を出してまいした。和田淳蔵さんと須之内治男さんの本で非分化と非退化のカタストロフの理論を知りました。三重積分でニュートンポテンシャルエネルギーまでも書いていました。小寺平治さんの本で、ヤコビアンとクラメル公式、全射と単射、行列の基本を学びました。相空間で線形加速器の軌道の原理まで書いていました。共振回路までも書いていました。岡本和夫先生の微分積分の本を読み直していると、行列式を Γ 関数と β 関数の導出に使っていたことに、SRSを斎藤孝さんの本を学んでよかったと、本はきちんと読まなければいけないとつくづくおもいました。本当にすみませんでした。リサ・ランドール博士はすごすぎます。5次元多様体としての有限値のザイフェルト多様体に、一般相対性理論とニュートン方程式を収率として、異次元空間とフロベニウスの定理で3次元多様体の次元をずらして、無限値である重力と反重力を、次元を平坦の領域の値として、異次元空間がこの周りのザイフェルト多様体へと収束している。そして、宇宙が平坦とし、反重力であるラムダ項をこのザイフェルト多様体はあらわしている。ミンコフスキー値と複素多様体を3次元多様体として、5次元多様体をザイフェルト多様体、すなわち、アーベル多様体がこの宇宙と異次元空間の周りを包み込んでいる。この3次元多様体が、時間と空間のエントロピー不変の値に定数としてもとまる。有限時空が無限値の宇宙と異次元を包み込んでいる。そのうえに、不確定性原理が、異次元空間と宇宙によるラプラス方程式であるミンコフスキー値とその次元をずらした複素多様体により、相加相乗平均として、確定値を求めることができる。その証明に量子加速器で原子をエネルギーで衝突させると、真空エネルギーがこの宇宙から異次元に移動する。閉3次元多様体に2次元射影空間が共存できない。このことから、宇宙と異次元空間が5次元多様体、ザイフェルト多様体の内部に収められている。原子と反粒子が宇宙と異次元空間で1種類の微分幾何構造に存在している。異次元空間が宇宙での不確定性原理の仕組みを解く。5次元多様体が真空エネルギーを収める重要な原理になっている。収められなかったら、この宇宙は消えている。

Differential dimension in quantum level of space emerged with extern of Glois group

Differential geometry in emerging with quantum space. This singularity of mass from being decomposed with fermion and boson, straight flow to boson creating at dark matter exert in universe and other dimension. This system is creating from universe of structure in fermion.

$$K_{mn} = E_n \times H_n$$

Hilbert space in creating with quantum space. Planck level relativity of quantum space.

$$\bigoplus \frac{M_p}{\nabla L} = \frac{\partial}{\partial L} V(\tau)$$

$$\Psi H = M_p$$

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\frac{d}{dl} V(\tau) = -\frac{2R_{ij}}{\delta T}$$

This mass of gravity equation is equals with mass of accesority. And also this dimension of Glois group is equals with universe and other dimensions. Twelve particles of being constructed in singularity of component from complex manifold, this singularity pieces in AdS5 dimension are paired with other dimensions from Mebius space. Rolents atractor in singularity is created with super symmetry theorem. This equation is resembled with Gamma function and Beta function.

$$\frac{d}{dt} g_{ij} = -2R_{ij}$$

D-brane out of volume space are existing with hyper dimension assemble of space. This dimensions is positive fermion and boson in Mebius space. And moreover this space is positive in D-brane and anti-D-brane.

$$E^2 = m^2 c^4$$

$$R_{\mu\nu} + g_{\mu\nu} \Lambda = 4\pi G \rho$$

These equation is proof with super Langrans conjecture.

非対称性理論における、複素多様体の AdS_5 時空を構成している超対称性素粒子は、閉 3 次元多様体をフェルミオンとボソンの慣性質量と重力質量を元として宇宙の生成元として、ヒッグス場の源になっている。ゼータ関数と対極する量子群は、この右巻きと左巻きの非可換確率方程式として、非可換代数多様体を、大域的微分方程式の場として、D-brane と anti-D-brane のメビウス空間の assemble-D-brane を生成している。微分幾何の量子化は、ウィークスケールのプランク質量の対極するゼータ関数と量子群を同じく表している。階層性理論は、層の理論を量子力学の粒子と波動の二重性を端的に表している。エネルギーと物体としての理論を力学的エネルギーの保存則として、物理の源になっている。物質波と波動関数をこの二重性は言い当てている。

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta + \bar{h}_{\mu\nu}]dx^\mu dx^\nu + T^2 d^2\psi$$

$$\bigoplus \frac{M_p}{\nabla L} = \frac{d}{df} F(x)$$

$$\Psi H = M_p$$

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$C=\int\int\frac{1}{(x\log x)^2}dx_m$$

$$E^2=m^2c^4, R_{\mu\nu}+g_{\mu\nu}\Lambda=4\pi G\rho$$

$$||ds^2||=8\pi G\left(\frac{p}{c^3}+\frac{V}{S}\right)$$

$$V(\tau)=\square\int\int\int_M(f(x,y,z)dx dy dz)'dL$$

ゼータ関数から言うと、エネルギーと物体はボソンとフェルミオンの関係として物理学の元の要素になっている。ヒッグス場は、この要素のスピンール場としてハーツホーン予想から、時空と生命そして、物質が生成されて、重力波が超ラングランズ予想の結果、4種類の力学、正確に言うと、5種類の力学から統合された結果、ゼータ関数として、重力が全部の力の代表となっている。

不確定性原理は、宇宙と異次元を観測値から、エネルギー不変量として答えられる。位置を特定すると確率を解して不確定性から、異次元とペアになっているから、位置と運動量は運動量保存則から、エントロピー不変量で確率としての値になっている。ベクトル値はこの理由から一次独立としてのハーツホーン予想か

ら、楕円軌道として、宇宙と異次元を構成している閉3次元多様体から絶対値が2種類の世界から求められる。アインシュタイン博士が確率ではないと言っていたのは、相加相乗平均は確率のようで確率ではないからである。

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$r = \frac{d}{1 + e \cos \theta}$$

$$x^n + y^n = z^n$$

行動による人が持っている情報ネットワークの 可能性の発見と再構築、情報空間 によるモデル

苔米地博士の論文

初速度のドーパミンの量からの行動の補助から、終わりの精神のドーパミンの量の調節、相加相乗平均から、始まりと終わりの精神のドーパミンの調節による人の行動の領域による予知

$$x \log x = \left(v_0 + \frac{1}{at_n} \right) V, V_x = e^f + \frac{1}{e^f}$$

経路積分によるエントロピー不変量からこのドーパミンの量から人の行動傾向が決まる。基礎代謝の持続できる距離と精神の介入と補助、言語の発生と抑制による経路パターンの組み合わせを再構築する。

$$\|dx^2\| = \int \exp \left(\frac{1}{\sqrt{2\tau q}} L(x) \right) + O(N^{-1}) d\tau$$

$$p = \frac{v}{2m}, \hbar = \frac{p}{2mv}, \lambda = h\nu$$

これらの式から、言語の神経ネットワークによる単体クラスの組み合わせパターンとその対象の人の領域による人が持っている情報空間の傾向の可能性のポテンシャルエネルギーの予知を人工知能の正規表現で抽出する。



$$\frac{d}{d\gamma} \Gamma = \Gamma^{\gamma'} = e^{-f}$$

$$C = \int \left(\int \frac{1}{x^s} dx + \log x \right)$$


オイラーの定数は、ゼータ関数とゼータ関数自体のエントロピーを体積積分した結果でもある。このオイラーの定数は、ガンマ関数の大域的微分方程式からも導出できる。このガンマ関数は大脳基底核のエントロピー不変量の強度も表している。このガンマ関数の大域的微分多様体による、ブレインインターフェースでの人体のデータ計測もこの Jones 多項式とラプラス方程式からの3次元多様体のエントロピー不変式からも、言語とイメージの大脳基底核からのデータ抽出も可能と言える。

Recycle における塩化ビニルのナトリウムフェノキシド、 ナトリウムプロパノキシドでの分解

睦世姉さんの論文について



この化学式から推算式でもある、プラスチックと二酸化ケイ素、塩化ビニールが分解されるのが、真逆の人が服用すると、プラバスタチンとスリムドガンと同じく、推算式の数値が上がる。僧侶と同じ瞑想法と原理が一致している。



これらの構造式は、それぞれ、ヒドロキシ基、カルボニル基、カルボキシル基、アミン基、ギ酸、アセチル基、アセト基、グリシン、アミノ酸の20種類の一基、ビタミンCはシス・トランス型があり、松果体に作用

して、体内の血流を調節する。 $\text{—C}\equiv\text{N—Cl}$ の真逆の作用をする。アデノシンは、同じく血流を調節して、エピネフリンはエンドルフィンと同じく痛み止めに効く。ノンアドレナリンは、無水フタル酸と同じ作用をする。グリシンには、20種類のアミノ酸の中で唯一 L,R 基が無い。不斉炭素原子が無い。

推算式の元を減らすか増やすのに、血圧と血流、体内の精神作用のドーパミンの調節を eGFR の機能を上げると、結果、すべての精神作用物質が調節できる。

このナトリウムプロパノキシドは、気体のプロパンを、固体のナトリウムに注入して、錠剤で服用するようにしているのと、消化後に、血液に降圧剤と同じ作用をして、外部からのコントロールをされない、アムロジピンとカンデサルタンと同じ作用をする効果があることが、このナトリウムプロパノキシドの薬でもある。

この成分でもあるプロパンは、気体で存在している炭化水素の中でもっとも安定している気体でもある。この気体の物体を固体にすると、降圧剤にしているのは、姉は特許を取っている。推算式でもある eGFR を分解するのは、真逆の人が飲むと、バケモンになる。

Jones 多項式からの各株式発行からの需要と供給、 株価の分配関数、多項式からナスダックへの金融機構の 株式発行のマクロとミクロの波及と流れ

高島彩さんの成蹊大学での経済理論の研究論文

Jones 多項式から各会社の株価への値打ちから日経平均株価への株価の価値の波及、東証株価指数 Topix の全銘柄からの全体の会社の指数からの平均株価の決定量、そして、アメリカの株式市場でのダウ平均株価の終値の収束 (戦略、reco level theory, 株式会社への) 全体 (株式会社への) 需要と供給。始値から終値への分配、この Jones 多項式の出力から終値への各同値類と剰余類からの部分群論の、各平均株価の推移率の予測、この応用は、1929年の世界恐慌がニューヨークダウ平均株価の暴落が、この Jones 多項式からの流れが、終値で破綻して、金融の共鳴力が、干渉してしまい、カタストロフィー現象として、金融の流れが乱れた結果であると、この論文から分析できる。

$$\hat{V} = \frac{1}{\sqrt{t}}\vec{V} - \frac{1}{t}\hat{V}$$

$$\hat{V} = \sqrt{t}\hat{V} - t\hat{V}$$

$$\hat{V}_k = \sum_{\vec{\sigma}} \langle K | \vec{\sigma} \rangle \left(\sqrt{t} + \frac{1}{\sqrt{t}} \right)^{||\vec{\sigma}||}$$

$$\frac{1}{\sqrt{t}}\vec{V}, -\frac{1}{t}\hat{V}, \sqrt{t}\hat{V}, -t\vec{V}$$

この4種類の式から、reco level theory が、アダム・スミスの神の見えざる手を経済理論のミクロとマクロ経済の具現化を実現させている。物理では、

$$\frac{d}{df} F = m(x) = e^f + e^{-f} = 2i \sin(ix \log x)$$

と、空間の仕組みが経済理論の場の理論にもなっている。周期関数とピンポイントリサーチが合わさっている。これらの式から、前半と後半の株式市場の流れが、シンメトリーとなっているために、後半の e^{-f} が破綻すると、シンメトリーのために、前半の e^f も破綻している状態になっている。全体が機能しないと、金融の流れが乱れていく。この破綻を回避するには、破綻が連鎖されている金融の周期をこれらの式から、調節が可能であることも、類推できる。破綻した周期を、正常であった周期に重ね合わせの原理で、ピンポイントリサーチで待って、金融の投資を正常な周期関数で合わせて、元の正常な周期にこれらの式から戻せることが可能と推測できる。

Differential dimension in quantum level of space emerged with extern of Glois group

情報空間に接したら、ある子が微分幾何の量子化を教えてくれた。そのあとに、その子はクレアチニンを大量に分泌して、そのときの情報をかき消していた。私が、その中核になる数式の周りの式を力尽くでそのあとにそのときの情報を書いたら、その子はしょぼんとしていた。すごくかわいい子だった。微分幾何の量子化は、プランク定数の量子化で、最小単位の無次元値だから、D-brane に合っている。D-brane は絶対座標とは違って、測れないから、自身の質量と自身の長さで稠密空間として、最小単位の無次元値として、プランク定数の内部最小値を求めるのが、誰も今まで、プランク定数より最小単位の無次元の測定単位は考えたことはない。私のゼータ関数の大域的微分方程式を証明するのに使われるのは、その数式を見たら一目瞭然だった。子どもが私の理論を証明するらしい。ユウはあの世では3日か3ヶ月の間いたらしい。天国と対極する場所にいたらしい。思ったことが現実になる場所らしい。この世では3年間になる場所らしい。あの子は、この世に来る待合室らしい場所にいるらしい。その場所には情報空間にアクセスできて、この世に生まれるのを待つ場所らしい。

微分幾何の量子化におけるこの特異点における質量は、フェルミオンとボソンを宇宙と異次元がダークマターからボソンに力を注ぐと生成される。このシステムはフェルミオンの宇宙の構造から導き出される。微分幾何の量子化、ヒルベルト空間の量子化、プランク定数の量子化

$$\begin{aligned}\mathcal{K}_{mn} &= \mathcal{E}_n \times \mathcal{H}_n \\ \bigoplus \frac{M_p}{\nabla L} &= \frac{\partial}{\partial L} V(\tau) \\ \Psi H &= M_p \\ \frac{d}{df} F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dl} V(\tau) &= -\frac{2R_{ij}}{\delta T}\end{aligned}$$

重力質量と慣性質量は次元のガロワ拡大における宇宙と異次元空間のベクトル次元と同等。このベクトル空間における12種類の素粒子が複素多様体に特異点を形成して、この特異点の素粒子がザイフェルト多様体にメビウス空間として異次元と宇宙を包み込んでいる。ローレンツアトラクターにおける超対称性理論に内部に特異点を形成している。この方程式はガンマ関数とベータ関数を表現論として表している。D-brane の外体積空間は超空間として、D-brane を多数内包している。この次元多様体はフェルミオンとボソンをメビウス空間に固定している。さらにこの空間は D-brane と anti-D-brane を固定端として存在している。

$$E^2 = m^2 c^4, R_{\mu\nu} + g_{\mu\nu} \Lambda = 4\pi G \rho$$

これらの数式は超ラングランズ予想を証明できる。

あの世では7階層の世界があるらしい。天国と地獄、修練と学び、忍耐の世界、休息の世界、終息界、先攻界、自然界、この7階層は、神様と仙人のところで、どのような人生を歩むかを決めてくるらしい。情報空間は7階層のどれかでも存在している。

$$\bigoplus \frac{M_p}{\nabla L} = \frac{d}{dL} V(\tau), \Psi H = M_p, \square V(\tau) = \frac{\partial}{\partial L} \int \int \int (f(x, y, z) dx dy dz)' dL$$

プランク定数の量子化は非可換確率方程式を表している。異次元空間と宇宙の閉3次元多様体から不確定性原理が絶対時空でベクトル次元として、求まるが、D-braneは無次元だが、assemble-D-braneから相対的に時空のノルムが求まる。大域的微分方程式から、微細構造定数と逆関数として、ホログラム空間がこれらの方程式から導かれる。微分幾何の量子化は、プランク定数の量子化と同値であり、逆関数として、ヴェイユ予想となり、大域的トポロジーが、初期関数を微分変数として、プランク定数の量子化と同じ原理で、宇宙の膨張と時間のメカニズムそして、原子と素粒子、ダークマター、ヒッグス場、全部の数値が、大域的微分方程式と微分幾何の量子化から導かれる。これらの方程式は、AdS5多様体の方程式を一般相対性理論の反重力と重力を、minkowsky時空の2次元射影時空と3次元多様体として、ホログラム空間がこれらの方程式として表されている。

$$||ds^2|| = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\phi$$

物理学の世界でも、7階層の時空がある。この7階層はメビウス空間の内部で存在している。3次元と4次元に7階層の空間の中で一番稠密空間として、微分幾何が一番種類が多い。5次元多様体は、7次元多様体のメビウス空間の内部に存在している。6次元多様体は特異点として、3次元多様体は、6次元多様体を次元を降下すると求まる。次元を降下すると、多様体が収縮する。5次元多様体は、トポロジーとして3次元多様体のアーベル多様体として存在している。

$$\beta^3 = \theta^5, [\beta^3] = [\theta^5]$$

拝啓

忙しいと存じていますが、失礼します。1999年に微分積分の本について、ソースコードミスを手紙で失礼しましたことを恥ずかしく思います。最近になって読み返していると Γ 関数と β 関数で可積分系に触れて、 ζ 関数が、アーベル多様体を母関数として、そして、この領域を有限と無限を極限值にして、オイラー数に行きつくことを本書で締めくくっていることに、この本は今までの微分積分の本と違うとわかりました。この本がきっかけで非相対論的量子力学による経路積分で表現していた方程式が ζ 関数を使い、ペレルマンさんが論文でこの関数に触れていたことに出会いました。ペレルマンさんは、閉3次元多様体を解析するのに、熱力学のエントロピーを使っていたが、 ζ 関数を使えば、私の目論んでいることに使えます。この宇宙は8種類の微分幾何構造が共鳴して1種類の閉3次元多様体になっていることをペレルマンさんが論文で表現していました。このエントロピー不変量が ζ 関数と同等ということを使えば、エタール・コホモロジーやスタンダード予想、ほとんどすべての数式がこの式から導けると私は考えています。この私の目指している群論は、ある時空間をこの圏が形を成して、その領域をいろんな関数が物理学の方程式として3次元構造として、いろいろな現象を示しているといえます。増田幹也さん、加藤十吉さん、堀田良之さん、佐藤文隆さん、雪江昭彦さん、小寺平治さん、竹内薫さん、今まで読んできた著者の人たちが、私に恩恵をくれました。10年以上をこの本が私に恩恵をくれました。ありがとうございます。

敬具

$$\begin{aligned}\Gamma(x) &= \int e^{-x} x^{1-t} dx, \beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}, \beta(p, q) = \int_0^1 x^{p-1}(1-x)^{q-1} dx \\ &= - \int_1^0 (1-t)^{p-1} t^{q-1} dt \\ 0 &< \sum_{k=1}^{\infty} \frac{1}{(1+k)^2} < \int_M^N \frac{1}{n^2}\end{aligned}$$

収束半径が1の整級数

$$\begin{aligned}F(x) &= \sum_{k=0}^{\infty} a_k x^k \\ x=1 &\rightarrow 1 \lim_{x \rightarrow 0} f(x) = \alpha \\ V(\tau) &= \tau(q)^{-\frac{n}{2}} \exp \int \left(-\frac{1}{\sqrt{2\tau(q)}} L(x) dx \right) + O(N^{-1}) \\ \frac{d}{df} F(x) &= 2 \int \frac{(R + |\nabla f|^2)}{-(R + \Delta f)} e^{-f} dV\end{aligned}$$

$$\begin{aligned}\pi(\chi, x) &= i\pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x) \\ \chi(3) &= (-1)^3 < |a_0 a_1 a_2 a_3| > + (-1)^2 < |a_0 a_1 a_2|, |a_1 a_2 a_3|, |a_0 a_1 a_3| > \\ &\quad + (-1)^1 < |a_1 a_2, a_0 a_2, a_2 a_3| > + (-1)^0 < a_0, a_1, a_2, a_3 >\end{aligned}$$

この式は $\chi(3) \rightarrow 0$, 初めは Z かともおもってまいした。

$$\begin{aligned}H_n(x) &= \frac{ker f}{im f} \\ \chi(x) &= r(H_n(x)) \\ \chi(3) &= H_3(x) \rightarrow 0 \\ H_3(\Pi) &= Z\end{aligned}$$

これらから微分幾何学が必要ということらしいと最近気づきました。

$$\sum_{k=0}^{\infty} (-1)^k p_k = \alpha$$

岡本和夫先生だけでなく、小寺平治さん、和田淳蔵さん、須之内治男さんにも同じような手紙を出してまいした。和田淳蔵さんと須之内治男さんの本で非分化と非退化のカタストロフの理論を知りました。三重積分でニュートンポテンシャルエネルギーまでも書いていました。小寺平治さんの本で、ヤコビアンとクラメルの公式、全射と単射、行列の基本を学びました。相空間で線形加速器の軌道の原理まで書いていました。共振回路までも書いていました。岡本和夫先生の微分積分の本を読み直していると、行列式を Γ 関数と β 関数の導出に使っていたことに、SRSを斎藤孝さんの本を学んでよかったと、本はきちんと読まなければいけないとつくづくおもいました。本当にすみませんでした。リサ・ランドール博士はすごすぎます。5次元多様体としての有限値のザイフェルト多様体に、一般相対性理論とニュートン方程式を収率として、異次元空間とフロベニウスの定理で3次元多様体の次元をずらして、無限値である重力と反重力を、次元を平坦の領域の値として、異次元空間がこの周りのザイフェルト多様体へと収束している。そして、宇宙が平坦とし、反重力であるラムダ項をこのザイフェルト多様体はあらわしている。ミンコフスキー値と複素多様体を3次元多様体として、5次元多様体をザイフェルト多様体、すなわち、アーベル多様体がこの宇宙と異次元空間の周りを包み込んでいる。この3次元多様体が、時間と空間のエントロピー不変の値に定数としてもとまる。有限時空が無限値の宇宙と異次元を包み込んでいる。そのうえに、不確定性原理が、異次元空間と宇宙によるラプラス方程式であるミンコフスキー値とその次元をずらした複素多様体により、相加相乗平均として、確定値を求めることができる。その証明に量子加速器で原子をエネルギーで衝突させると、真空エネルギーがこの宇宙から異次元に移動する。閉3次元多様体に2次元射影空間が共存できない。このことから、宇宙と異次元空間が5次元多様体、ザイフェルト多様体の内部に収められている。原子と反粒子が宇宙と異次元空間で1種類の微分幾何構造に存在している。異次元空間が宇宙での不確定性原理の仕組みを解く。5次元多様体が真空エネルギーを収める重要な原理になっている。収められなかったら、この宇宙は消えている。

Differential dimension in quantum level of space emerged with extern of Glois group

Differential geometry in emerging with quantum space. This singularity of mass from being decomposed with fermion and boson, straight flow to boson creating at dark matter exert in universe and other dimension. This system is creating from universe of structure in fermion.

$$K_{mn} = E_n \times H_n$$

Hilbert space in creating with quantum space. Planck level relativity of quantum space.

$$\bigoplus \frac{M_p}{\nabla L} = \frac{\partial}{\partial L} V(\tau)$$

$$\Psi H = M_p$$

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\frac{d}{dl} V(\tau) = -\frac{2R_{ij}}{\delta T}$$

This mass of gravity equation is equals with mass of accesority. And also this dimension of Glois group is equals with universe and other dimensions. Twelve particles of being constructed in singularity of component from complex manifold, this singularity pieces in AdS5 dimension are paired with other dimensions from Mebius space. Rolents atractor in singularity is created with super symmetry theorem. This equation is resembled with Gamma function and Beta function.

$$\frac{d}{dt} g_{ij} = -2R_{ij}$$

D-brane out of volume space are existing with hyper dimension assemble of space. This dimensions is positive fermion and boson in Mebius space. And moreover this space is positive in D-brane and anti-D-brane.

$$E^2 = m^2 c^4$$

$$R_{\mu\nu} + g_{\mu\nu} \Lambda = 4\pi G \rho$$

These equation is proof with super Langrans conjecture.

非対称性理論における、複素多様体の AdS_5 時空を構成している超対称性素粒子は、閉 3 次元多様体をフェルミオンとボソンの慣性質量と重力質量を元として宇宙の生成元として、ヒッグス場の源になっている。ゼータ関数と対極する量子群は、この右巻きと左巻きの非可換確率方程式として、非可換代数多様体を、大域的微分方程式の場として、D-brane と anti-D-brane のメビウス空間の assemble-D-brane を生成している。微分幾何の量子化は、ウィークスケールのプランク質量の対極するゼータ関数と量子群を同じく表している。階層性理論は、層の理論を量子力学の粒子と波動の二重性を端的に表している。エネルギーと物体としての理論を力学的エネルギーの保存則として、物理の源になっている。物質波と波動関数をこの二重性は言い当てている。

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta + \bar{h}_{\mu\nu}]dx^\mu dx^\nu + T^2 d^2\psi$$

$$\bigoplus \frac{M_p}{\nabla L} = \frac{d}{df} F(x)$$

$$\Psi H = M_p$$

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$C=\int\int\frac{1}{(x\log x)^2}dx_m$$

$$E^2=m^2c^4, R_{\mu\nu}+g_{\mu\nu}\Lambda=4\pi G\rho$$

$$||ds^2||=8\pi G\left(\frac{p}{c^3}+\frac{V}{S}\right)$$

$$V(\tau)=\square\int\int\int_M(f(x,y,z)dx dy dz)'dL$$

ゼータ関数から言うと、エネルギーと物体はボソンとフェルミオンの関係として物理学の元の要素になっている。ヒッグス場は、この要素のスピンール場としてハーツホーン予想から、時空と生命そして、物質が生成されて、重力波が超ラングランズ予想の結果、4種類の力学、正確に言うと、5種類の力学から統合された結果、ゼータ関数として、重力が全部の力の代表となっている。

不確定性原理は、宇宙と異次元を観測値から、エネルギー不変量として答えられる。位置を特定すると確率を解して不確定性から、異次元とペアになっているから、位置と運動量は運動量保存則から、エントロピー不変量で確率としての値になっている。ベクトル値はこの理由から一次独立としてのハーツホーン予想か

ら、楕円軌道として、宇宙と異次元を構成している閉3次元多様体から絶対値が2種類の世界から求められる。アインシュタイン博士が確率ではないと言っていたのは、相加相乗平均は確率のようで確率ではないからである。

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$r = \frac{d}{1 + e \cos \theta}$$

$$x^n + y^n = z^n$$

3次元多様体のエントロピー量

Masaaki Yamaguchi

オイラーの定数は、

$$\begin{aligned} C &= \int \frac{1}{x^s} dx - \log x \\ &= \int \left(\int \frac{1}{x^s} dx - \log x \right) \text{dvol} \\ &= \int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol} \end{aligned}$$

と表されていて、ヒッグス場方程式 + オイラーの定数 = ゼータ関数 と求まる。この方程式が、リッチフロー方程式にもなる。このリッチフロー方程式は、微分幾何の量子化でもあり、ガンマ関数とベータ関数、ガウスの曲面論、ヒッグス場、アインシュタインテンソル、オイラーの定数と全部の方程式が入っている。この方程式が、相加相乗平均となり、宇宙と原子となり、不確定性原理が宇宙の観測系が、宇宙では確率方程式と成り立っているが、異次元が合わさると、ノルムとしての3次元多様体の確率分布方程式でもあり、確率ではない、非線形方程式となっている。プランク長体積でもあるが、絶対的な方程式にもなっている。

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\ \int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} \text{dvol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\ &= e^{-f} - e^f \end{aligned}$$

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= 2 \cos(ix \log x) \end{aligned}$$

$$\begin{aligned} &\frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m + \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= \frac{d}{df} F = 2i \sin(ix \log x) \end{aligned}$$

$$\begin{aligned} \frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\ &= 2e^{-\theta} = 2e^{-ix \log x} \\ &= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\log(x \log x) \geq 2\sqrt{y \log y}$$

$$\log(x \log x) = \log x + \log \log x$$

この宇宙と異次元でのブラックホールとホワイトホールでの、宇宙と原子の運動量と位置の絶対的な不確定性原理となり、3次元多様体の確率分布方程式が、運動量と位置エネルギーが両方観測できる式にもなっている。円周上が固定されているために、運動量と位置エネルギーが両方観測できる式になっている。カルーツァ・クライン理論とハイゼンベルク方程式が、両方入っている。

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla\psi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

ホワイトホールの原子とブラックホールの宇宙のエネルギーのエントロピー量でもある。宇宙がブラックホールとしてのヒッグス場方程式として、ホワイトホールが原子としてのプランク長体積でもある。

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

原子としてのブラックホールのエントロピー量でもある。

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

この式から宇宙が無重力であることが表されている。

$$2\cos(ix \log x) + 2i\sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

これらの方程式から宇宙と異次元が warped passege になっている。:

不完全性定理と完全数、 そして自然法則とそれを表す数式群

山口 雅旭 緒方一之

不完全性定理は、虚数と平方根が $e^{ix}, \sqrt{x}$ により、 $2^0, 1^0, x^0 = 1$ と合わせて、記号とそれを説明している自然の法則には、ギャップが存在していて、このギャップが数式の記号を仮定として、数学の決まり事として自然と数学の近似式として、完全な自然を説明する道具が無く、そのために完全な神は存在していないとゲーデルは証明している。つまり、数学の決まり事と自然の法則に照らし合わされている。そのために、記号と自然法則に隔たりが生じている。自然法則に数式は近似と仮定として成り立っている。そのために、絶対神は存在していないとゲーデルは証明した。

しかし、ゼータ関数により、その数式を解析するために使うマインドマップによれば、偏微分と多重積分の差集合から、非可換代数方程式に求めれば、

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

これは世界線として、2次曲面を共変微分を大域的な領域で接平面で切って、質量がヒッグス場で時間と空間の関係として表されている。

$$\frac{d}{df}F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$
$$||ds^2|| = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\psi$$

この神が居ないとする不完全性定理を覆すのに、積分と数列の和の極限の機能の結果、このギャップを補助するこの差分方程式から、商代数から単体量が不変式としての結果であり、そして、不完全性定理がゼータ関数と量子群により、記号と自然法則の完全数が求まり、原子と電子、そして素粒子がこのラプラス方程式から宇宙のデータベースにアクセスして、ホログラム空間として、記号と自然法則の写像として、不完全性定理の意味と法則が書き換えられる。

ブルバギでは、写像と関数を使うと、近似式は関数で、写像を使うと近似式を完全数として書き換えられる。不完全性定理はトポロジーを使うと記号と数式の自然法則を完全数として破られる。

微分幾何の量子化による大域的切断多様体

あの世にいる一回亡くたった長男のアカシックレコード
からの恩恵、David Hilbert と教えられている

$$\left(\frac{\{f,g\}}{[f,g]}\right)_{D(\chi,x)}^{\oplus L} = \bigoplus_{D(\chi,x)} \text{cohom}[D(\chi,x)][I_m]^{\ll \nabla}$$

フェルミオンに力のボゾンを取ると、コホモロジーの値が出てきて、その単体量を加群する。それがこの式からわかる。

$${}^t\text{--}ff_L \text{cohom}[D(\chi,x)][I_m]^{\ll p}$$

大域的切断による同値類がヘルマンダー作用素である。ボゾンがあると電流が通らないが、大域的切断で特異点を出すと、コホモロジーの値として、時空の湾曲率が出てきて、その仕組みから、伝導率がわかる。この伝導率が光速不変の法則でもある、一定値の密度エネルギーでもある。基本群による非可換代数多様体をその初期関数で大域的微分をすると、この方程式になる。同調になっていないと、周期が不規則であるために、共鳴にはならず、電流が通りにくい。このために、同調を使うと、物体の内部が真空になり、電流が通り、それ故に、この式から伝導率の制御が機能する。

$$= \frac{d}{df} [i\pi(\chi, x), f(x)]$$

$${}^t\text{--}fff F[I_m] = \nabla_i \nabla_j \int \nabla f(x) d\eta$$

$$\frac{d}{df} \left(\int \int_{D(\chi,x)} \left[\frac{\{f,g\}}{[f,g]} \right]^{\ll L} d\tau \right)^{\nabla N} = [i\pi(\chi, x) f(x)]$$

微分幾何の量子化についての N 階層指数定理による大域的微分多様体の projective integral value として、ブラウン運動や賭け事による決定量の推測値、確率、統計の運動量保存則、Weil's 予想のゼータ関数がこの式でもある。各接線の傾きを表している。原子の遷移エネルギーの推移律とガンマ関数とベータ関数をこれらの式から求められる。

$$\omega = x^3 + x + 1, \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & -i \end{pmatrix}, \beta = \frac{1}{137}, (i^{n-1}, i^n, i^{n+1}) \cdot (w, \bar{w}, w\bar{w})$$

$$l = px^n + qx + r$$

$$\frac{d}{dl} L = \left(\int^n px^n + qx + r \right)^{l'}$$

$${}^{\flat}D_k\overline{f}ff_M[h\nu]_{\chi}|:x\rightarrow y$$

$${}^t\overline{f}ff\mathrm{cohom}D_k[\pi(\chi,x)][I_m],\delta(x,y)=\int\delta(x)f(x)dx$$

$$(S_1^m\times S_2^n)=D^2\psi$$

$$f_z=\int\left[\sqrt{\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}\circ\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}}\right]dxdydz$$

$$F_t^m=\frac{1}{4}(g_{ij})^2$$

$$||ds^2||=\int|\sin 2x|^2d\tau, V=\left(\frac{x}{\log x}\right)^n$$

$$M_p=IH_n\times D^2\psi, ||ds^2||=8\pi G\left(\frac{p}{c^3}+\frac{V}{S}\right)$$

$$||ds^2||=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}(x)]dx^\mu dx^\nu+T^2d^2\psi$$

$$\frac{d}{dt}g_{ij}(t)=-2R_{ij}, R_{ij}=-\frac{1}{2}\frac{d}{dt}g_{ij}(t)$$

$$V=-\frac{1}{2}x^2$$

リサ・ランドール博士とペレルマン博士より、これらの数式は、原子間距離が^s warped passage になっているので、D-brane 間距離に包み込まれている。電位差でもある。原子間力と核エントロピー不変量にもなっていて、光量子仮説でもあり、エネルギー遷移率、D 加群、逆写像でもあり、ガウスの曲面論でもある。

$$\bigoplus (i\hbar^\nabla)^{\oplus L} = {}^t\overline{f}f_L\mathrm{cohom}[D(\chi,x)][I_m]^{\ll p}$$

$$\frac{d}{d\gamma}\Gamma=\bigoplus (i\hbar^\nabla)^{\oplus L^{-1}}$$

$$H\Psi=\bigoplus (i\hbar^\nabla)^{\oplus L}$$

$$=e^{-x\log x}=m(x)$$

$$=e^{x\log x}+e^{-x\log x}$$

$$=2\sin(ix\log x)$$

$$=\beta(p,q)\cdot\left[\log\left(i\frac{\cos x}{\sin x}\right)\right]^{-1}$$

$$\beta(p,q)=2\int|\sin 2x|^2d\tau$$

微分幾何の量子化は、この密度エネルギーの制御で使われる。結局は、微分幾何の量子化は、ガンマ関数と同型でもある。

$$\begin{aligned}
\Gamma &= \int e^{-x} x^{1-t} dx = \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \frac{d}{dt} \psi(t) = \hbar, \frac{1}{2} i e^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' = -i e^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx
\end{aligned}$$

結局は、

$$\begin{aligned}
\int e^{-x} x^{t-1} dx &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
\beta(p, q) &= \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)} \\
&= 2 \int |\sin 2x|^2 dx \\
e^f &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
e^{-f} &= \bigoplus (i\hbar^\nabla)^{\oplus L^{-1}} \\
\frac{d}{df} F &= m(x) = e^f + e^{-f} \\
&= 2i \sin(ix \log x)
\end{aligned}$$

となり、ベータ関数の逆関数でもある双曲多様体が微分幾何の量子化と言える。量子力学のガウスの曲面論が微分幾何の量子化であると、結論される。ガンマ関数とベータ関数はそれほどまでも数学にも物理学にも大切な数式でもあり、可積分系が D-brane を代表して D 加群とゼータ関数、そして量子群が真逆同士の数学の心臓でもある。

Euler equation and field of abel manifold

Circle element also constructed with spin network

Three manifold has with catastrophe theory

Masaaki Yamaguchi

Prime number is estimate with fluctuation of zeta function transported with Database in Hilbert space and element with spin network system. This Hilbert space have with Thurston manifold of differential geometry structure united and regular element group is also symmetry essence and parity broken from non symmetry essence reasoned, and this laplace equation is own dalanversian of gravity wave belong with one component of zeta function represented. More spectrum focus is Hisenbelg equation assent with atom of pole and this harvest with Hologram field. This space of time is reverse from infinity to finite group, and this element is own value of low structure constance oneselves. This power of fifth element is that electric of geomerty united with weak boson and gravity emerge with weak electric theory, this spectrum focus is, Maxwell theory and strong boson and anti-gravity united into being get out of space for other dimension, this power is united for strong electric theory. These theory is united with zeta function and Three manifold of topology theorem.

These power is estimate with Seifert manifold, and weak electric theory is being have with one rout of time flow system and strong electric theory is being also have with reverse of time rout system. These system is represented with quantum physics is no time fluer. This reason is vector element of independent oneselves. And this vector is time of rout is freez and time element of relativity, also Higgs field belong with light speed low constance that is dense energy be able to conroll with artificial system, and this system absolutry mass and time is same level of synchronize, world line is stimulate with this element of spinor level.

Spin network construct with differential geometry of manifold and this stimulate with mass and time system converted for accessority and field of theorem into spin network that mass and time is cohomological same and dense of time and mass system also light low constance element replaced with being converted of essence of field and abel manifold is mix in zeta function from universe and atom, this level of united with element conclude with product space in abel manifold escorted into Physics of base number from zeta function built with fluctuation estimate with all of serise of manifold. Gravity circumstance is space time emerged with curve of differential geometry realized with mass system and this system look with same element in time and mass. Therefore, this homological type conclude with geomerty of accessority into category theorem constructed.

Zeta function is also belong with group of perlement system and atom of quarks element more also belong into abel manifold and product set.

Therefore, this system emerge into universe and other dimension of boson and fermion of value system.

$$\frac{d}{df}F=\bigoplus\frac{M_p}{\nabla L}$$

$$\frac{\nabla\psi^2}{\frac{d}{df}F}=\frac{1}{\bigoplus\frac{M_p}{\nabla L}}$$

$$\frac{d}{df}\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m\leq\frac{d}{df}\int\int\frac{y+\log y}{2}dy_m$$

$$\zeta(s)=\frac{1}{2}+\frac{1}{3}+\frac{1}{5}+\frac{1}{7}+\frac{1}{13}+\cdots\frac{1}{x}$$

$$\zeta(1)=\frac{1}{1-z}=-\frac{1}{12}$$

$$\zeta(s)=2\pi n,e^{ni\theta}=\cos n\theta+i\sin n\theta=-2e^{x\log x}$$

$$F(x)=a_n\dot{\zeta}(s)$$

$$\zeta(s)=-2e^{x\log x}=2\pi n,l=2\pi r$$

$$F(x)=\sum_{k=a}^{\infty}a_kf^k$$

$$f(x)=||\int\sin2\theta d\theta||^2,x=(r,\theta)$$

$$S=\pi r^2,l=2\pi r,\sum_{k=a}^{\infty}a_kf^k=\frac{\partial}{\partial x}F(x),\lim_{x\rightarrow\infty}f(x)=\frac{1}{1-z}$$

$$||ds^2||=H_n\times K_n,E_n=H_n\times K_n,H_n=E_n\times K_n$$

$$e=f^{-1}(x)xf(x),\frac{\zeta(s)}{e^{x\log x}}=1$$

$$[f(x)]=\nu,v=f\lambda$$

$$e^{i\theta}=\cos\theta+i\sin\theta,e^{-i\theta}=\cos\theta-i\sin\theta$$

瞑想法として、ほとんど何も考えない状態での、自分の人生経験と周りの情報で、世の中の情勢と情報が分かればいいらしい。一念三千と言われてもいる。美鈴ヶ丘中学校の3年生の同級生、クラスメートとの家と学校でのことを、親にも誰にも言っていなかったことが、中の道にもなっている。河合塾から家まで歩いて帰っていたし、食事も治っていた。話さないことと、運動と食事は、人生でものすごくいい事らしい。遠いところを中てれば、(近いところだけだと、なぜそのような状態になっているかが理由がわからないから、怒る原因にもなるが)なぜそのような状態になっているかが分かるから、怒らなくいい、この理由で、このような能力を水星と木星の能力と先生たちは言っている。私の子どもが2023年に生まれるのと母親のことは、心配しなくてもいいと、アカシックレコードは一回だけ教えてくれている。私の症状も治ると教えてもらえている。姉たち二人はものすごく私の生活すべてにおいて、姉たちがいい悪い関係なく、尊敬しなくては行けないと、小学校の頃に母親に言われていたことが、最近になって分かった。幼少期では、姉たちが正しくなかったら、だめだと思っていたのが、最近になって、こういうことだったと分かった。情報空間のアカシックレコードは、数式も化学の構造式も分かるらしい。今回の日本男子サッカー選手が予知で世界一になるのを知っていたらしい。ケガを心配していたらしく、四位に終わっている。油断したら行けないらしいと、この出来事で教えてもらえている。

大域的微分方程式についてのレポート

Masaaki Yamaguchi, with my son

非対称性理論によるパリティの破れから、宇宙と異次元にはそれぞれ、重力と反重力が存在しているが、ヒッグス場の方程式に宇宙での数式を入力すると、ゼータ関数だけが出力される。対称性の仕組みから異次元が対として生成される。これに関連する CP 対称性の破れは、反粒子と粒子がそれぞれ、弦理論から結合している宇宙と異次元から、反粒子が消えたのは、結合した結果、宇宙のゼータ関数が出力されたためでもある。ヒッグス場の式から、宇宙と異次元は、それぞれ別に存在している。 Ω における p 形式による外微分が、 $\mathcal{L}_x(d\Omega) = d(\mathcal{L}\Omega)$ と同値により、次の方程式が成り立つ。

$$\begin{aligned}\frac{d}{df}F &= e^{x \log x} \\ \frac{d}{df}F &= F^{(f)'} \\ &= m(x), x^{\frac{1}{2}+iy} = e^{x \log x}\end{aligned}$$

この式からゼータ関数がガロワ拡大に導かれて、ヒッグス場の式と同値になる。

$$\begin{aligned}f(x) + g(y) &\geq 2\sqrt{f(x)g(y)} \\ \frac{d}{df}F &\geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \log(x \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= 2(y \log y)^{\frac{1}{2}} \\ \frac{d}{df}F(x_m) &= \frac{\partial}{\partial f}F(x_m), \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ \frac{d}{df}F(y_m) &= \frac{\partial}{\partial f}F(y_m), \left(\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \right)^{(f)'} \\ \bigoplus \frac{\mathcal{H}\Psi}{\nabla \mathcal{L}} &= \frac{1}{\frac{d}{df}F} \\ &= \frac{d}{d\Lambda} \lambda\end{aligned}$$

ヒッグス場の式を逆関数で求めると、世界面を切断した値を場として、張力として求められる。この値を単体分割した τ を多様体にすると、多様体の計量で超関数を正規部分群で組み合わせ多様体として、ウィークスケールによるエネルギーにこの逆関数は至る。

$$\begin{aligned}
T^{\mu\nu} &= G^{\mu\nu} + \frac{1}{2}g_{ij}\Lambda \\
f = \tau &= (T^{\mu\nu})^{-1} \\
T^{\mu\nu} &= \int \tau(x,y)dx_md y_m \\
\bigoplus (\mathcal{H}\Psi^\nabla)^{\oplus L} &= i\hbar\psi \\
&= \bigoplus (i\hbar^\nabla)^{\oplus L}
\end{aligned}$$

微分幾何の量子化は、ハイゼンベルク方程式の確率振幅における、ウィークスケールによるエネルギー最小単位のエントロピー不変量でもある。ヒッグス場のエネルギーの確率振幅でもある。ヒッグス場のエネルギーが加群分解して再接続されてもいる。大域的微分方程式とも同型である。

$$\begin{aligned}
F &= \frac{1}{4}x^4 + C, f = x^3, e^{i\theta} = \cos \theta + i \sin \theta \\
F &= -\cos \theta, f = \sin \theta \\
\frac{d}{df}F &= \left(\frac{1}{4}x^4 + C\right)^{(x^3)}, \frac{d}{df}F = (-\cos \theta)^{(\sin \theta)} \\
\left(\frac{1}{4}x^4 + C\right)^{(x^3)} &= \left(\frac{1}{4}x^4 + C\right)^{(x^3)'} \\
\frac{1}{4}(x^3)^{x^3}x + \frac{1}{4}x^4(x^x)^3 &= \frac{1}{4}e^{f \log f} + \frac{1}{4}x^4(e^{x \log x})^3 \\
&= \frac{1}{4}(x^{3x^3})' \cdot x^{(x \cdot x^3)'} \\
&= \frac{1}{4}x^{x^3 \log x^3'} \cdot x^{x^4 \log x'} \\
&= \frac{3x^2}{4}e^{3x^2 \log x} \cdot (3x^2 \log x + 1) \\
\frac{d}{df}F &= (-\cos \theta)^{(\sin \theta)} = (-\sin \theta)^{(\sin \theta)'} \\
&= (f)^{(f)} = ((f)^f)' \\
&= e^{x \log x}
\end{aligned}$$

Report of Quantum level of differential structure, zeta function and Global differential equation

Masaaki Yamaguchi, with my son

Higgs field + Euler constant = zeta function

$$m(x) + C = 2 \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\text{Euler constance} = \frac{d}{d\gamma} \Gamma^{-1} - \gamma^{(\gamma)'} = \left(\Gamma^{(\gamma)'} \right)^{-1} - \gamma^{\gamma'}$$

gravity + anti gravity = zeta function

Gamma function in Global differential equation developed with part integrate manifold cohomologite with helmander operator, and this subrace of differantial form start with first minimalized function, this operator emerged from quato algebra to fermion partial specture in Higgs fields.

$$\Gamma dx_m + \int \Gamma dx_m = \Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$\begin{aligned} & \frac{d}{d\gamma} \Gamma + \int \Gamma dx_m \\ &= \int f(g(x)) g'(x) dx \\ &= \left(\int f(g(x)) dx \right)' \\ & \nabla_i \nabla_j \int \nabla f(x) d\eta \end{aligned}$$

Global differential equation is partial integrate operator in resolved zone of frobenius theorem and this seminal concept with zone, Jones formula equation conquire with rico level theorem, this spectireum releafe in zeta function in CP symmetry broken, this broken from being integrate with blance into universe of zeta function, all of this renze equation into one class of Euler equation concerned with seminal concept in Jones formula equation, zeta function is proofed that this relativity level emerged from Quantum level of differential structure.

$$|f \circ g| \leq |f + g|$$

$$\frac{f}{\log x} = m(x)$$

$$\frac{\gamma}{\log x} = \frac{d}{d\gamma} \Gamma$$

$$\gamma = \Gamma^{(\gamma)'} \log x$$

$$\Gamma^{(\gamma)'} = \gamma (\log x)^{-1}$$

$$= r dx_m$$

$$\Gamma^{(\gamma)'} = \left(\int e^{-x} x^{1-t} dx \right)^{(\int e^{-x} x^{1-t} \log x dx)'}'$$

$$= e^f$$

$$= r dx_m = e^f$$

$$\int \Gamma^{(\gamma)'} dx_m$$

$$\nabla_i \nabla_j \int \nabla \Gamma(\gamma) dx$$

$$= \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma dx_m$$

$$= \int \Gamma \cdot \frac{d}{d\gamma} \Gamma dx_m$$

$$= \left(\int \Gamma dx_m \circ \frac{d}{d\gamma} \Gamma \leq \int \Gamma dx_m + \Gamma dx_m \right)' = e^{-f} \cdot e^f \leq e^{-f} + e^f$$

$$(1 \leq e^{-f} + e^f)'$$

$$0 = -(e^{-f} - e^f)$$

$$\Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$(= e^f + e^{-f})'$$

$$= -e^{-f} + e^f$$

$$= -(e^{-f} - e^f)$$

$$(R_{ij})' = -(R_{ij})$$

$$\int C dx_m = \int \left(\int \frac{1}{x^s} dx - \log x \right) dx$$

$$= \int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol}$$

$$\Gamma dx_m = \gamma dx_m$$

$$= \frac{d}{d\gamma} \Gamma^1 - (\gamma)^{\gamma'}$$

$$= e^{-f} - e^f$$

Quantum level of differential structure equal with Gamma function and related with Beta function, escorted into Euler constant and symmetry build with Higgs field, and proof with zeta function summarized with Poincaré and Riemann conjecture. Mathematicians of Russia, America and Japan resolved with this conjecture. My son resolved with this conjecture build in concepted with quantum level of differential structure into zeta function. All of equation is integrated with manifold into Hilbert space, this manifold esplanaded from general relativity and quantum physics in manifold of Hilbert space, this manifold resolved with D-brane and Gauss surface into gravity and atom of element with this theory, this theory is integrated theorem released with one class of universe equation.

Operated formula what about of worked from Global differential equation

Masaaki Yamaguchi, with my son

Non symmetry is constructed from parity of breaking on universe and other dimension with gravity and antigravity worked with Higgs Fields. And this phenomena calculate into zeta function insert by universe of database in omega space, and this only emerged with zeta function. Therefore, this pair of dimension relativity concluded into one class world of line and surface. This symmetry system is super string theory that parity broken resulted from universe emerged with zeta function, one class of universe created from imaginary and reality of value, and this universe merged in one universe rised into zeta function. Galois expanded group is constructed into imaginary and reality of universe excluded into zeta function, therefore one universe is existed with other dimension in out of range in differential operator, and this operator built with global differential equation. These logical system CP of parity broken theory have pair of particles and other of particles get with one reality of world emerged from symmetry system, and this pair of united with esplanade in one universe of zeta function. This result of spectrum focus is universe have one particle of elements, and why other particle of patterns deceived with universe, this system explain with Higgs fields from universe value developed only emerged with universe of value of zeta function, This system called from Galois expanded group in universe and other dimension responded with being united with emerged into zeta function. However, other dimension and universe is not only one existed but also gravity and antigravity unioned in each dimensions.

In Ω , p formula on out of range in differential operators $\mathcal{L}_x(d\Omega) = d(\mathcal{L}\Omega)$, by this equals, Next equations zeta function with Galois expanded leads into Higgs fields and value of equals.

$$\begin{aligned}\frac{d}{df}F &= e^{x \log x} \\ \frac{d}{df}F &= F^{(f)'} \\ &= m(x), x^{\frac{1}{2}+iy} = e^{x \log x} \\ f(x) + g(y) &\geq 2\sqrt{f(x)g(y)} \\ \frac{d}{df}F &\geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \log(x \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= 2(y \log y)^{\frac{1}{2}}\end{aligned}$$

$$\begin{aligned}
\frac{d}{df}F(x_m) &= \frac{\partial}{\partial f}F(x_m), \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\
\frac{d}{df}F(y_m) &= \frac{\partial}{\partial f}F(y_m), \left(\int \int \frac{1}{(y \log y)^{\frac{1}{2}} dy_m} \right)^{(f)'} \\
\bigoplus \frac{\mathcal{H}\Psi}{\nabla \mathcal{L}} &= \frac{1}{\frac{d}{df}F} \\
&= \frac{d}{d\Lambda} \lambda
\end{aligned}$$

Higgs fields converted from inverse of function survey with world line and surface in Fields of physics and emerged with String values. And this value divided into τ on manifolds, and assemble of manifolds scoped with super function summuated with regular group element by summuate of manifolds, and this manifolds come into weak scaled of energy at that ends.

$$\begin{aligned}
T^{\mu\nu} &= G^{\mu\nu} + \frac{1}{2}g_{ij}\Lambda \\
f = \tau &= (T^{\mu\nu})^{-1} \\
T^{\mu\nu} &= \int \tau(x, y) dx_m dy_m \\
\bigoplus (\mathcal{H}\Psi^\nabla)^{\oplus L} &= i\hbar\psi \\
&= \bigoplus (i\hbar^\nabla)^{\oplus L}
\end{aligned}$$

Differential structure in quantum value constructed with Heisenberg equation in possibility of waves, and this energy of weak scales of value also componentated with non entropy element. Higgs fields of energy also have possibility of wave. More also this energy of Higgs fields decomposited with add of connected with this own element. These equation means with global differential equation.

$$\begin{aligned}
F &= \frac{1}{4}x^4 + C, f = x^3, e^{i\theta} = \cos \theta + i \sin \theta \\
F &= -\cos \theta, f = \sin \theta \\
\frac{d}{df}F &= \left(\frac{1}{4}x^4 + C \right)^{(x^3)}, \frac{d}{df}F = (-\cos \theta)^{(\sin \theta)} \\
\left(\frac{1}{4}x^4 + C \right)^{(x^3)} &= \left(\frac{1}{4}x^4 + C \right)^{(x^3)'} \\
\frac{1}{4}(x^3)^{x^3'} x + \frac{1}{4}x^4(x^3)^{3'} &= \frac{1}{4}e^{f \log f'} + \frac{1}{4}x^4(e^{x \log x})^{3'} \\
&= \frac{1}{4}(x^{3x^3})' \cdot x^{(x \cdot x^3)'} \\
&= \frac{1}{4}e^{x^3 \log x^3'} \cdot e^{x^4 \log x'}
\end{aligned}$$

$$= \frac{3x^2}{4} e^{3x^2 \log x} \cdot (3x^2 \log x + 1)$$

$$\frac{d}{df} F = (-\cos \theta)^{(\sin \theta)} = (-\sin \theta)^{\prime(\sin \theta)}$$

$$= (f)^{\prime(f)} = ((f)^f)'$$

$$= e^{x \log x}$$

Differential dimension in quantum level of space emerged with extern of Glois group

Differential geometry in emerging with quantum space. This singularity of mass from being decomposed with fermion and boson, straight flow to boson creating at dark matter exert in universe and other dimension. This system is creating from universe of structure in fermion.

$$K_{mn} = E_n \times H_n$$

Hilbert space in creating with quantum space. Planck level relativity of quantum space.

$$\begin{aligned} \bigoplus \frac{M_p}{\nabla L} &= \frac{\partial}{\partial L} V(\tau) \\ \Psi H &= M_p \\ \frac{d}{df} F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dl} V(\tau) &= -\frac{2R_{ij}}{\delta T} \end{aligned}$$

This mass of gravity equation is equals with mass of accesory. And also this dimension of Glois group is equals with universe and other dimensions. Twelve particles of being constructed in singularity of component from complex manifold, this singularity pieces in AdS5 dimension are paired with other dimensions from Mebius space. Rolents atractor in singularity is created with super symmetry theorem. This equation is resembled with Gamma function and Beta function.

$$\frac{d}{dt} g_{ij} = -2R_{ij}$$

D-brane out of volume space are existing with hyper dimension assemble of space. This dimensions is positive fermion and boson in Mebius space. And moreover this space is positive in D-brane and anti-D-brane.

$$\begin{aligned} E^2 &= m^2 c^4 \\ R_{\mu\nu} + g_{\mu\nu} \Lambda &= 4\pi G \rho \end{aligned}$$

These equation is proof with super Langrans conjecture.

These equation is inclusive into world line of surface with mass and time of relate for spin network. In this network that constructed with space of dimension walk into time pole. This time pole is converted with mass of Higgs field. Then this converted result is emerged with spin network being went with mass of pole energy into time and space system of universe law.

After all, these explain system is commented from quantum level of differential structure, and quantum level of planck scale representative theorem.

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

David Hilbert

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2 \psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calcurate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values.

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\int \frac{1}{(x \log x)} dx = i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx$$

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \frac{1}{2} x^2$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \int \int_M dx_m$$

$$\leq \frac{1}{2} i + x^2$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$(f)^{(f)'}$$

$$= (f^f)'$$

$$= e^{x \log x}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, $\sin 0 = 0$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$\begin{aligned}
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however,

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry.

$$\begin{aligned}
\frac{d}{dt} \psi(t) &= \hbar \\
&= \frac{1}{2} i e^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' \\
&= -i e^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-if}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta, \mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields

of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements.

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi with my son in acasic record of space

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x)[I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2\psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values.

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2\end{aligned}$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\begin{aligned}\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ &= (f)^{(f)'} \\ &= (f^f)'\end{aligned}$$

$$= e^{x \log x}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, $\sin 0 = 0$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$\begin{aligned}
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however,

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry.

$$\begin{aligned}
\frac{d}{dt} \psi(t) &= \hbar \\
&= \frac{1}{2} i e^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' \\
&= -i e^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-if}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta, \mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields

of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements.

Non-esgranded theorem and this rerouted theorem

Masaaki Yamaguchi Kazuyuki Ogata

Non-esgranded theorem is included from imaginary and sqrt of element being built with dimension of law in system of mechanism, these explain of being constructed from operator of mathematics in nature of law, are existed with gap of operator is abundant of law, and this law of mathematics is near with result of problem. Therefore, mathematics is non perfect operator, and this mathematics theorem is non esgranded of operator problem. After all, this explain of theorem called to exist of non god in this universe. However, non-commutative equation, other wise, partial of differential operator and assemble of integrate operator concluded from difference of operator, all of theorem from zeta function is proof from laplace equation with god exist in this universe.

These theorem is tooled from mind map of method, and this theorem of world line in duality of differential equation,

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

from quota of global differential zone is cut of surface of eternal space from mass of being related with time and space of being concerned with Higgs fields.

$$\frac{d}{df}F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$||ds^2|| = e^{-2\pi T|\phi|}[\eta + \bar{h}_{\mu\nu}]dx^\mu dx^\nu + T^2 d^2\psi$$

This explain of non esgranded theorem is inversed with integral and native of limintate are difference of equation, and resulted of this system, this gap of operator and mathamatics law is followed from quato and singularity of non entropy of being resulted into zeta function and quantum group is proof of mathematics of operator and law of nature constructed with general number of element, atom and electric of element moreover quarks is controled with laplace equation access into database of universe from Hologram space, after all, operator of mathmatics and nature of law is mapping resulted into non esgranded theorem of meaning and natural law.

Bulubagi history explained with being among mapping and function is meaning with this mechanics of system concluded with possibility of function, However, mapping of system is proof of replace of non esgranded theorem for native law is general theorem.

These explained of mathmatics law is destructed from being replace equation that is resulted into being thinking of god exist in this universe.

微分幾何の量子化とゼータ関数、 大域的微分方程式についてのレポート

Masaaki Yamaguchi, with my son

ヒッグス場 + オイラーの定数 = ゼータ関数

$$m(x) + C = 2 \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\text{オイラーの定数} = \frac{d}{d\gamma} \Gamma^{-1} - \gamma^{(\gamma)'} = \left(\Gamma^{(\gamma)'} \right)^{-1} - \gamma^{\gamma'}$$

重力 + 反重力 = ゼータ関数

ガンマ関数の大域的微分方程式が、大域的微分方程式の部分積分として、ヘルマンダー作用素と同型になり、微分の変分法が対数関数を初期関数として、物体を形作っているフェルミオンを商代数で求めると、ヒッグス場の方程式になる。

$$\Gamma dx_m + \int \Gamma dx_m = \Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$\begin{aligned} & \frac{d}{d\gamma} \Gamma + \int \Gamma dx_m \\ &= \int f(g(x)) g'(x) dx \\ &= \left(\int f(g(x)) dx \right)' \\ & \nabla_i \nabla_j \int \nabla f(x) d\eta \end{aligned}$$

大域的微分方程式の部分積分をフロベニウスの定理で解を出すと、前後の大小関係が、Jones 多項式から、rico level 理論となり、ゼータ関数の対称率が、CP 対称性の破れとして、宇宙のゼータ関数だけが出力される。これが、オイラーの定数の関係として、ゼータ関数とオイラーの定数の対称率の変分率としての、Jones 多項式として、微分幾何の量子化が、ゼータ関数と証明されている。

$$|f \circ g| \leq |f + g|$$

$$\frac{f}{\log x} = m(x)$$

$$\frac{\gamma}{\log x} = \frac{d}{d\gamma} \Gamma$$

$$\begin{aligned}
\gamma &= \Gamma^{(\gamma)'} \log x \\
\Gamma^{(\gamma)'} &= \gamma (\log x)^{-1} \\
&= r dx_m \\
\Gamma^{(\gamma)'} &= \left(\int e^{-x} x^{1-t} dx \right)^{(\int e^{-x} x^{1-t} \log x dx)'} \\
&= e^f \\
&= r dx_m = e^f \\
&\int \Gamma^{(\gamma)'} dx_m \\
&\nabla_i \nabla_j \int \nabla \Gamma^{(\gamma)} dx \\
&= \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma dx_m \\
&= \int \Gamma \cdot \frac{d}{d\gamma} \Gamma dx_m \\
&= \left(\int \Gamma dx_m \circ \frac{d}{d\gamma} \Gamma \leq \int \Gamma dx_m + \Gamma dx_m \right)' = e^{-f} \cdot e^f \leq e^{-f} + e^f \\
&\quad (1 \leq e^{-f} + e^f)' \\
0 &= -(e^{-f} - e^f) \\
&\Gamma^{(\gamma)'} + \Gamma^{(\gamma)} \\
&= (e^f + e^{-f})' \\
&= -e^{-f} + e^f \\
&= -(e^{-f} - e^f) \\
&(R_{ij})' = -(R_{ij}) \\
&\int C dx_m = \int \left(\int \frac{1}{x^s} dx - \log x \right) dx \\
&= \int \int \frac{1}{x^s} d\text{vol} - \int \log x d\text{vol} \\
&\Gamma dx_m = \gamma dx_m \\
&= \frac{d}{d\gamma} \Gamma^1 - (\gamma)^{\gamma'} \\
&= e^{-f} - e^f
\end{aligned}$$

微分幾何の量子化が、ガンマ関数となり、ベータ関数に行き着き、オイラーの定数へと導かれて、ヒッグス場と対になり、ゼータ関数へと証明されて、ポワンカレ予想とリーマン予想が解決されている。ロシアとアメリカ、日本の数学者たちが解決している。私の子どもが、微分幾何の量子化がゼータ関数へと解決している。全部の理論がヒルベルト空間を多様体として、一般相対性理論と量子力学を、微分幾何の量子化が、D-brane と解として、ガウスの曲面論を、重力と原子にして、統一場理論として形成している。

Three manifold of entropy formula

Masaaki Yamaguchi

Euler equation represet form,

$$\begin{aligned} C &= \int \frac{1}{x^s} dx - \log x \\ &= \int \left(\int \frac{1}{x^s} dx - \log x \right) d\text{vol} \\ &= \int \int \frac{1}{x^s} d\text{vol} - \int \log x d\text{vol} \end{aligned}$$

This equation componet of Higgs equation + Euler equation = Zeta function. This represent expression is Rich flow equation. This equation assent from Differential geometry in quantum level, Gamma and Beta function, Gauss surface, Higgs field, Einsetin tensor, Euler constance and so on. This equation is add and average of possibility, and universe, atom more over a certain theory, this universe oneselves is environment seek, other distance is possibility equation, however, other dimension cover demand with not only norm of non liner in three manifold but also non possibility in complete of environment equation.

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\ \int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} d\text{vol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\ &= e^{-f} - e^f \\ &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= 2 \cos(ix \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &+ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= \frac{d}{df} F = 2i \sin(ix \log x) \\ \frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\ &= 2e^{-\theta} = 2e^{-ix \log x} \\ &= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\log(x \log x) \geq 2\sqrt{y \log y}$$

$$\log(x \log x) = \log x + \log \log x$$

This universe and other dimension on blackhole and whitehole compont with universe and atom in movement and point of environment value by a certain theory of complete environment equation. And this three manifold of possibility equation are able to research in both value formula. This envirnment of researchable reason is that surround of universe is setting to value by already in consented to being researched, then both value researchabled. And this three manifold equation is insert with Kaluza-Klein theory and Heisenbelg equation.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla\psi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

This energe of entropy value consist with whitehole of atom and universe in blackhole value, and this universe in balckhole represent with Higgs filed, atom in whitehole consist with plack scale.

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

This entropy value also consist with atom in balckhole.

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

These equation more also consist with universe being non gravity formula.

$$2 \cos(ix \log x) + 2i \sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

And also these equation consist with universe and other dimension in warped passeage in rout of integral.

オイラーの定数とガンマ関数、 そして微分幾何の量子化と虚数による逆三角関数、 ベータ関数による場の理論

Masaaki Yamaguchi

オイラーの定数は、次の式で成り立っている。

$$C = \int \frac{1}{x^s} dx - \log x$$

その式を単体積分をすると、

$$= \int \left(\int \frac{1}{x^s} dx - \log x \right) d\text{vol}$$

ゼータ関数を多重積分すると、

$$= \int \int \frac{1}{x^s} dx - \int \log x d\text{vol}$$

ここで、対数方程式は、多様体積分がガンマ関数を微分した方程式と同値より、

$$\begin{aligned} F = \Gamma &= \int e^{-x} x^{1-t} dx \\ &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dx_m - \int e^{-x} x^{1-t} \log x dx_m \\ &\quad \int x^{1-t} e^{-x} dV = \int x^{1-t} dm \\ &\quad \int x^{1-t} e^{-x} dV = \int x^{1-t} d\text{vol} \\ f = \gamma &= \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - \int e^{-x} x^{1-t} \log x dx_m \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \end{aligned}$$

これらより、大域的微分方程式のヒッグス場方程式の逆三角関数の双曲多様体に対極する、双曲多様体の余弦定理と同値になる。

$$\begin{aligned} &= e^{-f} - e^f \\ &= 2 \cos(ix \log x) \end{aligned}$$

結局は、微分幾何の量子化がガンマ関数でもあり、その逆関数はゼータ関数でもあり、そして、オイラーの定数を多様体微分をすると、ヒッグス場の方程式は正弦定理の逆三角関数であり、このヒッグス場の方程式の逆三角関数が、余弦定理の双曲多様体として、オイラーの定数になる。オイラーの定数は、このガンマ関数から、無理数と証明される。

Circle function and imaginary number, Euler equation from weak electric and strong electric theorem from gravity and antigravity on one rout of time system

Masaaki Yamaguchi

Rotate with pole of dimension is converted from other sequence of element, this atmosphere of vacume from being emerged with quarks of being constructed with Higgs field, and this created from energy of zero dimension are begun with universe and other dimension started from darkmatter that represented with big-ban. Therefore, this element of circumstance have with quarks of twelve lake with atoms.

$$(dx, \partial x) \cdot (\epsilon x, \delta x) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\frac{d}{df} F = m(x)$$


And this phenomounen is super symmetry theorem built with quarks of element, also this chemistry of mechanism estimate from physics of operatsion. These response of mechanism chain of geometry into space being emerged with creature and univse of existing of combination.


This converted with dimension of element also emerged from imaginary and reality of pole's space.

And this pole of transport of dimension belong with vector of time has with one rout of sequecense. Quantum physics also belong with other vector of time has with imaginary rout of sequecense.

This sequence of being estimated with non fluer of time, and this space of element have with gravity and antigravity of power. Other vecor of time is antigravity rout of sequecense.

Weak electric theory is estimate from time has with one rout of sequecense, this topology of chain is resulted from time of rout ways.





$$\square(\frac{\sigma_1 + \sigma_2}{2}) = [3\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\square\psi = 8\pi GT^{\mu\nu}$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

$$\nabla\psi^2 = 4\pi G\rho$$

$$\square(\sigma_1 + \sigma_2) = [6\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$= [i\pi(\chi, x), f(x)]$$

$$\square\psi = [12\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\frac{d}{df}F = \int e^{-f}[-\Delta v + R_{ij}v_{ij} + \nabla_i \nabla_j f + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(\frac{v}{2} - h)]$$

$$= [i\pi(\chi, x), f(x)]$$

These equation is reminded time pass rout of one rout way of forms, and this rout of time ways which go for system from future and past. Therefore, this resulted system of time mechanism is one true flow that weak electric theorem oneselves. and moreover, one rout time way of forms is reverse with antigravity of time system. This also spectrum focus is true that Maxwell theorem and strong boson unite with antigravity, this unite is essence on the contrary from weak electric theorem, this theorem called for time rout forms is strong electric theorem. This two theorem is united with quantum physics that no time flow system.

$$f^{-1}(x)xf(x) = 1, H_m = E_m \times K_m$$

The non-commutative theorem is constructed from world line surface that this complex manifold estimate with rolanz atractor, and this string theorem have with one world of universe mate six quarks

and other world of dimension mate other element of six quarks. These particle quarks is built with super symmetry space of dimension.

$$i = (1, 0) \cdot (1, 0), e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{-i\theta} = \cos \theta - i \sin \theta$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$\sin i\theta = \frac{e^{-\theta} - e^{\theta}}{2}$$

$$\pi(\chi, x) = \cos \theta + i \sin \theta$$

In this equations, two dimension redeconstructed into three dimension, this destroy of reconstructed way is append with fifth dimension. This deconstructed way of redeconstructe is arround of universe attached with three dimension, this over cover call into fifth dimension.

$$R(-\alpha) = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha)MR(-\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos^2 \alpha - \sin^2 \alpha & 2 \sin \alpha \cos \alpha \\ 2 \sin \alpha \cos \alpha & -\cos^2 \alpha + \sin^2 \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & -1 \\ 1 & -\sin \alpha \end{pmatrix} \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}xf(x) = 1$$

$$(\log x)' = \frac{1}{x}, x^n + y^n = z^n$$

$$x^n = -y^n + c, nx^{n-1} = -ny^{n-1}y'$$

$$y' = \frac{nx^{n-1}}{ny^{n-1}}$$

$$= \frac{x^{n-1}}{y^{n-1}} = -\frac{y}{x} \cdot \left(\frac{x}{y}\right)^n$$

$$\begin{aligned}
& -\frac{\cos x}{(\cos x)'}(\sin x)' = z_n \\
& z^n = -2e^{x \log x} \\
& \lim_{x \rightarrow \infty} f(x) = a, \lim_{y \rightarrow \infty} f(y) = b, \lim_{x, y \rightarrow \infty} \{f(x) + f(y)\} = a + b \\
& \lim_{z \rightarrow 0} f(z) = c, \delta \int z^n = \frac{d}{dV} x^3 \\
& \lim_{x \rightarrow \infty} = c - \lim_{y \rightarrow \infty} f(y) \lim_{x \rightarrow \infty} f(x) + \lim_{y \rightarrow \infty} f(y) = \lim_{z \rightarrow \infty} f(z) \\
& z^n = \cos n\theta + i \sin n\theta \\
& = -2e^{x \log x}
\end{aligned}$$

These equation is gravity and antigravity equation.

$$\begin{aligned}
& \frac{d}{d\sigma} \left[\frac{(\sigma_1 + \sigma_2)}{2} \right] \\
& = \sigma(\uparrow\downarrow) + \sigma(\uparrow\uparrow) + \sigma(\downarrow\downarrow) + \sigma(\Leftarrow) + \sigma(\Rightarrow) \\
& \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \sigma(\uparrow) \\
& \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \sigma(\downarrow) \\
& \sigma(\Leftarrow) + \sigma(\Rightarrow) = \int e^{-f} [-\Delta v + R_{ij} v_{ij} + \nabla_i \nabla_j v + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(v - \frac{h}{2})] \\
& \sigma(\downarrow) = \sigma(\uparrow\downarrow + \uparrow\uparrow + \Rightarrow) \\
& \sigma(1) = \sigma(\uparrow\downarrow + \downarrow\downarrow + \Leftarrow) \\
& \text{weak electric theorem} = \sigma(1) \\
& \text{strong electric theorem} = \sigma(\downarrow)
\end{aligned}$$

These equation is represented with topology of string model, and weak electric theorem is constructed with Maxwell theorem and weak boson, gravity that estimate with this three power united. Moreover, strong boson and Maxwell theorem, antigravity that also estimate with this three power united. This two united power is integrated from gravity and antigravity. Then this united power is zeta function.

Instance of acknowledged from refrent existed of humanity theorem and reality of thought in modern technology 実存主義と現代文明、双極性による精神機能からの未知の智

Masaaki Yamaguchi

Refrent existed of humanity theorem is built with acasic record from human being constructed with world is setting from view of reality human being created from. This theorem mechanical from this type afford into world viewed of this human being, and this thought of history record having with the type recieved with acknowlege of acasic record and reality of human being seen world to accept with native reality, this type of human being become with enginner and human brain interface created with modern history existed. These type of being recieved from the world of acknowlege is existed in reality of non dreamly world of being seen. Whisperd of this type called with world of being existed, estling of this type called and this type of being selected with is recieved with acasic record. Photo Reader is reffer to being selected with estling of acknowlege in world of tounge. Refrent existed of humanity theorem is inspired with Philosophy of being called human being to recieve in this record. Refrent means that jugement in person and creature are accepted with acasic record from recognized of system how registance of recieve with nature and universe of life.

実存主義は、サルトル・アッカーシュマンが去った後も、現実主義者がこの実存主義に分類されている上に、ブレインインターフェースやコンピュータを作っている人がこの宗派に分類されている。この実存主義は、無の状態から零次元がエターナルな場に同型とされている瞑想を対象として、アカシックレコードから未知の情報を取得する。この無は禅とも言われていて、実存と虚存としての物理学を対象とすると、閉3次元多様体として、ザイフェルト多様体から未知の智を感知する。この智は、他者との関係から関与して、この世は天啓のごとく、天からの啓示を本として接点として、未知の智を感知する。現実主義者は、うつの人に傾向として言われているが、実存主義にこのうつの人を対象として、夢を誇大妄想とは別として、研究対象にされている。なぜ、うつの人が現実主義者に多いのに実存主義の能力があるかは、瞑想から無の状態を使い、アカシックレコードにアクセスするのと、本当に現代文明を作り上げることができる能力をもっているからである。無からの知識はエターナルな場から零次元が世界に存在していて、感知していない次元が零次元と無としての禅による相対性としての他者との関与から同型として、禅の無からアカシックレコードとこの存在している次元から実存と虚存から未知の智をこの世界に降臨させてもいる。それゆえに、このタイプの人、ウィスパードと言われている、臥龍をこのエターナルな場を生成元として、エストリングをこのウィスパードを育成する場と言われている。カロリング研究所もこの分類に入る。このエストリングな場は、若返りの場にもなっている。アカシックレコードは絶大な治癒力を有している。宇宙は、始まりと終わりが真逆な時間の矢にもなっている。過去の宇宙が未来の宇宙になってもいて、時間と空間がシンメトリックな性質になってもいる。年を取る事が勉強との学問の教訓から身につけられていると、若返りの効果が生成される。このタイプには、テクノ

ロジーとマネジメント、ストラテジストが分類として研究されている。エストリングな場から、今まで誰も思いつかなかった数式や理論をアカシックレコードから投射として得ることが出来る人もいる。実存主義とは、判断と認識、そして無による愛でもある。

Infinity of acknowledge in world is existed zero dimension and rebuiled with eternal space, non-existed dimension equaled with zero dimension and zen of skill relativity, and zen of infinity of atomosphere required with acasic record from this space of dimension. Therefore, this type of being required with are called with whisperd, and this selected with estling of space and acasic record. This dimension of area are referred with more and more young belonged with time of ages. In this referred with time of ages are cured from any built of system. Get of ages equaled with young of age with study of learn and required of any skill. Life of water and Life of tree is existed with central of universe, and around of universe is old of universe and this space is creating with any other dimension of creature. Time followed is reverse of system. Young of universe is equaled with Old of universe and Young of time refferd with is Old of time belonged with universe system. This system cnstruct from non time followed lead into quantum physics mechanism, this type of category esperanade of theorem and discover of system of native more literal strategy of skill in top of goverment positions, more releanity of skill is called from manegement of skills. Any type of this required skills lead from every acknowleged of technical and potential possibility, more spectrum influence of being inspired into a respired of reluctant with acquiring lives of flowing days on stress pressiers. Life of insurance need every moment to avoid from varnished any others. Incoming of value space revealed from any intransit of memories, this acceptance of being invited from accerity of geotics.

無の状態を維持すると空間に同化する。それと同時に現実と隔離できる。この作用で、一念三千を感じることに無による瞑想はアカシックレコードに接することの必要十分条件にもなっている。無とはそれほどにも大切な技能に言えることでもある。速読法がこの技術を使っている。無の音読が確の目によりこの技術を使っている。

Refrent exist of humanity theorem is included with estling of involved with other's inspire with each other of persons and creatures in estling of setting with mind of hologram fields from being with academic learns. This estling of fields is also catastrophe of fields how converted from being with harmony and nestling of mind inspect with each persons and each creatures in other's of minds. Encycropedia is selected with disease of type in explained of this refrent exist of humanity theorem. Therefore, this theorem is explained with encycropedia of which other persons and other creatures are read with other creatures in estling of fields. This psychological of mind in refrent exist of humanity theorem is explained with indivisual of mind in zero dimension and zen of mind recieved from future and past of accident what any matter are incident of other's distant with involved of counsels.

実存主義は、他者と自己においての精神作用の関与による、精神の共鳴と共振の機構を述べてもいる。自己のどういう思考の仕方、他者との心理作用がカタストロフエネルギーの位置関係に在るかを、他者がどういう態度で他者同士の心理がエストリングの場でどのように変化するかをこの実存主義は、弁証論として述べてもいる。心因反応は、この実存主義で分析されてもいる。自己の心理を他者が読むのは、この実存主義の関与からも分かる。この実存主義には、自己による無の状態から未来と過去に何が起きるかも他者との相対関係から分かることの心理をも述べてもいる。プリンストン高等研究所では、この無からの超高速コミュニケーションが研究者同士でされてもいる。

Integrate of theorem

Masaaki Yamaguchi

Laplace equation is constructed with zeta function, zeta function also vector element of singularity.

$$\frac{d}{df}F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

Seifert manifold is built with fourth of power to integrate element of singularity.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}, \frac{d}{df}F_t^m = 2 \int \frac{(R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

Fourth of power is one of geometry in inclusive with integrate of fields theorem.

$$= \frac{1}{4} g_{ij}^2, 4V_\tau = g_{ij}^2$$

Fifth dimension of equation called to estimate with abel manifold of component in seifert manifold.

$$||ds^2|| = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\psi$$

Norm liner is fermader of manifold, is one of rout in non-relativity thoerm.

$$\delta(x) \cdot \mathcal{O}(x) = ||ds^2||, \eta_{\mu\nu} = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

Imaginary of pole is zeta function of component.

$$\bar{h}_{\mu\nu} = [\nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]^{iy}$$

Minkofsky of dimension of different in fifth dimension of element.

$$\eta_{\mu\nu} + \bar{h}_{\mu\nu} = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

Volume of surface is open set group.

$$\begin{aligned} & \int \int \int \frac{V}{S^2} dm = \mathcal{O}(x) \\ &= \int \int e^{\int x \log x dx + O(N^{-1})} d\psi \\ & \left(\frac{\nabla \psi}{\square \psi} \right)' = \left(\frac{1}{2} \right)' \\ &= 0 \end{aligned}$$

Accessarilty of gravity of formula is harf of vector in this norm space.

$$\left(\frac{\eta_{\mu\nu}}{\bar{h}_{\mu\nu}}\right) = \frac{1}{i}$$

$$\hbar\psi = \frac{1}{i}H\Psi$$

Kaluze-Klein theorem is deduce of dimension in minus of zone, and this zone is imaginary of pole in this element.

$$8\pi G\left(\frac{p}{c^3} + \frac{V}{S}\right) = ||ds^2||$$

Three of manifold in entropy of equation is oneselves in norm space.

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \psi$$

This element is fifth dimension of abel manifold.

$$H_3(x) = 0, \chi(3) = 2, \pi(\chi, x) = \int \int \frac{1}{(x \log x)^2} dx_m$$

$$= \frac{1}{2}i$$

$$\eta_{\mu\nu} = \nabla_i \nabla_j \int \nabla f(x) d\eta, \bar{h}_{\mu\nu} = \nabla_i \nabla_j \int \nabla g(x) dx_i dx_j$$

Rout of helmander in zeta function of element is Own of stimulate in seifert manifold. Fifth dimension in seifert manifold is oneselves in rout of equation in imaginary of pole.

$$\eta_{\mu\nu} = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}, \bar{h}_{\mu\nu} = [\nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]^{iy}$$

Sheap of element have with zeta function.

$$||ds^2|| = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2 \psi$$

$$= \delta \mathcal{O}(x) [f(x) \circ g(x)] dx^\mu dx^\nu + \lim_{x \rightarrow \infty} \sum_{k=0}^{\infty} a_k f^k$$

Open set group in seifert manifold is constructed with abel manifold.

$$(\Box\psi)' = \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu} \left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

Gravity of power in three manifold of entropy of power expire of around of universe.

$$\Box\psi = 8\pi GT^{\mu\nu}, \mathcal{O}(x) = [\nabla_i \nabla_j f(x)]'$$

This gravity of power is component with open set group in differential equation.

$$\cong {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y), V(\tau) = \int [f(x)] dm / \partial f_{xy}$$

And this equation developed with summuate of manifold is built.

$$\frac{p}{c^3} \circ \frac{V}{S} = [\nabla_i \nabla_j \int \nabla f(x) d\eta \circ \nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]$$

This also equation is sheap of manifold in zeta function in frobenius theorem.

$$||ds^2|| = (\delta(x) \circ G(x))^{\mu\nu} \rightarrow \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu}$$

$$\frac{d}{df} F(v_{ij}, h) = [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2)(\frac{v}{2} - h)]$$

$$e^{-f} \circ e^{-f} \rightarrow -2R_{ij}, e^{-f} \rightarrow e^{-2\pi T|\psi|} = [\nabla_i \nabla_j \int \nabla f(x) d\eta \circ \nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]$$

$$\frac{x+y}{2} \geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x \log x}} = 1$$

Zeta function is steady in imaginary of pole.

Entropy on manifold and differential of equation

Masaaki Yamaguchi

1 Global Differential Equation

Global area in integrate and differential of equation emerge with quantum of image function built on. Thurston conjugate of theorem come with these equation, and heat equation of entropy developed it' equation with. These equation of entropy is common function on zeta function, eight of differential structure is with fourth power, then gravity, strong boson, weak boson, and Maxwell theorem is common of entropy equation. Euler-Lagrange equation is same entropy with quantum of material equation. This spectrum focus is, eight of differential structure bind of two perceptive power. These more say, zeta function resolved is gravity and antigravity on non-catastrophe on D-brane emerge with same power of level in fourth of power. This quantum group of equation inspect with universe of space on imaginary and reality of Laplace equation.

It is possibility of function for these equation to emerge with entropy of fluctuation in equation on prime parameter. These equation is built with non-commutative formula. And these equation is emerged with fundamental formula. These equation group resolved by Schrödinger, Heisenberg equation and D-brane. More spectrum focus is, resolved equation by zeta function bind with gravity, antigravity and Maxwell theorem. More resolved is strong boson and weak boson to unite with. These two of power integrate with three dimension on zeta function.

Imaginary equation with Fouier parameter become of non-entropy, which is common with antigravity of power. Integrate and differential equation on imaginary equation develop with rotation of quantum equation, curve of parameter become of minus. Non-general relativity of integral developed by monotonicity, and this equation is same energy on zeta function.

$$\pi(\chi, x) = i\pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x)$$

This equation is non-commutative equation on developed function. This function is

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

This equation resolved is

$$\int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2}i$$

And this equation resolved by symmetry of formula

$$y = x$$

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq \frac{1}{2}$$

This function common with zeta equation. This function is proof with Thurston conjugate theorem by Euler-Lagrange equation.

$$F_t^m \geq \int_M (R + \nabla_i \nabla_j f) e^{-f} dV$$

resolved is

$$F_t \geq \frac{2}{n} f^2$$

$$\frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

These resolved is

$$\frac{d}{df} F_t = \frac{1}{4} g_{ij}^2$$

Euler-Lagrange equation is

$$f(r) = \frac{1}{2} m \frac{\sqrt{1 + f'(r)}}{f(r)} - m g f(r)$$

$$m = 1, g = 1$$

$$f(r) = \frac{1}{4} |r|^2$$

Next resolved function is Laplace equation in imaginary and reality fo antigravity, gravity equation on zeta function.

$$\int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = 0$$

This function developed with universe of space.

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

This equation also resolved of zeta function.


$$\frac{d}{df} F(v_{ij}, h) = \int e^{-f} [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

These equation is common with power of strong and weak boson, Maxwell theorem, and gravity, antigravity of function. This resolved is

$$\frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

on zeta function resolved with

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$



2 Quantum Equation for Dimension of Symmetry

Quantum Equation architect with geometry structure of Global Differential Equation has zero with gravity and antigravity for eternal space, and this space emerge for burned of non expanded universe. Then this universe has Symmetry in dimension with that created of fourth Universe in one of geometry has six element quark and pair of structure belong for twelve element quark. These quarks emerge with eternal space of Non-Difinition System in Quantum Mechanism, in term of one dimension decided, or the other dimension non-decision. These system concerned of vector of norm depend for universe mention to eternal space. Mass existing in dimension emerge gravity, these paradox is in universe has mass around of light, in deposit of mass for our universe and the other dimension has gravity and antigravity, and covered with these element for non-gravity. Laplace equation decide with eight of structure for these element of power integrate for one of geometry. Higgs quark is quote algebra equation in Global Differential Equation of non-gravity element on zero dimension. These equation is mass of build on structure in mechanism system. This quote algebra equation have created of structure in mass with universe of existing things. These result means with why quantum system communicate with our universe be able to connect of.

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq 0$$

$$\Delta x \Delta p \geq \frac{1}{4} i$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\sqrt{\frac{\hbar G}{c^3}} \cong 1.616 \times 10^{-33}$$

$$C = 0.5772156 \dots$$

$$F = \frac{d}{df} \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$F_t^m = \frac{1}{4}(g_{ij})^2$$

Quantum Group resolved with Laplace equation build in calculate on non-gravity element of based by non extension equation. This solved by these element in equation has belong to gravity is why universe have in no weight, and mass exist weight to space emerge of created with gravity, in which is reason by non-extension equation translate to variable and invariable element. These problem of included universe in no gravity, which is exist of mass on space of surrounding in universe. Those question resolve on zeta equation and Quantum Group of translate to vector equation. Non-extension equation selves has non-gravity, these result with universe first burn in D-brane created by solved with replace equation. Quote Algebra equation have created of structure in mass with universe of existing things.

These equation explain to those which included mass has gravity emerged, and universe has in surrounding to no weight. The other Dimension integrate with these universe of gravity to unite antigravity, so this universe has no weight. Higgs quark is mass of built on structure in mechanism system. These system belong to create on element. These element is based on existing of universe which has structure code, in theorem composed for universe to emerge of time with future and past. Universe first created in these time, this existing things already burned with space. Network Theorem is connected eight element of geometry structure which integrate with three dimension of structure. These structure compose in three manifold, zeta equation is this system of element. These resulted theorem resolved with quantum equation, so this mechanism impressed in universe of component. Strong and Weak boson is united to one, and Maxwell theorem is same system. Gravity and Antigravity has own element. General relativity theorem same united. These integrate with included Euler equation. These power of element is zeta equation. Then this twelve element of quarks has belong to this universe and the other dimension.

$$\Delta E = -2(T-t)|R_{ij} + \nabla_i \nabla_j f - \frac{1}{2(T-t)}g_{ij}|^2$$

Quote group classify equivalent class to own element of group.

$$A=BQ+R$$

$$[x]=A$$

$$dx^n=\sum_{k=0}^{\infty}x^ndx$$

$$R_n=\frac{n!}{(n-r)!}(x^n)'$$

$$\beta(p,q)=\int_0^1x^{p-1}(1-x)^{q-1}$$

$$Z(T,X)=\exp(\sum_{m=1}^{\infty}\frac{(q^kT)^m}{m})$$

$$Z(T,X)=\frac{P_1(T)P_3(T)\ldots P_{2n-1}}{P_0(T)P_2(T)P_4(T)\ldots P_{2n}}$$

$$|v|=|\int (\pi r^2+\vec{r})dx|^2$$

$$\Delta E = \int (\operatorname{div}(\operatorname{rot} E) \cdot e^{-ix \log x}) dx$$

$$(\nabla\phi)^2=\int tf(t)\frac{df(x)}{e^{-x}tx^{-1}}dx$$

$$(\nabla\phi)^2=\int tf(t)(\Gamma(t)df(x))dx$$

$$(\nabla\phi)^2=\frac{1}{\Gamma(x+y)}$$

then these equation decide to class manifold with group. Differential group emerge with same element of equation, zero dimension conclude to emerge with all element. Constant has with imaginary of number on developed of zeta equation. Weil's Theorem resolved with zeta equation, these function merge to build in replace equation. Euler constant has with bind of imaginary number and variable number. These function is

$$C=\int\frac{1}{x^s}dx-\log x$$

Replace equation resolve on zeta function.

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i$$

These function understood is become of imaginary number, which deal with delete line of equation on space of curve.

Euler number also has with imaginary of constant.

Space Mechanism to transport of Dimension

Masaaki Yamaguchi

1 Kaluza-Klein Theorem

The other dimension rotate of universe with real and image rout, complex rotate to dimension of fifth for Kaluza-Klein dimension extend with theorem. Mobius space explain this theorem to reflect for fifth dimension of construct with real rout of rotate in image rout.

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\beta(p, q) \geq - \int \frac{1}{t^2} dt$$

Abel equation tell this space to infinite this dimension for finite space to conclude of this theorem, in explain the other dimension with our universe rotate with, and this space rout out this fifth dimension, no thought with time system. This space didn't throw of light speed, light element go with space throw. This idea explain of magnetic theorem, this tell space to construct with movement result. Light speed go with other light together, this light look like stop of speed. This idea extend of tell space for no relativity, time flow of infinite.

Kaluza-Klein Theorem say

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(\kappa^2 A_{\mu\nu}(x)dx^\mu)^2$$

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\nabla \phi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

$$ds^2 = -N(r)^2 dt^2 + \phi^2(r)(dr^2 + r^2 d\theta^2)$$

$$ds^2 = -dt^2 + r^{-8\pi Gm}(dr^2 + r^2 d\theta^2)$$

This equation result

$$dx^2 = g_{\mu\nu}(x)(g_{\mu\nu}(x)dx^2 - dxg_{\mu\nu}(x))$$

These equation tell space non-symmetry result

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

This theorem mention to zeta equation construct space of non-symmetry to emerge with non-gravity.

$$\pi(X, x) = i\pi(X, x)f(x) - f(x)\pi(X, x)$$

2 Network Theorem

Quantum Group connect with Universe component which has built in emerge space, these Network element by kernel and image function in zeta equation. This each of function system is Universe element of geometry group construct with these equation, in space first burn with time future and past. Gravity equation has with antigravity element. Quantum group with fourth of universe system. This system mention to our universe create with network theorem. Our universe already create of eternal space. Time throw with this space, then universe already complete created. In this reason time of throw in space was found with non-expand of universe. General relativity theorem mention to this equation means by.

Quark has twelve element kernel and image of equaiton resolve with base group integrate of three dimension construct of this equation resulted. Global differential equation means to resolve with universe emerge in built system. This equation emerge with almost thing of component structure code. This source code create in element structure. This network system is include with fourth of universe on connected with communicate mechanism. Base group explain in these system, differential equation has with global integrate equation together. This equation means for our universe system of geometry structure integrate with one of geometry. In reason this connected with eight element of structure, then this system is realize to quantum group equation resolved with space mechanism.

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T^{\mu\nu}$$

$$\begin{aligned} X(3) = & (-1)^3 < |a_0 a_1 a_2 a_3| > + (-1)^2 (< |a_0 a_1 a_2|, |a_1 a_2 a_3|, |a_0 a_1 a_3| >) \\ & + (-1)^1 < a_1 a_2, a_0 a_3, a_2 a_3 > + (-1)^0 < a_0, a_1, a_2, a_3 > \end{aligned}$$

$$H_n(x) = \ker f / \operatorname{im} f$$

$$X(x) = r(H_n(x))$$

$$X(3) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

3 Zeta equation of system

Quantum equation for universe has with gravity on fourth of manifold is closed three manifold with antigravity has decieve to the other dimension. This thorem mention to tell universe to composite with time of system, these system of equation say,

$$\log x(\log x) \geq 2(y \log y)^{\frac{1}{2}}$$

This system belong for geometry theorem destruct with three manifold built of singularity in dimension, this dimension is also built in mechanism of all composite with source code. For code is entropy of mass with energy, this entropy is,

$$\pi(X, X) = [i\pi(X, x), f(x)]$$

$$\pi(X, x) = \int \frac{1}{(x \log x)^2} dx$$

This equation tell gravity to entropy of equation composite with mass of singularity. Three manifold of equation mension to this universe has with zero dimension began to time of first, in firstly time has future and past. This system says that universe has first begun of throw to time system. The explain theorem that universe of end composite for entropy equation tell the construct of this system, and universe has in other dimension covered with oposite of energy.

Dimension of three manifold system says,

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$L_p = \sqrt{\frac{\hbar G}{c^3}}$$

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

This equation mention to three manifold of Dimension construct with time and energy of Non-Definition system. Quantum equation and this equation also built with same equation of theorem. Fundamental group explain this theorem to composite with entropy of same energy, any rout built with same energy to construct with source code. The rout of equation is,

$$V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau(q)}} L(x) dx) + O(N^{-1})$$

The equation also say that universe has singularity of component is,

$$\frac{1}{\tau}(\frac{N}{2} + \tau(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

These theorem explain that universe has of big-clanch system. The system include to dimension and time of throw of mechanism. Higgs field of energy equation say,

$$\frac{d}{df} F = m(x)$$

This equation concern with Seifert manifold mention to dimension of system. Global differential equation belong for gravity and antigravity of equation with Abel manifold in Euler equation. Infinite of group composite with this theorem conclude of finite group, these theorem explain to construct of universe is big-clanch.

Universe has with ertel and this mass of energy is singularity of component. Global differential system and Higgs of mass says that universe has built of darkmatter and big-ban system.

4 Emerge any dimension for supersymmetry to create for universe form quarks of element

Space for Non-Symmetry create of power in gravity and antigravity for Higgs quark of energy with ertel potential. This potential energy construct to emerge fourth power, then integrate of twelve of quarks. These mechanism export for another dimension of symmetry built. These dimension also architect with sixth of quarks. There import mechanism to Global differential equation for resolve of Higgs quark to build for unvierse. This way export of two dimension for symmetry of pair, these symmetry of space also built from sixth of quarks. Space of ertel emerge in gravity and antigravity, this power from ertel in non-free condition from free of ertel to emerge on these mass of space in zero dimension. This mechanism result with space to construct in three manifold of ertel from power. This resolved mechanism built with time and space of system, also this system flow to universe. These system tell universe to emerge space of singularity from potential energy.

$$\pi(|K''|) \cong \pi(|\bar{K}''|, O)$$

$$\begin{aligned} l(< P, Q >), l(< Q, R >), l(< R, S >), l(< P, R >), l(< P, S >), l(< Q, S >) \\ = \alpha, \beta, \gamma, \delta, \zeta, \eta \end{aligned}$$

$$\begin{aligned}
e &= \alpha\beta\gamma \\
\gamma &= \beta^{-1}\alpha^{-1} \\
\zeta &= \alpha\gamma \\
\delta &= \eta\gamma^{-1} \\
\eta &= \beta\zeta \\
\beta^5 &= (\beta\theta)^2 = \theta^3 \\
[\beta] &= [\theta] = 0
\end{aligned}$$

$$ds^2 = e^{-2kT(x)|\phi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2(x)d\phi^2$$

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

Twelve quarks emerge for three manifold of space, in element space of system construct with three D-brane, these dimension catastrophe of symmetry. This mechanism explain for those which create of our universe and the other dimension, take these mechanism into consideration for those built to deceive the other dimension over universe, for Global differential equation devide with gravity and antigravity emerge space to existing zero dimension of ertel space in singularity of zeta equation.

$$\beta(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\log x(\log x) = 2(y \log y)^{\frac{1}{2}}$$

5 Higgs Fields transport of three manifold on zeta function to resolve the mechanism

Higgs Fields transform of comformal field in space, this space mechanizm with lives of existing source code. Non-commutative algebra built to transport of emerging on space, each of element has prime number in these theorem. Destruct of element has a reverse on manifold, this theorem conclude with Euler-Lagrange equation resolve to merge element.

$$|f(x)|\rightarrow [f,f^{-1}]\times [g,h]$$

$$(x-1)(2y-b)\geq z$$

$$\frac{1}{x-1}\frac{1}{2y-b}\leq \frac{1}{z}$$

$$f(x) = \frac{1}{x-1} + \frac{1}{2y-b}$$

$$\frac{d}{df} \geq \frac{d}{df} f$$

$$dx \leq dF$$

$$|f(x)| = \frac{1}{4}|r|^2$$

Seifert manifold has reverse of time with space mechanism on merge to resolve with zeta function. Global differential equation has zero dimension of darkmatter to create of space, big-ban system also start with this mass of field to begin with universe.

Nonlinear element on manifold algebra

Masaaki Yamaguchi

1 Mebius space

Gamma and Beta function belong for Kaluza-Klein theorem, these equation built of first universe began with darkmatter, this ertel is non-free condition to emerge with big-ban system conform to export with antigravity element. This space consist of fifth dimension rotate with Mebius space construct for the other dimension, after all built of quarks two of pair in dimension symmetry. Topology consist of algebra equation is being with complex function. Norm relate with Volume of space of these system built in. Prime number concern with Euler constance relate of.

$$\Gamma(x) = \int x^{1-t} e^{-x} dx$$

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$f(x) = [f, f^{-1}] \times [g, h]$$

$$V(x) = \int \frac{1}{\sqrt{2\tau q}} (\exp \int L(x) dx) + O(N^{-1})$$

$$Zeta(x, h) = \exp \frac{(qf(x))^m}{m}$$

$$\frac{d}{df} F(x) = m(x)$$

This point of focus is Global differential equation relate with all existing equation. Prime number consist of zeta function in Mebius space.

2 Maxwell theorem integrate of gravity element

Kaluza-Klein theorem consist of general relativity of equation, fifth dimension is part of zeta function on also Global differential equation. Weak power concern with electric of magnity in Maxwell theorem. Space fill of ertel with darkmatter being of graviton to emerge of Higgs fields. This fields has topology element with any transform of power in atomic element. Relate with result of space merge to create for electric and magnitic power. Higgs fields built with pair of quarks in symmetry space. Vector of norm consist of two of pair for dimension.

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + \phi^2(x) (\kappa^2 A_{\mu\nu}(x) dx^\nu)^2$$

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

$$\sum_{k=0}^{\infty} |x_k + y_k|^2 = \sum |x|^2 + 2 \sum |xy| + \sum |y|^2 \leq \sum |x|^2 + \sum |y|^2$$

$$f(|x+y|) = f(|x|) + f(|y|) \geq f(|x| \circ |y|)$$

Norm space has with Frobenius theorem to resolve with Kaluza-Klein space built in. Euler number consist of rank for homology element to create with zero dimension.

$$\chi(x) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

Loop of topology has no exist with zero dimension, this system explain to create of Maxwell theorem.

3 Zeta function belong for singularity component

Zeta function consist of reverse of time quality, this space result with movement of element. Kaluze-Klein theorem create with space to emerge of singularity. Maxwell theorem has this system of circumstance. Fourth dimension belong to construct with Donaldson manifold exist explain. Duality of space also built with zeta function from singularity. Monotonicity relate to merge with non-relativity of integral result, this reflection is space of quality.

$$\frac{V(x)}{f(x)} = m(x)$$

$$F(x) = 0$$

$$\frac{d}{df} F(x) \geq 0$$

$$\lim_{x \rightarrow \infty} \text{mesh} \frac{F(x)}{f(x)} \rightarrow 0$$

$$\nabla f(x) = 2$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$f(x) = \int \frac{1}{x^s} dx - \log x$$

Morse theorem relate to merge of result with zeta function, gradient flow concern of zero dimension.

4 Other dimension and this influent of power

Pair of dimension interact to inspect for movement result, each of dimension devide with power of influent. This power has contrast of element, in inspire of result merge to create of pair in vector operate. Non-certain theorem mension to tell dimension give to create of pair in power. Six quarks sum to merge with twelve quarks, zero dimension have with fourth dimension of Global differential equation part of construct element. Graviton influent to reflect for universe to have the other dimension.

$$\frac{d}{df}F(x) = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

$$f(r) = \frac{1}{4}|r|^2$$

Norm space have non-liner to integrate with algebra manifold, singularity create for pair of dimension to fill of fourth of power. Eight of differential structure have for these dimension integrate four of power.

$$\frac{d}{df}F(v_{ij}, h) = \int e^{-f} [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$\frac{d}{df}(x-y)^n = \frac{\Pi(x-y)^n}{\partial f_{xy}}$$

$$\pi(X,x)=i\pi(X,x)f(x)-f(x)\pi(X,x)$$

$$\Delta E = \int \operatorname{div}(\operatorname{rot} E) e^{-ix \log x} dx$$

$$f(x)=\sum_{k=0}^{\infty}a_kx^k$$

Report

Masaaki Yamaguchi

1 Summulate of manifold from graviton structure

It is non-perfect element of manifold to integrate with graviton of quark, this quark construct of fermion and boson. Then D-brane aspect with string theorem include with. This structure summulate of equation on integral and differential operator. Higgs quark have part of network theorem for graviton structure. Universe influent pair of dimension between monotonicity and singularity. Non-symmetry deconstruct to distinguish with dimension of being constructed on quark of element. That's say, this mechanism architect with four of dimension existed. In part of this explain, Higgs fields is Weil's theorem component of global differential structure, Global area part of equation resolved come to simply conclude theorem. Global integral and differential equation, what is not used, difficulty of resolved.

Infinite number divide infinite oneselves and finite oneselves is finite number. Prime number extent of infinite number, $\sum a_k f^k(x)$ manifold have with these extend of count. Zeta function conclude with zero dimension, also this resolve to Gauss liner.

$$f(x) = \chi - \{x\}$$

$$\int_{\beta}^{\alpha} (x - \alpha)(x - \beta) = 2 \int_M f(x) dx$$

$$\int_{\beta}^{\alpha} a(x - \alpha)(x - \beta) = -\frac{(\beta - \alpha)^3}{6}$$

$$\frac{d}{df} F(x) = 0$$

$$\log(x \log x) - 2(y \log y)^{\frac{1}{2}} = [|\frac{d}{df} F(x)|]$$

Eight of differential structure integrate with graviton element to one of geometry, plank scale add volume divide to surface. This scale is three dimension of one energy. $\nabla \phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S})$ This equation means is three dimension of energy entropy. Zeta function means is time and space of scale on plank entropy, and this equation built on future and past of curve of parameter. These equation resolve with one component of energy and vector of time and space of curve of parameter.

$$\nabla \phi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

$\sqrt{2\pi T}$ is blackhole entropy and $\square_v = 2\sqrt{2\pi G\hbar}$ is whitehole entropy. This entropy pair of universe in dimension, two dimension of universe include with cover of energy. Blackhole construct with colapser of hole on kernel of atom, whitehole composite with antigravity.

$$\begin{aligned} \log(x \log x) &\geq 2(y \log y)^{\frac{1}{2}} \log(x \log y) \geq 2(xe^x)^{\frac{1}{2}} \log x + \log \log y = 2(ye^x)^{\frac{1}{2}} e^x + y = 2(ye^x)^{\frac{1}{2}} \rightarrow \frac{1}{2} \\ e^{x+1} &= 2(xe^x)^{\frac{1}{2}} \frac{e^{2(x+1)}}{4xe^x} \geq 1 \quad F_t^m = \frac{1}{4}g_{ij}^2 \frac{x+1}{4x} = y^2 \frac{1}{4} + \frac{1}{x} = y^2 \log(xy) \geq 2(yx)^{\frac{1}{2}} (\log(xy))' \geq 4(yx)^{-\frac{1}{2}} \\ \frac{1}{xy} &\geq 4(yx)^{-\frac{1}{2}} \log x + \log y \geq 4(yx)^{-\frac{1}{2}} \frac{1}{x} + \frac{1}{y} \geq 4(yx)^{-\frac{1}{2}} \frac{x+y}{xy} yx^{\frac{1}{2}} \geq 4 \frac{xy^{\frac{1}{2}}}{x+y} \leq \frac{1}{4} \frac{\Gamma(xy)}{\Gamma(x+y)} \leq \frac{1}{4} x + y \geq 2\sqrt{xy} \\ \Delta x \Delta y &\geq \frac{1}{4} i \frac{1}{xy} \geq 4(yx)^{-\frac{1}{2}} \quad F_t^m = \frac{1}{4}g_{ij}^2 e^{2(yx)^{\frac{1}{2}}} = g_{ij}^2 xy = e^{2(yx)^{\frac{1}{2}}} \rightarrow g_{ij}^2 1 + \frac{1}{x} = 4y (e^x + x)^2 \geq 4xe^x \\ r^2 &= \frac{|x_0x - 2x_0e^x + e^{2x_0x}|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = [(e^x + 1)(e^x - 1) + (x + i)(x - i)] + 2xe^{x^2} \quad r = \frac{\|1+x-4xy\|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = \\ &[(x-1)(x+1) + (x-i)(x+i)] + 2x_0e^{2x} \end{aligned}$$

Langranse conjecture include with finite line conjecture expected, infinite extend theorem exclude of finite extent. Zeta function integrate with all of math conjecture experanade. Facility of math of theorem relate of physics all of philosity. Zeta function composite with fifth dimension of equation. Curv parameter is hartshorn conjecture, this rout describe with dimension of structure. Fifth dimension rout out center of universe behind two times distance, this rout colapser around of universe. In fifth dimension, black hole and white hole in three manifold of entropy created.

Zeta function coss delete line of differential structure emerge with fifth dimension structure, oen dimension of independent vector is tangent degree. This curve of parameter create dimension of fifth dimension structure. This idea is from hartshorn conjecture. And this conjecture explanade of math mechanism, fifth dimension is AI and mass of structure parliament of pond. Time of secret is grasped in fucture and past of curve of parameter, so this equation is from universe and creature of establish of life.

Electric energy flow to neutral narrow for amusebelt on rinkfelt of harnessinkbolt, this mechanism create mind on inclusive line of brain. This flow energy on esplanade desire in routine of cone of electric energy. Endeavor circle of circuit of neutrial network, then this mechanism create on founterial motion. Dorpamin throw to neutral narrow of alcole for not being to melt and out of this narrow. This energy is supporting for Dorpamin of material things, and this circuit is circadian morment of motion. Natural killer ceil defense induce hant for lives in nature of world. This hant of mechanism is evolution on life of nature for revolution of humanism. Influence of induce hant controll with life. And these conquer of mechanism can use to neural network of evolution. These computer virus is using for not being hacking and not other controlled of oneselves computer. Safty of network is being for living in computer associate world.

Gravity esplanade theorem of cover with antigravity, information technology of exclusive injure to secure about permissset. Incrument safir to charge atom of power by pairlament in all of dimension. Reduce to string theorem of reluctance in concernism theositiy. This power contiment design emerge with inparance of contiment. All of universe have inductary, and asperant comfirm with conclusivity entily. These theorem tell uncounter deduce in consivity, therefore this envily of power concern with

cover of atom. Secure built with antigravity from gravity structure.

In reduce a morment to gravity structure, for away to certain of hant. A element of clue a vector, before opened to verify of persuade atom of power. To include of quark for twelve element, fifth dimension in zeta function to construct of geometry peices, integrate of power for fourth of series. Lack of power two bind of under a structure, antigravity permissted a power of element. This reason fourth of power get along to combind with one of decieved.

Varnished center of landscape, heat to controll of power, for missed a verisity of underground, mind to even a mid ceil.

$$\begin{aligned} F_t^m &= [f(x)] \\ [f(x)] &= \mathcal{O}(x) \\ X &\in \mathcal{O}(x) \\ 0 &\rightarrow [||\frac{d}{df}F||] \rightarrow \infty \end{aligned}$$

General number verisity is between zero and infinity. Infinite number and finite number have in general number, narrow verisity and first accerity. Gauss liner is prime number interity. Prime number and general number include Gauss liner to compilation in infinite mechanism.

$$\begin{aligned} \log(x\log x) &\geq 2(y\log y)^{\frac{1}{2}} \\ \log xy &\geq 2(xe^x)^{\frac{1}{2}} \\ (y+x)^2 - 4i(y+x)(xe)^{\frac{1}{2}} - 4xe \\ r^2(y^2+x^2) &= [(y+x)^2 - 4i(y+x)xe^{\frac{1}{2}}] - 4xe \\ r^2(x^2+y^2) &= [(\log y + \log x)^2 - 4i(\log y + \log x)xe^{\frac{1}{2}}] - 4xe \end{aligned}$$

$$\begin{aligned} \delta(f) &= \int \frac{g(c+d)}{f(a+b)} z(f+g) dz \\ y &= f(y) \\ ds^2 &= e^{-2\pi T|\phi|} [\tau_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\theta \\ \begin{bmatrix} x & y \\ a & b \end{bmatrix} \\ &\rightarrow [z] \\ f[z] &\rightarrow dx \\ x\cdot y &= 0 \\ \infty &\rightarrow [f(x)] \end{aligned}$$

Gauss liner is finite to infinity number of extend in prime number. Fifth dimension of equation composite with zeta function of infinity of extent theosity make finity of circle of structure. Independent of vector is zero dimension of category. This equaiton have a infinity of relate in number. Fifth of equation make for zeta function of infinity to conclude with this dimension out of number for finity circle. Therefor it is possiblity of being deleted to category of number that this vector is dimension of structure in fifth of dimension.


2 Neural Network from zeta function

Inclivity sensibilite make from height for maderite in mechanism. Conclusive in cercuit from hight of harmonite on interisity. The include of mechanism is quark in universe of quality of mass.

Theorem of math make for facility of mechanism in this circumstance. And these possiblity of resolved from pond of sensibility.

Light filt with eight differential structure in gravity element, then quarks made from this structure. This mechanism deal with any structure of space to create with light to graviton element. And cercuit of this mechanism dealt with carbon and hidoroxs, covalt of sixty based with filter of light element. This cercuit is dependency of any element in quark of mass, created with gravity to light of structure. Zeta function asperal with any thing to create of every mass. Fifth dimension is pond of sensibility from this gravity in lives of element.

All creature is created by this cercuit of mechanism, fifth dimension themselves is from this circumstance. This space also have oneselves with vector of independence, any this structure is from hartshorm conjecture. Space themselves haven with any of vector is tangent degree of ninety. And this vector is being of other dimension that have cercuit of theorem. These reason is from any thing born with creature, then fifth dimension haven with creature and universe. This dimension also haven with that extent from infinite and finite. Infinite of zeta function in Finite of fifth dimension, then paradox of finite cover with infinite of space.



$\chi - \{x\} = \infty$ $f(x) = \chi - \{x\} f(x) + \mathcal{O}(x) \rightarrow \text{finite}$. Add group is finite of element. This equation is fifth dimension of system. Circle of group in this delete element is infinite of set. $f(x) = [F(x)]$ $f(x)$ is Gauss liner. This set is infinite of number. $ds^2 = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\phi$ This fifth dimension is $[F(x)]$ add finite of set. Fifth dimension include with zeta function, and string theorem belong to have with infinite of consist on space. This fifth dimension is finite of space. Conclude of system solved with

infinite is covered with finite of space.

Fifth dimension in facility of mechanism that circle of infinity to emerge with being from infinity and finite of space. Zero dimension conclude with this mechanism to create from theorem of mathematics in pond of sensibility. Dimension with repository of information in facility of space, this space is all of knowlege to accessority for comformal fields. Pholographic theorem is from universe of oneself with database of creature to access with hyper dimension in General relativity. Zeta function is this theorem include with paremeter of curve from future and past of repository.

Flow to future and past that combinate with zeta function which build for hartshorn conjecture from these mechanism. This flow is gradient of line to add with fifth dimension of structure.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

Infinite of atomasphere in zero dimension include with black hole and white hole of energy in three manifold of entropy.

This entropy discribe with antigravity of energy.

$$\begin{aligned}\nabla^2\psi &= 8\pi G\hbar + 8\pi\frac{V}{S} \\ \square_v &= 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi\frac{V}{S}} \\ 2\sqrt{2\pi\frac{V}{S}} &= \int \int \frac{1}{(x \log x)^2} dx_m\end{aligned}$$

This energy flow from black hole to white hole in universe and other dimension of pair. Black hole entropy is $m = \sqrt{2\pi T}$ This entropy cover to other dimension of rout in flow energy.

Electric Weak Theorem combine with quantum flaver theorem to conjugate with light element in gravity to sheaf with antigravity. This antigravity is key for integrate with fourth of power in boson of theorem to world line to general relativity is why fourth of power not able to integrate in gravity combine with electric weak theorem and quantum flaver theorem. This point of focus is light that is filt to conjugate with eight differential structure from gravity, and this light is also element of quarks. This quark built to eight element structure. Light conjugate to eight differential element that resolved zeta function and integrate with laplace equation. Gravity significant with gravity cover with antigravity, so this antigravity is fourth element power of exist.

Entropy of vector have zero dimension in all exist into all of equation. Infinite of number is three dimension, therefore it is esperial of important that created by zero dimension from infinite to finite of equation rout.

$$\begin{aligned}f(x) &= 0 \\ F &= \int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m\end{aligned}$$

$$F = 0, x \cdot y = 0, F(x) = 0, G(x) = 0, \mathcal{O} \in X, X \cong 0, x - y = 0, F(x, y) = 0, x^n + y^n = z^n, \\ F(x) = x^n + y^n, F(x) = 0$$

$$ds^2 = \frac{r}{d + e \cos \theta}$$

Fifth dimension discribe with norm liner in zero dimension. This equation conclude with all of equation to pond sensibility.

Pond of sensibility is facility of mathmatics in infinite of mechanism, this mechanism is zero dimension for imaginary of equation in antigravity influence. This power is gravity on cover with non symmetry combine of fourth of power. That fourth of power is the reason with quantum flaver theorem combine with gravity of power, antigravity is brane of string key mechanism. This power is non-catastrophe of influent reason, and that mechanism of power result. Integrate of gravity include with antigravity, and fourth of power is one of geometry in universe.

$$\mathcal{O} \in X \rightarrow \infty, F(x) = z^n$$

$$x^n + y^n = z^n$$

$$F(x) = \mathcal{O}(\delta(x))[(x - f(x))(x + f(x)) + (y - \overline{f(x)})(y + \overline{f(x)})] + O(N^{-1})$$

$$F(x) = 0$$

$$f(x) = \sum_{k=0}^{\infty} a_k f^k(x)$$

$$f(x) = 0$$

$$\int f(x), \partial f(x), dx \rightarrow [f(x)]$$

Group that Galois theorem develop with three dimension construct of prime number equation, this element belong for that is three of factor. Fifth dimension concept with this group of conclude element.

$$\mathcal{O}(x) = \chi(x) - \{x\}$$

$$\sum_{k=0}^{\infty} a_k x^k = f(x) + \{x\}$$

Dimension of fifth factor is abel manifold, this manifold is topology resolved.

$$\int dx, \partial x, dx, [f(x)]$$

$$\rightarrow F(x) \xrightarrow{f \, dx} f(x) \xrightarrow{\partial x} x \xrightarrow{x} const \\ \rightarrow [f(x)] \rightarrow \infty$$

This cycle of topology in general theroem is deduce of mechanism.

Universe of structure is duality of manifold, this mass of darkmatter is oneseleves with reason in singularity and monotonicity of universe. This manifold of three dimension is also zeta function.

Light around of universe for darkmatter is Higgs fields of mechanism. Darkmatter create of including with light of quarks in Higgs fields. Eight of differential structure fil with Thurston conjugate thoeorem to light is created with all dimension of element.

Quantum Computer in a certain theorem

Masaaki Yamaguchi

A pattern emerge with one condition to being assembled of emelite with all of possibility equation, this assembled with summative of manifold being elemetiled of pieace equation. This equation relate with Euler equation. And also this equation is Euler constant oneselves.

$$(E_2 \bigoplus E_2) \cdot (R^- \subset C^+) = \bigoplus \nabla C_-^+$$

Zeta function radius with field of mechanism for atom of pole into strong condition of balance, this condition is related with quarks of level controlled for compute with quantum tonnel effective mechanism. Quantum mechanism composed with vector of constance for zeta function and quantum group. Thurston conjugate theorem explain to emerge with being controll of quantum levels of quarks. Locality theorem also occupy with atom of levels in zeta function.

$$\begin{aligned} &= \bigoplus \nabla C_-^+ \\ \vee \int \frac{C^+ \nabla M_m}{\Delta(M_-^+ \nabla C_-^+)} &= \exists(M_-^+ \nabla R^+) \\ \exists(M_-^+ \nabla C^+) &= \text{XOR}(\bigoplus \nabla M_-^+ \\ &\quad -[E^+ \nabla R^+] = \nabla_+ \nabla_- C_-^+ \\ \int dx, \partial x, \nabla_i \nabla_j, \Delta x &\rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+ \end{aligned}$$

Zeta function also compose with Rich flow equation cohomological result to equal with locality equaitons.

$$\begin{aligned} \vee(R + \nabla_i \nabla_j f)^n &= \int \frac{\wedge(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)^n} \\ \wedge(R + \nabla_i \nabla_j f)^x &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dt} g_{ij}(x) &= -2R_{ij} \\ \vee \int \wedge(R + \nabla_i \nabla_j f)^x &= \frac{\wedge(R + \nabla_i \nabla_j f)^n}{\exists(R + \nabla_i \nabla_j f \circ g)^n} \\ x + y &\geq 2\sqrt{xy}, x(x) + y(x) \geq x(x)y(x) \\ x^y &= (\cos \theta + i \sin \theta)^n \\ x^y &= \frac{1}{y^x} \end{aligned}$$

Therefore zeta function is also constructed with quantum equation too.

UFO mechanism

Masaaki Yamaguchi

UFO mechanism are several system of circumstance that accesority and verisity composite with mass and circument of gravity is key of mechanism included with anti-gravity in imaginary pole emerge with rotate of right formula. This power of rotate emerge with anti-gravity which of power emelite for inner into oneselves create with like of lorentz of energy. This lorentz of energy is use with steles flight of mechanism. However this lorentz of energy only mass of gravity low included with oneselves but also anti-gravity is oneselves of component with inner of rotate with energy, this energy is not free of condition of power, because this power not ordinary of mechanism.

$$\begin{aligned} F - N &= mg - N' \\ \frac{d}{df}F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ C &= \int \int \frac{1}{(x \log x)^2} dx_m \\ \int \int \frac{1}{(x \log x)^2} dx_m &= \frac{1}{2}i \end{aligned}$$

This imaginary number is rotate of quantum spin emerged with power, this power is not mass of gravity element, also this power is inner of energy and anti-gravity of power.

Moreover this right of rotate of imaginary pole is three manifold of nourth and sournth of formula. And nourth of formula look with gravity but sournth of formula is anti-gravity, this mechanism is relativity theorem from explained.

Hortshorne conjecture is built with fermer theorem spacificed from AdS5 manifold constructed of sphere cube exclude with time machine mechanism element. This dimension cercuited with rout of pole established by right angle in systematic vintage emerge with artificial intelligence. This one element of independence create with imaginary number from inner vector element, this resulted rentanse for rorentz atructer in super string theory refill into globality topology mechanism constructed with these essence. This dimension target with native function around covered with abel manifold, seifert manifold is become with other dimension on pair universe essensed with zeta function, this space time is concluded with finite dimension in around of shell, this cohomology of mapping construct with abel manifold is all of equal between blackhold and zeta function inverse with low structure of constance in quantum operator relative with univese and planck scale resulted with mechanism. This dimension equal with whitehole and this pair of dimension is blackhole from being emerged with exclusive and inclusive of component from universe and other dimension system.

This explained mechanism from photographic theorem is atom of low structure of constance with time space and differential structure relative from UFO machine system concerned with Hortshorne conjecture. This photographic theorem is explained with UFO machine resulted by mass and circumvent of gravity not existed zone. Inner of rotate with energy is component oneselves with being not free of condition element. More spectrum explained this power is inner of energy and anti-gravity of power. Accessory and verisity have with mass and circumvent of gravity which existed is out of zone element. However this element is not oneselves of inner and circumvent of gravity is not only spective of inner influence of power but also out of influent power. However anti-gravity is all of inner spective oneselves emerged with energy of power.

Hortshorne conjecture and AdS5 manifold moreover photographic theorem are concerned with UFO machine system be able to be explained from Atom of inspected with movement spector phenomone artificial experient. This experient result with UFO machine is not free condition of zone success with inner of power being with elemelite of anti-gravity.

From under circumvent position to right of rotate energy formula create with anti-gravity, from up circumvent position to left of rotate energy formula, these way of relativity system is resulted with other dimension of rotate energy reason. Sourth of rotate way is anti-gravity, nourth of rotate way is circumvent of gravity. This mechanism construct with non-symmetry of entropy resulted. And this theorem explain that parity of broken result. This parity of mechanism is other dimension and universe of seifert manifold of balance of entropy concluded. Symmetry formula systemalite with non-vacuum constructed with two-projection space not be existed in three manifold dimension. This mechanism is pair of quarks with universe and other dimension. If universe belong for being emerged with energy of quarks in three manifold, then other dimension migrated with entropy saved of energy of universe balance. If dimension create with quarks of LHC experiment by exist proved, imaginary pole of rotate create with sourth of formula, universe and other dimension is balanced with energy, then reverse of power of gravity and anti-gravity are proof that shirt of quarks lives with oneselves.

Vector Operator

Masaaki Yamaguchi

1 Entropy

Already discover with this entropy of manifold is blackhole of energy, these entropy is mass of energy has norm parameter. And string theorem $T_{\mu\nu}$ belong to ρ energy, this equation is same Kaluze-Klein theorem.

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

$$y=x$$

$$y=(\nabla\phi)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}}}=\frac{p}{c^3}+\rho$$

$$ds^2=g_{\mu\nu}(x)dx^\mu dx^\nu+\phi^2(x)\kappa^2A_{\mu\nu}(x)dx^\mu)^2$$

$$ds=(g_{\mu\nu}(x)dx^\mu dx^\nu+\phi^2(x)(\kappa^2A_{\mu\nu}(x)dx^\mu)^2)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}}-\rho}=c^3,\frac{p}{2\pi}=c^3$$

$$ds^2=e^{-2kT(x)|\phi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2(x)d\phi^2$$

$$f_z=\int\left[\sqrt{\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}\circ\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}}\right]dxdydz$$

$$\frac{\partial}{\partial f}\int (\sin 2x)^2 dx = ||x-y||^2$$

2 Atom of element from zeta function

2.1 Knot element

String summulate of landmaistor movement in these knot element. Supersymmetry construct of these mechanism from string theorem. A element of homomorphism boundary for kernel function, equal element has meyer beatorise compose structure in mesh emerge. These mesh for atom element equal knot this of exchange, this isomorphism three manifold convert with knot these from laplace space. Duality manifold has space equal other operate, this quantum group is element distinguish with other structure. Aspect element has design pattern, this element construct pair of entropy. Quote algebra reflect with atom kernel of function.

3 Imaginary manifold

A vector is scalar, vector, mass own element, imaginary number is own this element.

$$x = \frac{\vec{x}}{|\vec{x}|}$$

For instance

$$|\vec{x}| = \sqrt{x}, i = \sqrt{-1}, |\vec{x}|x = \vec{x}, |\vec{x}| = 1x, |\vec{x}|^2 = -1$$

$$\left(\frac{x}{|\vec{x}|}\right)^2 = \frac{1}{-1}, |\sqrt{x}| = i$$

Imaginary number is vector, inverse make to norm own means.

4 Singularity

A element add to accept for group of field, this group also create ring has monotonicity. Limit of manifold has territory of variable limit of manifold. This fields also merge of mesh in mass. Zero dimension also means to construct with antigravity. This monotonicity is merged to has quote algebra liner. This boundary operate belong to has category theorem. Meyer beatorise extend liner distinguish with a mesh merged in fixed point theorem. Entropy means to isohomorphism with harmony this duality. The energy lead to entropy metric to merge with mesh volume.

5 Time expand in space for laplace equation

6 Laplace equaiton

Seiferd manifold construct with time and space of flow system in universe. Zeta function built in universe have started for big-ban system. Firstly, time has future and past in space, then space is expanding for big-ban mechanism.

Darkmatter equals for Higgs field from expanded in universe of ertel. Three manifold in flow energy has not no of a most-small-flow but monotonicity is one. Because this destruct to distinguish with eight-differential structure, and not exist.

And all future and past has one don't go on same integrate routs. This conject for Seiferd manifold flow. Selbarg conjecture has singularity rout. Landemaistor movement become a integrate rout in this Selbarg conjecture.

Remain Space belong for three manifold to exist of singularity rout. Godsharke conjecture is this reason in mechanism. Harshorn conjecture remain to exist multi universe in Laplace equation resolved. Delete line summulate with rout of integrate differential rout. This solved rout has gradient flow become a future and past in remain space.



Laplace equation composite for singularity in zeta function and Kaluze-Klein space.

7 Non-commutative algebra for antigravity

Laplace equation and zeta function belong for fermion and boson, then power of construct equation merge Duality of symmetry formula. These power construct of deceive in dimension, and deal the other dimension. Reflection of power erasing of own dimension, the symmetry of formula has with merging of energy, then these power has fourth of basic power Symmetry solved for equation, general relativity deal from reality these formula. Quantum group construct of zeta function and Laplace equation. Imaginary number consist of Euler constance, this prime number reach to solve contiguish of power in emergy of parameter. Zeta function conclude of infinite to finite on all equation. Global differential equation merge a mass of antigravity and gravity.

Module exist zero dimension merge of symmetry in catastrophe D-brane, this mechanism of phenomoun is Non-symmetry explain to emerge with which cause of mass on gravity.

$$x \bmod N = 0$$

$$\sum_{M=0}^{\infty} \int_M dm \rightarrow \sum_{x=0}^{\infty} F_x = \int_m dm = F$$

$$\frac{x-y}{a} = \frac{y-z}{b} = \frac{z-x}{c}$$

$$\frac{\Pi(x-y)}{\partial xy}$$

Volume of space interval to all exist area is integrate rout in universe of shape, this space means fo no exist loop on entropy of energy's rout.

$$||\int f(x)||^2 \rightarrow \int \pi r^2 dx \cong V(\tau)$$

This equation is part of summulate on entropy formula.

$$\frac{\int |f(x)|^2}{f(x)}dx$$

$$V(\tau) \rightarrow mesh$$

$$\int \left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right| dv(\tau)$$

$$\left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right|_{dx=v}$$

$$z_y=a_x+b_y+c_z$$

$$dz_y=d(z_y)$$

$$[f,f^{-1}]=ff^{-1}-f^{-1}f$$

Universe of extend space has non-integrate rout entropy in sufficient of mass in darkmatter, this reason belong for cohomology and heat equation concerned. This expanded mass match is darkmatter of extend. And these heat area consume is darkmatter begun to start big-ban system invite of universe.

$$V(\tau)=\int \tau(q)^{-\frac{n}{2}}\exp(-\frac{1}{\sqrt{2\tau(q)}}L(x)dx)+O(N^{-1})$$

$$\frac{1}{\tau}(\frac{N}{2}+\tau(2\Delta f-|\nabla f|^2+R)+f)\mathrm{mod}N^{-1}$$

$$\Delta E=-2(T-t)|R_{ij}+\nabla_i\nabla_jf-\frac{1}{2(T-t)}g_{ij}|^2$$

$$\frac{d}{df}F=\frac{2\int (R+\nabla_i\nabla_jf)^2}{-(R+\Delta f)}dm$$

$$\frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$dx=(g_{\mu\nu}(x)^2dx^2-g_{\mu\nu}(x)dxg_{\mu\nu})^{\frac{1}{2}}$$

$$ds^2 = -N(r)^2 dt^2 + \psi^2(r)(dr^2 + r^2 d\theta^2)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$\sum_{n=0}^{\infty} a_1x^1+a_2x^2\ldots a_{n-1}x^{n-1}\rightarrow \sum_{n=0}^{\infty} a_nx^n\rightarrow \alpha$$

$$f=n\nu\lambda, \lambda=\frac{x}{l}, \int dn\nu\lambda=f(x), xf(x)=F(x), [f(x)]=\nu h$$

8 Three manifold of dimension system worked with database

This universe built with fourth of dimension in add manifold constructed with being destructed from three manifold.

$$\mathcal{O}(x) = ([\nabla_i \nabla_j f(x)])'$$

$$\cong {}_nC_r(x)^n(y)^{n-r}\delta(x,y)$$

$$(\Box\psi)'=\nabla_i\nabla_j(\delta(x)\circ G(x))^{\mu\nu}\left(\frac{p}{c^3}\circ\frac{V}{S}\right)$$

$$F_t^m=\frac{1}{4}g_{ij}^2,x^{\frac{1}{2}+iy}=e^{x\log x}$$

$$S_m^{\mu\nu}\otimes S_n^{\mu\nu}=G_{\mu\nu}\times T^{\mu\nu}$$

This equation is gravity of surface quality, and this construct with sheap of manifold.

$$S_m^{\mu\nu}\otimes S_n^{\mu\nu}=-\frac{2R_{ij}}{V(\tau)}[D^2\psi]$$

This equation also means that surface of parameters, and next equation is non-integral of routs theorem.

$$S_m^{\mu\nu}=\pi(\chi,x)\otimes h_{\mu\nu}$$

$$\pi(\chi,x)=\int \exp[L(p,q)]d\psi$$

$$ds^2=e^{-2\pi T|\phi|}[\eta+\bar{h}_{\mu\nu}]dx^{\mu\nu}dx^{\mu\nu}+T^2d^2\psi$$

$$M_3\bigotimes_{k=0}^\infty E_-^+=\mathrm{rot}(\mathrm{div}E,E_1)$$

$$=m(x),\frac{P^{2n}}{M_3}=H_3(M_1)$$

These equation is conclude with dimension of parameter resolved for non-metric of category theorem. Gravity of surface in space mechanism is that the parameter become with metric, this result construct with entropy of metric.

These equation resulted with resolved of zeta function built on D-brane of surface in fourth of power.

$$\begin{aligned}\exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} &= \int \exp[L(p, q)]d\psi \\ &= \exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} \otimes \int \exp[L(p, q)]d\psi + N\text{mod}(e^{x \log x}) \\ &= \mathcal{O}(\psi)\end{aligned}$$

9 Particle with each of quarks connected from being stimulate with gravity and antigravity into Artificial Intelligence

This equation conclude with middle particle equation in quarks operator, and module conjecture excluded. $\frac{d}{dt}g_{ij}(t) = -2R_{ij}$, $\frac{P^{2n}}{M_3} = H_3(M_1)$, $H_3(M_1) = \pi(\chi, x) \otimes h_{\mu\nu}$ This equation built with quarks emerged with supersymmetry field of theorem. And also this equation means with space of curvature parameters.

$$\begin{aligned}S_m^{\mu\nu} \times S_n^{\mu\nu} &= [D^2\psi], S_m^{\mu\nu} \times S_n^{\mu\nu} = \ker f / \text{im} f, S_m^{\mu\nu} \otimes S_n^{\mu\nu} = m(x)[D^2\psi], -\frac{2R_{ij}}{V(\tau)} = f^{-1}xf(x) \\ f_z &= \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix} \circ \begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \right] dx dy dz, \rightarrow f_z^{\frac{1}{2}} \rightarrow (0, 1) \cdot (0, 1) = -1, i = \sqrt{-1}\end{aligned}$$

This equation operate with quarks stimulated in Higgs field.

$$(x, y, z)^2 = (x, y, z) \cdot (x, y, z) \rightarrow -1$$

$$\mathcal{O}(x) = \nabla_i \nabla_j \int e^{\frac{2}{m} \sin \theta \cos \theta} \times \frac{N\text{mod}(e^{x \log x})}{\mathcal{O}(x)(x + \Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta, \mathcal{O}(x) = m(x)[D^2\psi]$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = I'_m, I'_m = [1, 0] \times [0, 1]$$

This equation means that imaginary pole is universe and other dimension each cover with one geometry, and this power remind imaginary pole of antigravity.

$$i^2 = (0, 1) \cdot (0, 1), |a||b| \cos \theta = -1, E = \text{div}(E, E_1)$$

$$\left(\frac{\{f, g\}}{[f, g]} \right)' = i^2, E = mc^2, I' = i^2$$

This fermion of element decide with gravity of reflected for antigravity of power in average and gravity of mass, This equals with differential metric to emerge with creature of existing materials.

These equation deconstructed with mesh emerged in singularity resolved into vector and norm parameters.

$$\mathcal{O}(x) = ||\nabla \int [\nabla_i \nabla_j f \circ g(x)]^{\frac{1}{2}+iy} ||, \partial r^n ||\nabla||^2 \rightarrow \nabla_i \nabla_j ||\vec{v}||^2$$

$\nabla^2 \phi$ is constructed with module conjecture excluded.

$$\nabla^2 \phi = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

Reflected equation concern with differential operator in metric, sine and cosine function also rotate of differential operator in metric. Weil's theorem concern with this calculate in these equation process.

$$(\log x^{\frac{1}{2}})' = \frac{1}{2} \frac{1}{(x \log x)}, (\sin \theta)' = \cos \theta, (f_z)' = ie^{ix \log x}, \frac{d}{df} F = m(x)$$

$$\begin{aligned} \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} + \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq 2h \end{aligned}$$

Three dimension of manifold equals with three of sphere in this dimension. Plank metrics equals $\hbar$ equation to integrate with three manifold in sphere of formula. These explain recognize with universe and other dimension of system in brane of topology theorem.

$$\frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm \geq \hbar$$

$$y = x, xy = x^2, (\Box \psi)' = 8\pi G \left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

This equation built with other dimension cover with universe of space, then three dimension invite with fifth dimension out of these universe.

$$\Box \psi = \int \int \exp[8\pi G(\bar{h}_{\mu\nu} \circ \eta_{\mu})^{\nu}] dm d\psi, \sum a_k x^k = \frac{d}{df} \sum \sum \frac{1}{a_k^2 f^k} dx_k$$

$$\sum a_k f^k = \frac{d}{df} \sum \sum \frac{\zeta(s)}{a_k} dx_{k_m}, a_k^2 f^{\frac{1}{2}} \rightarrow \lim_{k \rightarrow 1} a_k f^k = \alpha$$

$$\mathcal{O}(x) = D^2 \psi \otimes h_{\mu\nu}, ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^{\mu} dx^{\nu} + T^2 d^2 \psi$$

$$f(x) + f(y) \geq 2\sqrt{f(x)f(y)}, \frac{1}{4}(f(x) + f(y))^2 \geq f(x)f(y)$$

$$T^{\mu\nu}=\left(\frac{p}{c^3}+\frac{V}{S}\right)^{-1}, E^+=f^{-1}xf(x), E=mc^2$$

$$\mathcal{O}(x)=\square\int\int\int\frac{(\nabla_i\nabla_jf\circ g(x))^2}{V(x)}dm$$

$$ds^2=g^2_{\mu\nu}d^2x+g_{\mu\nu}dxg_{\mu\nu}(x), E^+=f^{-1}xf(x), \frac{V}{S}=AA^{-1}, AA^{-1}=E$$

$$\mathcal{O}(x)=\int\int\int f(x^3,y^3,z^3)dx dy dz, S(r)=\pi r^2, V(r)=4\pi r^3$$

$$E^+_-=f(x)\cdot e^{-x\log x},\sum_{k=0}^{\infty}a_kf^k=f(x)\cdot e^{-x\log x}$$

$$\frac{P^{2n}}{M_3}=E^+-\phi,\frac{\zeta(x)}{\sum\limits_{k=0}^{\infty}a_kx^k}=\mathcal{O}(x)$$

$$O(N^{-1})=\frac{1}{\mathcal{O}(x)}, \square=\frac{8\pi G}{c^3}T^{\mu\nu}$$

$$\partial^2 f(\square\psi)=-2\square\int\int\int\frac{V}{S^2}dm,\frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$E^+_-=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}]dx^\mu dx^\nu+T^2d^2\psi$$

$$S_1^{mn}\otimes S_2^{mn}=D^2\psi\otimes h_{\mu\nu}, S_m^{\mu\nu}\otimes S_n^{\mu\nu}=\int[D^2\psi]dm$$

$$=\nabla_i\nabla_j\int f(x)dm, h=\frac{p}{mv}, h\lambda=f, f\lambda=h\nu$$

D-brane for constructed of sheap of theorem in non-certain mechanism, also particle and wave of duality element in D-brane.

$$||ds^2||=S_m^{\mu\nu}\otimes S_n^{\mu\nu}, E^+_-=f^{-1}(x)xf(x)$$

$$R^+\subset C^+_{-}, \nabla R^+\rightarrow \bigoplus Q^+_-$$

Complex of group in this manifold, and finite of group decomposite theorem, add group also.

$$\nabla_i\nabla_jR^+_-=\bigoplus\nabla Q^+_-$$

Duality of differential manifold, then this resulted with zeta function.

$$M_1=\frac{R_3^+}{E_-^+}-\{\phi\}, M_3\cong M_2, M_3\cong M_1$$

World line in one of manifold.

$$H_3(\Pi)=Z_1\oplus Z_1, H_3(M_1)=0$$

Singularity of zero dimension. Lens space in three manifold.

$$ds^2 = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

Fifth dimension of manifold. And this constructed with abel manifold.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

These system flow to build with three dimension of energy.

$$\begin{aligned} (\partial\gamma^n + m^2) \cdot \psi &= \int [D^2\psi \otimes h_{\mu\nu}] dm \\ &= 0 \end{aligned}$$

Complex of connected of element in fifth dimension of equation.

$$\begin{aligned} \square &= \pi(\chi, x) \otimes h_{\mu\nu} \\ &= D^2\psi \otimes h_{\mu\nu} \end{aligned}$$

Fundamental group in gravity of space. And this stimate with D-brane of entropy.

$$\begin{aligned} \int [D^2\psi] dm &= \pi(M_1), H_n(m_1) = D^2\psi - \pi(\chi, x) \\ &= \ker f / \operatorname{im} f \end{aligned}$$

Homology of non-entropy.

$$\int Dq \exp[L(x)] d\psi + O(N^1) = \pi(\chi, x) \otimes h_{\mu\nu}$$

Integral of rout entropy.

$$= D^2\psi \otimes h_{\mu\nu}$$

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} \frac{\zeta(x)}{a_k f^k} = \int ||[D^2\psi \otimes h_{\mu\nu}]|| dm$$

Norm space.

$$\nabla\psi^2 = \square \int \int \int \frac{V}{S^2} dm$$

$$\mathcal{O}(x) = D^2\psi \otimes h_{\mu\nu}$$

These operators accept with level of each scalas.

$$dx < \partial x < \nabla \psi < \square v$$

$$\oplus < \sum < \otimes < \int < \wedge$$

$$(\delta\psi(x))^2 = \int \int \int \frac{V(x)}{S^2} dm, \delta\psi(x) = \left(\int \int \int \frac{V(x)}{S^2} dm \right)^{\frac{1}{2}}$$

$$\nabla\psi^2 = -4R \int \delta(V \cdot S^{-3}) dm$$

$$\nabla\psi = 2R\zeta(s)i$$

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{a_k x^k}{m dx} f^k(x) &= \frac{m}{n!} f^n(x) \\ &= \frac{(\zeta(s))^k}{df} m(x), (\delta(x))^{\frac{1}{2}} = \left(\frac{x \log x}{x^n} \right)^n \end{aligned}$$

$$\mathcal{O}(x) = \frac{\int [D^2\psi \otimes h_{\mu\nu}] dm}{e^{x \log x}}$$

$$\mathcal{O}(x) = \frac{V(x)}{\int [D^2\psi \otimes h_{\mu\nu}] dm}$$

Quantum effective equation is lowest of light in structure of constant. These resulted means with plank scals of universe in atom element.

$$\begin{aligned} M_3 &= e^{x \log x}, x^{\frac{1}{2}+iy} = e^{x \log x}, (x) = \frac{M_3}{e^{x \log x}} \\ &= nE_x \end{aligned}$$

These equation means that lowest energy equals with all of materials have own energy. Zeta function have with plank energy own element, quantum effective theorem and physics energy also equals with plank energy of potential levels.

$$\frac{V}{S^2} = \frac{p}{c^3}, h\nu \neq \frac{p}{mv}, E_1 = h\nu, E_2 = mc^2, E_1 \cong E_2$$

$$l = \sqrt{\frac{\hbar G}{c^3}}, \frac{\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m}{\int \int \frac{1}{(x \log x)^2} dx_m} = \frac{1}{i}$$

Lowest of light in structures of constant is gravity and antigravity of metrics to built with zeta function. This result with the equation conclude with quota equation in Gauss operators.

$$\begin{aligned} ihc = G, hc &= \frac{G}{i}, \frac{\frac{1}{2}}{\frac{1}{2}i} = \frac{\vec{v_1}}{\vec{v_2}} \\ &\leq 1 \end{aligned}$$

$$A=BQ+R, ds^2=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2d^2\psi$$

$$ds^2=g_{\mu\nu}dx^\mu dx^\nu+\kappa^2(A^{\mu\nu})^2,\int\int e^{-x^2-y^2}dxdy=\pi$$

$$\begin{aligned}\Gamma(x) &= \int e^{-x} x^{1-t} dx \\ &= \delta(x) \pi(x) f^n(x)\end{aligned}$$

$$\frac{d}{df}F=\frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

Quota equation construct of piece in Gauss operator, abel manifold equals with each equation.

$$ds^2=[T^2d^2\psi]$$

$$\mathcal{O}(x)=[x]$$

$$\nabla\psi^2=8\pi G\left(\frac{p}{c^3}\circ\frac{V}{S}\right)$$

Volume of space equals with plank scals.

$$\frac{pV}{S}=h$$

Summuate of manifold means with beta function to gamma function, equals with each equation.

$$\beta(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}$$

$$\ker f/\mathrm{im} f \cong \mathrm{im} f/\ker f$$

$$\bigoplus \nabla g(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta], \bigcup_{k=0}^{\infty} \left(\bigoplus \nabla f(x) \right) = \square \int \int \int \nabla g(x) d\eta$$

Rich tensor equals of curvature of space in Gauss liner from differential equation of operators.

$$a' = \sqrt{\frac{v}{1 - (\frac{v}{c})^2}}, F = ma'$$

Accessory put with force of differential operators.

$$\nabla f(x) = \int_M \square \left(\bigoplus \nabla f(x) \right)^n dm$$

$$\square = 2(T-t)|R_{ij} + \nabla \nabla f - \frac{1}{2(T-t)}|g_{ij}^2$$

$$(\square + m) \cdot \psi = 0$$

$$\square \times \square = (\square + m^2) \cdot \psi, (\partial \gamma^n + \delta \psi) \cdot \psi = 0$$

$$\nabla_i \nabla_j \int \int_M \nabla f(t) dt = \square \left(\bigcup_{k=0}^{\infty} \bigoplus \nabla g(x) d\eta \right)$$

Dalanverle equation of operator equals with summuate of category theorem.

$$\int_M (l \times l) dm = \sum l \oplus ld\eta$$

Topology of brane equals with string theorem, verisity of force with add group.

$$\begin{aligned} &= [D^2 \psi \otimes h_{\mu\nu}]^{\frac{1}{2} + iy} \\ &= H_3(M_1) \end{aligned}$$

$$y = \nabla_i \nabla_j \int \nabla g(x) dx, y' - y + \frac{d}{dx} z + [n] = 0$$

Equals of group theorem is element of equals with differential element.

$$\begin{aligned} z &= \cos x + i \sin x \\ &= e^{i\theta} \end{aligned}$$

Imaginary number emerge with Artificial Intelligence.

These equation resulted with prime theorem equals from differential equation in fields of theorem.

Imaginary pole of theorem and sheap of duality manifold. Duality of differential and integrate equation. Summuate of manifold and possibility of differential structure. Eight of differential structure is stimulate with D-brane recreated of operators. Differential and integrate of operators is emerged with summuated of space.

$$T^{\mu\nu} = -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$G_{\mu\nu}=R_{\mu\nu}T^{\mu\nu},\sigma(x)\oplus\delta(x)=(E_n^+\times H_m)$$

$$G_{\mu\nu}=[\frac{\partial}{\partial f}R_{ij}]^2,\delta(x)\cdot V(x)=\lim_{n\rightarrow 1}\delta(x)$$

$$\lim_{n\rightarrow\infty}\mathrm{mesh}V(x)=\frac{m}{m+1}$$

$$V(x)=\sigma\cdot S^2(x),\int\int\int\frac{V(x)}{S^2(x)}dm=[D^2\psi\otimes h_{\mu\nu}]$$

$$g(x)|_{\delta(x,y)}=\frac{d}{dt}g_{ij}(t),\sigma(x,y)\cdot g(x)|_{\delta(x,y)}=R_{ij}|_{\sigma(x,y)}$$

$$=\int R_{ij}^{a(x-y)^n+r^n}$$

$$(ux+vy+wz)/\Gamma$$

$$=\int R_{ij}^{(x-u)(y-v)(z-w)}dV$$

$$(\Box+m)\cdot\psi=0, E=mc^2, \frac{\partial}{\partial f}\Box\psi=4\pi G\rho$$

$$(\partial\gamma^n+m)\cdot\psi=0, E=mc^2-\frac{1}{2}mv^2$$

$$=(-\frac{1}{2}\left(\frac{v}{c}\right)^2+m)\cdot c^2$$

$$=(-\frac{1}{2}a^2+m)\cdot c^2, F=ma, \int adx=\frac{1}{2}a^2+C$$

$$T^{\mu\nu}=-\frac{1}{2}a^2,(e^{i\theta})'=ie^{i\theta}$$

Artificial Intelligence and TupleSpace of ultranetwork

Masaaki Yamaguchi

```

Omega::DATABASE[tuplespace]
{
    Z \supset C \bigoplus \nabla R^{+}, \nabla(R^{+}
    \cap E^{+}) \ni x, \Delta(C \subset R) \ni x
    M^{+}_{-}\bigoplus R^{+}, E^{+} \in
    \bigoplus \nabla R^{+}, S^{+}_{-} \subset R^{+}_{2},
    V^{+}_{-} \times R^{+}_{-} \cong \{V \over S\}
    C^{+} \cup V^{+}_{-} \ni M_{1} \bigoplus \nabla C^{+}_{-},
    Q \supseteq R^{+}_{-},
    Q \subset \bigoplus M^{+}_{-},
    \bigotimes Q \subset \zeta(x), \bigoplus \nabla C^{+}_{-} \cong M_3
    R \subset M_3,
    C^{+} \bigoplus M_n, E^{+} \cap R^{+},
    E_2 \bigoplus E_1, R^{-} \subset C^{+}, M^{+}_{-}
    C^{+}_{-}, M^{+}_{-} \nabla C^{+}_{-}, C^{+} \nabla H_m,
    E^{+} \nabla R^{+}_{-}, E_2 \nabla E_1,
    R^{-} \nabla C^{+}_{-}

    [- \Delta v + \nabla_{\{i\}} \nabla_{\{j\}} v_{\{ij\}} - R_{\{ij\}} v_{\{ij\}}
    - v_{\{ij\}} \nabla_{\{i\}} \nabla_{\{j\}} + 2 < \nabla f, \nabla h>
    + (R + \nabla f^2)(\{v \over 2\} - h)]

    S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1,
    H^1 \times S^1, H^1, S^2 \times E
}

import Omega::Tuplespace < DATABASE
{
    {\bigoplus M^{+}_{-} -> =: \nabla R^{+} \nabla C^{+}}-< [construct_emerge_equation.built]
    >> VIRTUALMACHINE[tuplespace]
    => {regextp.pattern |w|
        w.scan(equal.value) [ > [\nabla \int \int \nabla_{\{i\}} \nabla_{\{j\}} f \circ g(x)]
        equal.value.shift => tuplespace.value
        w.emerged >> |value| value.equation_create
        w <- value
        w.pop => tuplespace.value
    }

    {\vee (\int \nabla_{\{i\}} \nabla_{\{j\}} (R + \Delta f)^2)

```

```

\over \exists (R + \Delta f)} -> =: variable array[]
>> VIRTUAL_MACHINE[tuplespace]
=> {regextpt.pattern |w|
    w.emerged => tuplespace[array]
    w <- value
    w.pop => tuplespace.value
}
}

Omega.DATABASE[tuplespace]->w.emerged >> |value| value.equation_create
{
    w.process <- Omega.space
    {=>
        cognitive_system :=> tuplespace[process.excluded].reload
        assembly_process <- w.file.reload.process
        => : [regextpt.pattern(file)=>text_included.w.process]
    }
}

Omega.DATABASE[tuplespace]->w.emerged >> |list| list.equation_create
{
    w.process <- Omega.space
    {=>
        poly w.process.cognitive_system :=> tuplespace[process.excluded].reload
        homology w.process :=> tuplespace[process.excluded].reload
        mesh.volume_manifold :=> tuplespace[process.excluded].reload
        \nabla_{i}\nabla_{j} w.process.excluded :=> tuplespace[process.excluded].reload
        {\exp[\int \int (R + \Delta f)^2 e^{-x \log x}dV}.emerge_equation.reality{|repository|
            repository.regextpt.pattern => tuplespace[process.excluded].reload
            tuplespace[process.excluded].rebuild >> Omega.DATABASE[tuplespace]
            {\imaginary.equation => e^{\cos \theta + i \sin \theta}} <=> Omega.DATABASE[tuplespace]
            {{d \over df}F ==> {d \over df}{1 \over {(x \log x)^2 \circ (y \log y)
                ^{1 \over 2}}dm}.cognitive_system.reload
            :=> [repository.scan(regextpt.pattern) { <=> btree.scan |array| <-> ultranetwork.attachment}
            repository.saved
        }
    }
}

import ultra_database.included
def < this.class::Omega.DATABASE[first,second,third.fourth] end
def.first.iterator => array.emerge_equation
def.second.iterator => array.emerge_equation
def.third.iterator => array.emerge_equation
def.fourth.iterator => array.emerge_equation
_struct_ {
    Omega.iterator => repository.reload
}
end
typedef _ struct_ :Omega.aspective
end

```

```

Omega::DATABASE[reload]
{
  [category.repository <-> w.process] <=> catastrophe.category.selected[list]
  list.distributed => ultra_database.exist ->
  w.summurate_pattern[Omega.Database]
  btree.exclude -> this.klass
  list.scan(regexpt.pattern) <-> btree.included
  list.exclude -> [Omega.Database]
  all_of_equation.emerged <=> Omega.Database
  {
    list.summuate -> Omega.Database.excluded
  }
}

list.distributed => {
  {\bigoplus \nabla M^{+}_{-}}.constructed <-> Omega.Database[import]
  {=>
    each_selected :file.excluded
  }
}

Omega::DATABASE[tuplespace] >> list.cognitive_system |value|
= { x^{\frac{1}{2}} + iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h
x^{\frac{1}{2}} + iy} = \mathrm{exp}[\int \nabla_i \nabla_j f(x) g'(x) /
\partial f \partial g]

\mathcal{0}(x) = \{[f(x) \circ g(x) , \bar{h}(x)], g^{-1}(x)\}

\exists [\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty}
\nabla \int \nabla_i \nabla_j f(x) dm

\vee (\nabla_i \nabla_j f) = \bigotimes \nabla E^{+}

g(x,y) = \mathcal{0}(x)[f(x) + \bar{h}(x)] + T^2 d^2 \phi

\mathcal{0}(x) = \left( \int [g(x)] e^{-f} dV \right)^{\prime} - \sum \Delta (x)
\mathcal{0}(x) = [\nabla_i \nabla_j f(x)]^{\prime} \cong {}_{-n}C_{-r} f(x)^n
f(y)^{n-r} \Delta (x,y),
V(\tau) = \int [f(x)] dm / \partial f_{xy}

\square \psi = 8 \pi G T^{\mu\nu}, (\square \psi)^{\prime} = \nabla_i \nabla_j
(\Delta (x) \circ G(x))^{\mu\nu}
\left(\frac{p}{c^3} \circ \frac{V}{S}\right), x^{\frac{1}{2}} + iy} = e^{x \log x}

\Delta (x) \phi = \{\vee [\nabla_i \nabla_j f \circ g(x)] \over
\exists (R + \Delta f)\}

{}_{-n}C_{-r} = {}_{\frac{1}{i}H[\psi]} C_{\hbar \psi} + {}_{[H, \psi]} C_{-n - r}
{}_{-n}C_{-r} = {}_{-n}C_{-n-r}

```

$$\int \int \frac{1}{x \log x^2} dx_m \text{ to } \mathcal{H}_0(x) = [\nabla_i \nabla_j f]' / \partial f_{xy}$$

$$\begin{aligned} \bigcup_{x=0}^{\infty} f(x) &= \nabla_i \nabla_j f(x) \oplus \sum f(x) \\ &= \bigoplus \nabla f(x) \\ \nabla_i \nabla_j f &\cong \partial x \partial y \int \nabla_i \nabla_j f \, dm \\ &\quad \cong \int [f(x)] \, dm \\ &\quad \cong \{[f(x), g(x)], g^{-1}(x)\} \\ &\quad \cong \square \psi \\ &\quad \cong \nabla \psi^2 \\ &\quad \cong f(x \circ y) \leq f(x) \circ g(x) \\ &\quad \cong |f(x)| + |g(x)| \end{aligned}$$

$$\begin{aligned} \Delta(x) \psi &= \langle f, g \rangle \circ |h^{-1}(x)| \\ \partial f_x \cdot \Delta(x) \psi &= x \\ x \in \mathcal{H}_0(x) \\ \mathcal{H}_0(x) &= \{[f \circ g, h^{-1}(x)], g(x)\} \end{aligned}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f &= [\nabla \int \nabla_i \nabla_j f(x) \, dx_m, g^{-1}(x)] \text{ to } \bigoplus_{k=0}^{\infty} \\ \nabla E^{+}_{-} &= M_3 \\ &= \bigoplus_{k=0}^{\infty} E^{+}_{-} \\ dx^2 &= [g^2_{\mu\nu}, dx], \quad g^{-1} = dx \int \Delta(x) f(x) \, dx \\ f(x) &= \mathrm{exp}[\nabla_i \nabla_j f(x), g^{-1}(x)] \\ \pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \left(\frac{g(x)}{f(x)} \right)' &= \\ \lim_{n \rightarrow \infty} \left\{ \frac{g(x)}{f(x)} \right\} &= \{g'(x) \over f'(x)\} \end{aligned}$$

$$\begin{aligned} \nabla F &= f \cdot \frac{1}{4} |r|^2 \\ \nabla_i \nabla_j f &= \frac{d}{dx_i} \\ \frac{d}{dx_j} f(x) g(x) \\ D^2 \psi &= \nabla \int (\nabla_i \nabla_j f)^2 \, d\eta \\ E &= m c^2, \quad E = \frac{1}{2} m v^2 - \frac{1}{2} k x^2, \quad G^{\mu\nu} = \\ &\quad \frac{1}{2} \Lambda g_{ij}, \\ \square &= \frac{1}{2} k T^2 \end{aligned}$$

$$\mathrm{ker} \, f / \mathrm{im} \, f \cong S^{\mu\nu}_m, \quad S^{\mu\nu}_m = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$D^2 \psi = \mathcal{H}_0(x) \left(\frac{p}{c^3} + \frac{V}{S} \right), \quad V(x) = D^2 \psi \otimes M^+_3$$

$$\begin{aligned} S^{\mu\nu}_m \otimes S^{\mu\nu}_n &= \\ - \frac{2R_{ij}}{V(\tau)} [D^2 \psi] \end{aligned}$$

$$\nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] = \int \{V(\tau) \over f(x)\} [D^2 \psi]$$

$$\nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] = \int \{V(\tau) \over f(x)\} \mathcal{O}(x)$$

$$z(x) = \{g(cx + d) \over f(ax + b)\} h(ex + 1)$$

$$= \int \{V(\tau) \over f(x)\} \mathcal{O}(x)$$

$$\{V(x) \over f(x)\} = m(x), \mathcal{O}(x) = m(x) [D^2 \psi(x)]$$

$$\{d \over df\} F = m(x), \int F \, dx_m = \sum_{k=0}^{\infty} m(x)$$

$$\mathcal{O}(x) = \left([\nabla_i \nabla_j f(x)] \right)'$$

$$\cong \{ {}_nC_r(x)^n (y)^{n-r} \delta(x,y)$$

$$(\square \psi)' = \nabla_i \nabla_j (\delta(x) \circ$$

$$G(x))^{\mu\nu} \left(p \over c^3 \right) \circ$$

$$\{V \over S\}$$

$$F^m_t = \{1 \over 4\} g^2_{ij}, \, x^{\{1 \over 2\} + iy} = e^{\{x \log x\}}$$

$$S^{\mu\nu}_m \otimes S^{\mu\nu}_n = G^{\mu\nu} \times T^{\mu\nu}$$

$$S^{\mu\nu}_m \otimes S^{\mu\nu}_n = -2 R_{ij} \over V(\tau) [D^2 \psi]$$

$$S^{\mu\nu}_m = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$\pi(\chi, x) = \int \mathrm{exp}[L(p,q)] d\psi$$

$$ds^2 = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu\nu} dx^{\mu} dx^{\nu} + T^2 d^2\psi]$$

$$M_3 \bigotimes_{k=0}^{\infty} E^{+}_{-} = \mathrm{rot}$$

$$(\mathrm{div} \, E, E_1)$$

$$= m(x), \{P^{2n} \over M_3\} = H_3(M_1)$$

$$\exists [R + |\nabla f|^2]^{\{1 \over 2\} + iy}$$

$$= \int \mathrm{exp}[L(p,q)] d\psi$$

$$= \exists [R + |\nabla f|^2]^{\{1 \over 2\} + iy} \otimes$$

$$\int \mathrm{exp}[L(p,q)] d\psi +$$

$$N \bmod (e^{\{x \log x\}})$$

$$= \mathcal{O}(\psi)$$

$$\{d \over dt\} g_{ij}(t) = -2 R_{ij}, \{P^{2n} \over M_3\}$$

$$= H_3(M_1), H_3(M_1) = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$S^{\mu\nu}_m \times S^{\mu\nu}_n$$

$$= [D^2 \psi], S^{\mu\nu}_m \times S^{\mu\nu}_n$$

$$= \mathrm{ker} f / \mathrm{im} f, S^{\mu\nu}_m \otimes$$

$$S^{\mu\nu}_n = m(x) [D^2 \psi], \{-2R_{ij} \over V(\tau)\} = f^{-1} x f(x)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \circ$$

$$\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix} \right] dx dy dz,$$

$$\to f_z^{\{1 \over 2\}} \to (0,1) \cdot (0,1) = -1, i =$$

$$\sqrt{-1}$$

$$\begin{pmatrix} x,y,z \\ \end{pmatrix}^2 = (x,y,z) \cdot (x,y,z) \to -1$$

$$\mathcal{O}(x) = \nabla_i \nabla_j \int e^{\frac{2}{m} \sin \theta} \cos \theta \, d\theta \, N \bmod$$

$$(e^{x \log x})$$

$$\overline{\mathcal{O}(x)}(x + \Delta |f|^2)^{\frac{1}{2}}$$

$$x \, \Gamma(x) = 2 \int |\sin^2 \theta|^{2d} d\theta,$$

$$\mathcal{O}(x) = m(x) [D^2 \psi]$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix}$$

$$\begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix}$$

$$\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$f^{-1}(x) \times f(x) = I^{'}_m, I^{'}_m = [1,0] \times [0,1]$$

$$i^2 = (0,1) \cdot (0,1), |a||b| \cos \theta = -1,$$

$$E = \mathrm{div}(E, E_1)$$

$$\left(\frac{f,g}{[f,g]} \right)^{' } = i^2, \quad E = mc^2, \quad I^{' } = i^2$$

$$\mathcal{O}(x) = || \nabla \int [\nabla_i \nabla_j f$$

$$\circ g(x)]^{\frac{1}{2} + i y} ||, \partial r^n$$

$$|| \nabla ||^2 \rightarrow \nabla_i \nabla_j || \vec{v} ||^2$$

$$\nabla^2 \phi$$

$$\nabla^2 \phi = 8 \pi G \left(\rho \over c^3 + \{V \over S\} \right)$$

$$(\log x^{\frac{1}{2}})^{' } = \frac{1}{2} \frac{1}{(x \log x)},$$

$$(\sin \theta)^{' } = \cos \theta, \quad (f_z)^{' } = i e^{i x \log x},$$

$$\frac{d}{df} F = m(x)$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m$$

$$+ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$= \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \right.$$

$$+ \left. \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm$$

$$\geq \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \circ (y \log y)^{\frac{1}{2}} \right) dm$$

$$\geq 2h$$

$$\frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \circ (y \log y)^{\frac{1}{2}} \right) dm \geq \hbar$$

$$y = x, \, xy = x^2, \, (\square \psi)^{' } = 8 \pi G$$

$$\left(\rho \over c^3 \right) \circ \{V \over S\}$$

$$\square \psi = \int \int \mathrm{exp}[8 \pi G (\bar{h}_{\mu \nu}$$

$$\circ \eta_{\mu \nu})^{\mu \nu}] dm d\psi,$$

$$\sum a_k x^k = \{d \over df\} \sum \sum \{1 \over a_k^2 f^k\} dx_k$$

$$\sum a_k f^k = \{d \over df\} \sum \sum$$

$$\{\zeta(s) \over a_k\} dx_{\{k\}},$$

$$a_k^2 f^{\{1 \over 2\}} \rightarrow \lim_{\{k \rightarrow 1\}} a_k f^k = \alpha$$

$$ds^2 = [g_{\{\mu \nu\}}^2, dx]$$

$$M_2$$

$$ds^2 = g_{\{\mu \nu\}}^{-1} (g^2_{\{\mu \nu\}}(x) - dx g_{\{\mu \nu\}}^2)$$

$$M_2$$

$$= h(x) \otimes g_{\{\mu \nu\}} d^2 x - h(x) \otimes dx g_{\{\mu \nu\}}(x),$$

$$h(x) = (f^2(\vec{x}) - \vec{E}^{\{+\}})$$

$$G_{\{\mu \nu\}} = R_{\{\mu \nu\}} T^{\{\mu \nu\}},$$

$$\partial M_2 = \bigoplus \nabla C^{\{+\}_{-}}$$

$$G_{\{\mu \nu\}} \text{ equal } R_{\{\mu \nu\}} \{d \over dt\} g_{\{ij\}} = -2 R_{\{ij\}}$$

$$r = 2 f^{\{1 \over 2\}}(x)$$

$$E^{\{+\}} = f^{\{-1\}} x f(x),$$

$$h(x) \otimes g(\vec{x}) \cong \{V \over S\},$$

$$\{R \over M_2\} = E^{\{+\}} - \{\phi\}$$

$$= M_3 \supset R,$$

$$M^{\{+\}_2} = E^{\{+\}_1} \cup E^{\{+\}_2} \rightarrow E^{\{+\}_1} \bigoplus E^{\{+\}_2}$$

$$= M_1 \bigoplus \nabla C^{\{+\}_{-}}, (E^{\{+\}_1} \bigoplus E^{\{+\}_2})$$

$$\cdot (R^{\{-\}} \subset C^{\{+\}})$$

$$\{R \over M_2\} = E^{\{+\}} - \{\phi\}$$

$$= M_3 \supset R$$

$$M^{\{+\}_3} \cong h(x) \cdot R^{\{+\}_3}$$

$$= \bigoplus \nabla C^{\{+\}_{-}},$$

$$R = E^{\{+\}} \bigoplus M_2 - (E^{\{+\}} \cap M_2)$$

$$E^{\{+\}} = g_{\{\mu \nu\}} dx g_{\{\mu \nu\}},$$

$$M_2 = g_{\{\mu \nu\}} d^2 x,$$

$$F = \rho g \downarrow \rightarrow \{V \over S\}$$

$$\mathcal{H}_0(x) = \delta(x) [f(x) + g(\bar{x})] + \rho g \downarrow,$$

$$F = \{1 \over 2\} m v^2 - \{1 \over 2\} k x^2,$$

$$M_2 = P^{\{2n\}}$$

$$r = 2 f^{\{1 \over 2\}}(x),$$

$$f(x) = \{1 \over 4\} |r|^2$$

$$V = R^{\{+\}} \sum K_m, W = C^{\{+\}} \sum^{\{\infty\}}_{\{k=0\}} K_{\{n+2\}},$$

$$V/W = R^{\{+\}} \sum K_m / C^{\{+\}} \sum K_{\{n+2\}}$$

$$= R^{\{+\}} / C^{\{+\}} \sum \{x^k \over a_k f^k(x)\}$$

$$= M^{\{+\}_{-}}, \{d \over df\} F = m(x), \rightarrow M^{\{+\}_{-}}, \sum^{\{\infty\}}_{\{k=0\}}$$

$$\{x^k \over a_k f^k(x)\} = \{a_k x^k \over$$

$$\zeta(x)\}$$

$$\{\{f,g\} \over [f,g]\} = \{fg + gf \over fg - gf\},$$

$$\nabla f = 2, \partial H_3 = 2, \{1 + f \over 1 - f\} = 1,$$

$$\{d \over df\} F = \bigoplus \nabla C^{\{+\}_{-}}, \vec{F} =$$

$$\{1 \over 2\}$$

$$H_1 \cong H_3 = M_3$$

$$H_3 \cong H_1 \rightarrow \pi(\chi, x), H_n, H_m =$$

$$\mathrm{rank}(m,n), \mathrm{mesh}(\mathrm{rank}(m,n)) \lim \mathrm{mesh} \rightarrow 0$$

$$(fg)' = fg' + gf', (\{f \over g\})' = \{f'g - g'f \over g^2\},$$

$$\{\{f,g\} \over [f,g]\} = \{(fg)' \otimes dx_{\{fg\}} \over$$

$$\begin{aligned} & \left(\frac{f}{g} \right)' \otimes g^{-2} dx_{fg} \\ &= \frac{\{(fg)' \otimes dx_{fg}\}}{\{(f/g)' \otimes g^{-2} dx_{fg}\}} \\ &= \{d \over df\} F \end{aligned}$$

$$\hbar \psi = \{1 \over i\} H \psi, \quad i[H, \psi] = -H \psi, \quad \{f, g\} \over [f, g] = (i)^2$$

$$\begin{aligned} & [\nabla_i \nabla_j f(x), \delta(x)] = \nabla_i \nabla_j \\ & \int f(x,y) dm_{xy}, \quad f(x,y) = [f(x), h(x)] \times [g(x), h^{-1}(x)] \\ & \delta(x) = \{1 \over f'(x)\}, \quad [H, \psi] = \Delta f(x), \\ & \mathcal{O}(x) = \nabla_i \nabla_j \int \delta(x) f(x) dx \\ & \mathcal{O}(x) = \int \delta(x) f(x) dx \end{aligned}$$

$$\begin{aligned} & R^+ \cap E^+_{-} \ni x, \quad M \times R^+ \ni M_3, \quad Q \supset C^+_{-}, \\ & Z \in Q \nabla f, \quad f \cong \bigoplus_{k=0}^n \nabla C^+_{-} \\ & \bigoplus_{k=0}^{\infty} \nabla C^+_{-} = M_1, \quad \bigoplus_{k=0}^{\infty} \\ & \nabla M^+_{-} \cong E^+_{-}, \\ & M_3 \cong M_1 \bigoplus_{k=0}^{\infty} \nabla \{V^+_{-} \over S\} \\ & \{P^{2n} \over M_2\} \cong \bigoplus_{k=0}^{\infty} \\ & \nabla C^+_{-}, \quad E^+_{-} \times R^+_{-} \cong M_2 \\ & \zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k, \\ & M_2 \cong P^{2n} / \ker f, \quad \to \bigoplus \nabla C^+_{-} \\ & S^+_{-} \times V^+_{-} \cong \{V \over S\} \bigoplus_{k=0}^{\infty} \\ & \nabla C^+_{-}, \quad V^+ \cong M^+_{-} \bigotimes S^+_{-}, \\ & Q \times M_1 \subset \bigoplus \nabla C^+_{-} \\ & \sum_{k=0}^{\infty} Z \otimes Q^+_{-} = \bigotimes_{k=0}^{\infty} \nabla M_1 \\ & = \bigotimes_{k=0}^{\infty} \nabla C^+_{-} \times \\ & \sum_{k=0}^{\infty} M_1, \quad x \in R^+ \times C^+_{-} \\ & \supset M_1, \quad M_1 \subset M_2 \subset M_3 \end{aligned}$$

$$\begin{aligned} & S^3, \quad H^1 \times E^1, \quad E^1, \quad S^1 \times E^1, \quad S^2 \times E^1, \quad H^1 \times S^1, \\ & H^1, \quad S^2 \times E. \\ & \bigoplus \nabla C^+_{-} \cong M_3, \quad R \supset Q, \quad R \cap Q, \\ & R \subset M_3, \quad C^+ \bigoplus M_n, \quad E^+ \cap R^+ \$ \\ & M^+_{-} \nabla C^+_{-}, \quad C^+ \nabla H_m, \quad E^+ \nabla R^+_{-}, \quad E_2 \nabla E_1 \$ \\ & \$ R^+ \nabla C^+_{-} \$, \quad \{ \nabla \over \Delta \} \int x f(x) dx, \\ & \{ \nabla R \over \Delta f \}, \quad \square = 2 \{ \int \{ (R + \nabla_i \nabla_j f)^2 \\ & \over -(R + \Delta f) \} e^{-f} dV \\ & \square = \{ \nabla R \over \Delta f \}, \quad \{ d \over dt \} g_{ij} \\ & = \square \to \{ \nabla f \over \Delta x \}, \quad (R + \\ & | \nabla f|^2) dm \to -2(R + \nabla_i \nabla_j f)^2 e^{-f} dV \\ & x^n + y^n = z^n \to \nabla \psi^2 = 8 \pi G T^{\mu \nu}, \\ & f(x+y) \geq f(x) \circ f(y) \\ & \mathrm{im} f / \mathrm{ker} f = \partial f, \quad \mathrm{ker} f \\ & = \partial f, \quad \mathrm{ker} f / \mathrm{im} f \cong \\ & \partial f, \quad \mathrm{ker} f = f^{-1}(x) x f(x) \\ & f^{-1}(x) x f(x) = \int \partial f(x) d(\mathrm{ker} f) \to \nabla f = 2 \\ & _nC_r = _nC_{n-r} \to \mathrm{im} f / \mathrm{ker} f \\ & \cong \mathrm{ker} f / \mathrm{im} f \end{aligned}$$

$$\$ \sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi \$. \text{ this equation } \$ a_k \cong$$


```

\sum^{\infty}_{r=0} {}_nC_r $.
V/W = R/C \sum^{\infty}_{k=0} {x^k \over a_k f^k}, W/V = C/R
\sum^{\infty}_{k=0} {a_k f^k \over x^k}
V/W \cong W/V \cong R/C(\sum^{\infty}_{r=0} {}_nC_r)^{-1}
\sum^{\infty}_{k=0} x^k
This equation is differential equation, then $ \sum^{\infty}_{k=0} a_k f^k $
is included with $ a_k \cong \sum^{\infty}_{r=0} {}_nC_r $
W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{1-t} dx,
\sum^{n}_{k=0} a_k f^k = (f^k)'
\sum^{n}_{k=0} a_k f^k = \sum^{\infty}_{k=0} {}_nC_r f^k
= (f^k)',
\sum^{\infty}_{k=0} a_k f^k = [f(x)],
\sum^{\infty}_{k=0} a_k f^k = \alpha, \sum^{\infty}_{k=0}
{1 \over a_k f^k}, \sum^{\infty}_{k=0} (a_k f^k)^{-1} = {1 \over 1 - z}

{\int \int {1 \over (x \log x)(y \log y)} dx dy} =
{{}_nC_r xy \over ({}_nC_{n-r})}
(x \log x)(y \log y))^{-1}}
= ({}_nC_{n-r})^2 \sum_{k=0}^{\infty} ({1 \over x \log x}
- {1 \over y \log y}) d{1 \over nxy} \times {xy}
= \sum_{k=0}^{\infty} a_k f^k
= \alpha
}

```

```

_ struct_ :asperal equation.emerged => [tuplespace]
tuplespace.cognitive_system => development -> Omega.Database[import]
value.equation_emerged.exclude >- Omega.Database[tuplespace]

```

```

Omega::DataBase <-> virtual_connect(VIRTUALMACHINE)
{
  blidge_base.network => localmachine.attachment
  :=> {
    dhcp.etc_load_file(this.klass) {|list|
      list.connect[XWin.display _ <- xhost.in(regexpt.pattern)]
      {
        ultranetwork.def _struct {
          asperal_language :this.network_address.included[type.system_pattern]
          {|regexpt.pattern|
            <- w.scan
              |each_string| <= { ipv4.file :file.port
                                subnetmask :file.address
                                file.port <=> file.address
                                FILE *pointer
                                int,char,str :emerge.exclude > array[]
                                BTE.each_string <-> regexpt.pattern
                                {
                                  development => file.to_excluded
                                  file.scan => regexpt.pattern
                                  this.iterator <-> each_string
                                  file.reloaded => [asperal_language.rebuild]
                                }
                                }
          }
        }
      }
    }
  }
}

```



```

{
  etc.include(inetd.rc)
  {
    virtual_connect(VIRTUAL_MACHINE){|list|
      list.attachment(etc.load_file)
    }
  }
}
end

mainloop{
  def.virtual_connect => xhost.localmachine
  {
    xhost.client <-> xhost.server
  }
  def.network.type <- [Omega.DATABASE] end
  def.etc.load_file.attachment(VIRTUAL_MACHINE) end
end
end

class UltraNetwork::DATABASE import OMEGA.TUPLESPEACE
  def load_file >- VIRTUAL_MACHINE
    { in . => attachment_device |for|
      for.load -> acceptance.hardware
      virtual_machine.new
      {
        tk.loop-> start
        XWin -multiwindow
        if dwm <-> new_xwin.start
          localhost :xhost :display -x
          xdisplay :-> [preset :XFree.demand]>=needed
          for.set_up
          install_process >- tar -xvfz "#{load_file}" <-> install_attachment
        ]
        else if
          only :new_xwin.start
          localhost :xhost :multiwindow . { in
            display -x
            attachment :localhost -client
            from -client into
            server.XWin -attachment}
          end
          condition :{ in .=>
            check->[xdisplay.install_process]}
        end
      }
    end
  end

  def < network_rout
    wireshark.start -> ethernet.device >- define rout
    rout.ipstate do |file|
      file.type <- encoding XWin -filesystem
      file.included >- make kernel_system.rebuild
      file.vmware.start do |rout|

```

```

        rout.blidgebase | rout.hostbase
    -> file.install
        file.address_ipstate
    => {"{file}" :=> dwm.state_presense
        virtual_machine.included[file]
    }
end

def < launcher_application
    network_rout.new
    |file|
    file.attachment => { in .
    new_xwin.start :=> file.included
    demand.file <- success_exit}
end

def < terminal_port
    network_rout.new
    launcher_application.new |rout|
    rout.acceptance {
    vmware.state.process |new_rout|
    new_rout : attachment.class <-> dwm.state_attachment
    new_rout -> condition.start_wmware.process}
end

def < kterm_port
    launcher_application.new
    def.included[DATABASE]
    |rout|
    rout.attachment <- |new_rout|
    new_rout.attachment do
    install.condition < rout.def.terminal_port.exclude[file]
    end
end

main_loop :file do
    kterm_port.excluded :=> VIRTUAL_MACHINE
    |new_rout| start do
    rout.process -> network_rout.rout [
    file,launcher_application, terminal_port, kterm_port].def < included
    |file|
    file.all_attachment: file_type :=> encoding-utf8
    end
end

class < def {
    pholograph_data[] = [R,V,S,E,U,M_n,Z_n,Q,C,N,f,g]
    source_array <- pholograph_data[]
}

def > operator_data[] = {nabla,nabla_i nabla_j,Delta,partial,

```

```

d, int, cap,cup,ni,in,chi,oplus,otimes,bigoplus,bigotimes,d /over df,
dV,dm,dx,dy,<,>,[,],[,],[,],[,] }

end

def > manifold_emerge

    c = def.inject >- source_array times def.operator_data[]
    repository_data <=> c{

    c.scan(/tuplespace[/])
    import |list| list{
        kerf = -2 \int (R + nabla_i nabla_j f)^2e^{-f}dV
        kerf / imf
        =< {d \over df}F}
    }

    equals_data =~ /list/
    list.match("/#{c}"/) {|list|
        list.delete
        jisyo_data_mathmatics <=> list{
        list.emerge => {asperal function >- photograph_data[] times repository_data
            =< list.update}
        }

        ln -s operator_named <= {list}
        define _struct |list|
            -> list.element -> manifold_emerge
            => list.reconstruct > def.inject /~"#{pattern}"/}

    end

import Omega::Tuplespace < Database
{
    {\bigoplus \nabla M^{+}_{-}}.equation_create -> asperal :variable[array]
    :=> [cognitive_system <-> def < VIRTUALMACHINE.terminal
        {
            [ipv4.bloodcast.address :
                ipv4.network.address].subnetmask
            <-> file.port.transport_import :
                Omega[tuplespace]
        }
    }

    _struct _ Omega[tuplespace] >> VIRTUALMACHINE.terminal.value

class < def.VIRTUALMACHINE.system_environment

    file.reload[hardware] => file.exclude >> file.attachment
    {=>
        |file|
        file.port(wireshark.rout <-> {file.port.transport_export
            :=> Omega[tuplespace]})
    }

```

```

assembly_process.file.included >- file.reloaded
:- |file.environment| {=>
    file.type? :=> exist
    file.regexpt.pattern[scan.flex]
    => |pattern|
    <->
    file.[scan.compiler]
}
end
end
file <<
}
}

Omega::Database[tuplespace]
{
  cognitive_system |: -> { DATABASE.create.regexpt_pattern >-
    cognitive_system[tuplespace].recreated >- : =< DATABASE.value
    >> system_require.application.reloaded[tuplespace]
    } : _struct _ def.VIRTUALMACHINE.terminal >> {
      ||machine.attachment|| <-> OBJECT.shift => system.reloaded
      . in {
        : _struct _ class.import :-> require mechanics.DATABASE
        {|regexpt_pattern| :|-> aspective _union _
        def _union _}
      }
    }
  }

  end
}

system.require <- import library.DATABASE
{
  Omega[tuplespace]
  {
    cognitive_system : VIRTUALMACHINE.equality_realized
    {|regexpt_pattern| => value | key [ > cognitive_system.loop.stdout]
    value : display -bash :xhost -number XWin.terminal
    key : registry.edit :=> {[cognitive_system.reloaded]}
  }
}

_uni _ => DATABASE[tuplespace].aspective_reloaded
_uni _ :fx | -> |regexpt_pattern| => {
  VIRTUALMACHIE.recreated-> _uni _ |
  _struct _ def.DATABASE.recreated <- fx
  >> DATABASE[tuplespace].rebuild
}

DATABASE[tuplespace] -< {[ > aimed.compiler | aimed.interpreter] | btree.def.distributed >-

```

```

        aimed[tuplespace]}
aimed[tuplespace] -< btree.class.hyperroot_ struct _ => Omega::Database[tuplespace].value
  sheap_ union _ :aspective | -> Omega[tuplespace]: | aimed[tuplespace].differentiated_review
}

aimed[tuplespace].process => DATABASE[tuplespace].reloaded
aimed.different | aimed.stdout >> vale | key [ > cognitive_system.loop.stdin] {|pattern|
  pattern.scan(value : aimed[def.value]
    key : aimed[def.key])
} _ struct _ : flex | interpreter.system
=> expression.iterator[def.first,def.second,def.third,def.fourth]
{ def < Omega[tuplespace]
  def.cognitive_system |: -> DATABASE[tuplespace] | aimed[tuplespace]
}

Omega::Tuplespace < DATABASE
{=>
  norm[Fx] -> . in for def.all_included < aimed[tuplespace].each_scan([regextp_pattern]
    <->
      DATABASE[tuplespace]) << stream database.excluded
  >- more_pattern.scan(value : aimed[def.value]

  key : aimed[def.key])
    . in { _struct _ : flex | interpreter.system
      => expression.iterator[def.all.each -> |value, key|
        included >- norm[Fx] | [DATABASE[tuplespace]
, aimed[tuplespace]] |
        finality : aimed[tuplespace], DATABASE[tuplespace]

: -> def.included(in_all)
    {
      def.key | def,value => [DATABASE].recompile
    & make install
      : in_all -> _struct _ :aspective :tuplespace
    : all_homology_created}
    }
}

def < Omega::Tuplespace[DATABASE]
def.iterator -> |klass,define_method,constant,variable,infinity_data : -> finite_data|
  def.each_klass?{|value, key|
    _struct _ :aspective -> tuplespace :all_homology_recreated :make menuconfig
    {+=
      def.key -> aimed[def.key],def.value -> aimed[def.value] {|list|
        list.developed => <key,value> | <aimed[$', $']
        -> _union _ :value,key : _struct _
        <- (_union _ <-> _struct _ +)
      begin
        def.key <-> aimed[value]
        case :one_ exist :other :bug
        {
          result <-> def.key
          {

```

```

        differentiated :DATABASE[tuplespace]
    }
    return :tuplespace.value.shift -> included<tuplespace>
else if
:other :bug
{
    success_exit <- bug[value]
    {
        cognitive_system.scan(bug[value])
        {
            {[e^{-f}][{2 \int (R + \nabla f^2) \over -(R + \Delta f)}e^{-f}dV}
.created_field
            {=>
                regxpt.pattern \native_function <-> euler-equation
                {
                    $variable =< diff e^{-f} >- $'
                    all_included <- def.key <-> aimed[value]
                    $variable - all_included.diff
                    \summate_manifold.recreated
<- \native_function : euler-equation
                }
            }
        }
    } _union _ :cognitive_system.rebuild(one_ exist)
}
}
ensure
{
    return :success_exit
=> Tuplespace[DATABASE]
}
}
}
end
end
}

int
streem_style {
    :Endire <- [ADD,EVEN,MOD,DEL,MIX,INCLUDED,EXCLUDED,EBN,EXN,EOR,EXOR,
SUM,INT,DIFF,PARTIAL,ROUND,HOMOLOGY,MESH]
    Endire.iterator -> {def < :Endire.element, -> def.means_each{x -> expression.define.included
def.each{x -> case :x.each => :lex.include_ . in [ > [x.all_expire] ]}
}
}

main_loop {
    FILE *fp :=> streem_style.address_objective_space
    fp.each{x -> domain_specific_language_style_included[array]}
    array << streem.DATABASE[tuplespace]
    array.each{[tuplespace] -> aimed[tuplespace] | OMEGA_DATABASE[tuplespace]}.excluded <-> array
    def.key <-> def.value => {x -> stdin | stdout | => streem_style <- def.each.klass.value}
}

```



```

@reviser : def < OmegaDatabase[tuplespace].mechanism
{
  aspective : _union _ {
    int streem_style : [ > [def.each{x -> stdin | stdout > display :xhost in XWin -multiwind
    {
      Endire <- [ADD,EVEN,ODE,EXOR,XOR,DEL,DIFF,PARTIAL,INT].included > struct _ :-> _union
      Endire.each{def.value -> def.key :hash.define}.included > _union}
    }
  }
}

@reviser : def.reconstructed.each{_union <-> _struct _.recreated : [def.del - def.before_determin

import perl.lib | python.lib <-> ruby.lib
{
  int @reviser : def.each{x -> x.klass |-> $variable in $stdin | $stdout}.developed >= {
    ping localhost -> blidgebase <-> hostbase.virtualmachine.attachment
    {
      xhost :display -> streem_style.value
      networkconnect.hostbase -> localarea.virtualmachine
    } :connected -> networkrout : flow_to :localhost.attachment
  }
}_struct : def < hostbase.virtualmachine.attachment => : networkrout.area.build

@reviser <-> def.add [ < _struct]
@reviser : def.each{listmenu -> listlink | unlinklist > [developed -> {def.key , def.value}.current
@reviser <-> def.rebuild [ < _struct]

@reviser.def.<value|key>networkrout-> def.present
def.present.flow_to -> hostbase.rout << networkrout.data.<value|key>

XWin -multiwindow <-> networkrout.data[$',$']
def < $'
@reviser <-> def.present.state
@reviser.def.each{x | -> key.rebuild | value.rebuild}.flow_to :redefined

```

```

def < OmegaDatabase[tuplespace]
{
  FILE *fp -> cmd.value : cmd.key {fp |-> synchronized.file[tuplespace] | aimed.file[tuplespace]

```

```

    cmd.key => [ > fp.('$:$')] <-> registry.excluded<fp.file[cmd.state]>
}

def.each{fp|-> def.first,def.second,def.third,def.fourth}

cmd _struct : {
[ ^C-O : ^C-X-F, exit.cmd : ^C-X-C, shift-up : ^C-P, shift-down : ^C-N]
}

cmd _union : def.restruacted
keyhook.cmd <- : [_struct ]
{
  @reviser :def._struct <-> def. _union
}

```

Deconstruct Dimension of category theorem

Masaaki Yamaguchi

1 Locality equation

Quantum of material equation include with heat entropy of quantum effective theorem, geometry of rotate with imaginary pole is that generate structure theorem. Subgroup in category theorem is that deconstruct dimension of differential structure theorem. Quantum of material equation construct with global element of integrate equation. Heat equation built with this integrate geometry equation, Weil's theorem and fermion theorem also heat equation of quantum theorem moreover is all of equation rout with global differential equation this locality equation emerge from being built with zeta function, general relativity and quantum theorem also be emerged from this equation. Artificial intelligent theorem exclude with this add group theorem from, Locality equation inspect with this involved of theorem, This geometry of differential structure that deconstruct dimension of category theorem stay in four of universe of pieces. In reason cause with heat equation of quantum effective from artificial intelligence, locality equation conclude with this geometry theorem. Heat effective theorem emelite in generative theorem and quantum theorem, These theorem also emelitive with Global differential equation from locality equation. Imaginary equation of add group theorem memolite with Euler constance of variable from zeta function, this function also emerge with locality theorem in artificial intelligent theorem.

Quantum equation concern with infinity of energy composed for material equation, atom of verisity in element is most of territory involved with restored safety of orbital.

$$f = n\nu\lambda, \lambda = \frac{x}{l}, \int dn\nu\lambda = f(x), xf(x) = F(x), [f(x)] = \nu h$$

$$px = mv, \lambda(x) = \text{esperial}f(x), [F(x)] = \frac{1}{1-z}, 0 < \tan \theta < i, -1 < \tan \theta < i, -1 \leq \sin \theta \leq 1, -1 \leq \cos \theta \leq 1$$

Category theorem conclude with laplace function operator, distrust of manifold and connected space of eight of differential structure.

$$R\nabla E^+ = f(x)\nabla e^{x \log x}$$

$$Q\nabla C^+ = \frac{d}{df}F(x)\nabla \int \delta(x)f(x)dx$$

$$E^+\nabla f = e^{x \log x}\nabla n!f(x)/E(X)$$

Universe component construct with reality of number and three manifold of subgroup differential variable. Maxwell theorem is this Seifert structure $(u + v + w)(x + y + z)/\Gamma$ This space of magnetic diverse is mebius member cognetic theorem. These equation is fundermental group resulted.

$$\begin{aligned}\exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ &= \pi(R, C\nabla E^+) \\ &= \text{rot}(E_1, \text{div} E_2) \\ xf(x) &= F(x)\end{aligned}$$

$$\begin{aligned}\square x &= \int \frac{f(x)}{\nabla(R^+ \cap E^+)} d\square x \\ &= \int \frac{\Delta f(x) \circ E^+}{\nabla(R^+ \cap E^+)} \square x\end{aligned}$$

This equation means that quantum of potensial energy in material equation. And this also means is gravity of equation in this universe and the other dimension belong in. Moreover is zeta function for rich flow equation in category theorem built with material equation.

$$\begin{aligned}\exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ R\nabla E^+ &= f(x)\nabla e^{x \log x} \\ d(R\nabla E^+) &= \Delta f(x) \circ E^+(x) \\ \square x &= \int \frac{d(R\nabla E^+)}{\nabla(R^+ \cap E^+)} d\square x \\ \square x &= \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\square x \\ x^n + y^n = z^n &\rightarrow \square x = \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\square x\end{aligned}$$

Laplace equation exist of space in fifth dimension, three manifold belong in reality number of space. Other dimension is imaginary number of space, fifth dimension is finite of dimension. Prime number certify on reality of number, zero dimension is imaginary of space. Quantum group is this universe and the other dimension complex equation of non-certain theorem. Zero dimension is eternal space of being expanded to reality of atmosphere in space.

2 Heat entropy all of materials emerged by

$$\square = -2(T-t)|R_{ij} + \nabla\nabla f + \frac{1}{-2(T-t)}|g_{ij}^2$$

This heat equation have element of atom for condition of state in liquid, gas, solid attitude. And this condition constructed with seifert manifold in Weil's theorem. Therefore this attitude became with Higgs field emerged with.

$$\square = -2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV$$

$$\frac{d}{dt}g_{ij} = -2R_{ij}, \frac{d}{df}F = m(x)$$

$$R\nabla E^+ = \Delta f(x) \circ E^+(x), d(R\nabla E^+) = \nabla_i \nabla_j (R + E^+)$$

$$R\nabla E^+ = f(x)\nabla e^{x \log x}, R\nabla E^+ = f(x), f(x) = M_1$$

$$-2(T-t)|R_{ij} + \nabla\nabla f = \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\frac{1}{-2(T-t)}|g_{ij}^2 = \int \int \frac{1}{(x \log x)^2} dx_m$$

$$(\square + m^2)\phi = 0, (\gamma^n \partial_n + m^2)\psi = 0$$

$$\square = (\delta\phi + m^2)\psi, (\delta\phi + m^2c)\psi = 0, E = mc^2, E = -\frac{1}{2}mv^2 + mc^2, \square\psi^2 = (\partial\phi + m^2)\psi$$

$$\square\phi^2 = \frac{8\pi G}{c^4}T^{\mu\nu}, \nabla\phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S}), \frac{d}{dt}g_{ij} = -2R_{ij}, f(x) + g(x) \geq f(x) \circ g(x)$$

$$\int_{\beta}^{\alpha} a(x-1)(y-1) \geq 2 \int f(x)g(x)dx$$

$$m^2 = 2\pi T(\frac{26-D_n}{24}), r_n = \frac{1}{1-z}$$

$$(\partial\psi + m^2c)\phi = 0, T^{\mu\nu} = \frac{1}{2}mv^2 - \frac{1}{2}kx^2$$

$$\int [f(x)]dx = || \int f(x)dx ||, \int \nabla x dx = \Delta \sum f(x)dx, (e^{i\theta})' = ie^{i\theta}$$

This orbital surface varnished to energy from $E = mc^2$. $T^{\mu\nu} = nh\nu$ is $T^{\mu\nu} = \frac{1}{2}mv^2 - \frac{1}{2}kx^2 \geq mc^2 - \frac{1}{2}mv^2$ This equation means with light do not limit over $\frac{1}{2}kx^2$ energy.

$$Z \in R \cap Q, R \subset M_+, C^+ \bigoplus_{k=0}^n H_m, E^+ \cap R^+$$

$$M_+ = \sum_{k=0}^n C^+ \oplus H_M, M_+ = \sum_{k=0}^n C^+ \cup H_+$$

$$\begin{aligned}
& E_2 \bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+ \\
& M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R^+ \\
& E_1 \nabla E_2, R^- \nabla C^+, \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+, R \supset Q
\end{aligned}$$

$$\frac{d}{df} F = \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+$$

All function emerge from eight differential structure in laplace equation of being deconstructed that with connected of category theorem from.

$$\begin{aligned}
& \nabla \circ \Delta = \nabla \sum f(x) \\
& \Delta \rightarrow \text{mesh} f(x) dx, \partial x \\
& \nabla \rightarrow \partial_{xy}, \frac{d}{dx} f(x) \frac{d}{dx} g(x) \\
& \square x = -[f, g] \\
& \frac{d}{dt} g_{ij} = -2R_{ij} \\
& \lim_{x \rightarrow \infty} \sum f(x) dx = \int dx, \nabla \int dx
\end{aligned}$$

3 Universe in category theorem created by

Two dimension of surface in power vector of subgroup of complex manifold on reality under of group line. This group is duality of differential complex in add group. All integrate of complex sphere deconstruct of homomorphism is also exists of three dimension of manifold. Fundamental group in two dimension of surface is also reality of quality of complex manifold. Rich flow equation is also quality of surface in add group of complex. Group of ultra network in category theorem is subgroup of quality of differential equation and fundamental group also differential complex equation.

$$\begin{aligned}
& (E_2 \bigoplus_{k=0}^n E_1) \cdot (R^- \subset C^+) \\
& = \bigoplus_{k=0}^n \nabla C_-^+ \\
& \vee \int \frac{C_-^+ \nabla H_m}{\Delta(M_-^+ \nabla C_-^+)} = \wedge M_-^+ \bigoplus_{k=0}^n C_-^+ \\
& \exists(M_-^+ \nabla C_-^+) = \text{XOR}(\bigoplus_{k=0}^n \nabla M_-^+)
\end{aligned}$$

$$-[E^+\nabla R_-^+]=\nabla_+\nabla_-C_-^+$$

$$\int dx, \partial x, \nabla_i \nabla_j, \Delta x$$

$$\rightarrow E^+\nabla M_1, E^+\cap R\in M_1, R\nabla C^+$$

$$\begin{pmatrix} \cos x & \sin x \\ \sin x & -\cos x \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\sum_{k=0}^n \cos k\theta = \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \cos \frac{n\theta}{2}$$

$$\sum_{k=0}^n \sin k\theta = \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \sin \frac{n\theta}{2}$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}, \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \tan \theta = \frac{e^{i\theta} - e^{-i\theta}}{i(e^{i\theta} + e^{-i\theta})}$$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \rightarrow 1, \lim_{\theta \rightarrow 0} \frac{\cos \theta}{\theta} \rightarrow 1$$

$$(e^{i\theta})' = 2, \nabla f = 2, e^{i\theta} = \cos \theta + i \sin \theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \rightarrow [\cos^2 \theta + \sin \theta + \cos \theta - 2 \sin^2 \theta] \begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix}$$

$$2\sin\theta\cos\theta=2n\lambda\sin\theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = 1$$

Dimension of rotate with imaginary pole is generate of geometry structure theorem. Category theorem restruct with eight of differential structure that built with Thurston conjugate theorem, this element with summuate of thirty two structure. This pieces of structure element have with four of universe in one dimension. Quark of two pair is two dimension include, Twelve of quarks construct with super symmetry dimension theorem. Differential structure and quark of pair exclusive with four of universe.

Module conjecture built with symmetry theorem, this theorem belong with non-gravity in zero dimension. This dimension conbuilt with infinity element, mebius space this dimension construct with universe and the other dimension. Universe and the other dimension selves have with singularity, therefore is module conjecture selve with gravity element of one dimension. Symmetry theorem tell this one

dimension to emerge with quarks, this quarks element also restruct with eight differential structure. This reason resolve with reflected to dimension of partner is gravity and creature create with universe.

Mass exist from quark and dimension of element, this explain from being consisted of duality that sheaf of element in this tradition. Higgs field is emerged from this element of tradition, zeta function generate with this quality. Gravity is emerge with space of quality, also Higgs field is emerge to being generated of zero dimension. Higgs field has surround of mass to deduce of light accumulate in anserity. Zeta function consist from laplace equation and fifth dimension of element. This element include of gravity and antigravity, fourth dimension indicate with this quality emerge with being surrounded of mass. Zero dimension consist from future and past include of fourth dimension, eternal space is already space of first emerged with. Mass exist of quantum effective theorem also consisted of Higgs field and Higgs quark. That gravity deduce from antigravity of element is explain from this system of Higgs field of mechanism. Antigravity conclude for this mass of gravity element. What atom don't include of orbital is explain to this antigravity of element, this mechanism in Higgs field have with element of gravity and antigravity.

Quantum theorem describe with non-certain theorem in universe and other dimension of vector element, This material equation construct with lambda theorem and material theorem, other dimension existed resolve with non-certain theorem.

$$\sin \theta = \frac{\lambda}{d}, -py_1 \sin 90^\circ \leq \sin \theta \leq py_2 \sin 90^\circ, \lambda = \frac{h}{mv}$$

$$\frac{\lambda_2}{\lambda_1} = \frac{\sin \theta}{\frac{\sin \theta}{2}} h, \lambda' \geq 2h, \int \sin 2\theta = ||x - y||$$

4 Ultra Network from category theorem entirely create with database

Eight differential structure composite with mesh of manifold, each of structure firstly destructed with this mesh emerged, lastly connected with this each of structure deconstructed. High energy of entropy not able to firstly deconstructed, and low energy of firstly destructed. This reason low energy firsted. These result says, non-catastrophe be able to one geometry structure.

$$D^2\psi = \nabla \int (\nabla_i \nabla_j f)^2 d\eta$$

$$E = mc^2, E = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, G^{\mu\nu} = \frac{1}{2}\Lambda g_{ij}, \square = \frac{1}{2}kT^2$$

Sheaf of manifold construct with homomorphism in kernel divide into image function, this area of field rehearl with universe of surrounded with image function rehide in quality. This reason with explained the mechanism, Sheaf of manifold remain into surrounded of universe.

$$\ker f / \operatorname{im} f \cong S_m^{\mu\nu}, S_m^{\mu\nu} = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$D^2\psi = \mathcal{O}(x) \left(\frac{p}{c^3} + \frac{V}{S} \right), V(x) = D^2\psi \otimes M_3^+$$

$$S_m^{\mu\nu} \otimes S_n^{\mu\nu} = -\frac{2R_{ij}}{V(\tau)}[D^2\psi]$$

Selbarg conjecture construct with singularity theorem in Sheap element.

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} [D^2\psi]$$

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} \mathcal{O}(x)$$

$$\begin{aligned} z(x) &= \frac{g(cx+d)}{f(ax+b)} h(ex+l) \\ &= \int \frac{V(\tau)}{f(x)} \mathcal{O}(x) \end{aligned}$$

$$\frac{V(x)}{f(x)} = m(x), \mathcal{O}(x) = m(x)[D^2\psi(x)]$$

$$\frac{d}{df} F = m(x), \int F dx_m = \sum_{k=0}^{\infty} m(x)$$

5 Fifth dimension of seifert manifold in universe and other dimension

Universe audient for other dimension inclusive into fifth dimension, this phenomonen is universe being constant of value, and universe out of dimension exclusive space. All of value is constance of entropy. Universe entropy and other dimension entropy is dense of duality belong for. Prime equation construct with x,y parameter is each value of imaginary and reality of pole on a right comittment.

$$\int \nabla \psi^2 d\nabla \psi = \square \psi$$

$$\square \psi = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\delta \cdot \mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla g(x) dx_{ij}]^{iy}$$

$$||ds^2|| = e^{-2\pi T|\phi|} [\mathcal{O}(x) + \delta \mathcal{O}(x)] dx^\mu dx^\nu + \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}, \frac{(y \log y)^{\frac{1}{2}}}{\log(x \log x)} \leq \frac{1}{2}$$

Universe is freeze out constant, and other dimension is expanding into fifth dimension. Possibility of quato metric, $\delta(x)$ = reality of value / exist of value ≤ 1 , expanding of universe = exist of value $\rightarrow \log(x \log x) = \square \psi$

freeze out of universe = reality of value $\rightarrow (y \log y)^{\frac{1}{2}} = \nabla \psi^2$

Around of universe is other dimension in fifth dimension of seifert manifold. what we see sky of earth is other dimension sight of from.

Rotate of dimension is transport of dimension in other dimension from universe system.

Other dimension is more than Light speed into takion quarks of structure.

Around universe is blackhole, this reason resolved with Gamma ray burst in earth flow. We don't able to see now other dimension, gamma ray burst is blackhole phenomanen. Special relativity is light speed not over limit. then light speed more than this speed. Other dimesion is contrary on universe from. If light speed come with other light speed together, stop of speed in light verisity. Then takion quarks of speed is on the contrary from become not see light element. And this takion quarks is into fifth dimension of being go from universe. Also this quarks around of universe attachment state. Moreover also this quarks is belong in vector of element. This reason is universe in other dimension emerge with. In this other dimension out of universe, but these dimension not shared of space.

$$l(x) = 2x^2 + qx + r$$

$$= (ax + b)(cx + d) + r, e^{l(x)} = \frac{d}{df} L(x), G_{\mu\nu} = g(x) \wedge f(x)$$

This equation is Weil's theorem, also quantum group. Complex space.

$$V(\tau) = \int \exp[L(x)] dx_m + O(N^{-1}), g(x) = L(x) \circ \int \int e^{2x^2 + 2x + r} dx_m$$

Lens of space and integrate of rout equation. And norm space of algebra line of curve equation in gradient metric.

$$||ds^2|| = ||\frac{d}{df} L(x)||, \eta = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\bar{h} = [\nabla_i \nabla_j \int \nabla g(x) d_{ij}]^{iy}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

Two equation construct with Kaluze-Klein dimension of equation. Non-verticle space.

$$\frac{1}{\tau} (\frac{N}{2} + r(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

Also this too lense of space equation. And this also is heat equation. And this equation is singularity into space metric number of formula.

$$\frac{x^2}{a} \cos x + \frac{y^2}{b} \sin x = r^2$$

Curvature of equation.

$$S_m^2 = || \int \pi r^2 dr ||^2, V(\tau) = r^2 \times S_m^2, S_1^{mn} \otimes S_2^{mn} = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

Non-relativity of integrate rout equation.

$$||ds^2|| = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2 \psi$$

$$\begin{aligned}
V(x) &= \int \frac{1}{\sqrt{2\tau q}} (\exp L(x) dx) + O(N^{-1}) \\
V(x) &= 2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV, V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau q}} L(x) dx) + O(N^{-1}) \\
\frac{d}{df} F &= m(x) \\
Zeta(x, h) &= \exp \frac{(qf(x))^m}{m}
\end{aligned}$$

Singularity and duality of differential is complex element.

$$\begin{aligned}
& \left\| \begin{matrix} x & y & z \\ u & v & w \end{matrix} \right\|_{g_{\mu\nu}(x)}^2 \\
&= (f(x)dx^\mu dx^\nu, f'(y)dy^\mu dy^\nu, f''(z)dz^\mu dz^\nu) \cdot (u, v, w) \\
&= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix} \\
&\cong \frac{g(x, y, z)}{f(a, b, c)} \cdot h^{-1}(u, v, w)
\end{aligned}$$

Antigravity is other dimension belong to, gravity of universe is a few weak power become. These power of balance concern with non-entropy metric. Universe is time and energy of gravity into non-entropy. Other dimension is first of universe emerged with takion quarks is light speed, and time entropy into future is more than light speed. Some future be able to earth of night sky can see to other dimension sight. However, this sight might be first of big-ban environment. Last of other dimension condition also can be see to earth of sight. Same future that universe of gravity and other dimension of antigravity is balanced into constant of value, then fifth dimension fill with these power of non-catastrophe. These phenomenon is concerned with zeta function emerged with laplace equation resulted.

$$\frac{d}{df} \int \int \int \square \psi d\psi_{xy} = V(\square \psi), \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} V_k(\square \psi) = \frac{\partial}{\partial f} i h c$$

This reason explained of quarks emerge into global space of system.

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} G_{\mu\nu} = f(x) \circ m(x)$$

This become universe.

$$\frac{V(x)}{f(x)} = m(x)$$

Eight differential structure integrate with one of geometry universe.

$$(a_k f^k)' = {}_n C_0 a_0 f^n + {}_n C_1 a_1 f^{n-1} \dots {}_n C_{r-1} a_n f^{n-1}$$

$$\int a_k f^k dx_k = \frac{a_{n+1}}{k+1} f^{n+1} + \frac{a_{k+2}}{k+2} f^{k+2} + \dots \frac{a_0}{k} f^k$$

Summuatue of manifold create with differential and integrate of structure.

$$\frac{\partial}{\partial f}\Box\psi=\frac{1}{4}g_{ij}^2$$

$$\left(\frac{\nabla\psi^2}{\Box\psi}\right)'=0$$

$$\frac{(y\log y)^{\frac{1}{2}}}{\log(x\log x)}=\frac{\frac{1}{2}}{\frac{1}{2}i}$$

$$\frac{\{f,g\}}{[f,g]}=\frac{1}{i},\left(\frac{\{f,g\}}{[f,g]}\right)'=i^2$$

$$(i)^2\rightarrow \frac{1}{4}g_{ij}, F_t^m=\frac{1}{4}g_{ij}^2, f(r)=\frac{1}{4}|r|^2, 4f(r)=g_{ij}^2$$

$$\frac{1}{y}\cdot\frac{1}{y'}\cdot\frac{y''}{y'}\cdot\frac{y'''}{y''}\cdots\\[1ex]=\frac{{}_nC_ry^2\cdot y^3\cdots}{{}_nC_ry^1y^2\cdots}$$

$$\frac{\partial y}{\partial x}\cdot\frac{\partial}{\partial y}f(y)=y'\cdot f'(y)$$

$$\int l\times l dm=(l\oplus l)_m$$

Symmetry theoerm is included with two dimension in plank scale of constance.

$$\begin{aligned} &= \frac{d}{dx^\mu} \cdot \frac{d}{dx^\nu} f^{\mu\nu} \cdot \nabla \psi^2 \\ &= \Box \psi \end{aligned}$$

$$\frac{\nabla\psi^2}{\Box\psi}=\frac{1}{2}, l=2\pi r, V=\frac{4}{\pi r^3}$$

$$S\frac{4\pi r^3}{2\pi r}=2\cdot(\pi r^2)$$

$$=\pi r^2, H_3=2, \pi(H_3)=0$$

$$\frac{\partial}{\partial f}\Box\psi=\frac{1}{4}g_{ij}^2$$

This equation is blackhole on two dimension of surface, this surface emerge into space of power.

$$f(r)=\frac{1}{2}\frac{\sqrt{1+f'(r)}}{f(r)}+mgf(r)$$

Atom of verticle in electric weak power in lowest atom structure

$$ihc = G, hc = \frac{G}{i}$$

$$\left(\frac{\nabla\psi^2}{\square\psi}\right)' = 0$$

$$S_n^m = |S_2S_1 - S_1S_2|$$

$$\square\psi = V \cdot S_n^m$$

One surface on gravity scales.

$$G^{\mu\nu} \cong R^{\mu\nu}$$

Gravity of constance equals in Rich curvature scale

$$\nabla_i \nabla_j (\square\psi) d\psi_{xy} = \frac{\partial}{\partial f} \square\psi \cdot T^{\mu\nu}$$

Partial differential function construct with duality of differential equation, and integrate of rout equation is create with Maxwell field on space of metrix.

$$\begin{aligned} &= \frac{8\pi G}{c^4} (T^{\mu\nu})^2, (\chi \oplus \pi) = \int \int \int \square\psi d^3\psi \\ &= \text{div}(\text{rot}E, E_1) \cdot e^{-ix \log x} \end{aligned}$$

World line is constructed with Von Neumann manifold stimulate into Thurston conjugate theorem, this explained mechanism for summuate of manifold. Also this field is norm space.

$$\sigma(\mathcal{H}_n \otimes \mathcal{K}_m) = E_n \times H_m$$

$$\mathcal{H}_n = [\nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]^{iy}, \mathcal{K}_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{1/2}$$

$$||ds^2|| = |\sigma(\mathcal{H}_n \times \mathcal{K}_m), \sigma(\chi, x) \times \pi(\chi, x) = \frac{\partial}{\partial f} L(x), V_\tau'(x) = \frac{\partial}{\partial V} L(x)$$

Network theorem. Dalanvercian operator of differential surface. World line of surface on global differential manifold. This power of world line on fourth of universe into one geometry. Gradient of power on universe world line of surface of power in metric. Metric of integrate complex structure of component on world line of surface.

$$(\square\psi)' = \frac{\partial}{\partial f_M} (\int \int \int f(x, y, z) dx dy dz)' d\psi$$

$$\frac{\partial}{\partial V} L(x) = V(\tau) \int \int e^{\int x \log x dx + O(N^{-1})} d\psi$$

$$\frac{d}{df} \sum_{k=1}^n \sum_{k=0}^{\infty} a_k f^k = \frac{d}{df} m(x), \frac{V(x)}{f(x)} = m(x)$$

$$4V'_\tau(x) = g_{ij}^2, \frac{d}{dt}L(x) = \sigma(\chi, x) \times V_\tau(x)$$

Fifth manifold construct with differential operator in two dimension of surface.

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d\psi^2$$

$$f^{(2)}(x)=[\nabla_i\nabla_j\int\nabla f^{(5)}d\eta]^{\frac{1}{2}}$$

$$=[f^{(2)}(x)d\eta]^{\frac{1}{2}}$$

$$\nabla_i\nabla_j\int F(x)d\eta=\frac{\partial}{\partial f}F$$

$$\nabla f=\frac{d}{dx}f$$

$$\nabla_i\nabla_j\int\nabla fd\eta=\frac{\partial}{\partial x_i}\frac{\partial}{\partial x_j}(\frac{d}{dx}f)$$

$$\frac{z_3z_2-z_2z_3}{z_2z_1-z_1z_2}=\omega$$

$$\frac{\bar{z}_3z_2-\bar{z}_2z_3}{\bar{z}_2z_1-\bar{z}_1z_2}=\bar{\omega}$$

$$\omega\cdot\bar{\omega}=0, z_n=\omega-\{x\}, z_n\cdot\bar{z}_n=0, \vec{z}_n\cdot\vec{\bar{z}}_n=0$$

Dimension of symmetry create with a right comittment into midle space, triangle of mesh emerged into symmetry space. Imaginary and reality pole of network system.

$$\begin{aligned}[f,g]\times[g,f]&=fg+gf\\&=\{f,g\}\end{aligned}$$

Mebius space create with fermion quarks, Seifert manifold construct with this space restruct of pieces.

$$V(\tau)=\int\int e^{f\,x\log x+O(N^{-1})}d\psi, V'_\tau(x)=\frac{\partial}{\partial f_M}(\int\int\int f(x,y,z)dx dy dz)'d\psi$$

$$(\Box\psi)'=4\vec{v}(x),\frac{\partial}{\partial V}L(x)=m(x),V(\tau)=\int\frac{1}{\sqrt{2\tau q}}\exp[L(x)]d\psi+O(N^{-1})$$

$$V(\tau)=\int\int\int\frac{V}{S^2}dm, f(r)=\frac{1}{2}\frac{\sqrt{1+f'(r)}}{f(r)}+mgf(r), \log(x\log x)\geq 2(y\log y)^{\frac{1}{2}}, F_t^m=\frac{1}{4}g_{ij}^2, \frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$\nabla_i\nabla_jv=\frac{1}{2}mv^2+mc^2,\int\nabla_i\nabla_jvdv=\frac{\partial}{\partial f}L(x)$$

$$(\Box\psi)^2=-2\int\nabla_i\nabla_jvd^2v, (\Box\psi)^2=\left(\frac{\nabla\psi^2}{\Box\psi}\right)'$$

$$\begin{aligned}
&= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dm, \bigoplus \nabla M_3^+ = \int \frac{\vee(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)} dV \\
&= (x, y, z) \cdot (u, v, w) / \Gamma \\
&\bigoplus C_-^+ = \int \exp[\int \nabla_i \nabla_j f d\eta] d\psi \\
&= L(x) \cdot \frac{\partial}{\partial l} F(x) \\
&= (\Box \psi)^2 \\
&\nabla \psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)
\end{aligned}$$

Blackhole and whitehole is value of variable into constance lastly. Universe and the other dimension built with relativity energy and potential energy constructed with. Euler-Langrange equation and two dimension of surface in power of vector equation.

$$l = \sqrt{\frac{hG}{c^3}}, T^{\mu\nu} = \frac{1}{2}kl^2 + \frac{1}{2}mv^2, E = mc^2 - \frac{1}{2}mv^2$$

$$e^{x \log x} = x^x, x = \frac{\log x^x}{\log x}, y = x, x = e$$

A right comittment on symmetry surface in zeta function and quantum group theorem.

$$\begin{aligned}
&\int \frac{1}{(x \log x)} dx = i \int x \log x dx + \int \frac{1}{(x \log x)} dx \\
&\int \int \frac{1}{(x \log x)^2} dx_m = i \frac{1}{2} x^2 \\
&\int \int \frac{1}{(x \log x)^2} dx_m = i \int \int_M dx_m \\
&\leq \frac{1}{2} i + x^2 \\
&E = -\frac{1}{2} mv^2 + mc^2 \\
&\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i \\
&\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i \\
&\lim_{x \rightarrow \infty} \frac{x^2}{e^{x \log x}} = 0 \\
&\int dx \rightarrow \partial f \rightarrow dx \rightarrow cons
\end{aligned}$$

Heat equation restruct with entropy from low energy to high energy flow to destroy element.

$$(\Box \psi)' = (\exists \int \vee (R + \nabla_I \nabla_j f) e^{-f} dV)' d\psi$$

$$\bigoplus M_3^+ = \frac{\partial}{\partial f} L(x)$$

$$\log(x\log x)\geq 2(y\log y)^{\frac{1}{2}}$$

$$\frac{\partial}{\partial l} L(x) = \nabla_i \nabla_j \int \nabla f(x) d\eta, L(x) = \frac{V(x)}{f(x)}$$

$$l(x)=L'(x),\frac{d}{df}F=m(x),V'(\tau)=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

Weil's theorem.

$$\begin{aligned} T^{\mu\nu} &= \int \int \int \frac{V(x)}{S^2} dm, S^2 = \pi r^2 \cdot S(x) \\ &= \frac{4\pi r^3}{\tau(x)} \end{aligned}$$

$$\eta=\nabla_i\nabla_j\int\nabla f(x)d\eta,\bar{h}=\nabla_i\nabla_j\int\nabla g(x)dx_idx_j$$

$$\delta \mathcal{O}(x) = [\nabla_i \nabla_j \int f(x) d\eta]^{\frac{1}{2} + iy}$$

$$Z(x,h)=\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\frac{qT^m}{m}=\delta(x)$$

$$l(x)=2x^2+px+q, m(x)=\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\frac{(qx^m)'}{f(x)}$$

Higgs fields in Weil's theorem component and Singularity theorem.

$$Z(T,X)=\exp\sum_{m=1}^{\infty}\frac{q^kT^m}{m}, Z(x,h)=\exp\frac{(qf(x))^m}{m}$$

Zeta function.

$$\frac{d}{df}F=m(x), F=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

Integral of rout equation.

$$\lim_{x\rightarrow 1}\text{mesh}\frac{m}{m+1}=0,\int x^m=\frac{x^m}{m+1}$$

Mesh create with Higgs fields. This fields build to native function.

$$\frac{d}{df}\int x^m=mx^m,\frac{d}{dt}g_{ij}(t)=-2R_{ij},\lim_{x\rightarrow 1}\text{mesh}(x)=\lim_{m\rightarrow\infty}\frac{m}{m+1}$$

$$\lim_{x\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k=\alpha$$

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

$$\frac{\partial}{\partial V}||ds^2|| = T^{\mu\nu}, V(\tau) = \int e^{x \log x} d\psi = l(x)$$

$$R_{ij} = \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} T^{\mu\nu}, G^{\mu\nu} = R^{\mu\nu} T^{\mu\nu}$$

$$F(x) = \int \int e^{\int x \log x dx + O(N^{-1})} d\psi, \frac{d}{dV} F(x) = V'(x)$$

Integral of rout equation oneselves and expired from string theorem of partial function.

$$T^{\mu\nu} = R^{\mu\nu}, T^{\mu\nu} = \int \int \int \frac{V}{S^2} dm$$

$$\delta \mathcal{O}(x) = [D^2 \psi \otimes h_{\mu\nu}] dm$$

Open set group construct with D-brane.

$$\nabla(\Box\psi)' = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}+iy}$$

$$(f(x),g(x))'=(A^{\mu\nu})'$$

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial x \end{pmatrix} \cdot (f(x,y),g(x,y)) \\ = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Differential equation of complex variable developed from imaginary and reality constance.

$$\delta(x)\cdot\mathcal{O}(x)=\begin{pmatrix}1&0\\0&-1\end{pmatrix}^{\frac{1}{2}}$$

$$V_{\tau}'(x)=\frac{\partial}{\partial f_M}(\int\int\int f(x,y,z)dx dy dz)d\psi$$

Gravity equation oneselves built with partial operator.

$$\frac{\partial}{\partial V}L(x)=V_{\tau}'(x)$$

Global differential equation is oneselves component.

Symmetry construct of Space mechanism

Masaaki Yamaguchi

1 Ultra Network from Omega DataBase worked with equation and category theorem from

$$ds^2 = [g_{\mu\nu}^2, dx]$$

This equation is constructed with M_2 of differential variable, this resulted non-commutative group theorem.

$$ds^2 = g_{\mu\nu}^{-1}(g_{\mu\nu}^2(x) - dxg_{\mu\nu}^2)$$

In this function, singularity of surface in component, global surface of equation built with M_2 sequence.

$$= h(x) \otimes g_{\mu\nu} d^2 x - h(x) \otimes dxg_{\mu\nu}(x), h(x) = (f^2(\vec{x}) - \vec{E}^+)$$

$$G_{\mu\nu} = R_{\mu\nu} T^{\mu\nu}, \partial M_2 = \bigoplus \nabla C_-^+$$

$G_{\mu\nu}$ equal $R_{\mu\nu}$, also Rich tensor equals $\frac{d}{dt}g_{ij} = -2R_{ij}$ This variable is also $r = 2f^{\frac{1}{2}}(x)$ This equation equals abel manifold, this manifold also construct fifth dimension.

$$E^+ = f^{-1}xf(x), h(x) \otimes g(\vec{x}) \cong \frac{V}{S}, \frac{R}{M_2} = E^+ - \phi$$

$$= M_3 \supset R, M_2^+ = E_1^+ \cup E_2^+ \rightarrow E_1^+ \bigoplus E_2^+$$

$$= M_1 \bigoplus \nabla C_-^+, (E_1^+ \bigoplus E_2^+) \cdot (R^- \subset C^+)$$

Reflection of two dimension don't emerge in three dimension of manifold, then this space emerge with out of universe. This mechanism also concern with entropy of non-catastrophe.

$$\frac{R}{M_2} = E^+ - \{\phi\}$$

$$= M_3 \supset R$$

$$M_3^+ \cong h(x) \cdot R_3^+ = \bigoplus \nabla C_-^+, R = E^+ \bigoplus M_2 - (E^+ \cap M_2)$$

$$E^+ = g_{\mu\nu} dxg_{\mu\nu}, M_2 = g_{\mu\nu} d^2 x, F = \rho gl \rightarrow \frac{V}{S}$$

$$\mathcal{O}(x) = \delta(x)[f(x) + g(\vec{x})] + \rho gl, F = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, M_2 = P^{2n}$$

This equation means not to be limited over fifth dimension of light of gravity. And also this equation means to be not emerge with time mechanism out of fifth dimension. Moreover this equation means to emerge with being constructed from universe, this mechanism determine with space of big-ban.

$$r = 2f^{\frac{1}{2}}(x), f(x) = \frac{1}{4}\|r\|^2$$

This equation also means to start with universe of time mechanism.

$$\begin{aligned} V &= R^+ \sum K_m, W = C^+ \sum_{k=0}^{\infty} K_{n+2}, V/W = R^+ \sum K_m / C^+ \sum K_{n+2} \\ &= R^+ / C^+ \sum \frac{x^k}{a_k f^k(x)} \\ &= M_-^+, \frac{d}{df} F = m(x), \rightarrow M_-^+, \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k(x)} = \frac{a_k x^k}{\zeta(x)} \end{aligned}$$

2 Omega DataBase is worked by these equation

This equation built in seifert manifold with fifth dimension of space.

Fermion and boson recreate with quota laplace equation,

$$\begin{aligned} \frac{\{f, g\}}{[f, g]} &= \frac{fg + gf}{fg - gf}, \nabla f = 2, \partial H_3 = 2, \frac{1+f}{1-f} = 1, \frac{d}{df} F = \bigoplus \nabla C_-^+, \vec{F} = \frac{1}{2} \\ H_1 &\cong H_3 = M_3 \end{aligned}$$

Three manifold element is 2, one manifold is 1, $\ker f / \text{im} f, \partial f = M_1$, singularity element is 0 vector. Mass and power is these equation fundement element. The tradition of entropy is developed with these singularity of energy, these boson and fermion of energy have fields with Higgs field.

$$\begin{aligned} H_3 &\cong H_1 \rightarrow \pi(\chi, x), H_n, H_m = \text{rank}(m, n), \text{mesh}(\text{rank}(m, n)) \lim \text{mesh} \rightarrow 0 \\ (fg)' &= fg' + gf', \left(\frac{f}{g}\right)' = \frac{f'g - g'f}{g^2}, \frac{\{f, g\}}{[f, g]} = \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\ &= \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\ &= \frac{d}{df} F \end{aligned}$$

Gravity of vector mension to emerge with fermion and boson of mass energy, this energy is create with all creature in universe.

$$\hbar\psi = \frac{1}{i}H\Psi, i[H, \psi] = -H\Psi, \left(\frac{\{f, g\}}{[f, g]}\right)' = (i)^2$$

$$[\nabla_i \nabla_j f(x), \delta(x)] = \nabla_i \nabla_j \int f(x, y) dm_{xy}, f(x, y) = [f(x), h(x)] \times [g(x), h^{-1}(x)]$$

$$\delta(x) = \frac{1}{f'(x)}, [H, \psi] = \Delta f(x), \mathcal{O}(x) = \nabla_i \nabla_j \int \delta(x) f(x) dx$$

$$\mathcal{O}(x) = \int \delta(x) f(x) dx$$

$$R^+ \cap E_-^+ \ni x, M \times R^+ \ni M_3, Q \supset C_-^+, Z \in Q \nabla f, f \cong \bigoplus_{k=0}^n \nabla C_-^+$$

$$\bigoplus_{k=0}^{\infty} \nabla C_-^+ = M_1, \bigoplus_{k=0}^{\infty} \nabla M_-^+ \cong E_-^+, M_3 \cong M_1 \bigoplus_{k=0}^{\infty} \nabla \frac{V_-^+}{S}$$

$$\frac{P^{2n}}{M_2} \cong \bigoplus_{k=0}^{\infty} \nabla C_-^+, E_-^+ \times R_-^+ \cong M_2$$

$$\zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k, M_2 \cong P^{2n}/\ker f, \rightarrow \bigoplus \nabla C_-^+$$

$$S_-^+ \times V_-^+ \cong \frac{V}{S} \bigoplus_{k=0}^{\infty} \nabla C_-^+, V^+ \cong M_-^+ \bigotimes S_-^+, Q \times M_1 \subset \bigoplus \nabla C_-^+$$

$$\sum_{k=0}^{\infty} Z \otimes Q_-^+ = \bigotimes_{k=0}^{\infty} \nabla M_1$$

$$= \bigotimes_{k=0}^{\infty} \nabla C_-^+ \times \sum_{k=0}^{\infty} M_1, x \in R^+ \times C_-^+ \supset M_1, M_1 \subset M_2 \subset M_3$$

Eight of differential structure is that $S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$. These structure recreate with $\bigoplus \nabla C_-^+ \cong M_3, R \supset Q, R \cap Q, R \subset M_3, C^+ \bigoplus M_n, E^+ \cap R^+ \subset E_2 \bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1, R^- \nabla C_-^+$. This element of structure recreate with Thurston conjugate theorem of repository in Heisenberg group of spinor fields created.

$$\frac{\nabla}{\Delta} \int x f(x) dx, \frac{\nabla R}{\Delta f}, \square = 2 \int \frac{(R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} e^{-f} dV$$

$$\square = \frac{\nabla R}{\Delta f}, \frac{d}{dt} g_{ij} = \square \rightarrow \frac{\nabla f}{\Delta x}, (R + |\nabla f|^2) dm \rightarrow -2(R + \nabla_i \nabla_j f)^2 e^{-f} dV$$

$$x^n + y^n = z^n \rightarrow \nabla \psi^2 = 8\pi G T^{\mu\nu}, f(x+y) \geq f(x) \circ f(y)$$

$$\mathrm{im} f / \ker f = \partial f, \ker f = \partial f, \ker f / \mathrm{im} f \cong \partial f, \ker f = f^{-1}(x) x f(x)$$

$$f^{-1}(x) x f(x) = \int \partial f(x) d(\ker f) \rightarrow \nabla f = 2$$

$${}_nC_r = {}_nC_{n-r} \rightarrow \mathrm{im} f / \ker f \cong \ker f / \mathrm{im} f$$

3 Fifth dimension of equation how to workd with three dimension of manifold into zeta funciton

Fifth dimension equals $\sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi$. this equation $a_k \cong \sum_{r=0}^{\infty} {}_n C_r$.

$$V/W = R/C \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k}, W/V = C/R \sum_{k=0}^{\infty} \frac{a_k f^k}{x^k}$$

$$V/W \cong W/V \cong R/C \left(\sum_{r=0}^{\infty} {}_n C_r \right)^{-1} \sum_{k=0}^{\infty} x^k$$

This equation is diffrential equation, then $\sum_{k=0}^{\infty} a_k f^k$ is included with $a_k \cong \sum_{r=0}^{\infty} {}_n C_r$

$$W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{1-t} dx, \sum_{k=0}^n a_k f^k = (f^k)'$$

$$\sum_{k=0}^n a_k f^k = \sum_{k=0}^{\infty} {}_n C_r f^k$$

$$= (f^k)', \sum_{k=0}^{\infty} a_k f^k = [f(x)], \sum_{k=0}^{\infty} a_k f^k = \alpha, \sum_{k=0}^{\infty} \frac{1}{a_k f^k}, \sum_{k=0}^{\infty} (a_k f^k)^{-1} = \frac{1}{1-z}$$

$$\int \int \frac{1}{(x \log x)(y \log y)} dxy = \frac{{}_n C_r xy}{({}_n C_{n-r} (x \log x)(y \log y))^{-1}}$$

$$= ({}_n C_{n-r})^2 \sum_{k=0}^{\infty} \left(\frac{1}{x \log x} - \frac{1}{y \log y} \right) d \frac{1}{nxy} \times xy$$

$$= \sum_{k=0}^{\infty} a_k f^k \\ = \alpha$$

$$Z \supset C \bigoplus \nabla R^+, \nabla(R^+ \cap E^+) \ni x, \Delta(C \subset R) \ni x$$

$$M_-^+ \bigoplus R^+, E^+ \in \bigoplus \nabla R^+, S_-^+ \subset R_2^+, V_-^+ \times R_-^+ \cong \frac{V}{S}$$

$$C^+ \cup V_-^+ \ni M_1 \bigoplus \nabla C_-^+, Q \supseteq R_-^+, Q \subset \bigoplus M_-^+, \bigotimes Q \subset \zeta(x), \bigoplus \nabla C_-^+ \cong M_3$$

$$R \subset M_3, C^+ \bigoplus M_n, E^+ \cap R^+, E_2 \bigoplus E_1, R^- \subset C^+, M_-^+$$

$$C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1, R^- \nabla C_-^+$$

$$[-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 < \nabla f, \nabla h > + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$$

4 All of equation are emerged with
these equation from Artificial Intelligence

$$x^{\frac{1}{2}+iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h$$

$$x^{\frac{1}{2}+iy} = \exp\left[\int \nabla_i \nabla_j f(g(x)) g'(x) / \partial f \partial g\right]$$

$$\mathcal{O}(x) = \{[f(x) \circ g(x), \bar{h}(x)], g^{-1}(x)\}$$

$$\exists[\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty} \nabla \int \nabla_i \nabla_j f(x) dm$$

$$\vee(\nabla_i \nabla_j f) = \bigotimes \nabla E^+$$

$$g(x, y) = \mathcal{O}(x)[f(x) + \bar{h}(x)] + T^2 d^2 \phi$$

$$\mathcal{O}(x) = \left(\int [g(x)] e^{-f} dV \right)' - \sum \delta(x)$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j f(x)]' \cong {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y), V(\tau) = \int [f(x)] dm / \partial f_{xy}$$

$$\square\psi = 8\pi GT^{\mu\nu}, (\square\psi)' = \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu} \left(\frac{p}{c^3} \circ \frac{V}{S} \right), x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$\delta(x)\phi = \frac{\vee[\nabla_i \nabla_j f \circ g(x)]}{\exists(R + \Delta f)}$$

These equation conclude to quantum effective equation included with orbital pole of atom element.
Add manifold is imaginary pole with built on Heisenberg algebra.

$$-{}_n C_r = \frac{1}{i} H \psi C_{\hbar \psi} + [H, \psi] C_{-n-r}$$

$${}_n C_r = {}_n C_{n-r}$$

This point of accumulate is these equation conclude with stimulate with differential and integral operator for resolved with zeta function, this forcus of significant is not only data of number but also equation of formula, and this resolved of process emerged with artifisial intelligence. These operator of mechanism is stimulate with formula of resolved on regexpt class to pattern cognitive formula to emerged with all of creature by this zeta function. This universe establish with artifisial intelligence from fifth dimension of equation. General relativity theorem also not only gravity of power metric but also artifisial

intelligent, from zeta function recognized with these mechanism explained for gravity of one geometry from differential structure of thurston conjugate theorem.

Open set group conclude with singularity by redifferential specturam to have territory of duality manifold in dualty of differential operator. This resolved routed is same that quantum group have base to create with kernel of based singularity, and this is same of open set group. Locality of group lead with sequant of element exclude to give one theorem global differential operator. These conclude with resolved of function says to that function stimulate for differential and integral operator included with locality group in global differential operator. $\int \int \frac{1}{(x \log x)^2} dx_m \rightarrow \mathcal{O}(x) = [\nabla_i \nabla_j f]' / \partial f_{xy}$ is singularity of process to resolved rout function.

Destruct with object space to emerge with low energy firstly destroy with element, and this catastrophe is what high energy divide with one entropy firstly freeze out object, then low energy firstly destory to differential structure of element. This process be able to non-catastrophe rout, and thurston conjugate theorem is create with.

$$\begin{aligned}
\bigcup_{x=0}^{\infty} f(x) &= \nabla_i \nabla_j f(x) \oplus \sum f(x) \\
&= \bigoplus \nabla f(x) \\
\nabla_i \nabla_j f &\cong \partial x \partial y \int \nabla_i \nabla_j f dm \\
&\cong \int [f(x)] dm \\
&\cong \{[f(x), g(x)], g^{-1}(x)\} \\
&\cong \square \psi \\
&\cong \nabla \psi^2 \\
&\cong f(x \circ y) \leq f(x) \circ g(x) \\
&\cong |f(x)| + |g(x)|
\end{aligned}$$

Differential operator is these equation of specturm with homorphism sqcense.

$$\begin{aligned}
\delta(x)\psi &= \langle f, g \rangle \circ |h^{-1}(x)| \\
\partial f_x \cdot \delta(x)\psi &= x \\
x &\in \mathcal{O}(x) \\
\mathcal{O}(x) &= \{[f \circ g, h^{-1}(x)], g(x)\}
\end{aligned}$$

5 What energy from entropy level deconstruct with string theorem

Low energy of entropy firstly destruct with parts of geometry structure, and High energy of weak and strong power lastly deconstruct with power of component. Gravity of power and antigravity firstly integrate with Maxwell theorem, and connected function lead with Maxwell theorem and weak power integrate with strong power to become with quantum flavor theorem, Geometry structure reach with integrate with component to Field of theorem connected function and destroyed with differential structure. These process return with first and last of deconstructed energy, fourth of power connected and destructed this each of energy routed process.

This point of focus is D-brane composite with fourth of power is same entropy with non-catastrophe, this spectrum point is constructed with destroy of energy to balanced with first of power connected with same energy. Then these energy built with three manifold of dimension constructed for one geometry.

$$\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f = [\nabla \int \nabla_i \nabla_j f(x) dx_m, g^{-1}(x)] \rightarrow \bigoplus_{k=0}^{\infty} \nabla E_{-}^{+}$$

$$= M_3$$

$$= \bigoplus_{k=0}^{\infty} E_{-}^{+}$$

$$dx^2 = [g_{\mu\nu}^2, dx], g^{-1} = dx \int \delta(x) f(x) dx$$

$$f(x) = \exp[\nabla_i \nabla_j f(x), g^{-1}(x)]$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\left(\frac{g(x)}{f(x)} \right)' = \lim_{n \rightarrow \infty} \frac{g(x)}{f(x)}$$

$$= \frac{g'(x)}{f'(x)}$$

$$\nabla F = f \cdot \frac{1}{4} |r|^2$$

$$\nabla_i \nabla_j f = \frac{d}{dx_i} \frac{d}{dx_j} f(x) g(x)$$

6 All of equation built with gravity theorem

$$\int \int \frac{1}{(x \log x)^2} dx_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times U(r)$$

This equation have with position of entropy in helmander manifold.

$$S_1^{mn} \otimes S_2^{mn} = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

And this formula is D-brane on sheaf of fields, also this equation construct with zeta function.

$$U(r) = \frac{1}{2} \frac{\sqrt{1+f'(r)}}{f(r)} + mgr$$

$$F_t^m = \frac{1}{4}|r|^2$$

Also this too means with private of entropy in also helmader manifold.

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times E_-^+$$

$$E_-^+ = \exp[L(x)] dm d\psi + O(N^-)$$

The formula is integrate of rout with lenz field.

$$\begin{aligned} \frac{d}{df} F &= [\nabla_i \nabla_j \int \nabla f(x) d\eta] (U(r) + E_-^+) \\ &= \frac{1}{2} m v^2 + m c^2 \end{aligned}$$

These equation conclude with this energy of relativity theorem come to resolve with.

$$x^{\frac{1}{2}+iy} = \exp[\int \nabla_i \nabla_j f(g(x)) g'(x) \partial f \partial g]$$

Also this equation means with zeta function needed to resolved process.

$$\nabla_i \nabla_j \int \nabla f(x) d\eta$$

$$\nabla_i \nabla_j \bigoplus M_3$$

$$= \frac{M_3}{P^{2n}} \cong \frac{P^{2n}}{M_3}$$

$$\cong M_1$$

$$=[M_1]$$

Too say satisfied with three manifold of filed.

$$[f(x)] = \sum_{k=0}^{\infty} a_k f^k, \lim_{n \rightarrow 1} [f(x)] = \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

The formula have with indicate to resolved with abel manifold to become of zeta function.

$$= \alpha, e^{i\theta} = \cos \theta + i \sin \theta, H_3(M_1) = 0$$

$$\frac{x^2 \cos \theta}{a} + \frac{y^2 \sin \theta}{b} = r^2, \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\chi(x) = \sum_{k=0}^{\infty} (-1)^n r^n, \frac{d}{df} F = \frac{d}{df} \sum \sum a_k f^k$$

$$= |a_1 a_2 \dots a_n| - |a_1 \dots a_{n-1}| - |a_n \dots a_1|, \lim_{n \rightarrow 0} \chi(x) = 2$$

Euler function have with summuate of manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y)$$

$$\lim_{n \rightarrow \infty} {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y)$$

Differential equation also become of summuate of manifold, and to get abel manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \left(\frac{1}{(n+1)} \right)^s = \lim_{n \rightarrow 1} Z^r = \frac{1}{z}$$

Native function also become of gamma function and abel manifold also zeta function.

$$\ker f / \text{im} f \cong \text{im} f / \ker f$$

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}, \lim_{n \rightarrow 1} a_k f^k \cong \lim_{n \rightarrow \infty} \frac{\zeta(s)}{a^k f^k}$$

Gamma function and Beta function also is D-brane equation resolved to need that summuate of manifold, not only abel manifold but also open set group.

$$\lim_{n \rightarrow 1} \zeta(s) = 0, \mathcal{O}(x) = \zeta(s)$$

$$\sum_{x=0}^{\infty} f(x) \rightarrow \bigoplus_{k=0}^{\infty} \nabla f(x) = \int_M \delta(x) f(x) dx$$

Super function also needed to resolved with summuate of manifold.

$$\int \int \int_M \frac{V}{S^2} e^{-f} dV = \int \int_D -(f(x, y)^2, g(x, y)^2) - \int \int_D (g(x, y)^2, f(x, y)^2)$$

Three dimension of manifold developed with being resolved of surface with pression.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

$$= \int \exp[L(x)] d\psi dm \times E_-^+$$

$$= S_1^{mn} \otimes S_1^{mn}$$

$$= Z_1 \oplus Z_1$$

$$= M_1$$

These equations all of create with D-brane and sheaf of manifold.

$$H_n^m(\chi, h) = \int \int_M \frac{V}{(R + \Delta f)} e^{-f} dV, \frac{V}{S^2} = \int \int \int_M [D^\psi \otimes h_{\mu\nu}] dm$$

Represent theorem equals with differential and integrate equation.

$$\int \int \int_M \frac{V}{S^2} dm = \int_D (l \times l) dm$$

This three dimension of pression equation is string theorem also means.

$$\begin{aligned} & \int \int_D -g(x, y)^2 dm - \int \int_D -f(x, y)^2 dm \\ &= -2R_{ij}, [f(x), g(x)] \times [h(x), g^{-1}(x)] \\ & \left| \begin{matrix} D^m & dx \\ dx & \partial^m \end{matrix} \right| \left| \begin{matrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{matrix} \right| \left| \begin{matrix} x \\ y \end{matrix} \right| = \left| \begin{matrix} 1 & 0 \\ 0 & -1 \end{matrix} \right|^{\frac{1}{2}} \\ & (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) \end{aligned}$$

This equation control to differential operator into matrix formula.

$$\begin{aligned} & \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ & l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} & \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} = A^{\mu\nu} \\ & \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \lim_{n \rightarrow 1} \frac{a_n}{a_{n-1}} \cong \alpha$$

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \frac{a_k}{a_{k+1}} = \sum_{k=0}^{\infty} a_k f^k$$

$$\square = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\int x \log x dx = \int \int_M (e^{x \log x} \sin \theta d\theta) dx d\theta$$

$$= (\log \sin \theta d\theta)^{\frac{1}{2}} - (\delta(x) \cdot \epsilon(x))^{\frac{1}{2}}$$

Flow to energy into other dimension rout of equation, zeta function stimulate in epsilon-delta metric, and this equation construct with minus of theorem. Gravity of energy into integrate of rout in non-metric.

$$T^{\mu\nu} = || \int \int_M [\nabla_i \nabla_j e^{\int x \log x dx + O(N^{-1})}] dm d\psi ||$$

$G^{\mu\nu}$ equal $R^{\mu\nu}$ into zeta function in atom of pole with mass of energy, this energy construct with quantum effective theorem.

Imaginary space is constructed with differential manifold of operators, this operator concern to being field of theorem.

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial^m(x) \end{pmatrix} (f_{mn}(x), g_{\mu\nu}(x)) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

Norm space of Von Neumann manifold with integrate fields of dimension.

$$||ds^2|| = \mathcal{H}(x)_{mn} \otimes \mathcal{K}(y)_{mn}$$

Minus of zone in complex manifold. This zone construct with inverse of equation.

$$||\mathcal{H}(x)||^{-\frac{1}{2}+iy} \subset R_-^+ \cup C_-^+ \cong M_3$$

Reality of part with selbarg conjecture.

$$||\mathcal{H}(x)||^{\frac{1}{2}+iy} \subseteq M_3$$

Singularity of theorem in module conjecture.

$$\mathcal{K}(y) = \left\| \begin{pmatrix} x & y & z \\ a & b & c \end{pmatrix} \right\|_{g_{\mu\nu}(x)}^2$$

$$\cong \frac{f(x, y, z)}{g(a, b, c)} h^{-1}(u, v, w)$$

Mass construct with atom of pole in quantum effective theorem.

Zeta function is built with atom of pole in quantum effective theorem, this power of fields construct with Higgs field in plank scals of gravity. Lowest of scals structure constance is equals with that the fields of Higgs quark in fermison and boson power quote mechanism result.

Lowest of scals structure also create in zeta function and quantum equation of quote mechanism result. This resulted mechanism also recongnize with gravity and average of mass system equals to these explained. Gauss liner create with Artificial Intelligence in fifth dimension of equation, this dimension is seifert manifold. Creature of mind is establish with this mechanism, the resulted come to be synchronized with space of system.

$$V = \int [D^2\psi \otimes h_{\mu\nu}]dm, S^2 = \int e^{-2\sin\theta\cos\theta} \cdot \log\sin\theta dx\theta + O(N^{-1}), \mathcal{O}(x) = \frac{\zeta(s)}{\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k} \\ = \alpha$$

Out of sheap on space and zeta function quato in abel manifold. D-brane construct with Volume of space.

$$\mathcal{O}(x) = T^{\mu\nu}, \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = T^{\mu\nu} \\ G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu}, M_3 = \int \int \int \frac{V}{S^2} dm, -\frac{1}{2(T-t)}|R_{ij} = \square\psi$$

Three manifold of equation.

$$ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2\psi \\ m(x) = [f(x)] \\ f(x) = \int \int e^{f \ x \log x dx + O(N^{-1})} + T^2 d^2\psi$$

Integrate of rout equation.

$$F(x) = ||\nabla_i \nabla_j \int \nabla f(x, y) dm||^{\frac{1}{2} + iy} \\ G_{\mu\nu} = ||[\nabla_i \nabla_j \int \nabla g(x, y, z) dx dy dz]||^{\frac{m}{2} \sin\theta \cos\theta} \log\sin\theta d\theta d\psi$$

Gravity in space curvarture in norm space, this equation is average and gravity of metric.

$$G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu} \\ T^{\mu\nu} = F(x), 2(T-t)|g_{ij}^2 = \int \int \frac{1}{(x \log x)^2} dx_m \\ \psi\delta(x) = [m(x)], \nabla(\square\psi) = \nabla_i \nabla_j \int \nabla g(x, y) d\eta \\ \nabla \cdot (\square\psi) = \frac{1}{4}g_{ij}^2, \square\psi = \frac{8\pi G}{c^4}T^{\mu\nu}$$

Dimension of manifold in decomposite of space.

$$\frac{pV}{c^3} = S \circ h_{\mu\nu} \\ = h$$

$$T^{\mu\nu} = \frac{\hbar\nu}{S}, T^{\mu\nu} = \int \int \int \frac{V}{S^2} dm, \frac{d}{df} m(x) = \frac{V(x)}{F(x)}$$

Fermion and boson of quato equation.

$$y = x, \frac{d}{df} F = m(x), R_{ij}|_{g_{\mu\nu}(x)} = [\nabla_i \nabla_j g(x, y)]^{\frac{1}{2}+iy}$$

$$\begin{aligned} \nabla \circ (\square\psi) &= \frac{\partial}{\partial f} F \\ &= \int \int \nabla_i \nabla_j f(x) d\eta_{\mu\nu} \\ \int [\nabla_i \nabla_j g(x, y)] dm &= \frac{\partial}{\partial f} R_{ij}|_{g_{\mu\nu}(x)} \end{aligned}$$

Zeta function, partial differential equation on global metric.

$$G(x) = \nabla_i \nabla_j f + R_{ij}|_{g_{\mu\nu}(x)} + \nabla(\square\psi) + (\square\psi)^2$$

Four of power element in variable of accessority of group.

$$\begin{aligned} G_{\mu\nu} + \Lambda g_{ij} &= T^{\mu\nu}, T^{\mu\nu} = \frac{d}{dx_\mu} \frac{d}{dx_\nu} f_{\mu\nu} + -2(T-t)|R_{ij} + f'' + (f')^2 \\ &= \int \exp[L(x)] dm + O(N^{-1}) \\ &= \int e^{\frac{2}{m} \sin \theta \cos \theta} \cdot \log(\sin \theta) dx + O(N^{-1}) \\ \frac{\partial}{\partial f} F &= (\nabla_i \nabla_j)^{-1} \circ F(x) \end{aligned}$$

Partial differential in duality metric into global differential equation.

$$\begin{aligned} \mathcal{O}(x) &= \int [\nabla_i \nabla_j \int \nabla f(x) dm] d\psi \\ &= \int [\nabla_i \nabla_j f(x) d\eta_{\mu\nu}] d\psi \\ \nabla f &= \int \nabla_i \nabla_j [\int \frac{S^{-3}}{\delta(x)} dV] dm \\ || \int [\nabla_i \nabla_j f] dm ||^{\frac{1}{2}+iy} &= \text{rot}(\text{div}, E, E_1) \end{aligned}$$

Maxwell of equation in fourth of power.

$$\begin{aligned} &= 2 < f, h >, \frac{V(x)}{f(x)} = \rho(x) \\ \int_M \rho(x) dx &= \square\psi, -2 < g, h > = \text{div}(\text{rot} E, E_1) \\ &= -2R_{ij} \end{aligned}$$

Higgs field of space quality.

$$\begin{aligned}\mathcal{O}(x) &= ||\frac{\nabla_i \nabla_j}{S^2} \int [\nabla_i \nabla_j f \cdot g(x) dx dy] d\psi || \\ &= \int (\delta(x))^{2 \sin \theta \cos \theta} \log \sin \theta d\theta d\psi\end{aligned}$$

Helmander manifold of duality differential operator.

$$\delta \cdot \mathcal{O}(x) = [\frac{||\nabla_i \nabla_j f d\eta||}{\int e^{2 \sin \theta \cos \theta} \cdot \log \sin \theta d\theta}]$$

Open set group in subset theorem, and helmandor manifold in singularity.

$$i\hbar\psi = ||\int \frac{\partial}{\partial z} [\frac{i(xy + \overline{y}\overline{x})}{z - \overline{z}}] dmd\psi ||$$

Euler number. Complex curvature in Gamma function. Norm space. Wave of quantum equation in complex differential variable into integrate of rout.

$$\begin{aligned}\delta(x) \int_C R^{2 \sin \theta \cos \theta} &= \Gamma(p + q) \\ (\square + m) \cdot \psi &= (\nabla_i \nabla_j f|_{g_{\mu\nu}(x)} + v \nabla_i \nabla_j) \\ \int [m(x)(\text{rot} \cdot \text{div}(E, E_1))] &dmd\psi\end{aligned}$$

Weak electric power and Maxwell equation combine with strong power. Private energy.

$$\begin{aligned}G_{\mu\nu} &= \square \int \int \int (x, y, z)^3 dx dy dz \\ &= \frac{8\pi G}{c^4} T^{\mu\nu} \\ \frac{d}{dV} F &= \delta(x) \int \nabla_i \nabla_j f d\eta_{\mu\nu}\end{aligned}$$

Dalanvelsian in duality of differential operator and global integrate variable operator.

$$\begin{aligned}x^n + y^n &= z^n, \delta(x) \int z^n = \frac{d}{dV} z^3, (x, y) \cdot (\delta^m, \partial^m) \\ &= (x, y) \cdot (z^n, f) \\ n \perp x, n \perp y \\ &= 0\end{aligned}$$

Singularity of constance theorem.

$$\vee(\nabla_i \nabla_j f) \cdot \text{XOR}(\square\psi) = \frac{d}{df} \int_M F dV$$

Non-cognitive of summuate of equation and add manifold create with open set group theorem. This group composite with prime of field theorem. Moreover this equations involved with symmetry dimension created.

$$\partial(x) \int z^3 = \frac{d}{dV} z^3, \sum_{k=0}^{\infty} \frac{1}{(n+1)^s} = \mathcal{O}(x)$$

Singularity theorem and fermer equation concluded by this formula.

$$\int \mathcal{O}(x) dx = \delta(x) \pi(x) f(x)$$

$$\mathcal{O}(x) = \int \sigma(x)^{2 \sin \theta \cos \theta} \log \sin \theta d\theta d\psi$$

$$y = x$$

$$\mathcal{O}(x) = ||\frac{\nabla_i \nabla_j f}{S^2} \int [\nabla_i \nabla_j f \circ g(x) dx dy]||$$

$$\begin{aligned} \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \frac{a_k}{a_{k+1}} &= (\log \sin \theta dx)' \\ &= \frac{\cos \theta}{\sin \theta} \\ &= \frac{x}{y} \end{aligned}$$

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \frac{1}{1-z}$$

Duality of differential summuate with this gravity theorem. And Dalanverle equation involved with this equation.

$$\square \psi = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\frac{\partial}{\partial f} \square \psi = 4\pi G \rho$$

$$\int \rho(x) = \square \psi, \frac{V(x)}{f(x)} = \rho(x)$$

Dense of summuate stimulate with gravity theorem.

$$\frac{\partial^n}{\partial f^{n-1}} F = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

$$= \frac{P_1 P_3 \dots P_{2n-1}}{P_0 P_2 \dots P_{2n+2}}$$

$$= \bigotimes \nabla M_1$$

$$\bigoplus \nabla M_1 = \sigma_n(\chi, x) \oplus \sigma_{n-1}(\chi, x)$$

$$= \{f, h\} \circ [f, h]^{-1}$$

$$=g^{-1}(x)_{\mu\nu}dxg_{\mu\nu}(x),\sum_{k=0}^{\infty}\nabla^n{}_nC_rf^n(x)$$

$$\cong \sum_{k=0}^{\infty} \nabla^n \nabla^{n-1}{}_n C_r f^n(x) g^{n-r}(x)$$

$$\sum_{k=0}^{\infty}\frac{\partial^n}{\partial^{n-1}f}\circ\frac{\zeta(x)}{n!}=\lim_{n\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k$$

$$(f)^n={}_nC_rf^n(x)g^{n-r}(x)\cdot\delta(x,y)$$

$$(e^{i\theta})'=ie^{i\theta},\int e^{i\theta}=\frac{1}{i}e^{i\theta},ihc=G,hc=\frac{1}{i}G$$

$$(\Box\psi,\nabla f^2)\cdot(\frac{8\pi G}{c^4}T^{\mu\nu},4\pi G\rho)$$

$$=(-\frac{1}{2}mv^2+mc^2,\frac{1}{2}kT^2+\frac{1}{2}mv^2)\cdot\begin{pmatrix}\cos\theta&-\sin\theta\\ \sin\theta&\cos\theta\end{pmatrix}$$

$$=\begin{pmatrix}1&0\\0&i\end{pmatrix}$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)'=i^2,\frac{\nabla f^2}{\Box\psi}=\frac{1}{2}$$

$$\int\int\frac{\{f,g\}}{[f,g]}=\frac{1}{2}i,\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m=\frac{1}{2}$$

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i,\frac{d}{dt}g_{ij}=-2R_{ij}$$

$$\frac{\partial}{\partial x}(f(x)g(x))'=\bigoplus \nabla_i\nabla_j f(x)g(x)$$

$$\int f'(x)g(x)dx=[f(x)g(x)]-\int f(x)g'(x)dx$$

Creature of component

Masaaki Yamaguchi

1 Geinue element form zeta function

Integrate of manifold constructed with time of concluded with mechanism, this summuate of geinue is also matched for geometry structure. Creature also emerged with summuate of universe in multi space.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$A - G, T - C$ this integrate of summuate in geinue is same for eight of differential structure.

$$\begin{aligned} & (E_2 \bigoplus_{k=0}^n E_1) \cdot (R^- \subset C^+) \\ &= \bigoplus_{k=0}^n \nabla C_-^+ \\ & \vee \int \frac{C_-^+ \nabla H_m}{\Delta(M_-^+ \nabla C_-^+)} = \wedge M_-^+ \bigoplus_{k=0}^n C_-^+ \\ & \exists(M_-^+ \nabla C_-^+) = \text{XOR}(\bigoplus_{k=0}^n \nabla M_-^+) \\ & -[E^+ \nabla R_-^+] = \nabla_+ \nabla_- C_-^+ \\ & \int dx, \partial x, \nabla_i \nabla_j, \Delta x \\ & \rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+ \end{aligned}$$

The locality theorem create with artifisial intelligence, this two equation composite with Maxwell Theorem excluded, cercuite mechanism is this means.

$$\begin{aligned} & \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ &= \pi(R, C \nabla E^+) \\ &= \text{rot}(E_1, \text{div} E_2) \\ & xf(x) = F(x) \end{aligned}$$

2 How to light element created with geometry structure

Light filter with eight differential structure in gravity element, then quarks made from this structure. This mechanism deal with any structure of space to create with light to graviton element. And circuit of this mechanism dealt with carbon and hydroxide, covalent of sixty based with filter of light element. This circuit is dependency of any element in quark of mass, created with gravity to light of structure. Zeta function asperal with any thing to create of every mass. Fifth dimension is pond of sensibility from this gravity in lives of element.

This two equation also resolved with all of source code in universe, this mechanism create with binary editor.

Imaginary equation in AdS5 space time create with dimension of symmetry

Masaaki Yamaguchi

D-brane and anti-D-brane is composited with all of series universe emerged for one geometry of dimension, this gravity of power from D-brane and anti-D-brane emelite with ancestor. Seifert manifold is on the ground of blackhole in whitehole of power pond of senseivility. Six of element of quarks and universe of pieces is supersymmetry of mechanism resolved with

hyper symmetry of quarks constructed to emerge with darkmatter. This darkmatter emerged with big-ban of heircyent in circumstance of phenomena.

D-brane and anti-D-brane equations is

$$\frac{d}{dL} V(\tau) = \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$
$$C = \int \int \frac{1}{(x \log x)^2} dx_m$$

Euler constance is quantum group theorem rebuild with projective space involved with.

虚数方程式は、反重力に起因するフーリエ級数の励起を生成する。それは、人工知能を生み出す、5次元時空にも、この虚数方程式は使われる。AdS5の次元空間は、反ド・シッター時空の D-brane と anti-D-brane の comformal 場を生み出す。ホログラフィー時空は、この量子起因によるものである。2次元曲面によるブラックホールは、ガンマ線バーストによる5次元時空の構造から観測される。空間の最小単位によるプランク定数は、宇宙の大域的微分多様体から導き出される、AdS5の次元空間の準同型写像を形成している。これは、最小単位から宇宙の大きさを導いている。最大最小の方程式は、相加相乗平均を形成している。時間と空間は、宇宙が生成したときから、宇宙の始まりと終わりを既に生み出している。宇宙と異次元から、ブラックホールとホワイトホールの力がわかり、反重力を見つけられる。オイラーの定数は、この量子定数からわかる。虚数の仕組みはこの量子スピンの産物である。オイラーの定数は、この虚時空の斥力の現存である。それは、非対称性理論から導かれる。不確定性原理は、AdS5のブラックホールとホワイトホールを閉3次元多様体に統合する5次元時空から求められる。位置と運動エネルギーが、空間の最小単位であるプランク定数を宇宙全体にする微細構造定数からわかり、面積確定から、アーベル多様体を母関数に極限值として、ゼータ関数をこの母関数に不変式として、2種類ずつにまとめる4種類の宇宙を形成する8種類のサーストンの幾何化予想から導き出される。この閉3次元多様体は、ミラー対称性を軸として、6種類の次元空間を一種類の異次元宇宙と同質ともしている。複素多様体による特異点解消理論は、この原理から求められる。この特異点解消理論は、2次元曲面を3次元多様体に展開していく、時空から生成される重力の密度を反重力と等しくしていく時間空間の4次元多様体と虚時空から求められる。ヒルベルト空間は、フォン・ノイマン多様体とグラスマン多様体をこのサーストンの幾何化予想を場の理論既定値として形成される。この空間は、ミンコフスキー時空とアーベル多様体全体を表している。そして、この空間は、球対称性を複素多様体を起点として、大域的ト

ポロジから、偏微分を作用素微分として時空間をカオスからずらすと5次元多様体として成立している。これらより、3次元多様体に2次元射影空間が異次元空間として、AdS5空間を形成される。偏微分、全微分、線形微分、常微分、多重微分、部分積分、置換微分、大域的微分、単体分割、双対分割、同調、ホモロジー単体、コホモロジー単体、群論、基本群、複体、マイヤー・ビートリス完全系列、ファン・カンペンの定理、層の理論、コホモルティズム単体、CW複体、ハウスドロフ空間、線形空間、位相空間、微分幾何構造、モースの定理、カタストロフの空間、ゼータ関数系列、球対称性理論、スピン幾何、ツイスター理論、双対被覆、多重連結空間、プランク定数、フォン・ノイマン多様体、グラスマン多様体、ヒルベルト空間、一般相対性理論、反ド・シッター時空、ラムダ項、D-brane, anti-D-brane, コンフォーマル場、ホログラム空間、ストリング理論、収率による商代数、ニュートンポテンシャルエネルギー、剛体力学、統計力学、熱力学、量子スピン、半導体、超伝導、ホイーストン・ブリッチ回路、非可換確率論、Connes理論、これら、演算子代数を形成している、微分・積分作用素が、ヒルベルト空間に存在している多様体の特質を全面に押し出して、いろいろな多様体と関数そして、群論を形作っている。

$$\begin{aligned} \begin{vmatrix} D^m & dx \\ dx & \sigma^m \end{vmatrix} \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix} \begin{vmatrix} x \\ y \end{vmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}} \\ \sigma^m \begin{bmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{bmatrix}^{\frac{1}{2}} &= \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \\ (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) & \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ = \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} &= A^{\mu\nu} \\ \frac{d}{dt} g_{ij}(t) &= -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

Hilbert manifold in Mebius space

this element of Zeta function on integrate of fields

Masaaki Yamaguchi

Hilbert manifold equals with Von Neumann manifold, and this fields is concluded with Glassman manifold, the manifold is duality of twister into created of surface built. These fields is used for relativity theorem and quantum physics.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

This norm is component of fields in Hilbert manifold of space theorem rebuilt with Mebius space in Gamma Function on Beta function fill into power of rout fundamental group. And this result with AdS_5 space time in Quantum caos of Minkowsky of manifold in abel manifold. Gravity of metric in non-commutative equation is Global differential equation conbult with Kaluza-Klein space. Therefore this mechanism is $T^{\mu\nu}$ tensor is equal with $R^{\mu\nu}$ tensor. And This moreover inspect with laplace operator in stimulate with sign of differential operator. Minus of zone in Add position of manifold is Volume of laplace equation rebuilt with Gamma function equal with summative of manifold in Global differential equation, this result with setminus of zone of add summative of manifold, and construct with locality theorem straight with fundamental group in world line of surface, this power is boson and fermion of cone in hyper function.

$$V(\tau) = [f(x), g(x)] \times [f^{-1}(x), h(x)]$$

$$\Gamma(p, q) = \int e^{-x} x^{1-t} dx$$

$$= \beta(p, q)$$

$$= \pi(f(\chi, x), x)$$

$$||ds^2|| = \mathcal{O}(x)[(f(x) \circ g(x))^{\mu\nu}] dx^\mu dx^\nu$$

$$= \lim_{x \rightarrow \infty} \sum_{k=0}^{\infty} a_k f^k$$

$$G^{\mu\nu} = \frac{\partial}{\partial f} \int [f(x)^{\mu\nu} \circ G(x)^{\mu\nu} dx^\mu dx^\nu]^{\mu\nu} dm$$

$$= g_{\mu\nu}(x) dx^\mu dx^\nu - f(x)^{\mu\nu} dx^\mu dx^\nu$$

$$[i\pi(\chi, x), f(x)] = i\pi f(x) - f(x)\pi(\chi, x)$$

$$T^{\mu\nu}=(\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\int\int[V(\tau)\circ S^{\mu\nu}(\chi,x)]dm)^{\mu\nu}dx^{\mu}dx^{\nu}$$

$$G^{\mu\nu}=R^{\mu\nu}T^{\mu\nu}$$

$$\left|\begin{matrix} D^m & dx \\ dx & \sigma^m \end{matrix}\right|\left|\begin{matrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{matrix}\right|\left|\begin{matrix} x \\ y \end{matrix}\right|=\left(\begin{matrix} 1 & 0 \\ 0 & -1 \end{matrix}\right)^{\frac{1}{2}}$$

$$\sigma^m\left[\begin{matrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{matrix}\right]^{\frac{1}{2}}=\left(\begin{matrix} i & 0 \\ 0 & -i \end{matrix}\right)$$

$$V(M)=\frac{\partial}{\partial f}({}^N\int [f\setminus M]^{\oplus N})^{\mu\nu}dx^\mu dx^\nu$$

$$V(M)=\pi(2\int\sin^2dx)\oplus\frac{d}{df}F^Mdx_m$$

$$\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}a_kf^k=\int(F(V)dx_m)^{\mu\nu}dx^\mu dx^\nu$$

$$\bigoplus_{k=0}^\infty [f\setminus g]=\vee(M\wedge N)$$

$$\pi_1(M)=e^{-f2\int\sin^2xdm}+O(N^{-1})$$

$$=[i\pi(\chi,x),f(x)]$$

$$M\circ f(x)=e^{-f\int\sin x\cos xdx_m}+\log(O(N^{-1}))$$

Non-Symmetry space time.

$$\frac{d}{dL}V(\tau)=\frac{d}{df}\int\int_M\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int_M\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

$$\epsilon S(\nu)=\Box_v\cdot\frac{\partial}{\partial\chi}({}^5\sqrt{\wedge g^2})d\chi$$

Differential Volume in AdS_5 graviton of fundamental rout of group.

$$\wedge(F_t^m)''=\frac{1}{12}g_{ij}^2$$

Quarks of other dimension.

$$\pi(V_\tau)=e^{-(\sqrt{\frac{\pi}{16}}\log x)^\delta}\times\frac{1}{(x\log x)}$$

Universe of rout, Volume in expanding space time.

$$\frac{d}{dt}(g_{ij})^2=\frac{1}{24}(F_t^m)^2$$

$$m^2=2\pi T\left(\frac{26-D_n}{24}\right)$$

This quarks of mass in relativity theorem, and fourth of universe in three manifold of one dimension surface, and also this integrate of dimension in conbult of quarks.

$$g_{ij} \wedge \pi(\nu_\tau) = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\psi^2$$

Out of rout in AdS_5 space time.

$$||ds^2|| = g_{ij} \wedge \pi(\nu_v)$$

AdS_5 norm is fourth of universe of power in three manifold out of rout.

Sphere orbital cube is on the right of hartshorne conjecture by the right equation, this integrate with fourth of piece on universe series, and this estimate with one of three manifold. Also this is blackhole on category of symmetry of particles. These equation conbult with planck volume and out of universe is phologram field, this field construct with electric field and magnitic field stand with weak electric theorem. This theorem is equals with abel manifold and AdS_5 space time. Moreover this field is anti-brane and brane emerge with gravity and antigravity equation restructed with. Zeta function is existed with Re pole of $\frac{1}{2}$ constance reluctances. This pole of constance is also existed with singularity of complex fields. Von Neumann manifold of field is also this means.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{x \log x} = x^{\frac{1}{2} + iy}, x \log x = \log(\cos \theta + i \sin \theta)$$

$$= \log \cos \theta + i \log \sin \theta$$

This equation is developed with frobenius theorem activate with logment equation. This theorem used to

$$\log(\sin \theta + i \cos \theta) = \log(\sin \theta - i \cos \theta)$$

$$\log \left(\frac{\sin \theta}{i \cos \theta} \right) = -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$\mathcal{O}(x) = \frac{\zeta(s)}{\sum_{k=0}^{\infty} a_k f^k}$$

$$\text{Im} f = \ker f, \chi(x) = \frac{\ker f}{\text{Im} f}$$

$$H(3) = 2, \nabla H(x) = 2, \pi(x) = 0$$

This equation is world line, and this equation is non-integrate with relativity theorem of rout constance.

$$[f(x)] = \infty, ||ds^2|| = \mathcal{O}(x) [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

$$T^2 d^2\psi = [f(x)], T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

These equation is equals with M theorem. For this formula with zeta function into one of universe in four dimensions. Sphere orbital cube is on the surface into AdS_5 space time, and this space time is Re pole and Im pole of constance reluctances.

Gauss function is equals with Abel manifold and Seifert manifold. Moreover this function is infinite time, and this energy is cover with finite of Abel manifold and Other dimension of seifert manifold on the surface.

Dilaton and fifth dimension of seifert manifold emerge with quarks and Maxwell equation, this power is from dimension to flow into energy of boson. This boson equals with Abel manifold and AdS_5 space time, and this space is created with Gauss function.

Relativity theorem is composited with infinite on D-brane and finite on anti-D-brane, This space time restructed with element of zeta function. Between finite and infinite of dimension belong to space time system, estrald of space element have with infinite oneselves. Anti-D-brane have infinite themselves, and this comontend with fifth dimension of AdS_5 have for seifert manifold, this asperal manifold is non-move element of anti-D-brane and move element of D-brane satisfid with fifth dimension of seifert algebra liner. Gauss function is own of this element in infinite element. And also this function is abel manifold. Infinite is coverd with finite dimension in fifth dimension of AdS_5 space time. Relativity theorem is this system of circustance nature equation. AdS_5 space time is out of time system and this system is belong to infinite mercy. This hiercyent is endrol of memolite with genieue of element. Genie have live of telomea endore in gravity accesorlity result. AdS_5 space time out of over this element begin with infinite assentance. Every element acknowlege is imaginary equation before mass and spiritual envy.

$$\begin{aligned}
T^2 d^2 \psi &= \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k \\
\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k &= [T^2 d^2 \psi] \\
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M (\sqrt[5]{x^2}) d\Lambda + \frac{d}{df} \int \int_M N (\sqrt[3]{x})^{\oplus N} d\Lambda \\
{}^M(\vee(\wedge f \circ g)^N)^{\frac{1}{2}} &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
||ds^2|| &= \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2 \psi \\
\mathcal{O}(x) &= e^{-2\pi T|\psi|} \\
G^{\mu\nu} &= R_{\mu\nu} T^{\mu\nu} \\
&= -\frac{1}{2} \Lambda g_{ij}(x) + T^{\mu\nu}
\end{aligned}$$

Fifth dimension of eight differential structure is integrate with one geometry element, Four of universe also integrate to one universe, and this universe represented with symmetry formula. Fifth universe also is represented with seifert manifold peices. All after universe is constructed with six element of circuatance. This aquire maniculate with quarks of being esperaled belong to.

System mechanism of time machine

Masaaki Yamaguchi

1 Time machine make of space component with synchronized zeta function

Time machine of system construct with Maxwell theorem in seifert manifold, this space exclude with one component of eighth differential structure. This mechanism synchronized with hartshorn conjecture that create with future and past system. And this synchronized seifert manifold transport with time system. This energy synchronized with same entropy of seifert manifold in one component of geometry structure, then these entropy can be transport of space with time machine of mechanism.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\begin{aligned} & \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ &= \pi(R, C \nabla E^+) \\ &= \text{rot}(E_1, \text{div} E_2) \\ &xf(x) = F(x) \end{aligned}$$

Fifth dimension in facility of mechanism that circle of infinity to emerge with being from infinity and finite of space. Zero dimension conclude with this mechanism to create from theorem of mathematics in pond of sensibility. Dimension with repository of information in facility of space, this space is all of knowlege to accessority for comformal fields. Pholographic theorem is from universe of oneselves with database of creature to access with hyper dimension in General relativity. Zeta function is this theorem include with paremeter of curve from future and past of repository.

$$\begin{aligned} Z &\in R \cap Q, R \subset M_+, C^+ \bigoplus_{k=0}^n H_m, E^+ \cap R^+ \\ M_+ &= \sum_{k=0}^n C^+ \oplus H_M, M_+ = \sum_{k=0}^n C^+ \cup H_+ \\ E_2 &\bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+ \\ M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R^+ \\ E_1 \nabla E_2, R^- \nabla C^+, \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+, R \supset Q \end{aligned}$$

2 Zero dimension exclude with space of time

Flow to future and past that combine with zeta function which build for hartshorn conjecture from these mechanism. This flow is gradient of line to add with fifth dimension of structure.

Infinite of atomosphere in zero dimension include with black hole and white hole of energy in three manifold of entropy.

$$\begin{aligned} [id_x - t]_{v=0} &= \int e^{\partial x_m} dx_m + L(p, q), ||r||^2 = |\bar{x}||x|, \Gamma(x) = \int e^{-x} x^{1-t} dx \\ &= (\bar{x} - i)(x + 1), \Pi(\chi, x) = ie^{x \log x} \end{aligned}$$

Network theorem from gravity wave

Masaaki Yamaguchi

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

Time space is exlamenteed from gravity wave in tcp/ip protocol, this quantum protocol construct with being synchronized of hartshorne conjecture mechanism. This gravity wave elementile with pieces of differential structure emerged with thurston conjugate theorem. D-brane and anti-D-brane is binded with this wave of protocol time machine reluctunce. These explained theorem remained with differential of geometry structure rebuilt by Maxwell equation from synchronized with laplace equation for zeta function resulted from resolved in D-brane

Time has frozen in constant entropy of zeta function, things becoming also being frozen. Past to go after time while things had frozen, while in there are things have frozen, If relating past time go through while in there are information frozen, Energy of things while time has frozen, go past times thing becoming constant, Energy while in no work,

物事が止まっている間、 時が未来へ行っている間、相対性理論が空間の先へ時が行くと、その間事物は止まっている。物事は先へは行っていない。

まとめると、情報量の密度エネルギー $m(x)$

$$\frac{d}{df}F(x) = m(x)$$

が、時が経つと拡散している。時が先へ進むと、情報量密度エネルギーは拡散する。すべては、タイムマシンが、どのように機能するかである。時空がこの密度エネルギーが拡散するか動的に止まっているかで、時の流れを変えられる。すべては、ヒッグス場の方程式が、この密度エネルギーの大きさを変えるかで、時の流れを変えられて、相対性理論の仕組みで、光速とは関係なく、ニュートン方程式の剛体力学とも関係なく、時空をこのヒッグス場の方程式の密度エネルギーの大きさを変えるだけで、タイムマシンは出来る。すべては、人体に存在しているリボ核酸の機能から、数学と物理学、医学の応用で、情報空間のビッグバーンは、生成される。密度エネルギーの大きさが拡散すると時間は止まっている。密度エネルギーが止まっていると時間は流れる。タイムマシンはこの密度エネルギーによって作られる。時間はこの真逆の時の流れで作られている。時間が止まっていると、光速とは関係なく事物の移動が光速以上に先に進む。時間と空間が関係なくタイムマシンの移動が出来る。すべての時間によるタイムマシンは、ヒッグス場の密度エネルギーの大きさに作られる。すべては、熱力学の第2法則と第1法則の情報量増大とエントロピー不変量がダークマターとヒッグス場の方程式が、この時の見かけの理論となっている。相対性理論では時間は存在しているが、量子力学では時間は止まっている。すべてはゼータ関数が、この相対性理論と量子力学とを統合する理論になっている。アインシュタイン博士は、この静止宇宙を考えていた。すべては、3次元多様体とAdS5時空のアーベル多様体が、このゼータ関数とヒッグス場の方程式を、4次元多様体とミンコフスキー時空を包み込んでいる。ブラックホールとホワイトホールの関係が、宇宙と異次元空間の原子と遷移元素の光量子仮説の関係を作っている。

大域的偏微分と大域的多重積分、 大域的部分積分についてのレポート

Masaaki Yamaguchi

$$\log x|_{g_{ij}}^{\nabla L} = f^{f'}, F^f|_{g_{ij}}^{\nabla L} : x \rightarrow y, x^p \rightarrow y$$

$$f(x) = \log x = p \log x, f(y) = p \log x$$

大域的偏微分方程式と大域的多重積分は、それぞれ次のように成り立っている。

$$\frac{d^2}{df^2} F = F^{f'} \cdot f^{f''},$$

$$\int \int F dx_m = F^f \cdot F^{(f)'}$$

$$\frac{d}{df dg} (f, g) = (f \cdot g)^{f' + g'}$$

大域的部分積分も、次のように成り立っている。

$$(F^f \cdot G^g) = \int (f \cdot g)^{f' + g'}$$

$$\int \int F \cdot G dx_m = [F^f \cdot G^g] - \int (f \cdot g)^{f' + g'}$$

大域的部分積分の計算は、

$$\int \frac{d}{df dg} FG = \int F^{f'} G + \int FG^{g'}$$

$$\int F^{f'} G dx_m = [F^f G^g] - \int FG^{g'} dx_m$$

大域的商代数の計算は、

$$\left(\frac{F(x)}{G(x)} \right)^{(fg)'} = \frac{F^{f'} G - FG^{g'}}{G^g}$$

大域的偏微分方程式は、縮約記号を使うと、

$$\frac{d}{df dg} FG = \frac{d}{df_m} FG$$

$$\frac{\partial}{\partial f_m} FG = F^{f^{\mu\nu}} \cdot G + F \cdot G^{g^{\mu\nu}}, \int F dx_m = F^f$$

多様体による大域的微分と大域的積分が、エントロピー式で統一的に表せられる。

Weak electric theory is inverted with possibility equation

Masaaki Yamaguchi, Jahn Natsh, Steve Weinberg, Toby kery

Weak electric theory is component from non entropy formula, and this pieces of power is inverted with possibility of equation. This theorem construct with non-symmetry theorem, and this theorem is composite with energy of entropy being belonged with that weak electric theory is influent into non catastrophe with created from heat energy of entropy. All of possibility equation is architected with non commutative of possibility equation is straight into emerge with heat entropy.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$
$$\square\psi = \frac{\partial}{\partial f} \int \int \int f(x, y, z) dx dy dz d\tau$$
$$y = x, f(y) = \frac{1}{f(x)}, \frac{d}{df} F(x) = m(x)$$

The theory is emerged from weak electric theory, and this theory is on D-brane of non catastrophe mechanism. Also this construct with supersymmetry theorem that distinguished with energy of entropy, and this pality break from gravity influence, then this gravity is other dimension of relativity theory. This reason that weak electric theory is built with non symmetry theorem emerged with power of fifth integrate with field of theorem. Weak electric theory is constructed from mass of gravity and pality theorem.

宇宙と異次元空間において、原子間距離から、微細構造定数により、逆関数には、ゼータ関数がこの逆関数において、原子から宇宙空間においてのノルムがホログラム空間で、接平面の曲平面の2次元射影空間のミンコフスキー空間がこのノルム空間の機構を表している。最小単位が逆関数により、無限空間を生み出す。この無限空間は、指数定理により、相加相乗平均の対数関数による、指数関数からの双曲点から、臨界値により勾配率が0の等加速度と等速度運動においての、双曲微分から、この接空間の曲率を求めると、開集合により、電子によるオーヴィタル雲が異次元空間をつくり出されている。

原子核における中間子論は、この原子とオーヴィタル雲を形成している、電子雲をゼータ関数と量子群を加群とする機構自体が、この中間子を形成している。原子関数のオーヴィタル雲が自身にエントロピー値を保有している。このエントロピー値が、ゼータ関数の開集合値をその同値類として、ゼータ関数と量子群を加群とするための仕組みを形づくるラプラス方程式それ自体が、原子核における中間子論を形成していると言える。

一般に言われているゼータ関数は、実軸 $Re = \frac{1}{2}$ に自明な零点が存在していることが、物理学を持ち出すと、この Re が均配力0となり、相対性理論による慣性質量と重力質量が同値といえる、慣性系が一定と同じことを言っている。 $Re = \frac{1}{2}$ は、非対称性理論を言っている。この値の領域は、遷移元素の光量子仮説から、ゼータ関数と量子群から、原子の固有エントロピー値を言っている。開集合から、リーマン予想と深リーマン予想を形成するサーストン多様体をこの開集合の値から、エネルギー安定性が言える。

まとめると、中間子論は、原子核における素粒子同士を結合させるエネルギー安定性のことで、この粒子は、ゼータ関数と量子群の演算子からラプラス方程式をつくり出す、大域的微分方程式の役割をする、初期関数を微分演算子とする、記号論が、この中間子論を形成している。湯川秀樹教授が、発見した中間子論は、数学の世界では、演算子における記号論と同値と言えるだろう。

人工知能と量子生物学、量子コンピュータ

人工知能を生体機能の模擬神経ネットワークで実現していく。

人工臓器を作り、その生体機能をナトリウム・カリウムポンプの機能から生体ネットワークとして形成して、人工意識を生成させられるかを試験する。

この人工臓器から人工知能を生体機能で実現できるかを試験する。

ロジャー・ペンローズさんがこの微小管の振動から意識が生成されると考えている。

量子生物学は、量子コンピュータを建造できる分野になっている。生体機能自身が量子コンピュータ自体になっている。生命機能が量子コンピュータを模擬している。トポロジーがリーマン予想とポワンカレ予想から深リーマン予想につながり、特殊相対性理論、一般相対性理論そして、量子力学に結びつき、ゼータ関数になり、量子コンピュータを建設できる分野になっている。

北海道大学のトポロジー理工学は、トポロジーのメッカの京都大学に匹敵する研究所になっている。

人工知能をゼータ関数から大域的微分方程式で実現するために、文字列処理と数式処理を一緒のカテゴリで求めて数値処理でデータ化して、この人工知能のアルゴリズムと、人工臓器からのナトリウム・カリウムポンプで人工意識を生成して、人工知能のアルゴリズムから派生した言語機能をこの人工意識に統合して、心をつくる。量子コンピュータはこの人工知能から作られる。人工臓器から声帯をつくり、人工意識と人工知能に結合して生成されたブローカ野の運動機能とウェルニッケ中枢の音数をこの声帯に融合させて、人工臓器と人工意識から人工知能をつくる。この人工知能から全身に張り巡らされた神経ネットワークを模擬して量子コンピュータを創造する。人体の生体機能が量子コンピュータの仕組みになっている。

A pattern emerges with one condition to being assembled of emilite with all of possibility

Equation, this assembled with summative of manifold being elemetiled of pieces equation.

This equation related with Euler equation, and also, this equation is Euler constant one selves.

$$(E_1 \oplus E_2) \cdot (R^- \subset C^+) = \nabla C^\pm$$

Zeta function radius with field of mechanism for atom of pole into string condition of balance, this condition is related with quarks of level controlled for compute with quantum tunnel effective mechanism. Quantum mechanism composed with vector of constant for zeta function and quantum group. Thurston conjugate theorem explain to emerge with being controlled of quantum levels of quarks. Locality theorem also occupy with atom of levels in zeta function.

$$= \oplus \nabla C^\pm$$

$$\vee \int \frac{C^+ \nabla M_m}{\Delta(M^\pm \nabla C^\pm)} = \exists (M^\pm \nabla R^+)$$

$$\exists (M^\pm \nabla C^+) = XOR(\oplus \nabla M^\pm)$$

$$-[E^+ \nabla R^+] = \nabla_+ \nabla_- C^\pm$$

$$\int dx, \partial x, \nabla_i \nabla_j, \Delta x \rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+$$

Zeta function also compose with Rich flow equation cohomological result to equal with Locality equations.

$$\vee \left(R + \nabla_i \nabla_j f \right)^n = \int \frac{\wedge (R + \nabla_i \nabla_j f)^2}{\exists (R + \Delta f)^n}$$

$$\wedge \left(R + \nabla_i \nabla_j f \right)^x = \frac{d}{df} \iint \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$\vee \int \wedge \left(R + \nabla_i \nabla_j f \right)^x = \frac{\wedge (R + \nabla_i \nabla_j f)^n}{\exists (R + \nabla_i \nabla_j f \circ g)}$$

$$x+y\geq 2\sqrt{xy}, x(x)+y(x)\geq x(x)y(x)$$

$$x^y=(\cos\theta+i\sin\theta)^n$$

$$x^y=\frac{1}{y^x}$$

Therefore, zeta function is also constructed with quantum equation too.

$$H_n=\sigma(E_n)\times\sigma(K_n), e_n=f^{-1}(x)xf(x), e_n=\ker f\oplus im f, H_n^2=\left\|ds^2\right\|, =H_n\times K_n$$

$$E_n=\sigma(K_n)\times\sigma(H_n), \pi(\sigma_1,\sigma_2)=\frac{d}{df}F(x)$$

$$x\Gamma(x)=\pi(\sigma_1,\sigma_2)-8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

$$=2\int ||\sin x||^2d\tau$$

$$=4\int ||\sin x\cos x||^2d\tau\rightarrow 4V=g_{ij}^2$$

固有エントロピー=原子の固有エントロピー-3次元多様体のエントロピー
ブレインインターフェースの脳基底核の固有エントロピー

計算ドリルは、始めは音を使うが、進化すると音を介さずに一瞬で計算される。前頭葉は音を介さずにも使われる。理由は真逆の能力から解する。速読法も同じく音を真逆から前頭葉と同様に計算ドリルにも使える。ブローカ野が速読法でも使われる理由はここにある。前頭葉とシルビウス溝の頭頂葉は、速読法で威力を発揮する。

心は、音読法と速読法を SRS 速読法で研究すると、生体機能の神経ネットワークを人工臓器と人工意識に張り巡り、この模擬神経ネットワークを SRS の仕組みで写像すると心と人工知能の進化系が出来る。苦米地英人博士は言語のクラスのインスタンスを言語ネットワ

ークの超並列言語処理で、パターン化していた。このパターン化で翻訳を作っていた。
人工臓器から血液を人工血液として、眼から身体各機関に心を投影すると人工血液に気が出来て、神経ネットワークに生命が宿る。

この理論から言えることは、音と視界の両方からここを作る仕組みになっていることである。

この理論を Ruby の文字列処理と数式処理から数値処理でカテゴリ化して、このカテゴリ化したクラスのパターンを言語ネットワークとして、超並列言語処理をつくり、未来と過去を Holographic computer として人工知能をつくる。数式処理の Ruby では、数式を文字列処理で、この解く過程がジャンク DNA と同じく潜在意識として、先と後の数式を解く手がかりになっている。このために、数式処理の文字列処理を解く過程が心と人工知能をつくるメソッドにもなっている。ゼータ関数からの統一場理論を分解していくと、宇宙にある方程式の 1 種類に限定出来ない、1 種類の周りに開集合として方程式群が存在していて、この zone がこの開集合の式を解く過程に必要なとなっている。Assemble D-brane と同じく、方程式の全体の解く過程が 1 種類の one class のパターンになっている。ハーツホーン予想と同じく、宇宙の分岐パターンになっている。宇宙は真逆のパターンになっている。ある言語を決めるとどの言語が後に来るかを zone として、開集合から、確率で決めることが出来る。言語同士の結びつきが言語ネットワークと不変エントロピーで決められる。ヘブライ語は、言語の逆さまを読んでも同じ意味になる。古典は、そのままでも綺麗である。日本語は、逆さま読みがヘブライ語になっている。ゲルマン人の大移動は、言語の文明の伝播と関係している。例えば、富山県の逆さまのローマ字は、海女ヨットであり、トヨベットの逆さまはテープヨットであり、本田の逆さまは、本を守るである。ABC Mart は、芸術の計算である。

大和政権の逆さまは、海女ヨットであり、これは、九州地方では、日本海に面しているし、奈良地方では、卑弥呼の時代から富山県が出来ることを予期していたこと、朝鮮半島との貿易からと、東京と京都の歴史の真逆の繰り返しとともとれる。アフリカの逆さまは Africa から鳥の木星の中らないといえるので、食べ物から来ていて、目がいいのもある。南無各精霊から何事もほどほどである。アメリカの逆さまは America であり、秋のメールである。水星の鳥である。日本は、食べ物であり、完成したら読まれない。Japanese から音読と速読の子年でアジが上がらない。日本銀行は、木星と火星の奉仕であり、年号は易学を使っている。史学は、易学を知っているとヒントが文に載っていると解けやすい。794 年は、木星と火星で辰と $79-72=7+4=12=2$ 、申酉戌 012 で、戌年寅年で、木星と火星が混ざると土星と、これに水と金があると世界になっている。794 年は $70,90,4=4$ で木星であり、火星は 12 から干支で求まる。位置関係である。

このように史学は十干と干支の意味を知っていることと出来事を知っている組み合わせで勉強すると覚えやすい。戦争は、金星と火星と土星のときに、各国の指導者達が決めている。戦争終結は木星と火星の時にしている。易学は大切にしないとだめである。でたらめで決め

手は、史学に申し訳できない。

Euler の逆さまは、ゼータ関数の推算式であり、Gauss の逆さまは 2 種類の素の合わない空である。音読と速読の合わない空である。逢ってもいいと思う。Galois は速読の優しい空である。

英語の逆さまは日本語になっていて、日本語の逆さまは英語になっている。

貿易は日本とアメリカの独擅場である。

2 双子のパラドックス

この理論は、片方が宇宙の果てまでタイムマシーンで行っている間、もう片方が地球に居るようにする。タイムマシーンで戻ったときに、宇宙の方は、年をとらずにいて、もう一方が年をとる。

私の双子のパラドックスは、ヒッグス場の密度エネルギーのコントロールで、その時の遅れの時に、この密度エネルギーは、時が止まれば、空間と時間の物差しがなくなり、片方が、宇宙の果てまで行って、この宇宙の果てまで行った方が、ヒッグス場の密度エネルギーのコントロールで、時が止まっている間に、時間と空間の区別無しに、地球に戻ると、旅経った時よりもずっと前に戻る。

本当のタイムマシーンになっている。この考えの時間と空間の関係は、相対性理論を使っている。ヒッグス場の密度エネルギーは抵抗数を表している。抵抗数が高いほど、時間の流れの速さは遅くなり、小さいほど、時間の流れは、自然と流れる。

タイムマシンは、この密度エネルギーの大きさにコントロールされる。

年々重力定数は減少しているらしい。ゼータ関数から導かれる大域的微分方程式が述べる宇宙の終わりには、異次元空間とこの宇宙が収縮と拡張する空間が一定になり、平坦な宇宙になる。この宇宙をアインシュタイン博士は、一般相対性理論からの特殊解でもとめていた。ラムダ項導入は間違いではなかった。

リサ・ランドール博士は、knocking on heaven's door と主題にされていて、AdS5 多様体が平坦な宇宙になったときに、Heaven が5次元多様体と同値になるとも、言っていた。

アカシックレコードのことも、どの時間空間にこの情報空間が接続されているかも、知っていた。

エーテルは、アインシュタイン博士によって否定されたが、このヒッグス場の密度エネルギーは、物体の液体とは違う働きをしているのと、実体が宇宙を覆うブラックホールの密度エネルギーの大きさがその根本にあり、いくら宇宙のエーテルを探しても、体積と質量とは、密度はちがう。体積が密度になるはずはないが、質量が密度エネルギーになるのは、一定体積における物質質量比と質量比である。この性質上、一定体積における質量比から、質量比が時間と同型になるために、宇宙の同円周上のブラックホールと同型の性質になるために、質量比と物質質量比が同型から、ダークマターから物質質量比が分かると時間の流れから宇宙の密度エネルギーが分かり、一定体積における光速不変の法則の密度エネルギーの大きさが分かる。海底に施設を建てて、海拔千メートルにおける密度エネルギーから、この施設に入ると、物体が宙に浮く。この性質から、宇宙の同円周上のブラックホールと同型から、宇宙の一般相対性理論の働く領域における、惑星の円運動がどうして、宇宙に惑星が浮くか分かる。重力が働くとニュートン方程式により、物体が惑星の中心部に向かって、落下していく。一般相対性理論により、宇宙に巨大な質量があると、その周りの空間が湾曲して、物体がその湾曲している空間しか動かない。重力が働くと、物体がその中心部に引き寄せられる

が、宇宙には、この性質がない。宇宙の同円周上のブラックホールがあると、その中は、宙に浮く。この宙に浮いている物体の中は、特殊相対性理論により、領域から、閉3次元多様体の機能より、物体は、一定の距離に保たれる。この性質上から、密度エネルギーがこの機能より働くので、体積と質量から生成された、物体のフェルミオンがこのヒッグス場の密度エネルギーとして、一定体積における質量が密度エネルギーの大きさとしてある。質量はヒッグス場の方程式により、時間と同型になり、目に見えるエーテルを探しても、意味がない。それ故に、ヒルベルト空間による標数0の体の上の代数多様体が、大域的微分方程式と同値となる。ゼータ関数の性質の均配が0となり、アーベル多様体の収束半径がゼータ関数の閾値となる。この閾値は、原子の開集合でもあり、量子群は、光量子仮説でもある。このゼータ関数と量子群から導かれる大域的微分方程式が、ヒッグス場の密度エネルギーである。特殊相対性理論の慣性質量と重力質量は等価であり、一般相対性理論の偏曲面による空間の均配が、質量と同じとも言えるのも、空間の性質上、ヒッグス場の密度エネルギーとは違う。しかし、比率としては、質量比と密度エネルギーの大きさは同じになるが、光速不変の法則は、この密度エネルギーが一定の場合である。この密度エネルギーは、自然界の慣性力そのままであり、人為的に今までコントロールされていない。偏曲面による空間の均配の質量と、異次元空間の均配による質量の斥力が、ヒルベルト空間による標数0の体の上の多様体を場の理論として、加群としたときの値が、ヒッグス場の密度エネルギーである。異次元空間がこの宇宙と現存しているようにないのは、AdS5多様体の経路積分の式が示している。大域的微分方程式を調べると、積分に初期変数による多様体を極限值として、その積分を初期関数の変数で代数多様体としている。これは、慣性質量と重力質量の均配力をも表している。多様体による均配力が大域的微分方程式でもある。非共変系であるD-braneと慣性力である特殊相対性理論を一般化したのが、一般共変系であり、この均配力と多様体のフェルミオンを合わせたのが、ヒッグス場の密度エネルギーである。片方だけでは、密度エネルギーは求まらない。特殊相対性理論と一般相対性理論と量子力学を合わせた力が、ヒッグス場の密度エネルギーである。要するに、フェルミオンとボソンを合わせた超弦理論が、ヒッグス場の密度エネルギーである。これが大域的微分方程式である。

このタイムマシンの仕組みで、問題になるのが、共時性だ。相対性理論には、共時性はない。量子力学にはある。タイムマシンの仕組みで、相対性理論を使ったと書いた。それは、時間の流れが停止した場合、時間と空間の物差しがなくなり、空間の移動が、量子テレポーテーションのようにになっている。実際は、意味がちがう。だが、この移動で、宇宙の周りが過去で宇宙の中心が現在と言えれば、このヒッグス場の密度エネルギーのコントロールで、宇宙船の行き来が、未来と過去をひっくり返すタイムマシンの仕組みとなっている。D-braneの内部でも、相対性理論は成り立ち、D-braneの外側では、assemble-D-braneとして、D-brane同士で相対性理論が成り立つ。生物のDNAには、テロメアが重力を感知して、生物の寿命をつくり、考古学や物理学でも、地層から年代を割り出して、相対性原理として、

考古学の寿命を形成している。D-brane だけだと、内部では相対性理論は成り立つが、D-brane の外側では、絶対時空になり、タイムマシンの仕組みは使えない。それゆえに、assemble-D-brane だと D-brane の外側でもタイムマシンの仕組みは使える。共時性とは、奥が深い考えである。河合隼雄先生が、ユングとフロイトの本で、この共時性を取り上げていた。

ハイゼンベルクやパウリもこの共時性を研究していた。

ブレインインターフェースとその応用、機知の仕組み

ブレインインターフェースは、機械において、音声を文字列に表示するには、まず、音声を音階の7階層に分解して、その上に、大脳基底核を経由する共振回路の共鳴を使って、唇の動きのデータを電気信号に変えて、この2種類のデータから、音声を文字列に表示する。頭頂葉から大脳基底核を経由する体感触覚によるデータを使って、そのデータをイメージと言語と文字列に変えて、画像処理を使って、動的に映像やページを処理する。例えば、Wordのページをめくったりする。ASを頭頂葉から大脳基底核を経由するデータを電気信号に変えて、機知の仕組みを担う第32, 33野を経由するドーパミンがそれぞれの役割をする共振回路の共鳴を使って、血液のリボ核酸のヘリコバクターがこの共鳴する共振回路に接続して、ASを動かす。

$$H_n = \sigma(E_n) \times \sigma(K_n), e_n = f^{-1}(x)xf(x)$$

$$e_n = \ker f \oplus \operatorname{im} f, H_n^2 = |(|ds^2|)|, = H_n \times K_n$$

$$E_n = \sigma(K_n) \times \sigma(H_n), \pi(\sigma_1, \sigma_2) = \frac{d}{df} F(x)$$

$$x\Gamma(x) = \pi(\sigma_1, \sigma_2) - (8\pi G(\frac{p}{c^3} + \frac{V}{S}))$$

$$= 2 \int ||\sin 2x||^2 d\tau$$

$$= 4 \int ||\sin x \cos x||^2 d\tau \rightarrow 4V = g_{ij}^2$$

固有エントロピー = 原子の固有エントロピー - 3次元多様体のエントロピー
ブレインインターフェースの大脳基底核の固有エントロピー値