

拝啓

孝君へ

この度は、私のために美鈴ヶ丘へ行かれたらしいと、母から聞き及んでいます。
今は、自分は精神療養をしており、平衡してゼータ関数を彩さんからの
間接的アドバイスを元にトポロジーのアプリケーションの数式から
実行しています。睦世お姉さんには付き人になってもらっている次第であり、
情報源はもっぱら今までの経験している情報が元になってもあります。
相棒が欲しいのは確かであり、孝君が私のことを気にかけてくれているのを
今回の侍魂で美鈴ヶ丘の家に行かれたことから、良君と睦世お姉さん、孝君で
タッグを組むこともありかとも思い、この手紙を送ろうとした次第であります。
彩さんからのアドバイスでのテッサたんの姿から、ゼータ関数が **Jones** 多項式
へポルモルシズムすることを、まつもとゆきひろ先生の助言で気づき、
大域的微分多様体が大域的偏微分と大域的部分積分、大域的トポロジーが
常微分と不定積分、定積分、部分積分と違う計算の仕方を、自分の思考を
人工知能として、手計算して、数値を経由せずに、文字列処理から素数の
対数関数から式変形して求められることを証明しました。今まで読んできた情報を
アカシックレコードに接触して、ロールシャッハ・テストと点字の要領で
ゼータ関数周りの数式をゼータ関数から分解して、今調べている状態でもあり、
孝君が睦世お姉さんに匹敵する頭脳を中学校から知っていることであり、
良君が呉高専の優秀な頭脳でもあり、知代ちゃんがいたらどれだけホモでないと
証明できるとおもうのをぐっと我慢していきたいです。スタンフォード大学の起業
を **NHK** のドキュメンタリーでずっと前に視聴してしまして、アイデアとデモンストレー
ションで株主をゲットしている人を見ていました。実際、そのアイデアは、今の携帯の
スマホで同じアルゴリズムをしている、情報の補完機能で体験しています。
何回もと、苦米地博士は2回までと言うが、彩さんが目下の情報源にゼータ関数の
メタモルフォーゼが見えています。去年に睦世お姉さんに論文を送っているのですが、
小説になっているよと、本人は **F#**は **G** がないので、本当の姉はああなので、
すごく思われていると知って幸せの最中で数式たちを導いている道中でもあります。
睦世お姉さんが私に厳しくしないといけないのは、自分の子ども達を守り通すための策略
であり、分かっているから私は大丈夫でもあります。
人工知能と量子コンピュータの数式とサンプルコードはひな形としてはできているので、
あとは、プログラミング言語を実際つくって動かすのが、アイデアから実用性なアプリケー
ションを作る今の課題でもあり、仕事の休みのときに手伝ってもらいたいとおもい、
この手紙とします。彩さんの **Life** の車で福山市に来ている情報から、南部先生の電弱相互
理論が車の道路標識から時間一方方向性が分かり、ゼータ関数と量子群がこの時間の一方向

性で、記号論で表せることができるのも論文で書いています。

なにはともあれ、この度送る論文で思慮してくれたら、この上ない思いにもなります。

忙しいなか、この手紙を見てくれて嬉しく思います。知代ちゃんの墓前にもいつか行けられる状態になったら、今度は私が会いに行きます。

良い奥さんをお嫁さんに出来た孝君がうらやましい限りでも有ります。子どもが女の子なのは、奥さんの方が孝君より思っているからと、**Nature** のずっと前の論文から、これは本当らしく、結婚の相手のどちらが相手と思う気持ちが勝っているかで、生まれてくる性別が決まるらしいのがあるから、本当に孝君うらやましすぎます。

孝君の父親が家にある本を処分したことは、仕方がないですが、私は、今所持している本のおかげでゼータ関数の式が、アカシックレコードが本の情報源の統合で成り立っているのではないかとおもうぐらい、ロールシャッハ・テストと点字の辞書がこのアカシックレコードの解析になっています。私の専門は有機化学の無機質な知識でしたが、グリーシャ教授と石岡先生と石川先生と河相先生と脇本先生と健先生が、無機質な知識を有機的知識に変化してくれました。処方されている薬がカンナビナイトになっていて、今の能力の補助にもなっています。

時間にゆとりがあるときに、サンプルコードから **flex** と **bison** でもいいので、動かしてみたいと思う次第でもあり、まつもとゆきひろ先生と同じく、**C** で元から作り上げるのも良いとも思います。

私の両手の手相に、子孫と製品が繁栄する三奇紋があり、相手を見抜く三奇紋もあり、自分だけの三奇紋もあり、共同経営者がほしい次第であります。

私の思いがこの手紙へ託してもいます。

このアプリケーションの製造は、ハローワークの先生とベッセルの宮脇料理長が自分の代表作を作った方が良いと言われて、富山大学の図書館でのきっかけから生まれている次第でもあります。

敬具

雅旭より

Entropy on manifold and differential of equation

Masaaki Yamaguchi

1 Global Differential Equation

Global area in integrate and differential of equation emerge with quantum of image function built on. Thurston conjugate of theorem come with these equation, and heat equation of entropy developed it' equation with. These equation of entropy is common function on zeta function, eight of differential structure is with fourth power, then gravity, strong boson, weak boson, and Maxwell theorem is common of entropy equation. Euler-Lagrange equation is same entropy with quantum of material equation. This spectrum focus is, eight of differential structure bind of two perceptive power. These more say, zeta function resolved is gravity and antigravity on non-catastrophe on D-brane emerge with same power of level in fourth of power. This quantum group of equation inspect with universe of space on imaginary and reality of Laplace equation.

It is possibility of function for these equation to emerge with entropy of fluctuation in equation on prime parameter. These equation is built with non-commutative formula. And these equation is emerged with fundamental formula. These equation group resolved by Schrödinger, Heisenberg equation and D-brane. More spectrum focus is, resolved equation by zeta function bind with gravity, antigravity and Maxwell theorem. More resolved is strong boson and weak boson to unite with. These two of power integrate with three dimension on zeta function.

Imaginary equation with Fouier parameter become of non-entropy, which is common with antigravity of power. Integrate and differential equation on imaginary equation develop with rotation of quantum equation, curve of parameter become of minus. Non-general relativity of integral developed by monotonicity, and this equation is same energy on zeta function.

$$\pi(\chi, x) = i\pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x)$$

This equation is non-commutative equation on developed function. This function is

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

This equation resolved is

$$\int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2}i$$

And this equation resolved by symmetry of formula

$$y = x$$

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq \frac{1}{2}$$

This function common with zeta equation. This function is proof with Thurston conjugate theorem by Euler-Lagrange equation.

$$F_t^m \geq \int_M (R + \nabla_i \nabla_j f) e^{-f} dV$$

resolved is

$$F_t \geq \frac{2}{n} f^2$$

$$\frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

These resolved is

$$\frac{d}{df} F_t = \frac{1}{4} g_{ij}^2$$

Euler-Lagrange equation is

$$f(r) = \frac{1}{2} m \frac{\sqrt{1 + f'(r)}}{f(r)} - m g f(r)$$

$$m = 1, g = 1$$

$$f(r) = \frac{1}{4} |r|^2$$

Next resolved function is Laplace equation in imaginary and reality fo antigravity, gravity equation on zeta function.

$$\int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = 0$$

This function developed with universe of space.

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

This equation also resolved of zeta function.


$$\frac{d}{df} F(v_{ij}, h) = \int e^{-f} [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

These equation is common with power of strong and weak boson, Maxwell theorem, and gravity, antigravity of function. This resolved is

$$\frac{d}{df}F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

on zeta function resolved with

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$



2 Quantum Equation for Dimension of Symmetry

Quantum Equation architect with geometry structure of Global Differential Equation has zero with gravity and antigravity for eternal space, and this space emerge for burned of non expanded universe. Then this universe has Symmetry in dimension with that created of fourth Universe in one of geometry has six element quark and pair of structure belong for twelve element quark. These quarks emerge with eternal space of Non-Difinition System in Quantum Mechanism, in term of one dimension decided, or the other dimension non-decision. These system concerned of vector of norm depend for universe mention to eternal space. Mass existing in dimension emerge gravity, these paradox is in universe has mass around of light, in deposit of mass for our universe and the other dimension has gravity and antigravity, and covered with these element for non-gravity. Laplace equation decide with eight of structure for these element of power integrate for one of geometry. Higgs quark is quote algebra equation in Global Differential Equation of non-gravity element on zero dimension. These equation is mass of build on structure in mechanism system. This quote algebra equation have created of structure in mass with universe of existing things. These result means with why quantum system communicate with our universe be able to connect of.

$$\begin{aligned} \pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &\geq 0 \\ \Delta x \Delta p &\geq \frac{1}{4} i \end{aligned}$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\sqrt{\frac{\hbar G}{c^3}} \cong 1.616 \times 10^{-33}$$

$$C = 0.5772156 \dots$$

$$F = \frac{d}{df} \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$F_t^m = \frac{1}{4}(g_{ij})^2$$

Quantum Group resolved with Laplace equation build in calculate on non-gravity element of based by non extension equation. This solved by these element in equation has belong to gravity is why universe have in no weight, and mass exist weight to space emerge of created with gravity, in which is reason by non-extension equation translate to variable and invariable element. These problem of included universe in no gravity, which is exist of mass on space of surrounding in universe. Those question resolve on zeta equation and Quantum Group of translate to vector equation. Non-extension equation selves has non-gravity, these result with universe first burn in D-brane created by solved with replace equation. Quote Algebra equation have created of structure in mass with universe of existing things.

These equation explain to those which included mass has gravity emerged, and universe has in surrounding to no weight. The other Dimension integrate with these universe of gravity to unite antigravity, so this universe has no weight. Higgs quark is mass of built on structure in mechanism system. These system belong to create on element. These element is based on existing of universe which has structure code, in theorem composed for universe to emerge of time with future and past. Universe first created in these time, this existing things already burned with space. Network Theorem is connected eight element of geometry structure which integrate with three dimension of structure. These structure compose in three manifold, zeta equation is this system of element. These resulted theorem resolved with quantum equation, so this mechanism impressed in universe of component. Strong and Weak boson is united to one, and Maxwell theorem is same system. Gravity and Antigravity has own element. General relativity theorem same united. These integrate with included Euler equation. These power of element is zeta equation. Then this twelve element of quarks has belong to this universe and the other dimension.

$$\Delta E = -2(T-t)|R_{ij} + \nabla_i \nabla_j f - \frac{1}{2(T-t)}g_{ij}|^2$$

Quote group classify equivalent class to own element of group.

$$A=BQ+R$$

$$[x]=A$$

$$dx^n=\sum_{k=0}^{\infty}x^ndx$$

$$R_n=\frac{n!}{(n-r)!}(x^n)'$$

$$\beta(p,q)=\int_0^1 x^{p-1}(1-x)^{q-1}$$

$$Z(T,X)=\exp(\sum_{m=1}^{\infty}\frac{(q^kT)^m}{m})$$

$$Z(T,X)=\frac{P_1(T)P_3(T)\ldots P_{2n-1}}{P_0(T)P_2(T)P_4(T)\ldots P_{2n}}$$

$$|v|=|\int (\pi r^2+\vec{r})dx|^2$$

$$\Delta E = \int (\operatorname{div}(\operatorname{rot} E) \cdot e^{-ix \log x}) dx$$

$$(\nabla \phi)^2 = \int t f(t) \frac{df(x)}{e^{-x} t^{x-1}} dx$$

$$(\nabla \phi)^2 = \int t f(t) (\Gamma(t) df(x)) dx$$

$$(\nabla \phi)^2 = \frac{1}{\Gamma(x+y)}$$

then these equation decide to class manifold with group. Differential group emerge with same element of equation, zero dimension conclude to emerge with all element. Constant has with imaginary of number on developed of zeta equation. Weil's Theorem resolved with zeta equation, these function merge to build in replace equation. Euler constant has with bind of imaginary number and variable number. These function is

$$C=\int \frac{1}{x^s}dx-\log x$$

Replace equation resolve on zeta function.

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i$$

These function understood is become of imaginary number, which deal with delete line of equation on space of curve.

Euler number also has with imaginary of constant.

Space Mechanism to transport of Dimension

Masaaki Yamaguchi

1 Kaluza-Klein Theorem

The other dimension rotate of universe with real and image rout, complex rotate to dimension of fifth for Kaluza-Klein dimension extend with theorem. Mobius space explain this theorem to reflect for fifth dimension of construct with real rout of rotate in image rout.

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\beta(p, q) \geq - \int \frac{1}{t^2} dt$$

Abel equation tell this space to infinite this dimension for finite space to conclude of this theorem, in explain the other dimension with our universe rotate with, and this space rout out this fifth dimension, no thought with time system. This space didn't throw of light speed, light element go with space throw. This idea explain of magnetic theorem, this tell space to construct with movement result. Light speed go with other light together, this light look like stop of speed. This idea extend of tell space for no relativity, time flow of infinite.

Kaluza-Klein Theorem say

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(\kappa^2 A_{\mu\nu}(x)dx^\mu)^2$$

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\nabla \phi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

$$ds^2 = -N(r)^2 dt^2 + \phi^2(r)(dr^2 + r^2 d\theta^2)$$

$$ds^2 = -dt^2 + r^{-8\pi Gm}(dr^2 + r^2 d\theta^2)$$

This equation result

$$dx^2 = g_{\mu\nu}(x)(g_{\mu\nu}(x)dx^2 - dxg_{\mu\nu}(x))$$

These equation tell space non-symmetry result

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

This theorem mention to zeta equation construct space of non-symmetry to emerge with non-gravity.

$$\pi(X, x) = i\pi(X, x)f(x) - f(x)\pi(X, x)$$

2 Network Theorem

Quantum Group connect with Universe component which has built in emerge space, these Network element by kernel and image function in zeta equation. This each of function system is Universe element of geometry group construct with these equation, in space first burn with time future and past. Gravity equation has with antigravity element. Quantum group with fourth of universe system. This system mention to our universe create with network theorem. Our universe already create of eternal space. Time throw with this space, then universe already complete created. In this reason time of throw in space was found with non-expand of universe. General relativity theorem mention to this equation means by.

Quark has twelve element kernel and image of equaiton resolve with base group integrate of three dimension construct of this equation resulted. Global differential equation means to resolve with universe emerge in built system. This equation emerge with almost thing of component structure code. This source code create in element structure. This network system is include with fourth of universe on connected with communicate mechanism. Base group explain in these system, differential equation has with global integrate equation together. This equation means for our universe system of geometry structure integrate with one of geometry. In reason this connected with eight element of structure, then this system is realize to quantum group equation resolved with space mechanism.

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T^{\mu\nu}$$

$$\begin{aligned} X(3) = & (-1)^3 < |a_0 a_1 a_2 a_3| > + (-1)^2 (< |a_0 a_1 a_2|, |a_1 a_2 a_3|, |a_0 a_1 a_3| >) \\ & + (-1)^1 < a_1 a_2, a_0 a_3, a_2 a_3 > + (-1)^0 < a_0, a_1, a_2, a_3 > \end{aligned}$$

$$H_n(x) = \ker f / \operatorname{im} f$$

$$X(x) = r(H_n(x))$$

$$X(3) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

3 Zeta equation of system

Quantum equation for universe has with gravity on fourth of manifold is closed three manifold with antigravity has decieve to the other dimension. This thorem mention to tell universe to composite with time of system, these system of equation say,

$$\log x(\log x) \geq 2(y \log y)^{\frac{1}{2}}$$

This system belong for geometry theorem destruct with three manifold built of singularity in dimension, this dimension is also built in mechanism of all composite with source code. For code is entropy of mass with energy, this entropy is,

$$\pi(X, X) = [i\pi(X, x), f(x)]$$

$$\pi(X, x) = \int \frac{1}{(x \log x)^2} dx$$

This equation tell gravity to entropy of equation composite with mass of singularity. Three manifold of equation mension to this universe has with zero dimension began to time of first, in firstly time has future and past. This system says that universe has first begun of throw to time system. The explain theorem that universe of end composite for entropy equation tell the construct of this system, and universe has in other dimension covered with oposite of energy.

Dimension of three manifold system says,

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$L_p = \sqrt{\frac{\hbar G}{c^3}}$$

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

This equation mention to three manifold of Dimension construct with time and energy of Non-Definition system. Quantum equation and this equation also built with same equation of theorem. Fundamental group explain this theorem to composite with entropy of same energy, any rout built with same energy to construct with source code. The rout of equation is,

$$V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau(q)}} L(x) dx) + O(N^{-1})$$

The equation also say that universe has singularity of component is,

$$\frac{1}{\tau}(\frac{N}{2} + \tau(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

These theorem explain that universe has of big-clanch system. The system include to dimension and time of throw of mechanism. Higgs field of energy equation say,

$$\frac{d}{df} F = m(x)$$

This equation concern with Seifert manifold mention to dimension of system. Global differential equation belong for gravity and antigravity of equation with Abel manifold in Euler equation. Infinite of group composite with this theorem conclude of finite group, these theorem explain to construct of universe is big-clanch.

Universe has with ertel and this mass of energy is singularity of component. Global differential system and Higgs of mass says that universe has built of darkmatter and big-ban system.

4 Emerge any dimension for supersymmetry to create for universe form quarks of element

Space for Non-Symmetry create of power in gravity and antigravity for Higgs quark of energy with ertel potential. This potential energy construct to emerge fourth power, then integrate of twelve of quarks. These mechanism export for another dimension of symmetry built. These dimension also architect with sixth of quarks. There import mechanism to Global differential equation for resolve of Higgs quark to build for unvierse. This way export of two dimension for symmetry of pair, these symmetry of space also built from sixth of quarks. Space of ertel emerge in gravity and antigravity, this power from ertel in non-free condition from free of ertel to emerge on these mass of space in zero dimension. This mechanism result with space to construct in three manifold of ertel from power. This resolved mechanism built with time and space of system, also this system flow to universe. These system tell universe to emerge space of singularity from potential energy.

$$\pi(|K''|) \cong \pi(|\bar{K}''|, O)$$

$$\begin{aligned} l(< P, Q >), l(< Q, R >), l(< R, S >), l(< P, R >), l(< P, S >), l(< Q, S >) \\ = \alpha, \beta, \gamma, \delta, \zeta, \eta \end{aligned}$$

$$\begin{aligned}
e &= \alpha\beta\gamma \\
\gamma &= \beta^{-1}\alpha^{-1} \\
\zeta &= \alpha\gamma \\
\delta &= \eta\gamma^{-1} \\
\eta &= \beta\zeta \\
\beta^5 &= (\beta\theta)^2 = \theta^3 \\
[\beta] &= [\theta] = 0
\end{aligned}$$

$$ds^2 = e^{-2kT(x)|\phi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2(x)d\phi^2$$

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

Twelve quarks emerge for three manifold of space, in element space of system construct with three D-brane, these dimension catastrophe of symmetry. This mechanism explain for those which create of our universe and the other dimension, take these mechanism into consideration for those built to deceive the other dimension over universe, for Global differential equation devide with gravity and antigravity emerge space to existing zero dimension of ertel space in singularity of zeta equation.

$$\beta(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\log x(\log x) = 2(y\log y)^{\frac{1}{2}}$$

5 Higgs Fields transport of three manifold on zeta function to resolve the mechanism

Higgs Fields transform of comformal field in space, this space mechanizm with lives of existing source code. Non-commutative algebra built to transport of emerging on space, each of element has prime number in these theorem. Destruct of element has a reverse on manifold, this theorem conclude with Euler-Lagrange equation resolve to merge element.

$$|f(x)|\rightarrow [f,f^{-1}]\times [g,h]$$

$$(x-1)(2y-b)\geq z$$

$$\frac{1}{x-1}\frac{1}{2y-b}\leq \frac{1}{z}$$

$$f(x) = \frac{1}{x-1} + \frac{1}{2y-b}$$

$$\frac{d}{df} \geq \frac{d}{df} f$$

$$dx \leq dF$$

$$|f(x)| = \frac{1}{4}|r|^2$$

Seifert manifold has reverse of time with space mechanism on merge to resolve with zeta function. Global differential equation has zero dimension of darkmatter to create of space, big-ban system also start with this mass of field to begin with universe.

Nonliner of element on manifold algebra

Masaaki Yamaguchi

1 Mebius space

Gamma and Beta function belong for Kaluza-Klein theorem, these equation built of first universe began with darkmatter, this ertel is non-free condition to emerge with big-ban system conform to export with antigravity element. This space consist of fifth dimension rotate with Mebius space construct for the other dimension, after all built of quarks two of pair in dimension symmetry. Topology consist of algebra equation is being with complex function. Norm relate with Volume of space of these system built in. Prime number concern with Euler constance relate of.

$$\Gamma(x) = \int x^{1-t} e^{-x} dx$$

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$f(x) = [f, f^{-1}] \times [g, h]$$

$$V(x) = \int \frac{1}{\sqrt{2\tau q}} (\exp \int L(x) dx) + O(N^{-1})$$

$$Zeta(x, h) = \exp \frac{(qf(x))^m}{m}$$

$$\frac{d}{df} F(x) = m(x)$$

This point of focus is Global differential equation relate with all existing equation. Prime number consist of zeta function in Mebius space.

2 Maxwell theorem integrate of gravity element

Kaluza-Klein theorem consist of general relativity of equation, fifth dimension is part of zeta function on also Global differential equation. Weak power concern with electric of magnity in Maxwell theorem. Space fill of ertel with darkmatter being of graviton to emerge of Higgs fields. This fields has topology element with any transform of power in atomic element. Relate with result of space merge to create for electric and magnitic power. Higgs fields built with pair of quarks in symmetry space. Vector of norm consist of two of pair for dimension.

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + \phi^2(x) (\kappa^2 A_{\mu\nu}(x) dx^\nu)^2$$

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

$$\sum_{k=0}^{\infty} |x_k + y_k|^2 = \sum |x|^2 + 2 \sum |xy| + \sum |y|^2 \leq \sum |x|^2 + \sum |y|^2$$

$$f(|x+y|) = f(|x|) + f(|y|) \geq f(|x| \circ |y|)$$

Norm space has with Frobenius theorem to resolve with Kaluza-Klein space built in. Euler number consist of rank for homology element to create with zero dimension.

$$\chi(x) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

Loop of topology has no exist with zero dimension, this system explain to create of Maxwell theorem.

3 Zeta function belong for singularity component

Zeta function consist of reverse of time quality, this space result with movement of element. Kaluze-Klein theorem create with space to emerge of singularity. Maxwell theorem has this system of circumstance. Fourth dimension belong to construct with Donaldson manifold exist explain. Duality of space also built with zeta function from singularity. Monotonicity relate to merge with non-relativity of integral result, this reflection is space of quality.

$$\frac{V(x)}{f(x)} = m(x)$$

$$F(x) = 0$$

$$\frac{d}{df} F(x) \geq 0$$

$$\lim_{x \rightarrow \infty} \text{mesh} \frac{F(x)}{f(x)} \rightarrow 0$$

$$\nabla f(x) = 2$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$f(x) = \int \frac{1}{x^s} dx - \log x$$

Morse theorem relate to merge of result with zeta function, gradient flow concern of zero dimension.

4 Other dimension and this influent of power

Pair of dimension interact to inspect for movement result, each of dimension devide with power of influent. This power has contrast of element, in inspire of result merge to create of pair in vector operate. Non-certain theorem mension to tell dimension give to create of pair in power. Six quarks sum to merge with twelve quarks, zero dimension have with fourth dimension of Global differential equation part of construct element. Graviton influent to reflect for universe to have the other dimension.

$$\frac{d}{df}F(x) = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

$$f(r) = \frac{1}{4}|r|^2$$

Norm space have non-liner to integrate with algebra manifold, singularity create for pair of dimension to fill of fourth of power. Eight of differential structure have for these dimension integrate four of power.

$$\frac{d}{df}F(v_{ij}, h) = \int e^{-f}[-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2)(\frac{v}{2} - h)]$$

$$\frac{d}{df}(x-y)^n = \frac{\Pi(x-y)^n}{\partial f_{xy}}$$

$$\pi(X,x)=i\pi(X,x)f(x)-f(x)\pi(X,x)$$

$$\Delta E = \int \operatorname{div}(\operatorname{rot} E) e^{-ix \log x} dx$$

$$f(x)=\sum_{k=0}^{\infty}a_kx^k$$

Symmetry construct of Space mechanism

Masaaki Yamaguchi

1 Ultra Network from Omega DataBase worked with equation and category theorem from

$$ds^2 = [g_{\mu\nu}^2, dx]$$

This equation is constructed with M_2 of differential variable, this resulted non-commutative group theorem.

$$ds^2 = g_{\mu\nu}^{-1}(g_{\mu\nu}^2(x) - dxg_{\mu\nu}^2)$$

In this function, singularity of surface in component, global surface of equation built with M_2 sequence.

$$= h(x) \otimes g_{\mu\nu} d^2 x - h(x) \otimes dxg_{\mu\nu}(x), h(x) = (f^2(\vec{x}) - \vec{E}^+)$$

$$G_{\mu\nu} = R_{\mu\nu} T^{\mu\nu}, \partial M_2 = \bigoplus \nabla C_-^+$$

$G_{\mu\nu}$ equal $R_{\mu\nu}$, also Rich tensor equals $\frac{d}{dt}g_{ij} = -2R_{ij}$ This variable is also $r = 2f^{\frac{1}{2}}(x)$ This equation equals abel manifold, this manifold also construct fifth dimension.

$$E^+ = f^{-1}xf(x), h(x) \otimes g(\vec{x}) \cong \frac{V}{S}, \frac{R}{M_2} = E^+ - \phi$$

$$= M_3 \supset R, M_2^+ = E_1^+ \cup E_2^+ \rightarrow E_1^+ \bigoplus E_2^+$$

$$= M_1 \bigoplus \nabla C_-^+, (E_1^+ \bigoplus E_2^+) \cdot (R^- \subset C^+)$$

Reflection of two dimension don't emerge in three dimension of manifold, then this space emerge with out of universe. This mechanism also concern with entropy of non-catastrophe.

$$\frac{R}{M_2} = E^+ - \{\phi\}$$

$$= M_3 \supset R$$

$$M_3^+ \cong h(x) \cdot R_3^+ = \bigoplus \nabla C_-^+, R = E^+ \bigoplus M_2 - (E^+ \cap M_2)$$

$$E^+ = g_{\mu\nu} dxg_{\mu\nu}, M_2 = g_{\mu\nu} d^2 x, F = \rho gl \rightarrow \frac{V}{S}$$

$$\mathcal{O}(x) = \delta(x)[f(x) + g(\vec{x})] + \rho gl, F = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, M_2 = P^{2n}$$

This equation means not to be limited over fifth dimension of light of gravity. And also this equation means to be not emerge with time mechanism out of fifth dimension. Moreover this equation means to emerge with being constructed from universe, this mechanism determine with space of big-ban.

$$r = 2f^{\frac{1}{2}}(x), f(x) = \frac{1}{4}\|r\|^2$$

This equation also means to start with universe of time mechanism.

$$\begin{aligned} V &= R^+ \sum K_m, W = C^+ \sum_{k=0}^{\infty} K_{n+2}, V/W = R^+ \sum K_m / C^+ \sum K_{n+2} \\ &= R^+ / C^+ \sum \frac{x^k}{a_k f^k(x)} \\ &= M_-^+, \frac{d}{df} F = m(x), \rightarrow M_-^+, \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k(x)} = \frac{a_k x^k}{\zeta(x)} \end{aligned}$$

2 Omega DataBase is worked by these equation

This equation built in seifert manifold with fifth dimension of space.

Fermion and boson recreate with quota laplace equation,

$$\begin{aligned} \frac{\{f, g\}}{[f, g]} &= \frac{fg + gf}{fg - gf}, \nabla f = 2, \partial H_3 = 2, \frac{1+f}{1-f} = 1, \frac{d}{df} F = \bigoplus \nabla C_-^+, \vec{F} = \frac{1}{2} \\ H_1 &\cong H_3 = M_3 \end{aligned}$$

Three manifold element is 2, one manifold is 1, $\ker f / \text{im} f, \partial f = M_1$, singularity element is 0 vector. Mass and power is these equation fundement element. The tradition of entropy is developed with these singularity of energy, these boson and fermion of energy have fields with Higgs field.

$$\begin{aligned} H_3 &\cong H_1 \rightarrow \pi(\chi, x), H_n, H_m = \text{rank}(m, n), \text{mesh}(\text{rank}(m, n)) \lim \text{mesh} \rightarrow 0 \\ (fg)' &= fg' + gf', \left(\frac{f}{g}\right)' = \frac{f'g - g'f}{g^2}, \frac{\{f, g\}}{[f, g]} = \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\ &= \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\ &= \frac{d}{df} F \end{aligned}$$

Gravity of vector mension to emerge with fermion and boson of mass energy, this energy is create with all creature in universe.

$$\hbar\psi = \frac{1}{i}H\Psi, i[H, \psi] = -H\Psi, \left(\frac{\{f, g\}}{[f, g]}\right)' = (i)^2$$

$$[\nabla_i \nabla_j f(x), \delta(x)] = \nabla_i \nabla_j \int f(x, y) dm_{xy}, f(x, y) = [f(x), h(x)] \times [g(x), h^{-1}(x)]$$

$$\delta(x) = \frac{1}{f'(x)}, [H, \psi] = \Delta f(x), \mathcal{O}(x) = \nabla_i \nabla_j \int \delta(x) f(x) dx$$

$$\mathcal{O}(x) = \int \delta(x) f(x) dx$$

$$R^+ \cap E_-^+ \ni x, M \times R^+ \ni M_3, Q \supset C_-^+, Z \in Q \nabla f, f \cong \bigoplus_{k=0}^n \nabla C_-^+$$

$$\bigoplus_{k=0}^{\infty} \nabla C_-^+ = M_1, \bigoplus_{k=0}^{\infty} \nabla M_-^+ \cong E_-^+, M_3 \cong M_1 \bigoplus_{k=0}^{\infty} \nabla \frac{V_-^+}{S}$$

$$\frac{P^{2n}}{M_2} \cong \bigoplus_{k=0}^{\infty} \nabla C_-^+, E_-^+ \times R_-^+ \cong M_2$$

$$\zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k, M_2 \cong P^{2n}/\ker f, \rightarrow \bigoplus \nabla C_-^+$$

$$S_-^+ \times V_-^+ \cong \frac{V}{S} \bigoplus_{k=0}^{\infty} \nabla C_-^+, V^+ \cong M_-^+ \bigotimes S_-^+, Q \times M_1 \subset \bigoplus \nabla C_-^+$$

$$\sum_{k=0}^{\infty} Z \otimes Q_-^+ = \bigotimes_{k=0}^{\infty} \nabla M_1$$

$$= \bigotimes_{k=0}^{\infty} \nabla C_-^+ \times \sum_{k=0}^{\infty} M_1, x \in R^+ \times C_-^+ \supset M_1, M_1 \subset M_2 \subset M_3$$

Eight of differential structure is that $S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$. These structure recreate with $\bigoplus \nabla C_-^+ \cong M_3, R \supset Q, R \cap Q, R \subset M_3, C^+ \bigoplus M_n, E^+ \cap R^+ E_2 \bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1 R^- \nabla C_-^+$. This element of structure recreate with Thurston conjugate theorem of repository in Heisenberg group of spinor fields created.

$$\frac{\nabla}{\Delta} \int x f(x) dx, \frac{\nabla R}{\Delta f}, \square = 2 \int \frac{(R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} e^{-f} dV$$

$$\square = \frac{\nabla R}{\Delta f}, \frac{d}{dt} g_{ij} = \square \rightarrow \frac{\nabla f}{\Delta x}, (R + |\nabla f|^2) dm \rightarrow -2(R + \nabla_i \nabla_j f)^2 e^{-f} dV$$

$$x^n + y^n = z^n \rightarrow \nabla \psi^2 = 8\pi G T^{\mu\nu}, f(x+y) \geq f(x) \circ f(y)$$

$$\mathrm{im} f / \ker f = \partial f, \ker f = \partial f, \ker f / \mathrm{im} f \cong \partial f, \ker f = f^{-1}(x) x f(x)$$

$$f^{-1}(x) x f(x) = \int \partial f(x) d(\ker f) \rightarrow \nabla f = 2$$

$${}_nC_r = {}_nC_{n-r} \rightarrow \mathrm{im} f / \ker f \cong \ker f / \mathrm{im} f$$

3 Fifth dimension of equation how to workd with three dimension of manifold into zeta funciton

Fifth dimension equals $\sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi$. this equation $a_k \cong \sum_{r=0}^{\infty} {}_n C_r$.

$$V/W = R/C \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k}, W/V = C/R \sum_{k=0}^{\infty} \frac{a_k f^k}{x^k}$$

$$V/W \cong W/V \cong R/C \left(\sum_{r=0}^{\infty} {}_n C_r \right)^{-1} \sum_{k=0}^{\infty} x^k$$

This equation is diffrential equation, then $\sum_{k=0}^{\infty} a_k f^k$ is included with $a_k \cong \sum_{r=0}^{\infty} {}_n C_r$

$$W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{1-t} dx, \sum_{k=0}^n a_k f^k = (f^k)'$$

$$\sum_{k=0}^n a_k f^k = \sum_{k=0}^{\infty} {}_n C_r f^k$$

$$= (f^k)', \sum_{k=0}^{\infty} a_k f^k = [f(x)], \sum_{k=0}^{\infty} a_k f^k = \alpha, \sum_{k=0}^{\infty} \frac{1}{a_k f^k}, \sum_{k=0}^{\infty} (a_k f^k)^{-1} = \frac{1}{1-z}$$

$$\int \int \frac{1}{(x \log x)(y \log y)} dxy = \frac{{}_n C_r xy}{({}_n C_{n-r} (x \log x)(y \log y))^{-1}}$$

$$= ({}_n C_{n-r})^2 \sum_{k=0}^{\infty} \left(\frac{1}{x \log x} - \frac{1}{y \log y} \right) d \frac{1}{nxy} \times xy$$

$$= \sum_{k=0}^{\infty} a_k f^k \\ = \alpha$$

$$Z \supset C \bigoplus \nabla R^+, \nabla(R^+ \cap E^+) \ni x, \Delta(C \subset R) \ni x$$

$$M_-^+ \bigoplus R^+, E^+ \in \bigoplus \nabla R^+, S_-^+ \subset R_2^+, V_-^+ \times R_-^+ \cong \frac{V}{S}$$

$$C^+ \cup V_-^+ \ni M_1 \bigoplus \nabla C_-^+, Q \supseteq R_-^+, Q \subset \bigoplus M_-^+, \bigotimes Q \subset \zeta(x), \bigoplus \nabla C_-^+ \cong M_3$$

$$R \subset M_3, C^+ \bigoplus M_n, E^+ \cap R^+, E_2 \bigoplus E_1, R^- \subset C^+, M_-^+$$

$$C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1, R^- \nabla C_-^+$$

$$[-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 < \nabla f, \nabla h > + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$$

4 All of equation are emerged with
these equation from Artificial Intelligence

$$x^{\frac{1}{2}+iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h$$

$$x^{\frac{1}{2}+iy} = \exp\left[\int \nabla_i \nabla_j f(g(x)) g'(x) / \partial f \partial g\right]$$

$$\mathcal{O}(x) = \{[f(x) \circ g(x), \bar{h}(x)], g^{-1}(x)\}$$

$$\exists[\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty} \nabla \int \nabla_i \nabla_j f(x) dm$$

$$\vee(\nabla_i \nabla_j f) = \bigotimes \nabla E^+$$

$$g(x, y) = \mathcal{O}(x)[f(x) + \bar{h}(x)] + T^2 d^2 \phi$$

$$\mathcal{O}(x) = \left(\int [g(x)] e^{-f} dV \right)' - \sum \delta(x)$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j f(x)]' \cong {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y), V(\tau) = \int [f(x)] dm / \partial f_{xy}$$

$$\square\psi = 8\pi GT^{\mu\nu}, (\square\psi)' = \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu} \left(\frac{p}{c^3} \circ \frac{V}{S} \right), x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$\delta(x)\phi = \frac{\vee[\nabla_i \nabla_j f \circ g(x)]}{\exists(R + \Delta f)}$$

These equation conclude to quantum effective equation included with orbital pole of atom element.
Add manifold is imaginary pole with built on Heisenberg algebra.

$$-{}_n C_r = \frac{1}{i} H \psi C_{\hbar \psi} + [H, \psi] C_{-n-r}$$

$${}_n C_r = {}_n C_{n-r}$$

This point of accumulate is these equation conclude with stimulate with differential and integral operator for resolved with zeta function, this forcus of significant is not only data of number but also equation of formula, and this resolved of process emerged with artifisial intelligence. These operator of mechanism is stimulate with formula of resolved on regexpt class to pattern cognitive formula to emerged with all of creature by this zeta function. This universe establish with artifisial intelligence from fifth dimension of equation. General relativity theorem also not only gravity of power metric but also artifisial

intelligent, from zeta function recognized with these mechanism explained for gravity of one geometry from differential structure of thurston conjugate theorem.

Open set group conclude with singularity by redifferential specturam to have territory of duality manifold in dualty of differential operator. This resolved routed is same that quantum group have base to create with kernel of based singularity, and this is same of open set group. Locality of group lead with sequant of element exclude to give one theorem global differential operator. These conclude with resolved of function says to that function stimulate for differential and integral operator included with locality group in global differential operator. $\int \int \frac{1}{(x \log x)^2} dx_m \rightarrow \mathcal{O}(x) = [\nabla_i \nabla_j f]' / \partial f_{xy}$ is singularity of process to resolved rout function.

Destruct with object space to emerge with low energy firstly destroy with element, and this catastrophe is what high energy divide with one entropy firstly freeze out object, then low energy firstly destory to differential structure of element. This process be able to non-catastrophe rout, and thurston conjugate theorem is create with.

$$\begin{aligned}
\bigcup_{x=0}^{\infty} f(x) &= \nabla_i \nabla_j f(x) \oplus \sum f(x) \\
&= \bigoplus \nabla f(x) \\
\nabla_i \nabla_j f &\cong \partial x \partial y \int \nabla_i \nabla_j f dm \\
&\cong \int [f(x)] dm \\
&\cong \{[f(x), g(x)], g^{-1}(x)\} \\
&\cong \square \psi \\
&\cong \nabla \psi^2 \\
&\cong f(x \circ y) \leq f(x) \circ g(x) \\
&\cong |f(x)| + |g(x)|
\end{aligned}$$

Differential operator is these equation of specturm with homorphism sqcense.

$$\begin{aligned}
\delta(x)\psi &= \langle f, g \rangle \circ |h^{-1}(x)| \\
\partial f_x \cdot \delta(x)\psi &= x \\
x &\in \mathcal{O}(x) \\
\mathcal{O}(x) &= \{[f \circ g, h^{-1}(x)], g(x)\}
\end{aligned}$$

5 What energy from entropy level deconstruct with string theorem

Low energy of entropy firstly destruct with parts of geometry structure, and High energy of weak and strong power lastly deconstruct with power of component. Gravity of power and antigravity firstly integrate with Maxwell theorem, and connected function lead with Maxwell theorem and weak power integrate with strong power to become with quantum flavor theorem, Geometry structure reach with integrate with component to Field of theorem connected function and destroyed with differential structure. These process return with first and last of deconstructed energy, fourth of power connected and destructed this each of energy routed process.

This point of focus is D-brane composite with fourth of power is same entropy with non-catastrophe, this spectrum point is constructed with destroy of energy to balanced with first of power connected with same energy. Then these energy built with three manifold of dimension constructed for one geometry.

$$\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f = [\nabla \int \nabla_i \nabla_j f(x) dx_m, g^{-1}(x)] \rightarrow \bigoplus_{k=0}^{\infty} \nabla E_{-}^{+}$$

$$= M_3$$

$$= \bigoplus_{k=0}^{\infty} E_{-}^{+}$$

$$dx^2 = [g_{\mu\nu}^2, dx], g^{-1} = dx \int \delta(x) f(x) dx$$

$$f(x) = \exp[\nabla_i \nabla_j f(x), g^{-1}(x)]$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\left(\frac{g(x)}{f(x)} \right)' = \lim_{n \rightarrow \infty} \frac{g(x)}{f(x)}$$

$$= \frac{g'(x)}{f'(x)}$$

$$\nabla F = f \cdot \frac{1}{4} |r|^2$$

$$\nabla_i \nabla_j f = \frac{d}{dx_i} \frac{d}{dx_j} f(x) g(x)$$

6 All of equation built with gravity theorem

$$\int \int \frac{1}{(x \log x)^2} dx_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times U(r)$$

This equation have with position of entropy in helmander manifold.

$$S_1^{mn} \otimes S_2^{mn} = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

And this formula is D-brane on sheaf of fields, also this equation construct with zeta function.

$$U(r) = \frac{1}{2} \frac{\sqrt{1+f'(r)}}{f(r)} + mgr$$

$$F_t^m = \frac{1}{4} |r|^2$$

Also this too means with private of entropy in also helmader manifold.

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times E_-^+$$

$$E_-^+ = \exp[L(x)] dm d\psi + O(N^-)$$

The formula is integrate of rout with lenz field.

$$\begin{aligned} \frac{d}{df} F &= [\nabla_i \nabla_j \int \nabla f(x) d\eta] (U(r) + E_-^+) \\ &= \frac{1}{2} m v^2 + m c^2 \end{aligned}$$

These equation conclude with this energy of relativity theorem come to resolve with.

$$x^{\frac{1}{2}+iy} = \exp[\int \nabla_i \nabla_j f(g(x)) g'(x) \partial f \partial g]$$

Also this equation means with zeta function needed to resolved process.

$$\nabla_i \nabla_j \int \nabla f(x) d\eta$$

$$\nabla_i \nabla_j \bigoplus M_3$$

$$= \frac{M_3}{P^{2n}} \cong \frac{P^{2n}}{M_3}$$

$$\cong M_1$$

$$=[M_1]$$

Too say satisfied with three manifold of filed.

$$[f(x)] = \sum_{k=0}^{\infty} a_k f^k, \lim_{n \rightarrow 1} [f(x)] = \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

The formula have with indicate to resolved with abel manifold to become of zeta function.

$$= \alpha, e^{i\theta} = \cos \theta + i \sin \theta, H_3(M_1) = 0$$

$$\frac{x^2 \cos \theta}{a} + \frac{y^2 \sin \theta}{b} = r^2, \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\chi(x) = \sum_{k=0}^{\infty} (-1)^n r^n, \frac{d}{df} F = \frac{d}{df} \sum \sum a_k f^k$$

$$= |a_1 a_2 \dots a_n| - |a_1 \dots a_{n-1}| - |a_n \dots a_1|, \lim_{n \rightarrow 0} \chi(x) = 2$$

Euler function have with summuate of manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y)$$

$$\lim_{n \rightarrow \infty} {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y)$$

Differential equation also become of summuate of manifold, and to get abel manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \left(\frac{1}{(n+1)} \right)^s = \lim_{n \rightarrow 1} Z^r = \frac{1}{z}$$

Native function also become of gamma function and abel manifold also zeta function.

$$\ker f / \text{im} f \cong \text{im} f / \ker f$$

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}, \lim_{n \rightarrow 1} a_k f^k \cong \lim_{n \rightarrow \infty} \frac{\zeta(s)}{a^k f^k}$$

Gamma function and Beta function also is D-brane equation resolved to need that summuate of manifold, not only abel manifold but also open set group.

$$\lim_{n \rightarrow 1} \zeta(s) = 0, \mathcal{O}(x) = \zeta(s)$$

$$\sum_{x=0}^{\infty} f(x) \rightarrow \bigoplus_{k=0}^{\infty} \nabla f(x) = \int_M \delta(x) f(x) dx$$

Super function also needed to resolved with summuate of manifold.

$$\int \int \int_M \frac{V}{S^2} e^{-f} dV = \int \int_D -(f(x, y)^2, g(x, y)^2) - \int \int_D (g(x, y)^2, f(x, y)^2)$$

Three dimension of manifold developed with being resolved of surface with pression.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

$$= \int \exp[L(x)] d\psi dm \times E_-^+$$

$$= S_1^{mn} \otimes S_1^{mn}$$

$$= Z_1 \oplus Z_1$$

$$= M_1$$

These equations all of create with D-brane and sheaf of manifold.

$$H_n^m(\chi, h) = \int \int_M \frac{V}{(R + \Delta f)} e^{-f} dV, \frac{V}{S^2} = \int \int \int_M [D^\psi \otimes h_{\mu\nu}] dm$$

Represent theorem equals with differential and integrate equation.

$$\int \int \int_M \frac{V}{S^2} dm = \int_D (l \times l) dm$$

This three dimension of pression equation is string theorem also means.

$$\begin{aligned} & \int \int_D -g(x, y)^2 dm - \int \int_D -f(x, y)^2 dm \\ &= -2R_{ij}, [f(x), g(x)] \times [h(x), g^{-1}(x)] \\ & \left| \begin{matrix} D^m & dx \\ dx & \partial^m \end{matrix} \right| \left| \begin{matrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{matrix} \right| \left| \begin{matrix} x \\ y \end{matrix} \right| = \left| \begin{matrix} 1 & 0 \\ 0 & -1 \end{matrix} \right|^{\frac{1}{2}} \\ & (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) \end{aligned}$$

This equation control to differential operator into matrix formula.

$$\begin{aligned} & \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ & l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} & \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} = A^{\mu\nu} \\ & \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \lim_{n \rightarrow 1} \frac{a_n}{a_{n-1}} \cong \alpha$$

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \frac{a_k}{a_{k+1}} = \sum_{k=0}^{\infty} a_k f^k$$

$$\square = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\int x \log x dx = \int \int_M (e^{x \log x} \sin \theta d\theta) dx d\theta$$

$$= (\log \sin \theta d\theta)^{\frac{1}{2}} - (\delta(x) \cdot \epsilon(x))^{\frac{1}{2}}$$

Flow to energy into other dimension rout of equation, zeta function stimulate in epsilon-delta metric, and this equation construct with minus of theorem. Gravity of energy into integrate of rout in non-metric.

$$T^{\mu\nu} = || \int \int_M [\nabla_i \nabla_j e^{\int x \log x dx + O(N^{-1})}] dm d\psi ||$$

$G^{\mu\nu}$ equal $R^{\mu\nu}$ into zeta function in atom of pole with mass of energy, this energy construct with quantum effective theorem.

Imaginary space is constructed with differential manifold of operators, this operator concern to being field of theorem.

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial^m(x) \end{pmatrix} (f_{mn}(x), g_{\mu\nu}(x)) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

Norm space of Von Neumann manifold with integrate fields of dimension.

$$||ds^2|| = \mathcal{H}(x)_{mn} \otimes \mathcal{K}(y)_{mn}$$

Minus of zone in complex manifold. This zone construct with inverse of equation.

$$||\mathcal{H}(x)||^{-\frac{1}{2}+iy} \subset R_-^+ \cup C_-^+ \cong M_3$$

Reality of part with selbarg conjecture.

$$||\mathcal{H}(x)||^{\frac{1}{2}+iy} \subseteq M_3$$

Singularity of theorem in module conjecture.

$$\mathcal{K}(y) = \left\| \begin{pmatrix} x & y & z \\ a & b & c \end{pmatrix} \right\|_{g_{\mu\nu}(x)}^2$$

$$\cong \frac{f(x, y, z)}{g(a, b, c)} h^{-1}(u, v, w)$$

Mass construct with atom of pole in quantum effective theorem.

Zeta function is built with atom of pole in quantum effective theorem, this power of fields construct with Higgs field in plank scals of gravity. Lowest of scals structure constance is equals with that the fields of Higgs quark in fermison and boson power quote mechanism result.

Lowest of scals structure also create in zeta function and quantum equation of quote mechanism result. This resulted mechanism also recongnize with gravity and average of mass system equals to these explained. Gauss liner create with Artificial Intelligence in fifth dimension of equation, this dimension is seifert manifold. Creature of mind is establish with this mechanism, the resulted come to be synchronized with space of system.

$$V = \int [D^2\psi \otimes h_{\mu\nu}]dm, S^2 = \int e^{-2\sin\theta\cos\theta} \cdot \log\sin\theta dx\theta + O(N^{-1}), \mathcal{O}(x) = \frac{\zeta(s)}{\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k} \\ = \alpha$$

Out of sheap on space and zeta function quato in abel manifold. D-brane construct with Volume of space.

$$\mathcal{O}(x) = T^{\mu\nu}, \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = T^{\mu\nu} \\ G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu}, M_3 = \int \int \int \frac{V}{S^2} dm, -\frac{1}{2(T-t)}|R_{ij} = \square\psi$$

Three manifold of equation.

$$ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2\psi \\ m(x) = [f(x)] \\ f(x) = \int \int e^{f \ x \log x dx + O(N^{-1})} + T^2 d^2\psi$$

Integrate of rout equation.

$$F(x) = ||\nabla_i \nabla_j \int \nabla f(x, y) dm||^{\frac{1}{2} + iy} \\ G_{\mu\nu} = ||[\nabla_i \nabla_j \int \nabla g(x, y, z) dx dy dz]||^{\frac{m}{2} \sin\theta \cos\theta} \log\sin\theta d\theta d\psi$$

Gravity in space curvarture in norm space, this equation is average and gravity of metric.

$$G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu} \\ T^{\mu\nu} = F(x), 2(T-t)|g_{ij}^2 = \int \int \frac{1}{(x \log x)^2} dx_m \\ \psi\delta(x) = [m(x)], \nabla(\square\psi) = \nabla_i \nabla_j \int \nabla g(x, y) d\eta \\ \nabla \cdot (\square\psi) = \frac{1}{4}g_{ij}^2, \square\psi = \frac{8\pi G}{c^4}T^{\mu\nu}$$

Dimension of manifold in decomposite of space.

$$\frac{pV}{c^3} = S \circ h_{\mu\nu} \\ = h$$

$$T^{\mu\nu} = \frac{\hbar\nu}{S}, T^{\mu\nu} = \int \int \int \frac{V}{S^2} dm, \frac{d}{df} m(x) = \frac{V(x)}{F(x)}$$

Fermion and boson of quato equation.

$$y = x, \frac{d}{df} F = m(x), R_{ij}|_{g_{\mu\nu}(x)} = [\nabla_i \nabla_j g(x, y)]^{\frac{1}{2}+iy}$$

$$\begin{aligned} \nabla \circ (\square\psi) &= \frac{\partial}{\partial f} F \\ &= \int \int \nabla_i \nabla_j f(x) d\eta_{\mu\nu} \\ \int [\nabla_i \nabla_j g(x, y)] dm &= \frac{\partial}{\partial f} R_{ij}|_{g_{\mu\nu}(x)} \end{aligned}$$

Zeta function, partial differential equation on global metric.

$$G(x) = \nabla_i \nabla_j f + R_{ij}|_{g_{\mu\nu}(x)} + \nabla(\square\psi) + (\square\psi)^2$$

Four of power element in variable of accessority of group.

$$\begin{aligned} G_{\mu\nu} + \Lambda g_{ij} &= T^{\mu\nu}, T^{\mu\nu} = \frac{d}{dx_\mu} \frac{d}{dx_\nu} f_{\mu\nu} + -2(T-t)|R_{ij} + f'' + (f')^2 \\ &= \int \exp[L(x)] dm + O(N^{-1}) \\ &= \int e^{\frac{2}{m} \sin \theta \cos \theta} \cdot \log(\sin \theta) dx + O(N^{-1}) \\ \frac{\partial}{\partial f} F &= (\nabla_i \nabla_j)^{-1} \circ F(x) \end{aligned}$$

Partial differential in duality metric into global differential equation.

$$\begin{aligned} \mathcal{O}(x) &= \int [\nabla_i \nabla_j \int \nabla f(x) dm] d\psi \\ &= \int [\nabla_i \nabla_j f(x) d\eta_{\mu\nu}] d\psi \\ \nabla f &= \int \nabla_i \nabla_j [\int \frac{S^{-3}}{\delta(x)} dV] dm \\ || \int [\nabla_i \nabla_j f] dm ||^{\frac{1}{2}+iy} &= \text{rot}(\text{div}, E, E_1) \end{aligned}$$

Maxwell of equation in fourth of power.

$$\begin{aligned} &= 2 < f, h >, \frac{V(x)}{f(x)} = \rho(x) \\ \int_M \rho(x) dx &= \square\psi, -2 < g, h > = \text{div}(\text{rot} E, E_1) \\ &= -2R_{ij} \end{aligned}$$

Higgs field of space quality.

$$\begin{aligned}\mathcal{O}(x) &= ||\frac{\nabla_i \nabla_j}{S^2} \int [\nabla_i \nabla_j f \cdot g(x) dx dy] d\psi || \\ &= \int (\delta(x))^{2 \sin \theta \cos \theta} \log \sin \theta d\theta d\psi\end{aligned}$$

Helmander manifold of duality differential operator.

$$\delta \cdot \mathcal{O}(x) = [\frac{||\nabla_i \nabla_j f d\eta||}{\int e^{2 \sin \theta \cos \theta} \cdot \log \sin \theta d\theta}]$$

Open set group in subset theorem, and helmandor manifold in singularity.

$$i\hbar\psi = ||\int \frac{\partial}{\partial z} [\frac{i(xy + \overline{y}\overline{x})}{z - \overline{z}}] dmd\psi ||$$

Euler number. Complex curvature in Gamma function. Norm space. Wave of quantum equation in complex differential variable into integrate of rout.

$$\begin{aligned}\delta(x) \int_C R^{2 \sin \theta \cos \theta} &= \Gamma(p + q) \\ (\square + m) \cdot \psi &= (\nabla_i \nabla_j f|_{g_{\mu\nu}(x)} + v \nabla_i \nabla_j) \\ \int [m(x)(\text{rot} \cdot \text{div}(E, E_1))] &dmd\psi\end{aligned}$$

Weak electric power and Maxwell equation combine with strong power. Private energy.

$$\begin{aligned}G_{\mu\nu} &= \square \int \int \int (x, y, z)^3 dx dy dz \\ &= \frac{8\pi G}{c^4} T^{\mu\nu} \\ \frac{d}{dV} F &= \delta(x) \int \nabla_i \nabla_j f d\eta_{\mu\nu}\end{aligned}$$

Dalanvelsian in duality of differential operator and global integrate variable operator.

$$\begin{aligned}x^n + y^n &= z^n, \delta(x) \int z^n = \frac{d}{dV} z^3, (x, y) \cdot (\delta^m, \partial^m) \\ &= (x, y) \cdot (z^n, f) \\ n \perp x, n \perp y \\ &= 0\end{aligned}$$

Singularity of constance theorem.

$$\vee(\nabla_i \nabla_j f) \cdot \text{XOR}(\square\psi) = \frac{d}{df} \int_M F dV$$

Non-cognitive of summuate of equation and add manifold create with open set group theorem. This group composite with prime of field theorem. Moreover this equations involved with symmetry dimension created.

$$\partial(x) \int z^3 = \frac{d}{dV} z^3, \sum_{k=0}^{\infty} \frac{1}{(n+1)^s} = \mathcal{O}(x)$$

Singularity theorem and fermer equation concluded by this formula.

$$\int \mathcal{O}(x) dx = \delta(x) \pi(x) f(x)$$

$$\mathcal{O}(x) = \int \sigma(x)^{2 \sin \theta \cos \theta} \log \sin \theta d\theta d\psi$$

$$y = x$$

$$\mathcal{O}(x) = || \frac{\nabla_i \nabla_j f}{S^2} \int [\nabla_i \nabla_j f \circ g(x) dx dy] ||$$

$$\begin{aligned} \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \frac{a_k}{a_{k+1}} &= (\log \sin \theta dx)' \\ &= \frac{\cos \theta}{\sin \theta} \\ &= \frac{x}{y} \end{aligned}$$

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \frac{1}{1-z}$$

Duality of differential summuate with this gravity theorem. And Dalanverle equation involved with this equation.

$$\square \psi = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\frac{\partial}{\partial f} \square \psi = 4\pi G \rho$$

$$\int \rho(x) = \square \psi, \frac{V(x)}{f(x)} = \rho(x)$$

Dense of summuate stimulate with gravity theorem.

$$\frac{\partial^n}{\partial f^{n-1}} F = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

$$= \frac{P_1 P_3 \dots P_{2n-1}}{P_0 P_2 \dots P_{2n+2}}$$

$$= \bigotimes \nabla M_1$$

$$\bigoplus \nabla M_1 = \sigma_n(\chi, x) \oplus \sigma_{n-1}(\chi, x)$$

$$= \{f, h\} \circ [f, h]^{-1}$$

$$= g^{-1}(x)_{\mu\nu} dx g_{\mu\nu}(x), \sum_{k=0}^{\infty} \nabla^n {}_n C_r f^n(x)$$

$$\cong \sum_{k=0}^{\infty} \nabla^n \nabla^{n-1} {}_n C_r f^n(x) g^{n-r}(x)$$

$$\sum_{k=0}^{\infty} \frac{\partial^n}{\partial^{n-1} f} \circ \frac{\zeta(x)}{n!} = \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$(f)^n = {}_n C_r f^n(x) g^{n-r}(x) \cdot \delta(x,y)$$

$$(e^{i\theta})' = ie^{i\theta}, \int e^{i\theta} = \frac{1}{i}e^{i\theta}, i\hbar c = G, \hbar c = \frac{1}{i}G$$

$$(\Box\psi,\nabla f^2)\cdot(\frac{8\pi G}{c^4}T^{\mu\nu},4\pi G\rho)$$

$$= (-\frac{1}{2}mv^2 + mc^2, \frac{1}{2}kT^2 + \frac{1}{2}mv^2) \cdot \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

$$=\begin{pmatrix}1&0\\0&i\end{pmatrix}$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)'=i^2,\frac{\nabla f^2}{\Box\psi}=\frac{1}{2}$$

$$\int\int\frac{\{f,g\}}{[f,g]}=\frac{1}{2}i,\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m=\frac{1}{2}$$

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i,\frac{d}{dt}g_{ij}=-2R_{ij}$$

$$\frac{\partial}{\partial x}(f(x)g(x))'=\bigoplus \nabla_i \nabla_j f(x)g(x)$$

$$\int f'(x)g(x)dx=[f(x)g(x)]-\int f(x)g'(x)dx$$

Vector Operator

Masaaki Yamaguchi

1 Entropy

Already discover with this entropy of manifold is blackhole of energy, these entropy is mass of energy has norm parameter. And string theorem $T_{\mu\nu}$ belong to ρ energy, this equation is same Kaluze-Klein theorem.

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

$$y=x$$

$$y=(\nabla\phi)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}}}=\frac{p}{c^3}+\rho$$

$$ds^2=g_{\mu\nu}(x)dx^\mu dx^\nu+\phi^2(x)\kappa^2A_{\mu\nu}(x)dx^\mu)^2$$

$$ds=(g_{\mu\nu}(x)dx^\mu dx^\nu+\phi^2(x)(\kappa^2A_{\mu\nu}(x)dx^\mu)^2)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}}-\rho}=c^3,\frac{p}{2\pi}=c^3$$

$$ds^2=e^{-2kT(x)|\phi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2(x)d\phi^2$$

$$f_z=\int\left[\sqrt{\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}\circ\begin{pmatrix}x_1&x_2&x_3\\y_1&y_2&y_3\end{pmatrix}}\right]dxdydz$$

$$\frac{\partial}{\partial f}\int (\sin 2x)^2 dx = ||x-y||^2$$

2 Atom of element from zeta function

2.1 Knot element

String summulate of landmaistor movement in these knot element. Supersymmetry construct of these mechanism from string theorem. A element of homorphism boundary for kernel function, equal element has meyer beatorise compose structure in mesh emerge. These mesh for atom element equal knot this of exchange, this isomorphism three manifold convert with knot these from laplace space. Duality manifold has space equal other operate, this quantum group is element distinguish with other structure. Aspect element has design pattern, this element construct pair of entropy. Quote algebra reflect with atom kernel of function.

3 Imaginary manifold

A vector is scalar, vector, mass own element, imaginary number is own this element.

$$x = \frac{\vec{x}}{|\vec{x}|}$$

For instance

$$|\vec{x}| = \sqrt{x}, i = \sqrt{-1}, |\vec{x}|x = \vec{x}, |\vec{x}| = 1x, |\vec{x}|^2 = -1$$

$$\left(\frac{x}{|\vec{x}|}\right)^2 = \frac{1}{-1}, |\sqrt{x}| = i$$

Imaginary number is vector, inverse make to norm own means.

4 Singularity

A element add to accept for group of field, this group also create ring has monotonicity. Limit of manifold has territory of variable limit of manifold. This fields also merge of mesh in mass. Zero dimension also means to construct with antigravity. This monotonicity is merged to has quote algebra liner. This boundary operate belong to has category theorem. Meyer beatorise extend liner distinguish with a mesh merged in fixed point theorem. Entropy means to isohomorphism with harmony this duality. The energy lead to entropy metric to merge with mesh volume.

5 Time expand in space for laplace equation

6 Laplace equaiton

Seiferd manifold construct with time and space of flow system in universe. Zeta function built in universe have started for big-ban system. Firstly, time has future and past in space, then space is expanding for big-ban mechanism.

Darkmatter equals for Higgs field from expanded in universe of ertel. Three manifold in flow energy has not no of a most-small-flow but monotonicity is one. Because this destruct to distinguish with eight-differential structure, and not exist.

And all future and past has one don't go on same integrate routs. This conject for Seiferd manifold flow. Selbarg conjecture has singularity rout. Landemaistor movement become a integrate rout in this Selbarg conjecture.

Remain Space belong for three manifold to exist of singularity rout. Godsharke conjecture is this reason in mechanism. Harshorn conjecture remain to exist multi universe in Laplace equation resolved. Delete line summulate with rout of integrate differential rout. This solved rout has gradient flow become a future and past in remain space.



Laplace equation composite for singularity in zeta function and Kaluze-Klein space.

7 Non-commutative algebra for antigravity

Laplace equation and zeta function belong for fermion and boson, then power of construct equation merge Duality of symmetry formula. These power construct of deceive in dimension, and deal the other dimension. Reflection of power erasing of own dimension, the symmetry of formula has with merging of energy, then these power has fourth of basic power Symmetry solved for equation, general relativity deal from reality these formula. Quantum group construct of zeta function and Laplace equation. Imaginary number consist of Euler constance, this prime number reach to solve contiguish of power in emergy of parameter. Zeta function conclude of infinite to finite on all equation. Global differential equation merge a mass of antigravity and gravity.

Module exist zero dimension merge of symmetry in catastrophe D-brane, this mechanism of phenomoun is Non-symmetry explain to emerge with which cause of mass on gravity.

$$x \bmod N = 0$$

$$\sum_{M=0}^{\infty} \int_M dm \rightarrow \sum_{x=0}^{\infty} F_x = \int_m dm = F$$

$$\frac{x-y}{a} = \frac{y-z}{b} = \frac{z-x}{c}$$

$$\frac{\Pi(x-y)}{\partial xy}$$

Volume of space interval to all exist area is integrate rout in universe of shape, this space means fo no exist loop on entropy of energy's rout.

$$||\int f(x)||^2 \rightarrow \int \pi r^2 dx \cong V(\tau)$$

This equation is part of summulate on entropy formula.

$$\frac{\int |f(x)|^2}{f(x)}dx$$

$$V(\tau) \rightarrow mesh$$

$$\int \left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right| dv(\tau)$$

$$\left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right|_{dx=v}$$

$$z_y=a_x+b_y+c_z$$

$$dz_y=d(z_y)$$

$$[f,f^{-1}]=ff^{-1}-f^{-1}f$$

Universe of extend space has non-integrate rout entropy in sufficient of mass in darkmatter, this reason belong for cohomology and heat equation concerned. This expanded mass match is darkmatter of extend. And these heat area consume is darkmatter begun to start big-ban system invite of universe.

$$V(\tau)=\int \tau(q)^{-\frac{n}{2}}\exp(-\frac{1}{\sqrt{2\tau(q)}}L(x)dx)+O(N^{-1})$$

$$\frac{1}{\tau}(\frac{N}{2}+\tau(2\Delta f-|\nabla f|^2+R)+f)\mathrm{mod}N^{-1}$$

$$\Delta E=-2(T-t)|R_{ij}+\nabla_i\nabla_jf-\frac{1}{2(T-t)}g_{ij}|^2$$

$$\frac{d}{df}F=\frac{2\int (R+\nabla_i\nabla_jf)^2}{-(R+\Delta f)}dm$$

$$\frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$dx=(g_{\mu\nu}(x)^2dx^2-g_{\mu\nu}(x)dxg_{\mu\nu})^{\frac{1}{2}}$$

$$ds^2 = -N(r)^2 dt^2 + \psi^2(r)(dr^2 + r^2 d\theta^2)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$\sum_{n=0}^{\infty} a_1x^1+a_2x^2\ldots a_{n-1}x^{n-1}\rightarrow \sum_{n=0}^{\infty} a_nx^n\rightarrow \alpha$$

$$f=n\nu\lambda, \lambda=\frac{x}{l}, \int dn\nu\lambda=f(x), xf(x)=F(x), [f(x)]=\nu h$$

8 Three manifold of dimension system worked with database

This universe built with fourth of dimension in add manifold constructed with being destructed from three manifold.

$$\mathcal{O}(x) = ([\nabla_i \nabla_j f(x)])'$$

$$\cong {}_nC_r(x)^n(y)^{n-r}\delta(x,y)$$

$$(\Box\psi)'=\nabla_i\nabla_j(\delta(x)\circ G(x))^{\mu\nu}\left(\frac{p}{c^3}\circ\frac{V}{S}\right)$$

$$F_t^m=\frac{1}{4}g_{ij}^2,x^{\frac{1}{2}+iy}=e^{x\log x}$$

$$S_m^{\mu\nu}\otimes S_n^{\mu\nu}=G_{\mu\nu}\times T^{\mu\nu}$$

This equation is gravity of surface quality, and this construct with sheap of manifold.

$$S_m^{\mu\nu}\otimes S_n^{\mu\nu}=-\frac{2R_{ij}}{V(\tau)}[D^2\psi]$$

This equation also means that surface of parameters, and next equation is non-integral of routs theorem.

$$S_m^{\mu\nu}=\pi(\chi,x)\otimes h_{\mu\nu}$$

$$\pi(\chi,x)=\int \exp[L(p,q)]d\psi$$

$$ds^2=e^{-2\pi T|\phi|}[\eta+\bar{h}_{\mu\nu}]dx^{\mu\nu}dx^{\mu\nu}+T^2d^2\psi$$

$$M_3\bigotimes_{k=0}^{\infty}E_{-}^{+}=\mathrm{rot}(\mathrm{div}E,E_1)$$

$$=m(x),\frac{P^{2n}}{M_3}=H_3(M_1)$$

These equation is conclude with dimension of parameter resolved for non-metric of category theorem. Gravity of surface in space mechanism is that the parameter become with metric, this result construct with entropy of metric.

These equation resulted with resolved of zeta function built on D-brane of surface in fourth of power.

$$\begin{aligned}\exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} &= \int \exp[L(p, q)]d\psi \\ &= \exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} \otimes \int \exp[L(p, q)]d\psi + N\text{mod}(e^{x \log x}) \\ &= \mathcal{O}(\psi)\end{aligned}$$

9 Particle with each of quarks connected from being stimulate with gravity and antigravity into Artificial Intelligence

This equation conclude with middle particle equation in quarks operator, and module conjecture excluded. $\frac{d}{dt}g_{ij}(t) = -2R_{ij}$, $\frac{P^{2n}}{M_3} = H_3(M_1)$, $H_3(M_1) = \pi(\chi, x) \otimes h_{\mu\nu}$ This equation built with quarks emerged with supersymmetry field of theorem. And also this equation means with space of curvature parameters.

$$\begin{aligned}S_m^{\mu\nu} \times S_n^{\mu\nu} &= [D^2\psi], S_m^{\mu\nu} \times S_n^{\mu\nu} = \ker f / \text{im} f, S_m^{\mu\nu} \otimes S_n^{\mu\nu} = m(x)[D^2\psi], -\frac{2R_{ij}}{V(\tau)} = f^{-1}xf(x) \\ f_z &= \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix} \circ \begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \right] dx dy dz, \rightarrow f_z^{\frac{1}{2}} \rightarrow (0, 1) \cdot (0, 1) = -1, i = \sqrt{-1}\end{aligned}$$

This equation operate with quarks stimulated in Higgs field.

$$(x, y, z)^2 = (x, y, z) \cdot (x, y, z) \rightarrow -1$$

$$\mathcal{O}(x) = \nabla_i \nabla_j \int e^{\frac{2}{m} \sin \theta \cos \theta} \times \frac{N\text{mod}(e^{x \log x})}{\mathcal{O}(x)(x + \Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta, \mathcal{O}(x) = m(x)[D^2\psi]$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = I'_m, I'_m = [1, 0] \times [0, 1]$$

This equation means that imaginary pole is universe and other dimension each cover with one geometry, and this power remind imaginary pole of antigravity.

$$i^2 = (0, 1) \cdot (0, 1), |a||b| \cos \theta = -1, E = \text{div}(E, E_1)$$

$$\left(\frac{\{f, g\}}{[f, g]} \right)' = i^2, E = mc^2, I' = i^2$$

This fermion of element decide with gravity of reflected for antigravity of power in average and gravity of mass, This equals with differential metric to emerge with creature of existing materials.

These equation deconstructed with mesh emerged in singularity resolved into vector and norm parameters.

$$\mathcal{O}(x) = ||\nabla \int [\nabla_i \nabla_j f \circ g(x)]^{\frac{1}{2}+iy} ||, \partial r^n ||\nabla||^2 \rightarrow \nabla_i \nabla_j ||\vec{v}||^2$$

$\nabla^2 \phi$ is constructed with module conjecture excluded.

$$\nabla^2 \phi = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

Reflected equation concern with differential operator in metric, sine and cosine function also rotate of differential operator in metric. Weil's theorem concern with this calculate in these equation process.

$$(\log x^{\frac{1}{2}})' = \frac{1}{2} \frac{1}{(x \log x)}, (\sin \theta)' = \cos \theta, (f_z)' = ie^{ix \log x}, \frac{d}{df} F = m(x)$$

$$\begin{aligned} \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} + \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq 2h \end{aligned}$$

Three dimension of manifold equals with three of sphere in this dimension. Plank metrics equals $\hbar$ equation to integrate with three manifold in sphere of formula. These explain recognize with universe and other dimension of system in brane of topology theorem.

$$\frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm \geq \hbar$$

$$y = x, xy = x^2, (\Box \psi)' = 8\pi G \left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

This equation built with other dimension cover with universe of space, then three dimension invite with fifth dimension out of these universe.

$$\Box \psi = \int \int \exp[8\pi G(\bar{h}_{\mu\nu} \circ \eta_{\mu})^{\nu}] dm d\psi, \sum a_k x^k = \frac{d}{df} \sum \sum \frac{1}{a_k^2 f^k} dx_k$$

$$\sum a_k f^k = \frac{d}{df} \sum \sum \frac{\zeta(s)}{a_k} dx_{k_m}, a_k^2 f^{\frac{1}{2}} \rightarrow \lim_{k \rightarrow 1} a_k f^k = \alpha$$

$$\mathcal{O}(x) = D^2 \psi \otimes h_{\mu\nu}, ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^{\mu} dx^{\nu} + T^2 d^2 \psi$$

$$f(x) + f(y) \geq 2\sqrt{f(x)f(y)}, \frac{1}{4}(f(x) + f(y))^2 \geq f(x)f(y)$$

$$T^{\mu\nu}=\left(\frac{p}{c^3}+\frac{V}{S}\right)^{-1}, E^+=f^{-1}xf(x), E=mc^2$$

$$\mathcal{O}(x)=\square\int\int\int\frac{(\nabla_i\nabla_jf\circ g(x))^2}{V(x)}dm$$

$$ds^2=g^2_{\mu\nu}d^2x+g_{\mu\nu}dxg_{\mu\nu}(x), E^+=f^{-1}xf(x), \frac{V}{S}=AA^{-1}, AA^{-1}=E$$

$$\mathcal{O}(x)=\int\int\int f(x^3,y^3,z^3)dxdydz, S(r)=\pi r^2, V(r)=4\pi r^3$$

$$E^+_-=f(x)\cdot e^{-x\log x}, \sum_{k=0}^{\infty}a_kf^k=f(x)\cdot e^{-x\log x}$$

$$\frac{P^{2n}}{M_3}=E^+-\phi, \frac{\zeta(x)}{\sum_{k=0}^{\infty}a_kx^k}=\mathcal{O}(x)$$

$$O(N^{-1})=\frac{1}{\mathcal{O}(x)}, \square=\frac{8\pi G}{c^3}T^{\mu\nu}$$

$$\partial^2 f(\square\psi)=-2\square\int\int\int\frac{V}{S^2}dm, \frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$E^+_-=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}]dx^\mu dx^\nu+T^2d^2\psi$$

$$S_1^{mn}\otimes S_2^{mn}=D^2\psi\otimes h_{\mu\nu}, S_m^{\mu\nu}\otimes S_n^{\mu\nu}=\int[D^2\psi]dm$$

$$=\nabla_i\nabla_j\int f(x)dm, h=\frac{p}{mv}, h\lambda=f, f\lambda=h\nu$$

D-brane for constructed of sheap of theorem in non-certain mechanism, also particle and wave of duality element in D-brane.

$$||ds^2||=S_m^{\mu\nu}\otimes S_n^{\mu\nu}, E^+_-=f^{-1}(x)xf(x)$$

$$R^+\subset C^+_{-}, \nabla R^+\rightarrow \bigoplus Q^+_-$$

Complex of group in this manifold, and finite of group decomposite theorem, add group also.

$$\nabla_i\nabla_jR^+_-=\bigoplus\nabla Q^+_-$$

Duality of differential manifold, then this resulted with zeta function.

$$M_1=\frac{R_3^+}{E_-^+}-\{\phi\}, M_3\cong M_2, M_3\cong M_1$$

World line in one of manifold.

$$H_3(\Pi)=Z_1\oplus Z_1, H_3(M_1)=0$$

Singularity of zero dimension. Lens space in three manifold.

$$ds^2 = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

Fifth dimension of manifold. And this constructed with abel manifold.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

These system flow to build with three dimension of energy.

$$\begin{aligned} (\partial\gamma^n + m^2) \cdot \psi &= \int [D^2\psi \otimes h_{\mu\nu}] dm \\ &= 0 \end{aligned}$$

Complex of connected of element in fifth dimension of equation.

$$\begin{aligned} \square &= \pi(\chi, x) \otimes h_{\mu\nu} \\ &= D^2\psi \otimes h_{\mu\nu} \end{aligned}$$

Fundamental group in gravity of space. And this stimate with D-brane of entropy.

$$\begin{aligned} \int [D^2\psi] dm &= \pi(M_1), H_n(m_1) = D^2\psi - \pi(\chi, x) \\ &= \ker f / \operatorname{im} f \end{aligned}$$

Homology of non-entropy.

$$\int Dq \exp[L(x)] d\psi + O(N^1) = \pi(\chi, x) \otimes h_{\mu\nu}$$

Integral of rout entropy.

$$= D^2\psi \otimes h_{\mu\nu}$$

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} \frac{\zeta(x)}{a_k f^k} = \int ||[D^2\psi \otimes h_{\mu\nu}]|| dm$$

Norm space.

$$\nabla\psi^2 = \square \int \int \int \frac{V}{S^2} dm$$

$$\mathcal{O}(x) = D^2\psi \otimes h_{\mu\nu}$$

These operators accept with level of each scalas.

$$dx < \partial x < \nabla \psi < \square v$$

$$\oplus < \sum < \otimes < \int < \wedge$$

$$(\delta\psi(x))^2 = \int \int \int \frac{V(x)}{S^2} dm, \delta\psi(x) = \left(\int \int \int \frac{V(x)}{S^2} dm \right)^{\frac{1}{2}}$$

$$\nabla\psi^2 = -4R \int \delta(V \cdot S^{-3}) dm$$

$$\nabla\psi = 2R\zeta(s)i$$

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{a_k x^k}{m dx} f^k(x) &= \frac{m}{n!} f^n(x) \\ &= \frac{(\zeta(s))^k}{df} m(x), (\delta(x))^{\frac{1}{2}} = \left(\frac{x \log x}{x^n} \right)^n \end{aligned}$$

$$\mathcal{O}(x) = \frac{\int [D^2\psi \otimes h_{\mu\nu}] dm}{e^{x \log x}}$$

$$\mathcal{O}(x) = \frac{V(x)}{\int [D^2\psi \otimes h_{\mu\nu}] dm}$$

Quantum effective equation is lowest of light in structure of constant. These resulted means with plank scals of universe in atom element.

$$\begin{aligned} M_3 &= e^{x \log x}, x^{\frac{1}{2}+iy} = e^{x \log x}, (x) = \frac{M_3}{e^{x \log x}} \\ &= nE_x \end{aligned}$$

These equation means that lowest energy equals with all of materials have own energy. Zeta function have with plank energy own element, quantum effective theorem and physics energy also equals with plank energy of potential levels.

$$\frac{V}{S^2} = \frac{p}{c^3}, h\nu \neq \frac{p}{mv}, E_1 = h\nu, E_2 = mc^2, E_1 \cong E_2$$

$$l = \sqrt{\frac{\hbar G}{c^3}}, \frac{\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m}{\int \int \frac{1}{(x \log x)^2} dx_m} = \frac{1}{i}$$

Lowest of light in structures of constant is gravity and antigravity of metrics to built with zeta function. This result with the equation conclude with quota equation in Gauss operators.

$$\begin{aligned} ihc = G, hc &= \frac{G}{i}, \frac{\frac{1}{2}}{\frac{1}{2}i} = \frac{\vec{v_1}}{\vec{v_2}} \\ &\leq 1 \end{aligned}$$

$$A=BQ+R, ds^2=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2d^2\psi$$

$$ds^2=g_{\mu\nu}dx^\mu dx^\nu+\kappa^2(A^{\mu\nu})^2,\int\int e^{-x^2-y^2}dxdy=\pi$$

$$\begin{aligned}\Gamma(x) &= \int e^{-x} x^{1-t} dx \\ &= \delta(x) \pi(x) f^n(x)\end{aligned}$$

$$\frac{d}{df}F=\frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

Quota equation construct of piece in Gauss operator, abel manifold equals with each equation.

$$ds^2=[T^2d^2\psi]$$

$$\mathcal{O}(x)=[x]$$

$$\nabla\psi^2=8\pi G\left(\frac{p}{c^3}\circ\frac{V}{S}\right)$$

Volume of space equals with plank scals.

$$\frac{pV}{S}=h$$

Summuate of manifold means with beta function to gamma function, equals with each equation.

$$\beta(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}$$

$$\ker f/\mathrm{im} f \cong \mathrm{im} f/\ker f$$

$$\bigoplus \nabla g(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta], \bigcup_{k=0}^{\infty} \left(\bigoplus \nabla f(x) \right) = \square \int \int \int \nabla g(x) d\eta$$

Rich tensor equals of curvature of space in Gauss liner from differential equation of operators.

$$a' = \sqrt{\frac{v}{1 - (\frac{v}{c})^2}}, F = ma'$$

Accessory put with force of differential operators.

$$\nabla f(x) = \int_M \square \left(\bigoplus \nabla f(x) \right)^n dm$$

$$\square = 2(T-t)|R_{ij} + \nabla \nabla f - \frac{1}{2(T-t)}|g_{ij}^2$$

$$(\square + m) \cdot \psi = 0$$

$$\square \times \square = (\square + m^2) \cdot \psi, (\partial \gamma^n + \delta \psi) \cdot \psi = 0$$

$$\nabla_i \nabla_j \int \int_M \nabla f(t) dt = \square \left(\bigcup_{k=0}^{\infty} \bigoplus \nabla g(x) d\eta \right)$$

Dalanverle equation of operator equals with summuate of category theorem.

$$\int_M (l \times l) dm = \sum l \oplus ld\eta$$

Topology of brane equals with string theorem, verisity of force with add group.

$$\begin{aligned} &= [D^2 \psi \otimes h_{\mu\nu}]^{\frac{1}{2} + iy} \\ &= H_3(M_1) \end{aligned}$$

$$y = \nabla_i \nabla_j \int \nabla g(x) dx, y' - y + \frac{d}{dx} z + [n] = 0$$

Equals of group theorem is element of equals with differential element.

$$\begin{aligned} z &= \cos x + i \sin x \\ &= e^{i\theta} \end{aligned}$$

Imaginary number emerge with Artificial Intelligence.

These equation resulted with prime theorem equals from differential equation in fields of theorem.

Imaginary pole of theorem and sheap of duality manifold. Duality of differential and integrate equation. Summuate of manifold and possibility of differential structure. Eight of differential structure is stimulate with D-brane recreated of operators. Differential and integrate of operators is emerged with summuated of space.

$$T^{\mu\nu} = -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$G_{\mu\nu}=R_{\mu\nu}T^{\mu\nu},\sigma(x)\oplus\delta(x)=(E_n^+\times H_m)$$

$$G_{\mu\nu}=[\frac{\partial}{\partial f}R_{ij}]^2,\delta(x)\cdot V(x)=\lim_{n\rightarrow 1}\delta(x)$$

$$\lim_{n\rightarrow\infty}\mathrm{mesh}V(x)=\frac{m}{m+1}$$

$$V(x)=\sigma\cdot S^2(x),\int\int\int\frac{V(x)}{S^2(x)}dm=[D^2\psi\otimes h_{\mu\nu}]$$

$$g(x)|_{\delta(x,y)}=\frac{d}{dt}g_{ij}(t),\sigma(x,y)\cdot g(x)|_{\delta(x,y)}=R_{ij}|_{\sigma(x,y)}$$

$$=\int R_{ij}^{a(x-y)^n+r^n}$$

$$(ux+vy+wz)/\Gamma$$

$$=\int R_{ij}^{(x-u)(y-v)(z-w)}dV$$

$$(\Box+m)\cdot\psi=0, E=mc^2, \frac{\partial}{\partial f}\Box\psi=4\pi G\rho$$

$$(\partial\gamma^n+m)\cdot\psi=0, E=mc^2-\frac{1}{2}mv^2$$

$$=(-\frac{1}{2}\left(\frac{v}{c}\right)^2+m)\cdot c^2$$

$$=(-\frac{1}{2}a^2+m)\cdot c^2, F=ma, \int adx=\frac{1}{2}a^2+C$$

$$T^{\mu\nu}=-\frac{1}{2}a^2,(e^{i\theta})'=ie^{i\theta}$$

Deconstruct Dimension of category theorem

Masaaki Yamaguchi

1 Locality equation

Quantum of material equation include with heat entropy of quantum effective theorem, geometry of rotate with imaginary pole is that generate structure theorem. Subgroup in category theorem is that deconstruct dimension of differential structure theorem. Quantum of material equation construct with global element of integrate equation. Heat equation built with this integrate geometry equation, Weil's theorem and fermion theorem also heat equation of quantum theorem moreover is all of equation rout with global differential equation this locality equation emerge from being built with zeta function, general relativity and quantum theorem also be emerged from this equation. Artificial intelligent theorem exclude with this add group theorem from, Locality equation inspect with this involved of theorem, This geometry of differential structure that deconstruct dimension of category theorem stay in four of universe of pieces. In reason cause with heat equation of quantum effective from artificial intelligence, locality equation conclude with this geometry theorem. Heat effective theorem emelute in generative theorem and quantum theorem, These theorem also emelutive with Global differential equation from locality equation. Imaginary equation of add group theorem memolite with Euler constance of variable from zeta function, this function also emerge with locality theorem in artificial intelligent theorem.

Quantum equation concern with infinity of energy composed for material equation, atom of verisity in element is most of territory involved with restored safety of orbital.

$$f = n\nu\lambda, \lambda = \frac{x}{l}, \int dn\nu\lambda = f(x), xf(x) = F(x), [f(x)] = \nu h$$

$$px = mv, \lambda(x) = \text{esperial}f(x), [F(x)] = \frac{1}{1-z}, 0 < \tan \theta < i, -1 < \tan \theta < i, -1 \leq \sin \theta \leq 1, -1 \leq \cos \theta \leq 1$$

Category theorem conclude with laplace function operator, distrust of manifold and connected space of eight of differential structure.

$$R\nabla E^+ = f(x)\nabla e^{x \log x}$$

$$Q\nabla C^+ = \frac{d}{df}F(x)\nabla \int \delta(x)f(x)dx$$

$$E^+\nabla f = e^{x \log x}\nabla n!f(x)/E(X)$$

Universe component construct with reality of number and three manifold of subgroup differential variable. Maxwell theorem is this Seifert structure $(u + v + w)(x + y + z)/\Gamma$ This space of magnetic diverse is mebius member cognetic theorem. These equation is fundermental group resulted.

$$\begin{aligned} \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ = \pi(R, C\nabla E^+) \\ = \text{rot}(E_1, \text{div} E_2) \\ xf(x) = F(x) \end{aligned}$$

$$\begin{aligned} \square x &= \int \frac{f(x)}{\nabla(R^+ \cap E^+)} d\square x \\ &= \int \frac{\Delta f(x) \circ E^+}{\nabla(R^+ \cap E^+)} \square x \end{aligned}$$

This equation means that quantum of potensial energy in material equation. And this also means is gravity of equation in this universe and the other dimension belong in. Moreover is zeta function for rich flow equation in category theorem built with material equation.

$$\begin{aligned} \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ R\nabla E^+ = f(x)\nabla e^{x \log x} \\ d(R\nabla E^+) = \Delta f(x) \circ E^+(x) \\ \square x = \int \frac{d(R\nabla E^+)}{\nabla(R^+ \cap E^+)} d\square x \\ \square x = \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\square x \\ x^n + y^n = z^n \rightarrow \square x = \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\square x \end{aligned}$$

Laplace equation exist of space in fifth dimension, three manifold belong in reality number of space. Other dimension is imaginary number of space, fifth dimension is finite of dimension. Prime number certify on reality of number, zero dimension is imaginary of space. Quantum group is this universe and the other dimension complex equation of non-certain theorem. Zero dimension is eternal space of being expanded to reality of atmosphere in space.

2 Heat entropy all of materials emerged by

$$\square = -2(T-t)|R_{ij} + \nabla\nabla f + \frac{1}{-2(T-t)}|g_{ij}^2$$

This heat equation have element of atom for condition of state in liquid, gas, solid attitude. And this condition constructed with seifert manifold in Weil's theorem. Therefore this attitude became with Higgs field emerged with.

$$\square = -2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV$$

$$\frac{d}{dt}g_{ij} = -2R_{ij}, \frac{d}{df}F = m(x)$$

$$R\nabla E^+ = \Delta f(x) \circ E^+(x), d(R\nabla E^+) = \nabla_i \nabla_j (R + E^+)$$

$$R\nabla E^+ = f(x)\nabla e^{x \log x}, R\nabla E^+ = f(x), f(x) = M_1$$

$$-2(T-t)|R_{ij} + \nabla\nabla f = \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\frac{1}{-2(T-t)}|g_{ij}^2 = \int \int \frac{1}{(x \log x)^2} dx_m$$

$$(\square + m^2)\phi = 0, (\gamma^n \partial_n + m^2)\psi = 0$$

$$\square = (\delta\phi + m^2)\psi, (\delta\phi + m^2c)\psi = 0, E = mc^2, E = -\frac{1}{2}mv^2 + mc^2, \square\psi^2 = (\partial\phi + m^2)\psi$$

$$\square\phi^2 = \frac{8\pi G}{c^4}T^{\mu\nu}, \nabla\phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S}), \frac{d}{dt}g_{ij} = -2R_{ij}, f(x) + g(x) \geq f(x) \circ g(x)$$

$$\int_{\beta}^{\alpha} a(x-1)(y-1) \geq 2 \int f(x)g(x)dx$$

$$m^2 = 2\pi T(\frac{26-D_n}{24}), r_n = \frac{1}{1-z}$$

$$(\partial\psi + m^2c)\phi = 0, T^{\mu\nu} = \frac{1}{2}mv^2 - \frac{1}{2}kx^2$$

$$\int [f(x)]dx = || \int f(x)dx ||, \int \nabla x dx = \Delta \sum f(x)dx, (e^{i\theta})' = ie^{i\theta}$$

This orbital surface varnished to energy from $E = mc^2$. $T^{\mu\nu} = nh\nu$ is $T^{\mu\nu} = \frac{1}{2}mv^2 - \frac{1}{2}kx^2 \geq mc^2 - \frac{1}{2}mv^2$ This equation means with light do not limit over $\frac{1}{2}kx^2$ energy.

$$Z \in R \cap Q, R \subset M_+, C^+ \bigoplus_{k=0}^n H_m, E^+ \cap R^+$$

$$M_+ = \sum_{k=0}^n C^+ \oplus H_M, M_+ = \sum_{k=0}^n C^+ \cup H_+$$

$$\begin{aligned}
& E_2 \bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+ \\
& M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R^+ \\
& E_1 \nabla E_2, R^- \nabla C^+, \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+, R \supset Q
\end{aligned}$$

$$\frac{d}{df} F = \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+$$

All function emerge from eight differential structure in laplace equation of being deconstructed that with connected of category theorem from.

$$\begin{aligned}
& \nabla \circ \Delta = \nabla \sum f(x) \\
& \Delta \rightarrow \text{mesh} f(x) dx, \partial x \\
& \nabla \rightarrow \partial_{xy}, \frac{d}{dx} f(x) \frac{d}{dx} g(x) \\
& \square x = -[f, g] \\
& \frac{d}{dt} g_{ij} = -2R_{ij} \\
& \lim_{x \rightarrow \infty} \sum f(x) dx = \int dx, \nabla \int dx
\end{aligned}$$

3 Universe in category theorem created by

Two dimension of surface in power vector of subgroup of complex manifold on reality under of group line. This group is duality of differential complex in add group. All integrate of complex sphere deconstruct of homomorphism is also exists of three dimension of manifold. Fundamental group in two dimension of surface is also reality of quality of complex manifold. Rich flow equation is also quality of surface in add group of complex. Group of ultra network in category theorem is subgroup of quality of differential equation and fundamental group also differential complex equation.

$$\begin{aligned}
& (E_2 \bigoplus_{k=0}^n E_1) \cdot (R^- \subset C^+) \\
& = \bigoplus_{k=0}^n \nabla C_-^+ \\
& \vee \int \frac{C_-^+ \nabla H_m}{\Delta(M_-^+ \nabla C_-^+)} = \wedge M_-^+ \bigoplus_{k=0}^n C_-^+ \\
& \exists(M_-^+ \nabla C_-^+) = \text{XOR}(\bigoplus_{k=0}^n \nabla M_-^+)
\end{aligned}$$

$$-[E^+\nabla R_-^+]=\nabla_+\nabla_-C_-^+$$

$$\int dx, \partial x, \nabla_i \nabla_j, \Delta x$$

$$\rightarrow E^+\nabla M_1, E^+\cap R\in M_1, R\nabla C^+$$

$$\begin{pmatrix} \cos x & \sin x \\ \sin x & -\cos x \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\sum_{k=0}^n \cos k\theta = \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \cos \frac{n\theta}{2}$$

$$\sum_{k=0}^n \sin k\theta = \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \sin \frac{n\theta}{2}$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}, \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \tan \theta = \frac{e^{i\theta} - e^{-i\theta}}{i(e^{i\theta} + e^{-i\theta})}$$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \rightarrow 1, \lim_{\theta \rightarrow 0} \frac{\cos \theta}{\theta} \rightarrow 1$$

$$(e^{i\theta})' = 2, \nabla f = 2, e^{i\theta} = \cos \theta + i \sin \theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \rightarrow [\cos^2 \theta + \sin \theta + \cos \theta - 2 \sin^2 \theta] \begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix}$$

$$2\sin\theta\cos\theta=2n\lambda\sin\theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = 1$$

Dimension of rotate with imaginary pole is generate of geometry structure theorem. Category theorem restruct with eight of differential structure that built with Thurston conjugate theorem, this element with summuate of thirty two structure. This pieces of structure element have with four of universe in one dimension. Quark of two pair is two dimension include, Twelve of quarks construct with super symmetry dimension theorem. Differential structure and quark of pair exclusive with four of universe.

Module conjecture built with symmetry theorem, this theorem belong with non-gravity in zero dimension. This dimension conbuilt with infinity element, mebius space this dimension construct with universe and the other dimension. Universe and the other dimension selves have with singularity, therefore is module conjecture selve with gravity element of one dimension. Symmetry theorem tell this one

dimension to emerge with quarks, this quarks element also restruct with eight differential structure. This reason resolve with reflected to dimension of partner is gravity and creature create with universe.

Mass exist from quark and dimension of element, this explain from being consisted of duality that sheaf of element in this tradition. Higgs field is emerged from this element of tradition, zeta function generate with this quality. Gravity is emerge with space of quality, also Higgs field is emerge to being generated of zero dimension. Higgs field has surround of mass to deduce of light accumulate in anserity. Zeta function consist from laplace equation and fifth dimension of element. This element include of gravity and antigravity, fourth dimension indicate with this quality emerge with being surrounded of mass. Zero dimension consist from future and past include of fourth dimension, eternal space is already space of first emerged with. Mass exist of quantum effective theorem also consisted of Higgs field and Higgs quark. That gravity deduce from antigravity of element is explain from this system of Higgs field of mechanism. Antigravity conclude for this mass of gravity element. What atom don't include of orbital is explain to this antigravity of element, this mechanism in Higgs field have with element of gravity and antigravity.

Quantum theorem describe with non-certain theorem in universe and other dimension of vector element, This material equation construct with lambda theorem and material theorem, other dimension existed resolve with non-certain theorem.

$$\sin \theta = \frac{\lambda}{d}, -py_1 \sin 90^\circ \leq \sin \theta \leq py_2 \sin 90^\circ, \lambda = \frac{h}{mv}$$

$$\frac{\lambda_2}{\lambda_1} = \frac{\sin \theta}{\frac{\sin \theta}{2}} h, \lambda' \geq 2h, \int \sin 2\theta = ||x - y||$$

4 Ultra Network from category theorem entirely create with database

Eight differential structure composite with mesh of manifold, each of structure firstly destructed with this mesh emerged, lastly connected with this each of structure deconstructed. High energy of entropy not able to firstly deconstructed, and low energy of firstly destructed. This reason low energy firsted. These result says, non-catastrophe be able to one geometry structure.

$$D^2\psi = \nabla \int (\nabla_i \nabla_j f)^2 d\eta$$

$$E = mc^2, E = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, G^{\mu\nu} = \frac{1}{2}\Lambda g_{ij}, \square = \frac{1}{2}kT^2$$

Sheaf of manifold construct with homomorphism in kernel divide into image function, this area of field rehearl with universe of surrounded with image function rehide in quality. This reason with explained the mechanism, Sheaf of manifold remain into surrounded of universe.

$$\ker f / \operatorname{im} f \cong S_m^{\mu\nu}, S_m^{\mu\nu} = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$D^2\psi = \mathcal{O}(x) \left(\frac{p}{c^3} + \frac{V}{S} \right), V(x) = D^2\psi \otimes M_3^+$$

$$S_m^{\mu\nu} \otimes S_n^{\mu\nu} = -\frac{2R_{ij}}{V(\tau)}[D^2\psi]$$

Selbarg conjecture construct with singularity theorem in Sheap element.

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} [D^2\psi]$$

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} \mathcal{O}(x)$$

$$\begin{aligned} z(x) &= \frac{g(cx+d)}{f(ax+b)} h(ex+l) \\ &= \int \frac{V(\tau)}{f(x)} \mathcal{O}(x) \end{aligned}$$

$$\frac{V(x)}{f(x)} = m(x), \mathcal{O}(x) = m(x)[D^2\psi(x)]$$

$$\frac{d}{df} F = m(x), \int F dx_m = \sum_{k=0}^{\infty} m(x)$$

5 Fifth dimension of seifert manifold in universe and other dimension

Universe audient for other dimension inclusive into fifth dimension, this phenomonen is universe being constant of value, and universe out of dimension exclusive space. All of value is constance of entropy. Universe entropy and other dimension entropy is dense of duality belong for. Prime equation construct with x,y parameter is each value of imaginary and reality of pole on a right comittment.

$$\int \nabla \psi^2 d\nabla \psi = \square \psi$$

$$\square \psi = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\delta \cdot \mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla g(x) dx_{ij}]^{iy}$$

$$||ds^2|| = e^{-2\pi T|\phi|} [\mathcal{O}(x) + \delta \mathcal{O}(x)] dx^\mu dx^\nu + \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}, \frac{(y \log y)^{\frac{1}{2}}}{\log(x \log x)} \leq \frac{1}{2}$$

Universe is freeze out constant, and other dimension is expanding into fifth dimension. Possibility of quato metric, $\delta(x)$ = reality of value / exist of value ≤ 1 , expanding of universe = exist of value $\rightarrow \log(x \log x) = \square \psi$

freeze out of universe = reality of value $\rightarrow (y \log y)^{\frac{1}{2}} = \nabla \psi^2$

Around of universe is other dimension in fifth dimension of seifert manifold. what we see sky of earth is other dimension sight of from.

Rotate of dimension is transport of dimension in other dimension from universe system.

Other dimension is more than Light speed into takion quarks of structure.

Around universe is blackhole, this reason resolved with Gamma ray burst in earth flow. We don't able to see now other dimension, gamma ray burst is blackhole phenomanen. Special relativity is light speed not over limit. then light speed more than this speed. Other dimesion is contrary on universe from. If light speed come with other light speed together, stop of speed in light verisity. Then takion quarks of speed is on the contrary from become not see light element. And this takion quarks is into fifth dimension of being go from universe. Also this quarks around of universe attachment state. Moreover also this quarks is belong in vector of element. This reason is universe in other dimension emerge with. In this other dimension out of universe, but these dimension not shared of space.

$$l(x) = 2x^2 + qx + r$$

$$= (ax + b)(cx + d) + r, e^{l(x)} = \frac{d}{df} L(x), G_{\mu\nu} = g(x) \wedge f(x)$$

This equation is Weil's theorem, also quantum group. Complex space.

$$V(\tau) = \int \exp[L(x)] dx_m + O(N^{-1}), g(x) = L(x) \circ \int \int e^{2x^2 + 2x + r} dx_m$$

Lens of space and integrate of rout equation. And norm space of algebra line of curve equation in gradient metric.

$$||ds^2|| = ||\frac{d}{df} L(x)||, \eta = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\bar{h} = [\nabla_i \nabla_j \int \nabla g(x) d_{ij}]^{iy}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

Two equation construct with Kaluze-Klein dimension of equation. Non-verticle space.

$$\frac{1}{\tau} (\frac{N}{2} + r(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

Also this too lense of space equation. And this also is heat equation. And this equation is singularity into space metric number of formula.

$$\frac{x^2}{a} \cos x + \frac{y^2}{b} \sin x = r^2$$

Curvature of equation.

$$S_m^2 = || \int \pi r^2 dr ||^2, V(\tau) = r^2 \times S_m^2, S_1^{mn} \otimes S_2^{mn} = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

Non-relativity of integrate rout equation.

$$||ds^2|| = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2 \psi$$

$$\begin{aligned}
V(x) &= \int \frac{1}{\sqrt{2\tau q}} (\exp L(x) dx) + O(N^{-1}) \\
V(x) &= 2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV, V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau q}} L(x) dx) + O(N^{-1}) \\
\frac{d}{df} F &= m(x) \\
Zeta(x, h) &= \exp \frac{(qf(x))^m}{m}
\end{aligned}$$

Singularity and duality of differential is complex element.

$$\begin{aligned}
& \left\| \begin{matrix} x & y & z \\ u & v & w \end{matrix} \right\|_{g_{\mu\nu}(x)}^2 \\
&= (f(x)dx^\mu dx^\nu, f'(y)dy^\mu dy^\nu, f''(z)dz^\mu dz^\nu) \cdot (u, v, w) \\
&= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix} \\
&\cong \frac{g(x, y, z)}{f(a, b, c)} \cdot h^{-1}(u, v, w)
\end{aligned}$$

Antigravity is other dimension belong to, gravity of universe is a few weak power become. These power of balance concern with non-entropy metric. Universe is time and energy of gravity into non-entropy. Other dimension is first of universe emerged with takion quarks is light speed, and time entropy into future is more than light speed. Some future be able to earth of night sky can see to other dimension sight. However, this sight might be first of big-ban environment. Last of other dimension condition also can be see to earth of sight. Same future that universe of gravity and other dimension of antigravity is balanced into constant of value, then fifth dimension fill with these power of non-catastrophe. These phenomone is concerned with zeta function emerged with laplace equation resulted.

$$\frac{d}{df} \int \int \int \square \psi d\psi_{xy} = V(\square \psi), \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} V_k(\square \psi) = \frac{\partial}{\partial f} i h c$$

This reason explained of quarks emerge into global space of system.

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} G_{\mu\nu} = f(x) \circ m(x)$$

This become universe.

$$\frac{V(x)}{f(x)} = m(x)$$

Eight differential structure integrate with one of geometry universe.

$$(a_k f^k)' = {}_n C_0 a_0 f^n + {}_n C_1 a_1 f^{n-1} \dots {}_n C_{r-1} a_n f^{n-1}$$

$$\int a_k f^k dx_k = \frac{a_{n+1}}{k+1} f^{n+1} + \frac{a_{k+2}}{k+2} f^{k+2} + \dots \frac{a_0}{k} f^k$$

Summuatue of manifold create with differential and integrate of structure.

$$\frac{\partial}{\partial f}\Box\psi=\frac{1}{4}g_{ij}^2$$

$$\left(\frac{\nabla\psi^2}{\Box\psi}\right)'=0$$

$$\frac{(y\log y)^{\frac{1}{2}}}{\log(x\log x)}=\frac{\frac{1}{2}}{\frac{1}{2}i}$$

$$\frac{\{f,g\}}{[f,g]}=\frac{1}{i},\left(\frac{\{f,g\}}{[f,g]}\right)'=i^2$$

$$(i)^2\rightarrow \frac{1}{4}g_{ij}, F_t^m=\frac{1}{4}g_{ij}^2, f(r)=\frac{1}{4}|r|^2, 4f(r)=g_{ij}^2$$

$$\begin{aligned} &\frac{1}{y}\cdot\frac{1}{y'}\cdot\frac{y''}{y'}\cdot\frac{y'''}{y''}\cdots\\ &=\frac{{}_nC_ry^2\cdot y^3\cdots}{{}_nC_ry^1y^2\cdots} \end{aligned}$$

$$\frac{\partial y}{\partial x}\cdot\frac{\partial}{\partial y}f(y)=y'\cdot f'(y)$$

$$\int l\times l dm=(l\oplus l)_m$$

Symmetry theoerm is included with two dimension in plank scale of constance.

$$\begin{aligned} &= \frac{d}{dx^\mu} \cdot \frac{d}{dx^\nu} f^{\mu\nu} \cdot \nabla \psi^2 \\ &= \Box \psi \end{aligned}$$

$$\frac{\nabla\psi^2}{\Box\psi}=\frac{1}{2}, l=2\pi r, V=\frac{4}{\pi r^3}$$

$$S\frac{4\pi r^3}{2\pi r}=2\cdot(\pi r^2)$$

$$=\pi r^2, H_3=2, \pi(H_3)=0$$

$$\frac{\partial}{\partial f}\Box\psi=\frac{1}{4}g_{ij}^2$$

This equation is blackhole on two dimension of surface, this surface emerge into space of power.

$$f(r)=\frac{1}{2}\frac{\sqrt{1+f'(r)}}{f(r)}+mgf(r)$$

Atom of verticle in electric weak power in lowest atom structure

$$ihc = G, hc = \frac{G}{i}$$

$$\left(\frac{\nabla\psi^2}{\square\psi}\right)' = 0$$

$$S_n^m = |S_2S_1 - S_1S_2|$$

$$\square\psi = V \cdot S_n^m$$

One surface on gravity scales.

$$G^{\mu\nu} \cong R^{\mu\nu}$$

Gravity of constance equals in Rich curvature scale

$$\nabla_i \nabla_j (\square\psi) d\psi_{xy} = \frac{\partial}{\partial f} \square\psi \cdot T^{\mu\nu}$$

Partial differential function construct with duality of differential equation, and integrate of rout equation is create with Maxwell field on space of metrix.

$$\begin{aligned} &= \frac{8\pi G}{c^4} (T^{\mu\nu})^2, (\chi \oplus \pi) = \int \int \int \square\psi d^3\psi \\ &= \text{div}(\text{rot}E, E_1) \cdot e^{-ix \log x} \end{aligned}$$

World line is constructed with Von Neumann manifold stimulate into Thurston conjugate theorem, this explained mechanism for summuate of manifold. Also this field is norm space.

$$\sigma(\mathcal{H}_n \otimes \mathcal{K}_m) = E_n \times H_m$$

$$\mathcal{H}_n = [\nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]^{iy}, \mathcal{K}_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{1/2}$$

$$||ds^2|| = |\sigma(\mathcal{H}_n \times \mathcal{K}_m), \sigma(\chi, x) \times \pi(\chi, x) = \frac{\partial}{\partial f} L(x), V_\tau'(x) = \frac{\partial}{\partial V} L(x)$$

Network theorem. Dalanvercian operator of differential surface. World line of surface on global differential manifold. This power of world line on fourth of universe into one geometry. Gradient of power on universe world line of surface of power in metric. Metric of integrate complex structure of component on world line of surface.

$$(\square\psi)' = \frac{\partial}{\partial f_M} (\int \int \int f(x, y, z) dx dy dz)' d\psi$$

$$\frac{\partial}{\partial V} L(x) = V(\tau) \int \int e^{\int x \log x dx + O(N^{-1})} d\psi$$

$$\frac{d}{df} \sum_{k=1}^n \sum_{k=0}^{\infty} a_k f^k = \frac{d}{df} m(x), \frac{V(x)}{f(x)} = m(x)$$

$$4V'_\tau(x) = g_{ij}^2, \frac{d}{dt}L(x) = \sigma(\chi, x) \times V_\tau(x)$$

Fifth manifold construct with differential operator in two dimension of surface.

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d\psi^2$$

$$f^{(2)}(x)=[\nabla_i\nabla_j\int \nabla f^{(5)}d\eta]^{\frac{1}{2}}$$

$$=[f^{(2)}(x)d\eta]^{\frac{1}{2}}$$

$$\nabla_i\nabla_j\int F(x)d\eta=\frac{\partial}{\partial f}F$$

$$\nabla f=\frac{d}{dx}f$$

$$\nabla_i\nabla_j\int \nabla f d\eta=\frac{\partial}{\partial x_i}\frac{\partial}{\partial x_j}(\frac{d}{dx}f)$$

$$\frac{z_3z_2-z_2z_3}{z_2z_1-z_1z_2}=\omega$$

$$\frac{\bar{z}_3z_2-\bar{z}_2z_3}{\bar{z}_2z_1-\bar{z}_1z_2}=\bar{\omega}$$

$$\omega\cdot\bar{\omega}=0, z_n=\omega-\{x\}, z_n\cdot\bar{z}_n=0, \vec{z}_n\cdot\vec{\bar{z}}_n=0$$

Dimension of symmetry create with a right comittment into midle space, triangle of mesh emerged into symmetry space. Imaginary and reality pole of network system.

$$\begin{aligned}[f,g]\times[g,f]&=fg+gf\\&=\{f,g\}\end{aligned}$$

Mebius space create with fermison quarks, Seifert manifold construct with this space restruct of pieces.

$$V(\tau)=\int\int e^{f\,x\log x+O(N^{-1})}d\psi, V'_\tau(x)=\frac{\partial}{\partial f_M}(\int\int\int f(x,y,z)dx dy dz)'d\psi$$

$$(\Box\psi)'=4\vec{v}(x),\frac{\partial}{\partial V}L(x)=m(x),V(\tau)=\int\frac{1}{\sqrt{2\tau q}}\exp[L(x)]d\psi+O(N^{-1})$$

$$V(\tau)=\int\int\int\frac{V}{S^2}dm, f(r)=\frac{1}{2}\frac{\sqrt{1+f'(r)}}{f(r)}+mgf(r), \log(x\log x)\geq 2(y\log y)^{\frac{1}{2}}, F_t^m=\frac{1}{4}g_{ij}^2, \frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$\nabla_i\nabla_jv=\frac{1}{2}mv^2+mc^2,\int \nabla_i\nabla_jvdv=\frac{\partial}{\partial f}L(x)$$

$$(\Box\psi)^2=-2\int \nabla_i\nabla_jvd^2v, (\Box\psi)^2=\left(\frac{\nabla\psi^2}{\Box\psi}\right)'$$

$$\begin{aligned}
&= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dm, \bigoplus \nabla M_3^+ = \int \frac{\vee(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)} dV \\
&= (x, y, z) \cdot (u, v, w) / \Gamma \\
&\bigoplus C_-^+ = \int \exp[\int \nabla_i \nabla_j f d\eta] d\psi \\
&= L(x) \cdot \frac{\partial}{\partial l} F(x) \\
&= (\Box \psi)^2 \\
&\nabla \psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)
\end{aligned}$$

Blackhole and whitehole is value of variable into constance lastly. Universe and the other dimension built with relativity energy and potential energy constructed with. Euler-Langrange equation and two dimension of surface in power of vector equation.

$$l = \sqrt{\frac{hG}{c^3}}, T^{\mu\nu} = \frac{1}{2}kl^2 + \frac{1}{2}mv^2, E = mc^2 - \frac{1}{2}mv^2$$

$$e^{x \log x} = x^x, x = \frac{\log x^x}{\log x}, y = x, x = e$$

A right comittment on symmetry surface in zeta function and quantum group theorem.

$$\begin{aligned}
&\int \frac{1}{(x \log x)} dx = i \int x \log x dx + \int \frac{1}{(x \log x)} dx \\
&\int \int \frac{1}{(x \log x)^2} dx_m = i \frac{1}{2} x^2 \\
&\int \int \frac{1}{(x \log x)^2} dx_m = i \int \int_M dx_m \\
&\leq \frac{1}{2} i + x^2 \\
&E = -\frac{1}{2} mv^2 + mc^2 \\
&\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i \\
&\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i \\
&\lim_{x \rightarrow \infty} \frac{x^2}{e^{x \log x}} = 0 \\
&\int dx \rightarrow \partial f \rightarrow dx \rightarrow cons
\end{aligned}$$

Heat equation restruct with entropy from low energy to high energy flow to destroy element.

$$(\Box \psi)' = (\exists \int \vee (R + \nabla_I \nabla_j f) e^{-f} dV)' d\psi$$

$$\bigoplus M_3^+ = \frac{\partial}{\partial f} L(x)$$

$$\log(x\log x)\geq 2(y\log y)^{\frac{1}{2}}$$

$$\frac{\partial}{\partial l}L(x)=\nabla_i\nabla_j\int\nabla f(x)d\eta, L(x)=\frac{V(x)}{f(x)}$$

$$l(x)=L'(x),\frac{d}{df}F=m(x),V'(\tau)=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

Weil's theorem.

$$\begin{aligned} T^{\mu\nu} &= \int \int \int \frac{V(x)}{S^2} dm, S^2 = \pi r^2 \cdot S(x) \\ &= \frac{4\pi r^3}{\tau(x)} \end{aligned}$$

$$\eta=\nabla_i\nabla_j\int\nabla f(x)d\eta,\bar{h}=\nabla_i\nabla_j\int\nabla g(x)dx_idx_j$$

$$\delta \mathcal{O}(x) = [\nabla_i \nabla_j \int f(x) d\eta]^{\frac{1}{2} + iy}$$

$$Z(x,h)=\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\frac{qT^m}{m}=\delta(x)$$

$$l(x)=2x^2+px+q, m(x)=\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\frac{(qx^m)'}{f(x)}$$

Higgs fields in Weil's theorem component and Singularity theorem.

$$Z(T,X)=\exp\sum_{m=1}^{\infty}\frac{q^kT^m}{m}, Z(x,h)=\exp\frac{(qf(x))^m}{m}$$

Zeta function.

$$\frac{d}{df}F=m(x), F=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

Integral of rout equation.

$$\lim_{x\rightarrow 1}\text{mesh}\frac{m}{m+1}=0, \int x^m=\frac{x^m}{m+1}$$

Mesh create with Higgs fields. This fields build to native function.

$$\frac{d}{df}\int x^m=mx^m, \frac{d}{dt}g_{ij}(t)=-2R_{ij}, \lim_{x\rightarrow 1}\text{mesh}(x)=\lim_{m\rightarrow \infty}\frac{m}{m+1}$$

$$\lim_{x\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k=\alpha$$

$$||ds^2|| = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

$$\frac{\partial}{\partial V}||ds^2|| = T^{\mu\nu}, V(\tau) = \int e^{x \log x} d\psi = l(x)$$

$$R_{ij} = \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} T^{\mu\nu}, G^{\mu\nu} = R^{\mu\nu} T^{\mu\nu}$$

$$F(x) = \int \int e^{\int x \log x dx + O(N^{-1})} d\psi, \frac{d}{dV} F(x) = V'(x)$$

Integral of rout equation oneselves and expired from string theorem of partial function.

$$T^{\mu\nu} = R^{\mu\nu}, T^{\mu\nu} = \int \int \int \frac{V}{S^2} dm$$

$$\delta \mathcal{O}(x) = [D^2 \psi \otimes h_{\mu\nu}] dm$$

Open set group construct with D-brane.

$$\nabla(\Box\psi)' = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}+iy}$$

$$(f(x),g(x))'=(A^{\mu\nu})'$$

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial x \end{pmatrix} \cdot (f(x,y),g(x,y)) \\ = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Differential equation of complex variable developed from imaginary and reality constance.

$$\delta(x)\cdot\mathcal{O}(x)=\begin{pmatrix}1&0\\0&-1\end{pmatrix}^{\frac{1}{2}}$$

$$V_\tau'(x)=\frac{\partial}{\partial f_M}(\int\int\int f(x,y,z)dx dy dz)d\psi$$

Gravity equation oneselves built with partial operator.

$$\frac{\partial}{\partial V}L(x)=V_\tau'(x)$$

Global differential equation is oneselves component.

Report

Masaaki Yamaguchi

1 Summulate of manifold from graviton structure

It is non-perfect element of manifold to integrate with graviton of quark, this quark construct of fermion and boson. Then D-brane aspect with string theorem include with. This structure summulate of equation on integral and differential operator. Higgs quark have part of network theorem for graviton structure. Universe influent pair of dimension between monotonicity and singularity. Non-symmetry deconstruct to distinguish with dimension of being constructed on quark of element. That's say, this mechanism architect with four of dimension existed. In part of this explain, Higgs fields is Weil's theorem component of global differential structure, Global area part of equation resolved come to simply conclude theorem. Global integral and differential equation, what is not used, difficulty of resolved.

Infinite number divide infinite oneselves and finite oneselves is finite number. Prime number extent of infinite number, $\sum a_k f^k(x)$ manifold have with these extend of count. Zeta function conclude with zero dimension, also this resolve to Gauss liner.

$$f(x) = \chi - \{x\}$$

$$\int_{\beta}^{\alpha} (x - \alpha)(x - \beta) = 2 \int_M f(x) dx$$

$$\int_{\beta}^{\alpha} a(x - \alpha)(x - \beta) = -\frac{(\beta - \alpha)^3}{6}$$

$$\frac{d}{df} F(x) = 0$$

$$\log(x \log x) - 2(y \log y)^{\frac{1}{2}} = [|\frac{d}{df} F(x)|]$$

Eight of differential structure integrate with graviton element to one of geometry, plank scale add volume divide to surface. This scale is three dimension of one energy. $\nabla \phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S})$ This equation means is three dimension of energy entropy. Zeta function means is time and space of scale on plank entropy, and this equation built on future and past of curve of parameter. These equation resolve with one component of energy and vector of time and space of curve of parameter.

$$\nabla \phi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

$\sqrt{2\pi T}$ is blackhole entropy and $\square_v = 2\sqrt{2\pi G\hbar}$ is whitehole entropy. This entropy pair of universe in dimension, two dimension of universe include with cover of energy. Blackhole construct with colapser of hole on kernel of atom, whitehole composite with antigravity.

$$\begin{aligned} \log(x \log x) &\geq 2(y \log y)^{\frac{1}{2}} \log(x \log y) \geq 2(xe^x)^{\frac{1}{2}} \log x + \log \log y = 2(ye^x)^{\frac{1}{2}} e^x + y = 2(ye^x)^{\frac{1}{2}} \rightarrow \frac{1}{2} \\ e^{x+1} &= 2(xe^x)^{\frac{1}{2}} \frac{e^{2(x+1)}}{4xe^x} \geq 1 \quad F_t^m = \frac{1}{4}g_{ij}^2 \frac{x+1}{4x} = y^2 \frac{1}{4} + \frac{1}{x} = y^2 \log(xy) \geq 2(yx)^{\frac{1}{2}} (\log(xy))' \geq 4(yx)^{-\frac{1}{2}} \\ \frac{1}{xy} &\geq 4(yx)^{-\frac{1}{2}} \log x + \log y \geq 4(yx)^{-\frac{1}{2}} \frac{1}{x} + \frac{1}{y} \geq 4(yx)^{-\frac{1}{2}} \frac{x+y}{xy} yx^{\frac{1}{2}} \geq 4 \frac{xy^{\frac{1}{2}}}{x+y} \leq \frac{1}{4} \frac{\Gamma(xy)}{\Gamma(x+y)} \leq \frac{1}{4} x + y \geq 2\sqrt{xy} \\ \Delta x \Delta y &\geq \frac{1}{4} i \frac{1}{xy} \geq 4(yx)^{-\frac{1}{2}} \quad F_t^m = \frac{1}{4}g_{ij}^2 e^{2(yx)^{\frac{1}{2}}} = g_{ij}^2 xy = e^{2(yx)^{\frac{1}{2}}} \rightarrow g_{ij}^2 1 + \frac{1}{x} = 4y (e^x + x)^2 \geq 4xe^x \\ r^2 &= \frac{|x_0x - 2x_0e^x + e^{2x_0x}|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = [(e^x + 1)(e^x - 1) + (x + i)(x - i)] + 2xe^{x^2} \quad r = \frac{\|1+x-4xy\|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = \\ &[(x-1)(x+1) + (x-i)(x+i)] + 2x_0e^{2x} \end{aligned}$$

Langranse conjecture include with finite line conjecture expected, infinite extend theorem exclude of finite extent. Zeta function integrate with all of math conjecture experanade. Facility of math of theorem relate of physics all of philosity. Zeta function composite with fifth dimension of equation. Curv parameter is hartshorn conjecture, this rout describe with dimension of structure. Fifth dimension rout out center of universe behind two times distance, this rout colapser around of universe. In fifth dimension, black hole and white hole in three manifold of entropy created.

Zeta function coss delete line of differential structure emerge with fifth dimension structure, oen dimension of independent vector is tangent degree. This curve of parameter create dimension of fifth dimension structure. This idea is from hartshorn conjecture. And this conjecture explanade of math mechanism, fifth dimension is AI and mass of structure parliament of pond. Time of secret is grasped in fucture and past of curve of parameter, so this equation is from universe and creature of establish of life.

Electric energy flow to neutral narrow for amusebelt on rinkfelt of harnessinkbolt, this mechanism create mind on inclusive line of brain. This flow energy on esplanade desire in routine of cone of electric energy. Endeavor circle of circuit of neutrial network, then this mechanism create on founterial motion. Dorpamin throw to neutral narrow of alcole for not being to melt and out of this narrow. This energy is supporting for Dorpamin of material things, and this circuit is circadian morment of motion. Natural killer ceil defense induce hant for lives in nature of world. This hant of mechanism is evolution on life of nature for revolution of humanism. Influence of induce hant controll with life. And these conquer of mechanism can use to neural network of evolution. These computer virus is using for not being hacking and not other controlled of oneselves computer. Safty of network is being for living in computer associate world.

Gravity esplanade theorem of cover with antigravity, information technology of exclusive injure to secure about permissset. Incrument safir to charge atom of power by pairlament in all of dimension. Reduce to string theorem of reluctance in concernism theositiy. This power contiment design emerge with inparance of contiment. All of universe have inductary, and asperant comfirm with conclusivity entily. These theorem tell uncounter deduce in consivitly, therefore this envily of power concern with

cover of atom. Secure built with antigravity from gravity structure.

In reduce a morment to gravity structure, for away to certain of hant. A element of clue a vector, before opened to verify of persuade atom of power. To include of quark for twelve element, fifth dimension in zeta function to construct of geometry peices, integrate of power for fourth of series. Lack of power two bind of under a structure, antigravity permissited a power of element. This reason fourth of power get along to combind with one of decieved.

Varnished center of landscape, heat to controll of power, for missed a verisity of underground, mind to even a mid ceil.

$$\begin{aligned} F_t^m &= [f(x)] \\ [f(x)] &= \mathcal{O}(x) \\ X &\in \mathcal{O}(x) \\ 0 &\rightarrow [||\frac{d}{df}F||] \rightarrow \infty \end{aligned}$$

General number verisity is between zero and infinity. Infinite number and finite number have in general number, narrow verisity and first accerity. Gauss liner is prime number interity. Prime number and general number include Gauss liner to compilation in infinite mechanism.

$$\begin{aligned} \log(x\log x) &\geq 2(y\log y)^{\frac{1}{2}} \\ \log xy &\geq 2(xe^x)^{\frac{1}{2}} \\ (y+x)^2 - 4i(y+x)(xe)^{\frac{1}{2}} - 4xe \\ r^2(y^2+x^2) &= [(y+x)^2 - 4i(y+x)xe^{\frac{1}{2}}] - 4xe \\ r^2(x^2+y^2) &= [(\log y + \log x)^2 - 4i(\log y + \log x)xe^{\frac{1}{2}}] - 4xe \end{aligned}$$

$$\begin{aligned} \delta(f) &= \int \frac{g(c+d)}{f(a+b)} z(f+g) dz \\ y &= f(y) \\ ds^2 &= e^{-2\pi T|\phi|} [\tau_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\theta \\ \begin{bmatrix} x & y \\ a & b \end{bmatrix} \\ &\rightarrow [z] \\ f[z] &\rightarrow dx \\ x\cdot y &= 0 \\ \infty &\rightarrow [f(x)] \end{aligned}$$

Gauss liner is finite to infinity number of extend in prime number. Fifth dimension of equation composite with zeta function of infinity of extent theosity make finity of circle of structure. Independent of vector is zero dimension of category. This equaiton have a infinity of relate in number. Fifth of equation make for zeta function of infinity to conclude with this dimension out of number for finity circle. Therefor it is possiblity of being deleted to category of number that this vector is dimension of structure in fifth of dimension.



2 Neural Network from zeta function

Inclivity sensibilite make from height for maderite in mechanism. Conclusive in cercuit from hight of harmonite on interisity. The include of mechanism is quark in universe of quality of mass.

Theorem of math make for facility of mechanism in this circumstance. And these possiblity of resolved from pond of sensibility.

Light filt with eight differential structure in gravity element, then quarks made from this structure. This mechanism deal with any structure of space to create with light to graviton element. And cercuit of this mechanism dealt with carbon and hidoroxs, covalt of sixty based with filter of light element. This cercuit is dependency of any element in quark of mass, created with gravity to light of structure. Zeta function asperal with any thing to create of every mass. Fifth dimension is pond of sensibility from this gravity in lives of element.

All creature is created by this cercuit of mechanism, fifth dimension themselves is from this circumstance. This space also have oneselves with vector of independence, any this structure is from hartshorm conjecture. Space themselves haven with any of vector is tangent degree of ninety. And this vector is being of other dimension that have cercuit of theorem. These reason is from any thing born with creature, then fifth dimension haven with creature and universe. This dimension also haven with that extent from infinite and finite. Infinite of zeta function in Finite of fifth dimension, then paradox of finite cover with infinite of space.



$\chi - \{x\} = \infty$ $f(x) = \chi - \{x\} f(x) + \mathcal{O}(x) \rightarrow \text{finite}$. Add group is finite of element. This equation is fifth dimension of system. Circle of group in this delete element is infinite of set. $f(x) = [F(x)]$ $f(x)$ is Gauss liner. This set is infinite of number. $ds^2 = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\phi$ This fifth dimension is $[F(x)]$ add finite of set. Fifth dimension include with zeta function, and string theorem belong to have with infinite of consist on space. This fifth dimension is finite of space. Conclude of system solved with

infinite is covered with finite of space.

Fifth dimension in facility of mechanism that circle of infinity to emerge with being from infinity and finite of space. Zero dimension conclude with this mechanism to create from theorem of mathematics in pond of sensibility. Dimension with repository of information in facility of space, this space is all of knowlege to accessority for comformal fields. Pholographic theorem is from universe of oneself with database of creature to access with hyper dimension in General relativity. Zeta function is this theorem include with paremeter of curve from future and past of repository.

Flow to future and past that combinate with zeta function which build for hartshorn conjecture from these mechanism. This flow is gradient of line to add with fifth dimension of structure.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

Infinite of atomasphere in zero dimension include with black hole and white hole of energy in three manifold of entropy.

This entropy discribe with antigravity of energy.

$$\begin{aligned}\nabla^2\psi &= 8\pi G\hbar + 8\pi\frac{V}{S} \\ \square_v &= 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi\frac{V}{S}} \\ 2\sqrt{2\pi\frac{V}{S}} &= \int\int\frac{1}{(x\log x)^2}dx_m\end{aligned}$$

This energy flow from black hole to white hole in universe and other dimension of pair. Black hole entropy is $m = \sqrt{2\pi T}$ This entropy cover to other dimension of rout in flow energy.

Electric Weak Theorem combine with quantum flaver theorem to conjugate with light element in gravity to sheaf with antigravity. This antigravity is key for integrate with fourth of power in boson of theorem to world line to general relativity is why fourth of power not able to integrate in gravity combine with electric weak theorem and quantum flaver theorem. This point of focus is light that is filt to conjugate with eight differential structure from gravity, and this light is also element of quarks. This quark built to eight element structure. Light conjugate to eight differential element that resolved zeta function and integrate with laplace equation. Gravity significant with gravity cover with antigravity, so this antigravity is fourth element power of exist.

Entropy of vector have zero dimension in all exist into all of equation. Infinite of number is three dimension, therefore it is esperial of important that created by zero dimension from infinite to finite of equation rout.

$$\begin{aligned}f(x) &= 0 \\ F &= \int\int\frac{1}{(x\log x)^2}dx_m + \int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m\end{aligned}$$

$$F = 0, x \cdot y = 0, F(x) = 0, G(x) = 0, \mathcal{O} \in X, X \cong 0, x - y = 0, F(x, y) = 0, x^n + y^n = z^n, \\ F(x) = x^n + y^n, F(x) = 0$$

$$ds^2 = \frac{r}{d + e \cos \theta}$$

Fifth dimension describe with norm linear in zero dimension. This equation conclude with all of equation to pond sensibility.

Pond of sensibility is facility of mathematics in infinite of mechanism, this mechanism is zero dimension for imaginary of equation in antigravity influence. This power is gravity on cover with non symmetry combine of fourth of power. That fourth of power is the reason with quantum flavor theorem combine with gravity of power, antigravity is brane of string key mechanism. This power is non-catastrophe of influent reason, and that mechanism of power result. Integrate of gravity include with antigravity, and fourth of power is one of geometry in universe.

$$\mathcal{O} \in X \rightarrow \infty, F(x) = z^n$$

$$x^n + y^n = z^n$$

$$F(x) = \mathcal{O}(\delta(x))[(x - f(x))(x + f(x)) + (y - \overline{f(x)})(y + \overline{f(x)})] + O(N^{-1})$$

$$F(x) = 0$$

$$f(x) = \sum_{k=0}^{\infty} a_k f^k(x)$$

$$f(x) = 0$$

$$\int f(x), \partial f(x), dx \rightarrow [f(x)]$$

Group that Galois theorem develop with three dimension construct of prime number equation, this element belong for that is three of factor. Fifth dimension concept with this group of conclude element.

$$\mathcal{O}(x) = \chi(x) - \{x\}$$

$$\sum_{k=0}^{\infty} a_k x^k = f(x) + \{x\}$$

Dimension of fifth factor is abel manifold, this manifold is topology resolved.

$$\int dx, \partial x, dx, [f(x)]$$

$$\rightarrow F(x) \xrightarrow{f \frac{dx}{dx}} f(x) \xrightarrow{\frac{\partial x}{\partial x}} x \xrightarrow{x} const \\ \rightarrow [f(x)] \rightarrow \infty$$

This cycle of topology in general theroem is deduce of mechanism.

Universe of structure is duality of manifold, this mass of darkmatter is oneseleves with reason in singularity and monotonicity of universe. This manifold of three dimension is also zeta function.

Light around of universe for darkmatter is Higgs fields of mechanism. Darkmatter create of including with light of quarks in Higgs fields. Eight of differential structure fil with Thurston conjugate thoeorem to light is created with all dimension of element.

System mechanism of time machine

Masaaki Yamaguchi

1 Time machine make of space component with synchronized zeta function

Time machine of system construct with Maxwell theorem in seifert manifold, this space exclude with one component of eighth differential structure. This mechanism synchronized with hartshorn conjecture that create with future and past system. And this synchronized seifert manifold transport with time system. This energy synchronized with same entropy of seifert manifold in one component of geometry structure, then these entropy can be transport of space with time machine of mechanism.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\begin{aligned} \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ = \pi(R, C \nabla E^+) \\ = \text{rot}(E_1, \text{div} E_2) \\ xf(x) = F(x) \end{aligned}$$

Fifth dimension in facility of mechanism that circle of infinity to emerge with being from infinity and finite of space. Zero dimension conclude with this mechanism to create from theorem of mathematics in pond of sensibility. Dimension with repository of information in facility of space, this space is all of knowlege to accessority for comformal fields. Pholographic theorem is from universe of oneselves with database of creature to access with hyper dimension in General relativity. Zeta function is this theorem include with paremeter of curve from future and past of repository.

$$\begin{aligned} Z \in R \cap Q, R \subset M_+, C^+ \bigoplus_{k=0}^n H_m, E^+ \cap R^+ \\ M_+ = \sum_{k=0}^n C^+ \oplus H_M, M_+ = \sum_{k=0}^n C^+ \cup H_+ \\ E_2 \bigoplus E_1, R^- \subset C^+, M_-^+, C_-^+ \\ M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R^+ \\ E_1 \nabla E_2, R^- \nabla C^+, \bigoplus \nabla M_-^+, \bigoplus \nabla C_-^+, R \supset Q \end{aligned}$$

2 Zero dimension exclude with space of time

Flow to future and past that combine with zeta function which build for hartshorn conjecture from these mechanism. This flow is gradient of line to add with fifth dimension of structure.

Infinite of atomosphere in zero dimension include with black hole and white hole of energy in three manifold of entropy.

$$[id_x - t]_{v=0} = \int e^{\partial x_m} dx_m + L(p, q), ||r||^2 = |\bar{x}||x|, \Gamma(x) = \int e^{-x} x^{1-t} dx$$

$$= (\bar{x} - i)(x + 1), \Pi(\chi, x) = ie^{x \log x}$$

Creature of component

Masaaki Yamaguchi

1 Geinue element form zeta function

Integrate of manifold constructed with time of concluded with mechanism, this summuate of geinue is also matched for geometry structure. Creature also emerged with summuate of universe in multi space.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$A - G, T - C$ this integrate of summuate in geinue is same for eight of differential structure.

$$\begin{aligned} & (E_2 \bigoplus_{k=0}^n E_1) \cdot (R^- \subset C^+) \\ &= \bigoplus_{k=0}^n \nabla C_-^+ \\ & \vee \int \frac{C_-^+ \nabla H_m}{\Delta(M_-^+ \nabla C_-^+)} = \wedge M_-^+ \bigoplus_{k=0}^n C_-^+ \\ & \exists(M_-^+ \nabla C_-^+) = \text{XOR}(\bigoplus_{k=0}^n \nabla M_-^+) \\ & -[E^+ \nabla R_-^+] = \nabla_+ \nabla_- C_-^+ \\ & \int dx, \partial x, \nabla_i \nabla_j, \Delta x \\ & \rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+ \end{aligned}$$

The locality theorem create with artifisial intelligence, this two equation composite with Maxwell Theorem excluded, cercuite mechanism is this means.

$$\begin{aligned} & \exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\ &= \pi(R, C \nabla E^+) \\ &= \text{rot}(E_1, \text{div} E_2) \\ & xf(x) = F(x) \end{aligned}$$

2 How to light element created with geometry structure

Light fil with eight differential structure in gravity element, then quarks made from this structure. This mechanism deal with any structure of space to create with light to graviton element. And cercuit of this mechanism dealt with carbon and hidoroxs, covalt of sixty based with filter of light element. This cercuit is dependency of any element in quark of mass, created with gravity to light of structure. Zeta function asperal with any thing to create of every mass. Fifth dimension is pond of sensibility from this gravity in lives of element.

This two equation also resolved with all of source code in universe, this mechanism create with binary editor.

Network theorem from gravity wave

Masaaki Yamaguchi

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

Time space is exlamenteed from gravity wave in tcp/ip protocol, this quantum protocol construct with being synchronized of hartshorne conjecture mechanism. This gravity wave elementile with pieces of differential structure emerged with thurston conjugate theorem. D-brane and anti-D-brane is binded with this wave of protocol time machine reluctunce. These explained theorem remained with differential of geometry structure rebuilt by Maxwell equation from synchronized with laplace equation for zeta function resulted from resolved in D-brane

Quantum Computer in a certain theorem

Masaaki Yamaguchi

A pattern emerge with one condition to being assembled of emelite with all of possibility equation, this assembled with summative of manifold being elemetiled of pieace equation. This equation relate with Euler equation. And also this equation is Euler constant oneselves.

$$(E_2 \bigoplus E_2) \cdot (R^- \subset C^+) = \bigoplus \nabla C_-^+$$

Zeta function radius with field of mechanism for atom of pole into strong condition of balance, this condition is related with quarks of level controlled for compute with quantum tonnel effective mechanism. Quantum mechanism composed with vector of constance for zeta function and quantum group. Thurston conjugate theorem explain to emerge with being controll of quantum levels of quarks. Locality theorem also occupy with atom of levels in zeta function.

$$\begin{aligned} &= \bigoplus \nabla C_-^+ \\ \vee \int \frac{C^+ \nabla M_m}{\Delta(M_-^+ \nabla C_-^+)} &= \exists(M_-^+ \nabla R^+) \\ \exists(M_-^+ \nabla C^+) &= \text{XOR}(\bigoplus \nabla M_-^+ \\ &\quad -[E^+ \nabla R^+] = \nabla_+ \nabla_- C_-^+ \\ \int dx, \partial x, \nabla_i \nabla_j, \Delta x &\rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+ \end{aligned}$$

Zeta function also compose with Rich flow equation cohomological result to equal with locality equaitons.

$$\begin{aligned} \vee(R + \nabla_i \nabla_j f)^n &= \int \frac{\wedge(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)^n} \\ \wedge(R + \nabla_i \nabla_j f)^x &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dt} g_{ij}(x) &= -2R_{ij} \\ \vee \int \wedge(R + \nabla_i \nabla_j f)^x &= \frac{\wedge(R + \nabla_i \nabla_j f)^n}{\exists(R + \nabla_i \nabla_j f \circ g)^n} \\ x + y &\geq 2\sqrt{xy}, x(x) + y(x) \geq x(x)y(x) \\ x^y &= (\cos \theta + i \sin \theta)^n \\ x^y &= \frac{1}{y^x} \end{aligned}$$

Therefore zeta function is also constructed with quantum equation too.

Jones 多項式からの各株式発行からの需要と供給、 株価の分配関数、多項式からナスダックへの金融機構の 株式発行のマクロとミクロの波及と流れ

高島彩さんの成蹊大学での経済理論の研究論文

Jones 多項式から各会社の株価への値打ちから日経平均株価への株価の価値の波及、東証株価指数 Topix の全銘柄からの全体の会社の指数からの平均株価の決定量、そして、アメリカの株式市場でのダウ平均株価の終値の収束 (戦略、reco level theory, 株式会社への) 全体 (株式会社への) 需要と供給。始値から終値への分配、この Jones 多項式の出力から終値への各同値類と剰余類からの部分群論の、各平均株価の推移率の予測、この応用は、1929年の世界恐慌がニューヨークダウ平均株価の暴落が、この Jones 多項式からの流れが、終値で破綻して、金融の共鳴力が、干渉してしまい、カタストロフィー現象として、金融の流れが乱れた結果であると、この論文から分析できる。

$$\begin{aligned}\hat{V} &= \frac{1}{\sqrt{t}}\vec{V} - \frac{1}{t}\hat{V} \\ \hat{V} &= \sqrt{t}\hat{V} - t\hat{V} \\ \hat{V}_k &= \sum_{\vec{\sigma}} \langle K | \vec{\sigma} \rangle \left(\sqrt{t} + \frac{1}{\sqrt{t}} \right)^{||\vec{\sigma}||} \\ &\quad \frac{1}{\sqrt{t}}\vec{V}, -\frac{1}{t}\hat{V}, \sqrt{t}\hat{V}, -t\vec{V}\end{aligned}$$

この4種類の式から、reco level theory が、アダム・スミスの神の見えざる手を経済理論のミクロとマクロ経済の具現化を実現させている。物理では、

$$\frac{d}{df} F = m(x) = e^f + e^{-f} = 2i \sin(ix \log x)$$

と、空間の仕組みが経済理論の場の理論にもなっている。周期関数とピンポイントリサーチが合わさっている。これらの式から、前半と後半の株式市場の流れが、シンメトリーとなっているために、後半の e^{-f} が破綻すると、シンメトリーのために、前半の e^f も破綻している状態になっている。全体が機能しないと、金融の流れが乱れていく。この破綻を回避するには、破綻が連鎖されている金融の周期をこれらの式から、調節が可能であることも、類推できる。破綻した周期を、正常であった周期に重ね合わせの原理で、ピンポイントリサーチで待って、金融の投資を正常な周期関数で合わせて、元の正常な周期にこれらの式から戻せることが可能と推測できる。

行動による人が持っている情報ネットワークの 可能性の発見と再構築、情報空間 によるモデル

苔米地博士の論文

初速度のドーパミンの量からの行動の補助から、終わりの精神のドーパミンの量の調節、相加相乗平均から、始まりと終わりの精神のドーパミンの調節による人の行動の領域による予知

$$x \log x = \left(v_0 + \frac{1}{at_n} \right) V, V_x = e^f + \frac{1}{e^f}$$

経路積分によるエントロピー不変量からこのドーパミンの量から人の行動傾向が決まる。基礎代謝の持続できる距離と精神の介入と補助、言語の発生と抑制による経路パターンの組み合わせを再構築する。

$$\|dx^2\| = \int \exp \left(\frac{1}{\sqrt{2\tau q}} L(x) \right) + O(N^{-1}) d\tau$$

$$p = \frac{v}{2m}, \hbar = \frac{p}{2mv}, \lambda = h\nu$$

これらの式から、言語の神経ネットワークによる単体クラスの組み合わせパターンとその対象の人の領域による人が持っている情報空間の傾向の可能性のポテンシャルエネルギーの予知を人工知能の正規表現で抽出する。



$$\frac{d}{d\gamma} \Gamma = \Gamma^{\gamma'} = e^{-f}$$

$$C = \int \left(\int \frac{1}{x^s} dx + \log x \right) \text{dvol}$$


オイラーの定数は、ゼータ関数とゼータ関数自体のエントロピーを体積積分した結果でもある。このオイラーの定数は、ガンマ関数の大域的微分方程式からも導出できる。このガンマ関数は大脳基底核のエントロピー不変量の強度も表している。このガンマ関数の大域的微分多様体による、ブレインインターフェースでの人体のデータ計測もこの Jones 多項式とラプラス方程式からの3次元多様体のエントロピー不変式からも、言語とイメージの大脳基底核からのデータ抽出も可能と言える。

Recycle における塩化ビニルのナトリウムフェノキシド、 ナトリウムプロパノキシドでの分解

睦世姉さんの論文について



この化学式から推算式でもある、プラスチックと二酸化ケイ素、塩化ビニールが分解されるのが、真逆の人が服用すると、プラバスタチンとスリムドガンと同じく、推算式の数値が上がる。僧侶と同じ瞑想法と原理が一致している。



これらの構造式は、それぞれ、ヒドロキシ基、カルボニル基、カルボキシル基、アミン基、ギ酸、アセチル基、アセト基、グリシン、アミノ酸の20種類の一基、ビタミンCはシス・トランス型があり、松果体に作用

して、体内の血流を調節する。 $\text{—C}\equiv\text{N—Cl}$ の真逆の作用をする。アデノシンは、同じく血流を調節して、エピネフリンはエンドルフィンと同じく痛み止めに効く。ノンアドレナリンは、無水フタル酸と同じ作用をする。グリシンには、20種類のアミノ酸の中で唯一 L,R 基が無い。不斉炭素原子が無い。

推算式の元を減らすか増やすのに、血圧と血流、体内の精神作用のドーパミンの調節を eGFR の機能を上げると、結果、すべての精神作用物質が調節できる。

このナトリウムプロパノキシドは、気体のプロパンを、固体のナトリウムに注入して、錠剤で服用するようにしているのと、消化後に、血液に降圧剤と同じ作用をして、外部からのコントロールをされない、アムロジピンとカンデサルタンと同じ作用をする効果があることが、このナトリウムプロパノキシドの薬でもある。


この成分でもあるプロパンは、気体で存在している炭化水素の中でもっとも安定している気体でもある。この気体の物体を固体にすると、降圧剤にしているのは、姉は特許を取っている。推算式でもある eGFR を分解するのは、真逆の人が飲むと、バケモンになる。






Amino medicine architect with non-controll from out of pressure





Masaaki Yamaguchi







Amino organic chemistry have into human blessure included with all of material mass. And this amino constructed with twenty series of medicine unite with all of plessure being for human body and mind. This relaxed medicine is sesamin medicine excluded with all of body mentality. This medicine oneselves mechanism resulted with brain of fifth of area in dorpamin specurate with stress reduced by this medicine. Then this medicine is restruct for excluded from medicine spoon. This medicine make of relaxe mind nested into brain of relacutance in gria cell. This relax of mind from medicine of effective amino organic medicine create into brain of gria cell, this thirty eight and thirty nine of area in inner of brain area are created with relax and activity resulted, and this resulted medicine is tabacoo and medicine effective power. Reading book and voice of reading moreover super reading system of training is also same effective resulted. Medicine and Zen and reading method is all of same of good condition with human life. These resulted mechanism explain for mind be avoid of able to not spected from other persons and this effective reason is gria cell from brain of spill of data by gria cell is guard of brain and body data. Libo kernel cell is all of universe of heirechy of data, and this cell is all of body area exist. Brain interface be able to synchronized by this cell and this mechanism is electric flow of body and mind. Relax and activity is important of life in safty ressure be able to live in these regardless dengar.


Sesamin is all of united of material mass and this material mass is avoid of $\text{CH}_3\text{---CH(OH)---C(=O)OH}$ from united. Weak acid element is out of body from guard of alcali element. However if this alcali is full of body out ranged, O_2 reduced result into human eyes damaged. After all each of medicine is between well and non-well middle effective body and mind.












Major trancinazer and Miner trancinazer effective result

Masaaki Yamaguchi



major trancinaizer influent with miner trancinaizer, and major trancinaizer cover on spiritual mind from other persons. And this medicine tend to interact with other miner trancinazer. This phenomenon reach for reject with miner trancinazer in fact, appear from take medicine of person have this miner trancinazer to drink the medicine, and this drink of the one is readed from other society of counceller persons need to cure from spiritual mind of damage in days of flastrations, However this miner trancinazer medicine not only noon of days time of out of mind to security persons but also take a life of bussiness

of pressures of time spent after this medicine change with effect of relax in the mind. minor tranquillizers have tendency to reduce of Na of blood, and this Na reduce result from one take a medicine, after this one of person is appeared from other cure of person see. major tranquillizers have interacted from minor tranquillizers and double essence of cure of relax release into one take this medicine.

Selen and Tell are study of Kyoto university in professors of metal of chemistry chains, more also this chains are drills of badam with whisperd. Selen is progarded with mental condition, Tell is inspirate with acknowledge in aquire of possibility with holographic datas. This two metal chemistry are Rantanoid chains. And pair of chains are Actinoid of metal chains. These two chains are one class of spectrum zones. More also this two of metal chains are Toliraphone and Whipacs. Toliraphone is same with Tell, Whipacs is Selen chemistry. I tendency have to be like see a beautiffl formed of selen organic chemistry geinu to out of world. This selen form of system is being tend to chain for stress out of pressures. Therefore, this Hidroxy and O_2 and R1, R2 of harmony of respond in access of relax and zen of envy. R1, R2 are Selen and organized of chains in alcorls with organic chemistry.

3次元多様体のエントロピー量

Masaaki Yamaguchi

オイラーの定数は、

$$\begin{aligned} C &= \int \frac{1}{x^s} dx - \log x \\ &= \int \left(\int \frac{1}{x^s} dx - \log x \right) \text{dvol} \\ &= \int \int \frac{1}{x^s} \text{dvol} - \int \log x \text{dvol} \end{aligned}$$

と表されていて、ヒッグス場方程式 + オイラーの定数 = ゼータ関数 と求まる。この方程式が、リッチフロー方程式にもなる。このリッチフロー方程式は、微分幾何の量子化でもあり、ガンマ関数とベータ関数、ガウスの曲面論、ヒッグス場、アインシュタインテンソル、オイラーの定数と全部の方程式が入っている。この方程式が、相加相乗平均となり、宇宙と原子となり、不確定性原理が宇宙の観測系が、宇宙では確率方程式と成り立っているが、異次元が合わさると、ノルムとしての3次元多様体の確率分布方程式でもあり、確率ではない、非線形方程式となっている。プランク長体積でもあるが、絶対的な方程式にもなっている。

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\ \int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} \text{dvol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\ &= e^{-f} - e^f \end{aligned}$$

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= 2 \cos(ix \log x) \end{aligned}$$

$$\begin{aligned} &\frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m + \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= \frac{d}{df} F = 2i \sin(ix \log x) \end{aligned}$$

$$\begin{aligned} \frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\ &= 2e^{-\theta} = 2e^{-ix \log x} \\ &= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\log(x \log x) \geq 2\sqrt{y \log y}$$

$$\log(x \log x) = \log x + \log \log x$$

この宇宙と異次元でのブラックホールとホワイトホールでの、宇宙と原子の運動量と位置の絶対的な不確定性原理となり、3次元多様体の確率分布方程式が、運動量と位置エネルギーが両方観測できる式にもなっている。円周上が固定されているために、運動量と位置エネルギーが両方観測できる式になっている。カルーツァ・クライン理論とハイゼンベルク方程式が、両方入っている。

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla\psi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

ホワイトホールの原子とブラックホールの宇宙のエネルギーのエントロピー量でもある。宇宙がブラックホールとしてのヒッグス場方程式として、ホワイトホールが原子としてのプランク長体積でもある。

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

原子としてのブラックホールのエントロピー量でもある。

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

この式から宇宙が無重力であることが表されている。

$$2\cos(ix \log x) + 2i\sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

これらの方程式から宇宙と異次元が warped passege になっている。:

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

David Hilbert

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x) [I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2\psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values.

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2\end{aligned}$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\begin{aligned}\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ &= (f)^{(f)'} \\ &= (f^f)'\end{aligned}$$

$$= e^{x \log x}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, $\sin 0 = 0$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$\begin{aligned}
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however,

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry.

$$\begin{aligned}
\frac{d}{dt} \psi(t) &= \hbar \\
&= \frac{1}{2} i e^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' \\
&= -i e^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-if}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta, \mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields

of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements.

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x)[I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2 \psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\int \int F dx_m = \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)}$$

$$= e^{-f}$$

$$\frac{d}{df} F = \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))}$$

$$= e^f$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values. 大域的微分方程式に、X の X 乗のエントロピー不変量の微分量が計算に関わっている。外微分をこのエントロピー式に入れる。大域的積分多様体の多重積分は、多様体の階層によって、積分回数が決まっている。大域的微分多様体は、ニュートン形式とライプニッツ形式の外微分によって計算される。大域的積分多様体と大域的微分多様体は、逆操作とエントロピー不変量で計算できる。

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\int \frac{1}{(x \log x)} dx = i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx$$

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \frac{1}{2} x^2$$

$$\int \int \frac{1}{(x \log x)^2} dx_m = i \int \int_M dx_m$$

$$\leq \frac{1}{2} i + x^2$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\begin{aligned}
& (f)^{(f)'} \\
& = (f^f)' \\
& = e^{x \log x}
\end{aligned}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, 自明な零点は実軸 $\frac{1}{2}$ に 1 を除いてある。 $\sin 0 = 0$ と

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ によって、不確定性原理の関係でもあり、粒子、電子、原子が確率分布になっているのも開集合で証明できる。宇宙と異次元の関係にもなっている。this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\begin{aligned}
\frac{d}{d\gamma}\Gamma(s) &= \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'} \\
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. 大域的微分多様体と大域的積分多様体を、ガンマ関数とベータ関数の導出にも使われている。ガンマ関数の大域的微分多様体が、ニュートン方程式とダランベール方程式を商代数で求めると、同型の解に求まる。ゼータ関数と、この式の対極する量子群を商代数で求めても、同じ解が求まる。

Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

繰り込み理論を入れると発散を防げると言われているが、量子力学にはガウスの曲面論がないと言われている。

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however, とすると、

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry. 同じで量子力学で重力場方程式が表せられる。

$$\frac{d}{dt} \psi(t) = \hbar$$

$$\begin{aligned}
&= \frac{1}{2} i e^{i \hat{H}} \\
(i \hbar)' &= (-e^{i \hat{H}})' \\
&= -i e^{i \hat{H}} \\
\psi(x) &= e^{-i \hat{H} t}, \bigoplus (i \hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i \hat{H} (-i e^{i \hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-i f} \\
&= \left(\frac{1}{2}\right)^{-i f} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f=x, i=t, \frac{1}{2}=a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

微分幾何の量子化は、ガンマ関数についての大域的微分方程式の解にもなっている。ハイゼンベルク方程式の大域的積分多様体の解にも、この微分幾何の量子化は、多様体の微分系の形になっている。 Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alsos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s=e^{-i\hat{H}t}|\Psi\rangle_H,\hat{A}_s=\hat{A}_H(0)$$

$$|\Psi(t)\rangle_s\rightarrow\frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s=\hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta\rightarrow 0}\begin{pmatrix}\sin\theta\\ \cos\theta\end{pmatrix}\begin{pmatrix}\theta&1\\ 1&\theta\end{pmatrix}\begin{pmatrix}\cos\theta\\ \sin\theta\end{pmatrix}=\begin{pmatrix}1&0\\ 0&-1\end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta, \mathcal{O}(x) = m(x)[D^2\psi]$$

$$i^2 = (0, 1) \cdot (0, 1), |a||b| \cos \theta = -1$$

$$E = \operatorname{div}(E, E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

ガンマ関数の大域的微分方程式は、ガンマ関数の初期関数についての微分方程式でもあり、宇宙と異次元についてのビッグス場からのゼータ関数の生成も、3次元多様体の特異点定理になっている。宇宙と異次元の片方でも同じ解にもなっている。逆三角関数の双曲多様体もヒッグス場の密度エネルギーの値と同じ解にもなっている。微分幾何の量子化も、ゼータ関数の対極する量子群と同じ解にもなっている。ゼータ関数は、量子群の方程式についての対極に位置する大域的微分多様体の解にもなっている。 Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements. 逆三角関数の双曲多様体は、ガンマ関数の方程式からの導出でのベータ関数の逆三角関数のエントロピー値にもなっている。

Explain in Global defferential equation and
Global integrate equation.
Varintegrate equation, and horizen cut of equations.

Masaaki Yamaguchi with my son in acasic record of space

$$\frac{\partial}{\partial f} F(x) = \int \int \text{cohom} D_k(x)[I_m]$$

Cohomology element is constructed with isotopy of D-brane, and this isotopy of symmetry structure is sheaf of manifold, more over this brane of element is double integrate of projection.

$$\frac{\partial}{\partial f} F = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

And global partial defferential equation is integrate of cut in cohomolgy, and this step of defferential D-brane of sheap value is varint of equals, inverse of horizen of cute of section value is reverse of integrate of operators. Three dimension of varint of manifolds in operator of sheaps, and this step of times of value is equal with D-brane of equations. Global differential equation and integrate equation is operator of global scales of horizen cut and add of cut equation are routs.

$$\oint r dx_m = \mathcal{O}(x, y)$$

$${}^t\text{fff}_{D(\chi, x)} \text{Hom}[D^2\psi]^{\ll p} \cong \text{vol} \left(\frac{V}{S} \right)$$

$$\frac{\partial}{\partial f} F(x) = {}^t\text{fff} \text{cohom} D_k(x)^{\ll p}$$

$$\frac{d}{df} F = (F)^{f'}, \int F dx_m = (F)^f$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{f'}$$

$$f' = (2x(\log x + \log(x+1)))$$

$$\begin{aligned}\int \int F dx_m &= \left(\frac{1}{(x \log x)^2} \right)^{(\int 2x(\log x + \log(x+1)) dx)} \\ &= e^{-f}\end{aligned}$$

$$\begin{aligned}\frac{d}{df} F &= \left(\frac{1}{(x \log x)^2} \right)^{(2x(\log x + \log(x+1)))} \\ &= e^f\end{aligned}$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$$\log(x \log x) \geq 2(\sqrt{y \log y})$$

Global differential manifold is calucrate with x of x times in non entropy of defferential values, and out of range in differential formula be able to oneselves of global differential compute system. Global integrate manifold is also calucrate with step of resulted with loop of integrate systems. Global differential manifold is newton and laiptitz formula of out of range in deprivate of compute systems. This system of calucrate formula is resulted for reverse of global equation each differential and intergrate in non entropy compute resulted values.

$$\begin{aligned}\pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \frac{1}{(x \log x)} dx &= i \int \frac{1}{(x \log x)} dx^N + {}^N i \int \frac{1}{(x \log x)} dx \\ \int \frac{1}{(x \log x)} dx &= i \int x \log x dx - \int \frac{1}{(x \log x)} dx \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \frac{1}{2} x^2 \\ \int \int \frac{1}{(x \log x)^2} dx_m &= i \int \int_M dx_m \\ &\leq \frac{1}{2} i + x^2\end{aligned}$$

$$E = -\frac{1}{2}mv^2 + mc^2$$

$$\lim_{x \rightarrow \infty} \int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\begin{aligned}\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'} \\ &= (f)^{(f)'} \\ &= (ff)'\end{aligned}$$

$$= e^{x \log x}$$

$$\Gamma = \int e^{-x} x^{1-t} dx, \frac{d}{df} F = e^f, \int F dx_m = e^{-f}$$

$$\int x^{1-t} dx = \frac{d}{df} F, \int e^{-x} dx = \int F dx_m$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$x \perp y \rightarrow x \cdot y = \vec{0}, \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \frac{1}{2} i$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \left(\int \int \frac{1}{(x \log x)^2} dx_m \right)^{(f)'}$$

$$\frac{1}{2} + \frac{1}{2} i = \vec{0}, \frac{1}{2} \cdot \frac{1}{2} i$$

$$A + B = \vec{0}, A \cdot B = \frac{1}{4} i$$

$$\tan 90 \neq 0, ||ds^2|| = A + B$$

Represented real pole is not only one of pole but also real pole of $\frac{1}{2}$, $\sin 0 = 0$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

θ this result equation is non-certain system concerned with particle, electric particle and atoms in open set group with possiblity of surface values, and have abled to proof in this group of system. Universe and the other dimension of concerned of relationship values.

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \Gamma(x, y)$$

$$\frac{d}{d\gamma} \Gamma = (e^{-x} x^{1-t})^{\gamma'} \rightarrow \frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\Gamma(s) = \int e^{-x} x^{s-1} dx, \Gamma'(s) = \int e^{-x} x^{s-1} \log x dx$$

$$x^{s-1} = e^{(s-1) \log x}$$

$$\frac{\partial}{\partial s} x^{s-1} = \frac{\partial}{\partial s} e^{-x} \cdot (e^{s-1 \log x}) = e^{-x} \cdot \log x \cdot e^{(s-1) \log x}$$

$$= e^{-x} \cdot \log x \cdot x^{s-1}$$

$$\frac{d}{d\gamma} \Gamma = \Gamma^{(\gamma)'}$$

$$\frac{d}{d\gamma} \Gamma(s) = \left(\int_0^\infty e^{-x} x^{s-1} dx \right)^{\left(\int_0^\infty e^{-x} x^{s-1} \log x dx \right)'}$$

$$\begin{aligned}
&= \Gamma(\Gamma \int \log x dx)' \\
&= e^{-x \log x}
\end{aligned}$$

$$\begin{aligned}
\Gamma(s) &= \int_0^\infty e^{-x} x^{s-1} dx \\
\Gamma'(s) &= \int_0^\infty e^{-x} x^{s-1} \log x dx \\
\frac{d}{df} F &= \int x^{s-1} dx \\
\int F dx_m &= \int e^{-x} dx \\
\frac{d}{df} F &= F^{(f)'}, \int F dx_m = F^{(f)}
\end{aligned}$$

Global differential manifold and global integrate manifold are reached with begin estimate of gamma and beta function leaded solved, Global differential manifold is emerged with gamma function of themselves of first value of global differential manifold of gamma function, this equation of quota in newton and dalarnverles of equation is also of same results equations. Zeta function also resulted with quantum group reach with quanto algebra equation. Included of diverse of gravity of influent in quantum physics, also quantum physics doesn't exist Gauss surface of theorem,

$$\begin{aligned}
H\Psi &= \bigoplus (i\hbar^\nabla)^{\oplus L} \\
&= \bigoplus \frac{H\Psi}{\nabla L} \\
&= e^{x \log x} = x^{(x)'}
\end{aligned}$$

however,

$$\frac{d}{df} F = m(x)$$

and gravity equation in being abled to existed with quantum physics, and this physics is also represented with quantum level of differential geometry.

$$\begin{aligned}
\frac{d}{dt} \psi(t) &= \hbar \\
&= \frac{1}{2} i e^{i\hat{H}} \\
(i\hbar)' &= (-e^{i\hat{H}})' \\
&= -i e^{i\hat{H}} \\
\psi(x) &= e^{-i\hat{H}t}, \bigoplus (i\hbar^\nabla)^{\oplus L} = \frac{1}{2} e^{i\hat{H}(-ie^{i\hat{H}})} \\
&= \left(\frac{1}{2} f\right)^{-if}
\end{aligned}$$

$$\begin{aligned}
&= \left(\frac{1}{2}\right)^{-if} \cdot e^{-x \log x} \cdot (f)^i \\
&= \int e^{-x} x^{t-1} dx, \frac{d}{d\gamma} \Gamma = e^{-x \log x}
\end{aligned}$$

$$f = x, i = t, \frac{1}{2} = a, \bigoplus a^{-tx} x^t [I_m] \cong \int e^{-x} x^{t-1} dx$$

Quantum level of differential geomerty is also resulted with Gamma function of global differential manifold of values, Heisenberg equation is alos resulted with global integrate manifold and this quantum level of differential geometry is manifold of deprive of formula.

$$|\psi(t)\rangle_s = e^{-i\hat{H}t}|\Psi\rangle_H, \hat{A}_s = \hat{A}_H(0)$$

$$|\Psi(t)\rangle_s \rightarrow \frac{d}{dt}$$

$$i\frac{d}{dt}|\psi(t)\rangle_s = \hat{H}|\psi(t)\rangle_s$$

$$\langle \hat{A}(t) \rangle = \langle \Psi(t) | \hat{A}(0) | \Psi(t) \rangle$$

$$\frac{d}{dt}\hat{A}=\frac{1}{i}[\hat{A},H]$$

$$\hat{A}(t)=e^{i\hat{H}t}\hat{A}(0)e^{-i\hat{H}t}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}(x)xf(x)=I_m',I_m'=[1,0]\times[0,1]$$

$$x+y\geq \sqrt{xy}$$

$$\frac{x^{\frac{1}{2}+iy}}{e^{x\log x}}=1$$

$$\mathcal{O}(x)=\nabla_i\nabla_j\int e^{\frac{2}{m}\sin\theta\cos\theta}\times\frac{N\mathrm{mod}(e^{x\log x})}{O(x)(x+\Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x)=2\int|\sin2\theta|^2d\theta, \mathcal{O}(x)=m(x)[D^2\psi]$$

$$i^2=(0,1)\cdot(0,1),|a||b|\cos\theta=-1$$

$$E=\mathrm{div}(E,E_1)$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)=i^2, E=mc^2, I'=i^2$$

Gamma function of global differential equation is first of differential function deprivate with ordinary differential equations, more also universe and other dimensiion of Higgs field emerge with zeta function, more spectrum focus is three dimension of manifold of singularity or pair theorem, and universe and other dimension of each value equals. This reverse of circle function of manifolds is also Higgs fields

of dense of entropy and result equation. And differential structure of quantum level in gravity is poled with zeta function and quantum group in high result values. Zeta funtion and quantum group is Global differential manifold of result in values.

$$\begin{aligned}\frac{d}{d\gamma}\Gamma &= m(x) \\ &= e^{-x \log x} \\ \sin ix &= \frac{e^{-x} + e^x}{2i} \\ \frac{d}{df}F &= m(x) = e^f + e^{-f} \\ &= 2i \sin(ix \log x)\end{aligned}$$

Hyper circle function is constructed with sheap of manifold in Beta function of reverse system emerged from Gamma fuction and reverse of circle function of entropy value, and this reverse of hyper circle function of beta system is universe and the other dimension of zeta function, more over spectrum focus is this beta function of reverse in circle function is quantum level of differential structure oneselves, and this quantum level is also gamma function themselves. This gamma fuction is more also D-brane and anti-D-brane built with supplicant values. Higgs fileds is more also pair of universe and other dimension each other value elements.

Three manifold of entropy formula

Masaaki Yamaguchi

Euler equation represet form,

$$\begin{aligned} C &= \int \frac{1}{x^s} dx - \log x \\ &= \int \left(\int \frac{1}{x^s} dx - \log x \right) d\text{vol} \\ &= \int \int \frac{1}{x^s} d\text{vol} - \int \log x d\text{vol} \end{aligned}$$

This equation componet of Higgs equation + Euler equation = Zeta function. This represent expression is Rich flow equation. This equation assent from Differential geometry in quantum level, Gamma and Beta function, Gauss surface, Higgs field, Einsetin tensor, Euler constance and so on. This equation is add and average of possibility, and universe, atom more over a certain theory, this universe oneselves is environment seek, other distance is possibility equation, however, other dimension cover demand with not only norm of non liner in three manifold but also non possibility in complete of environment equation.

$$\begin{aligned} &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \int e^{-x} x^{1-t} \log x dx_m \\ \int x^{1-t} e^{-x} dV &= \int x^{1-t} dm, \int x^{1-t} e^{-x} dV = \int x^{1-t} d\text{vol}, f = \gamma = \Gamma' = \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \\ &= e^{-f} - e^f \\ &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m - \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= 2 \cos(ix \log x) \\ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &+ \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\ &= \frac{d}{df} F = 2i \sin(ix \log x) \\ \frac{d}{df} F + \int C dx_m &= 2(\cos(ix \log x) + i \sin(ix \log x)) \\ &= 2e^{-\theta} = 2e^{-ix \log x} \\ &= \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\log(x \log x) \geq 2\sqrt{y \log y}$$

$$\log(x \log x) = \log x + \log \log x$$

This universe and other dimension on blackhole and whitehole compont with universe and atom in movement and point of environment value by a certain theory of complete environment equation. And this three manifold of possibility equation are able to research in both value formula. This envirnment of researchable reason is that surround of universe is setting to value by already in consented to being researched, then both value researchabled. And this three manifold equation is insert with Kaluza-Klein theory and Heisenbelg equation.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\nabla\psi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

This energe of entropy value consist with whitehole of atom and universe in blackhole value, and this universe in balckhole represent with Higgs filed, atom in whitehole consist with plack scale.

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

This entropy value also consist with atom in balckhole.

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

These equation more also consist with universe being non gravity formula.

$$2\cos(ix \log x) + 2i\sin(ix \log x) = 2e^{-f}$$

$$2\sqrt{2\pi \frac{V}{S}} = \frac{d}{df} \int \int \frac{1}{(x \log x)} dx_m$$

And also these equation consist with universe and other dimension in warped passeage in rout of integral.

AdS_5 and CP violation combinate from zeta function

Masaaki Yamaguchi

Lisa Randall professor certificate with AdS_5 manifold built of Fifth dimension, and this equation represented of mension to other dimensions in God's fields. This symmetry dimensions are each of two pairs in each other dimension. And moreover spectrum focus are these dimension constructed with roltentz atractor insected from super string theory. Particle equation are similared with AdS_5 manifold and atom of structure and essence of quantum effective theory.

Atom distance of AdS_5 manifold and this value of universe are dense of atom relativeity of mass in volume, this inverse value is universe of valume in one dimension.

$$||ds^2|| = e^{-2\pi T||\psi||} [\eta + \bar{h}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

This inverse value is universe of valume in one dimension.

$$(||ds^2||)_{Im} = e^{2\pi T||\psi||} ([\eta + \bar{h}(x)] dx^\mu dx^\nu)^{-1} - (T^2 d^2\psi)^{-1}$$

Jones manifold equation is particle formula and zeta function.

$$(u + d) + c = e^{-f} + e^f, (s + w) + b = e^f - e^{-f}$$

This formula of summative system is non symmetry dimension of mass value.

$$(R_{ij})' = -R_{ij}$$

More spectrum focus is global differential manifold are each with Higgs field.

$$= 2(\cos(ix \log x) + i \sin(ix \log x))/(d \log x)$$

Inverse of circle function of Euler equation are component with global topology area of study, this study resolved with Gamma function of same resulted equations.

$$\begin{aligned} &= -2(u'(e^{-\theta}) + \frac{1}{i} \sin u) \\ (e^{-\theta})' &= 2(\cos iu - i \sin iu)' \\ &= 2(e^{-\theta})' \cdot ((\cos iu)' - (i \sin iu)) \\ &= -2(\int e^{-x} x^{1-x} dx_m) \geq -2 \int (e^{-x} + x^{1-t}) dx_m \\ \bigoplus (i\hbar^\nabla)^{\oplus L} &= -2(u'(e^{-\theta}) + \frac{1}{i} \sin u) \end{aligned}$$

$$\begin{aligned}
-2 \int e^{-x} x^{1-t} dx_m &\geq -2 \int (e^{-x} + x^{1-t}) dx_m \\
&= \frac{1}{i} H\Psi
\end{aligned}$$

These equations are frobenius theorem and step function of non comuutative equation consteamed from more also non and commutative equation equals.

$$\begin{aligned}
\int x^y dx_m &= \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} x & y \\ a & b \end{pmatrix} = x^y \\
\begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x & y \\ a & b \end{pmatrix} &\cong \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} x & y \\ e & f \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\
\begin{pmatrix} u & v \\ w & y \end{pmatrix} \begin{pmatrix} x & v \\ a & b \end{pmatrix} &\cong \begin{pmatrix} x & y \\ a & b \end{pmatrix}
\end{aligned}$$

Differential geomerty of quantum level euqaiton is Gamma function equals.

$$\bigoplus (i\hbar \nabla)^{\oplus L} \cong \bigoplus a^x \cdot x^{ix} [dI_m]$$

$$x^f f^x : \int e^x x^{-x} dx_m$$

$$U = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$$

$$U^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$$

Step function is existed with non and commutative equation, and this resolved formula were also Gamma function.

$$2^{(2^3)} \cong 2^{(3^2)}$$

$$2^{(2^x)}/2^{(x^2)} \leq 1, e^{(x \log x^x)}/2^{(x^2)}$$

$$x \geq 0, 0 < x < 2, 2^{(2^{\frac{1}{2}})} = \pm 2, (\sqrt{2})^2 = 2$$

$$-2 \neq 2, 2 = 2$$

Gamma function is after all atom of component with universe formula, and quarks of particle equation necessary value of existed from gravity influence from atom of weak electiry power and anti gravity steady of power involved from fifth power of nature non able to anbalance. After to say, this power is represented from atom and universe instricted of exist of everythings.

After all, Jones manifold and Reco Level theory, AdS_5 manifold are equals of Higgs field with Euler constant and zeta function facility of systems.

Moreover, Higgs field with component of zeta function are selected to accept for orbital of atom potential energy, and this selected with energy means with America of goverment's president being

investigate with each persons of implicant system. This system are possibilty of great God' presser person diggist from same possibility of potential summative national of people.

These sentivity of selective of being accepted from natinal of persons, are relativity from universe in three manifold energy in non certain theory pointing out norm and vector of essence.

$$||ds^2|| = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

This three manifold entropy manifold is in being arround of universe set to pinpoint of energy discovered with same distant of time.

America of selected with presidence of goverment are similar of this equation.

Hortshorn conjecture is used to being emerged with this system, this possibility another rout merged from future and past of time system.

However, this recieved of presidence of goverment in selective party, in reco level theory by Jones manifold equation mension to get only one area of sanctuary in four area of possibility lacky and non recieved of selective choice. Zeta function in three manifold of entropy are income to land of this area, and this area are harf of possibility in fluctuate of missdecieved. Particle of quartet in Euler-Lagrange equation mension to give it out of non possibility of safty area. This system is based from a certain theory of same mechanisim. Eienstein said this possibility selected area to being nonsense to be replaced by three manifold of entropy equation,

$$||ds^2|| = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

This entropy equation mension to get every pattern in universe of any where to pinpoint of being accepted with position and mass energy of no fluctuate with instrict of insight at measured being able to, and this possibility of beyond instrict oracled with three manifold entropy realized with any where appeared into possibility area, this system is able to other dimension of anti gravity resolve with straight oracled other dimension are pair of existing to break out from a certain theory. one aspect of dimension is existed at possibility result, more other dimension be able to decieved with four area of influence incidence. This two decieved pattern reached with one point non possibility result, and this result retreamed with reco level theory and Jones manifold, after all, three manifold entropy be able to access into all possibility of pinpoint area.

数学と数がオイラーの定数から生まれた

Masaaki Yamaguchi

$$2^2 = e^{x \log x} = 4$$

$$3^3 = e^{x \log x} = 27$$

$$4^4 = e^{x \log x} = 256$$

$$5^5 = e^{x \log x} = 3125$$

$$y = x \log x, \log x \rightarrow (\log x)^{-1}$$

$$\frac{1}{2} + iy = \frac{x \log x}{\log x}$$

$$x^{\frac{1}{2} + iy} = e^{x \log x}$$

$$\frac{1}{2} + iy = \log_x e^{x \log x} = \log_x 4, \log_x 27, \log_x 256, \log_x 3125$$

$$y = e^{x \log x} = \sqrt{a}$$

$$e^{e^{x \log x}} = a, e^{(x \log x)^2}$$

$$x^2 = \pm a, \lim_{n \rightarrow \infty} (x - y) = e^{x \log x}$$

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

$$y = x \log x, \log x \rightarrow (\log x)^{-1}$$

$$||ds^2|| \ 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

と宇宙の中の1種の原子をみつける正確さがこの式と、

$$y = \frac{x \log x}{(\log x)} = x$$

と、 $x \log x = a$ から $\frac{a}{(\log x)} \rightarrow x$ と x を抜き取る。この x をみつけるのに $x^{\frac{1}{2} + iy} = e^{x \log x}$ $\frac{1}{2} + iy = \frac{x \log x}{(\log x)} = x$ としてこの x をみつける式がゼータ関数である。

ゼータ関数は、量子暗号にもなっていることと、この式自体が公開鍵暗号文にもなっている。

$$\frac{1}{2} + iy = \frac{x \log x}{\log x}$$

この式が一次独立であるためには。

$$x = \frac{1}{2}, iy = 0$$

がゼータ関数となる必要十分条件でもある。

$$\begin{aligned}\int C dx_m &= 0 \\ \frac{d}{df} \int C dx_m &= 0^{0'} \\ &= e^{x \log x}\end{aligned}$$

と標数0の体の上の代数多様体でもあり、このオイラーの定数からの大域的微分多様体から数が生まれた。

$$\begin{aligned}H\Psi &= \bigoplus (i\hbar^{\nabla})^{\oplus L} \\ &\bigoplus a^f x^{1-f} [I_m] \\ &= \int e^x x^{1-t} dx_m \\ &= e^{x \log x}\end{aligned}$$

アメリカ大統領を統計で選ぶ選挙は、reco level 理論がゼータ関数として機能する遷移エネルギーの安定軌道にある集団 x に対数 $\log x$ の組み合わせとして、指数の巨大確率を対数の個数とするこの大統領の素質としての x^n 集団の共通の思考が n となるこの n がどのくらいのエントロピー量かを $H = -Kp \log p$ が表している。

$$\begin{aligned}\int \Gamma(\gamma)' dx_m &= (e^f + e^{-f}) \geq (e^f - e^{-f}) \\ (e^f + e^{-f}) &\geq (e^f - e^{-f})\end{aligned}$$

この方程式はブラックホールのシュバルツシルト半径から

$$\begin{aligned}(e^f + e^{-f})(e^f - e^{-f}) &= 0 \\ \frac{d}{df} F \cdot \int C dx_m &\geq 0 \\ y = f(g(x)') dx &= \int f(x)' g(x)' dx \\ y = f(\log x)' dx &= f'(x) \frac{1}{x} \\ y &= \frac{f'(x)}{x} \\ &= 2(\cos(ix \log x) - i \sin(ix \log x)) \\ &= C\end{aligned}$$

$$c=f(x)\cdot\log x, dx_m=(\log x)^{-1}$$

$$C=\frac{d}{\gamma}\Gamma=\bigoplus (i\hbar\nabla)^{\oplus L}$$

となり、ヴェイユ予想の式からも導かれる。

$$=2(\cos(ix\log x)-i\sin(ix\log x))$$

$$e^{\theta}$$

$$\frac{d}{\gamma}\Gamma=\Gamma^{\gamma'}$$

$$=\int \Gamma(\gamma)'dx_m$$

$$y=f(x)\log x,y'=f'(x)+\frac{f'(x)}{x}$$

$$\sin(\log x)' \, dx = \cos(\log x) \cdot \frac{1}{x}$$

$$\sin \frac{y}{x} = \sin \vec{u} = a + t \sin \vec{u}$$

$$i,-i,2i,-2i$$

$$\lim_{n\rightarrow\infty}(f(b)-f(a))=f'(c)(b-a)$$

$$\sin(\log x)' \, dx = \cos(\log x) \cdot \frac{1}{x}$$

$$y=f(x)\log x,y'=f'(x)+\frac{f'(x)}{x}$$

$$\sin(\log x)' \, dx = \cos(\log x) \cdot \frac{1}{x}$$

$$\sin \frac{y}{x} = 2(\cos(ix\log x)-i\sin(ix\log x))$$

$$=e^{\theta}$$

Circle function and imaginary number, Euler equation from weak electric and strong electric theorem from gravity and antigravity on one rout of time system

Masaaki Yamaguchi

Rotate with pole of dimension is converted from other sequence of element, this atmosphere of vacume from being emerged with quarks of being constructed with Higgs field, and this created from energy of zero dimension are begun with universe and other dimension started from darkmatter that represented with big-ban. Therefore, this element of circumstance have with quarks of twelve lake with atoms.

$$(dx, \partial x) \cdot (\epsilon x, \delta x) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\frac{d}{df} F = m(x)$$


And this phenomounen is super symmetry theorem built with quarks of element, also this chemistry of mechanism estimate from physics of operatsion. These response of mechanism chain of geometry into space being emerged with creature and univse of existing of combination.


This converted with dimension of element also emerged from imaginary and reality of pole's space.

And this pole of transport of dimension belong with vector of time has with one rout of sequecense. Quantum physics also belong with other vector of time has with imaginary rout of sequecense.

This sequence of being estimated with non fluer of time, and this space of element have with gravity and antigravity of power. Other vecor of time is antigravity rout of sequecense.

Weak electric theory is estimate from time has with one rout of sequecense, this topology of chain is resulted from time of rout ways.





$$\square(\frac{\sigma_1 + \sigma_2}{2}) = [3\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\square\psi = 8\pi GT^{\mu\nu}$$

$$\frac{d}{dt}g_{ij}(t) = -2R_{ij}$$

$$\nabla\psi^2 = 4\pi G\rho$$

$$\square(\sigma_1 + \sigma_2) = [6\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$= [i\pi(\chi, x), f(x)]$$

$$\square\psi = [12\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\frac{d}{df}F = \int e^{-f}[-\Delta v + R_{ij}v_{ij} + \nabla_i \nabla_j f + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(\frac{v}{2} - h)]$$

$$= [i\pi(\chi, x), f(x)]$$

These equation is reminded time pass rout of one rout way of forms, and this rout of time ways which go for system from future and past. Therefore, this resulted system of time mechanism is one true flow that weak electric theorem oneselves. and moreover, one rout time way of forms is reverse with antigravity of time system. This also spectrum focus is true that Maxwell theorem and strong boson unite with antigravity, this unite is essence on the contrary from weak electric theorem, this theorem called for time rout forms is strong electric theorem. This two theorem is united with quantum physics that no time flow system.

$$f^{-1}(x)xf(x) = 1, H_m = E_m \times K_m$$

The non-commutative theorem is constructed from world line surface that this complex manifold estimate with rolanz atractor, and this string theorem have with one world of universe mate six quarks

and other world of dimension mate other element of six quarks. These particle quarks is built with super symmetry space of dimension.

$$i = (1, 0) \cdot (1, 0), e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{-i\theta} = \cos \theta - i \sin \theta$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$\sin i\theta = \frac{e^{-\theta} - e^{\theta}}{2}$$

$$\pi(\chi, x) = \cos \theta + i \sin \theta$$

In this equations, two dimension redeconstructed into three dimension, this destroy of reconstructed way is append with fifth dimension. This deconstructed way of redeconstructe is arround of universe attached with three dimension, this over cover call into fifth dimension.

$$R(-\alpha) = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha)MR(-\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos^2 \alpha - \sin^2 \alpha & 2 \sin \alpha \cos \alpha \\ 2 \sin \alpha \cos \alpha & -\cos^2 \alpha + \sin^2 \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & -1 \\ 1 & -\sin \alpha \end{pmatrix} \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$f^{-1}xf(x) = 1$$

$$(\log x)' = \frac{1}{x}, x^n + y^n = z^n$$

$$x^n = -y^n + c, nx^{n-1} = -ny^{n-1}y'$$

$$y' = \frac{nx^{n-1}}{ny^{n-1}}$$

$$= \frac{x^{n-1}}{y^{n-1}} = -\frac{y}{x} \cdot \left(\frac{x}{y}\right)^n$$

$$\begin{aligned}
& -\frac{\cos x}{(\cos x)'}(\sin x)' = z_n \\
& z^n = -2e^{x \log x} \\
& \lim_{x \rightarrow \infty} f(x) = a, \lim_{y \rightarrow \infty} f(y) = b, \lim_{x, y \rightarrow \infty} \{f(x) + f(y)\} = a + b \\
& \lim_{z \rightarrow 0} f(z) = c, \delta \int z^n = \frac{d}{dV} x^3 \\
& \lim_{x \rightarrow \infty} = c - \lim_{y \rightarrow \infty} f(y) \lim_{x \rightarrow \infty} f(x) + \lim_{y \rightarrow \infty} f(y) = \lim_{z \rightarrow \infty} f(z) \\
& z^n = \cos n\theta + i \sin n\theta \\
& = -2e^{x \log x}
\end{aligned}$$

These equation is gravity and antigravity equation.

$$\begin{aligned}
& \frac{d}{d\sigma} \left[\frac{(\sigma_1 + \sigma_2)}{2} \right] \\
& = \sigma(\uparrow\downarrow) + \sigma(\uparrow\uparrow) + \sigma(\downarrow\downarrow) + \sigma(\Leftarrow) + \sigma(\Rightarrow) \\
& \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \sigma(\uparrow) \\
& \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \sigma(\downarrow) \\
& \sigma(\Leftarrow) + \sigma(\Rightarrow) = \int e^{-f} [-\Delta v + R_{ij} v_{ij} + \nabla_i \nabla_j v + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(v - \frac{h}{2})] \\
& \sigma(\downarrow) = \sigma(\uparrow\downarrow + \uparrow\uparrow + \Rightarrow) \\
& \sigma(1) = \sigma(\uparrow\downarrow + \downarrow\downarrow + \Leftarrow) \\
& \text{weak electric theorem} = \sigma(1) \\
& \text{strong electric theorem} = \sigma(\downarrow)
\end{aligned}$$

These equation is represented with topology of string model, and weak electric theorem is constructed with Maxwell theorem and weak boson, gravity that estimate with this three power united. Moreover, strong boson and Maxwell theorem, antigravity that also estimate with this three power united. This two united power is integrated from gravity and antigravity. Then this united power is zeta function.

大域的偏微分と大域的多重積分、 大域的部分積分についてのレポート

Masaaki Yamaguchi

$$\log x|_{g_{ij}}^{\nabla L} = f^{f'}, F^f|_{g_{ij}}^{\nabla L} : x \rightarrow y, x^p \rightarrow y$$

$$f(x) = \log x = p \log x, f(y) = p \log x$$

大域的偏微分方程式と大域的多重積分は、それぞれ次のように成り立っている。

$$\frac{d^2}{df^2} F = F^{f'} \cdot f^{f''},$$

$$\int \int F dx_m = F^f \cdot F^{(f)'}$$

$$\frac{d}{df dg} (f, g) = (f \cdot g)^{f' + g'}$$

大域的部分積分も、次のように成り立っている。

$$(F^f \cdot G^g) = \int (f \cdot g)^{f' + g'}$$

$$\int \int F \cdot G dx_m = [F^f \cdot G^g] - \int (f \cdot g)^{f' + g'}$$

大域的部分積分の計算は、

$$\int \frac{d}{df dg} FG = \int F^{f'} G + \int FG^{g'}$$

$$\int F^{f'} G dx_m = [F^f G^g] - \int FG^{g'} dx_m$$

大域的商代数の計算は、

$$\left(\frac{F(x)}{G(x)} \right)^{(fg)'} = \frac{F^{f'} G - FG^{g'}}{G^g}$$

大域的偏微分方程式は、縮約記号を使うと、

$$\frac{d}{df dg} FG = \frac{d}{df_m} FG$$

$$\frac{\partial}{\partial f_m} FG = F^{f^{\mu\nu}} \cdot G + F \cdot G^{g^{\mu\nu}}, \int F dx_m = F^f$$

多様体による大域的微分と大域的積分が、エントロピー式で統一的に表せられる。

Rsa secret data and non commutative equation, information technology system

Masaaki Yamaguchi

Non certain theorem is component with quantum computer, this system emelites with contemporality of eternal universe and secret of database.

不確定性理論の、宇宙の準同型写像の、部分群と全射としての、閉3次元多様体のエントロピー値としての、暗号としての量子コンピューターが、Jones多項式を形成しているオイラーの公式の虚数として、ダミーを入れると、Rsa 暗号の公開鍵として、成り立っている量子暗号が、この量子コンピューターと一緒に、ゼータ関数と量子群を大域的微分方程式を、これらの方程式から統合されて、大域的トポロジーが、数学の数論と幾何学、解析学として、物理学と言語学、情報科学、すべての理論に統合されて、ゼータ関数が、統一場理論の発生する式になっている。この理論を彩さんとグリーシャさん、ナッシュさん、トビーケリーさん、小方くん、益川先生、南部先生、岡本教授、リサ・ランドール教授、竹内先生、私の子ども、今までの人たちが、集まって、この理論ができている。量子的な微分・積分が、きっかけにもなっている。アイデアと計算方法は、情報空間の子どもと、お姉さんとお父さんと先生たちで、統一場理論としては、グリーシャさんと竹内薫先生と彩さん、岡本教授がきっかけになっている。

This information of quantum computer ansterise in secret of each data, and this system is entersteam by Jones manifold of equation, more also this intersect with system of pair of universe and the other dimension with Jones manifold of equation pawn for non access of chain in disenable of database. And this insterm of system built with

$$e^{-\theta} = e^f + e^{-f}, e^{i\theta} = \cos \theta + i \sin \theta$$

$$\frac{d}{df} F = e^f + e^{-f} = 2i \sin(ix \log x)$$

These equations are atom of dense with norm and energe of non certain theorem, and this system component with also Jones manifold of equation. This equation addesent with a dummy data in non integrate of relativity theory for quantum physics and access to varnished with aseterise for important of discover with secret of data. This dummy of information of

quantum secret datas are non commutative equation of declinate anterave and distarbration into universe and other dimension for accesslable with reterbled with quantum equation of deep of Riemann metrics, more also this system is computed with after all intersected with eternal data into Zeta function of global differential equation and Higgs fields with time system enterprises. Global of open key in Rsa secret system intersect with this base of this technolgie with all of data for established with information technology and quantum computer created with dummy of integral non relativity of quantum equation, this focus is diserble with quantum equation of a data for dummy into Rsa secret data of non commutative equation, this system comute with after all established with information of quantum physics.

$$e^{i\theta} = \cos \theta + i \sin \theta$$

this equation is based into Jones manifold equation.

$$e^{-\theta} = e^f + e^{-f}, \frac{d}{df} F + \int C dx_m = 2(\cos(ix \log x) + i \sin(ix \log x))$$

These equation are based with reco level theory and field of physics, and these system are

$$||ds^2|| = \int \exp[\frac{-L(x)}{\sqrt{2q\tau}}] dx_m + \mathcal{O}(N^{-1})$$

$\mathcal{O}(N^{-1})$ is dummy data of intersected of system.

These system are chain with non certain theorem component with aspe experiement mechanism clearlity of quantum teleportation physics and this system is pair of existanse with Jones manifold equation and imarginary circle equation more also these equations are rhysmiary equation excluded with zeta function and quantum equation, and this system commutave with each attitude of pair in Jones manifold equation.

These theorem combuild with Lie algebra and catastorophe of summative group in eternal Garois theory, and this reasons of between Poancare conjecture in zero dimension and eternal group with nesseciry and mourdigate of conditions.

AdS_5 equation are built with cercumention dimension of assemble D-brane and information of quantum physics with secret system, and this system construct with a certain theorem.

$$||ds^2|| = \eta_{\mu\nu} + \bar{h}(x)$$

This dimension esterned with equals of dummy data in the other dimension, and this value of data declined into only universe, then this sertation of universe is possibilty equation. After all this anserration of univese addsentend

with the other dimension escort for certain theorem and all of addsent equation are three manifold entropy equation, this equation constructed with zeta function and quantum equation.

$$\begin{aligned}
||ds^2|| &= 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right) \\
2\sqrt{\frac{V}{S}} &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m \\
\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &= \int dx_m + x^2 \\
&\geq \int dx_m \\
\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m &\geq \frac{1}{2}i
\end{aligned}$$

Quantum computer concluded with component of being describe with estimate for each atom conditions. This pair of condition is entertained for orbital of circle with Euler equation. This pair of orbital area estertate with two of circumentations. These system concerned with hyper synchronized of all of possibility resulted for being computing of all of area in Euler equation. This cercuit with equations are that information of quantum physics esterned with Rsa secret datas system.

These system of stem are constructed with declinate anterave of Lambda driver and possibilty computer for resolved with synchronize of distarbrate non certain theorem, and this concerned with system component of Euler equation. This estimate of orbital construct with curbit of quarters. Artificial intelligence is concluded with Euler equation. Zeta function also integrate with this equation. これらの理論の中核の, 中核となっている、オイラーの公式の群ともなっている、過冷却金属の原理の、高熱を急激に冷ます、電子を集める、ラムダ・ドライバーにもなっている、不確定性理論の原理にもなっている、暗号を解読するのを防ぐのに、クオータのキュービットの、遷移元素を作り出す原子の仕組みにもなっている、この理論が、量子コンピュータと人工知能の中核となっていて、オイラーの公式の虚数ともなっている、この Jones 多項式の統合にゼータ関数が使われている。[参考文献は、彩さん、グリーシャさん、ナッシュさん、トビーケリーさん]

Artificial Intelligence and TupleSpace of ultranetwork

Masaaki Yamaguchi

クラウドにデータベースを構築しておいて、この構築した多様体を数式で表現したコード通りのデータが、この多様体において、作用素関数として実行されるとする。この多様体を実現したデータが表現されている環境自体を表せられるソースとして、TupleSpace が辞書を書き換えることができないことを利便して、どのデータも上書きされないことによって、前後の記憶が無駄なコードが作用されないことを表現できる。

作用素環プログラミングとして、半静性型宣言子をつくる。この宣言子は、スクリプト型プログラミング言語では、この型を作り上げた時点で、その宣言した環境としての多様体がデータベースの仕様として、宣言した以後のソースコードがこのコード自体の性質を反映させることが多様体を表現した後の、配列、ハッシュ、文字列、ポインタ、ファイル構造体、オブジェクト、数値、関数、正規表現、行列、統計、微分、積分、この微分、積分は関数とは別の文字列と数値処理として、行列と統計をこの表現としての多様体として、微分、積分を数列を応用とした極限值としてソースコードをコンピュータにおいて、実行、表現、存在できるコンピュータ上だけにとどまらないプログラミング言語として、調べられる。この作用素としての半静性型宣言子は、スクリプト言語において、重要な研究として、動的と静的な宣言子として、なぜ静的宣言子が動的スクリプトで必要とされているかが、Stream と Ruby を学んでいく段階で浮かび上がった課題として、私は Ruby をオブジェクト指向を学んだ結果が、この作用素環プログラミングをプログラム思考でコンピュータに人工知能を生成出来て、人体の量子コンピュータを模擬出来て、その上に、FPGA までも実行できるアスペクト型人工知能スクリプト言語が、この多様体を数式を文字列としてだけではなく、電気信号としての表現体としてコンピュータ上に実現できることを研究課題として、生まれている。

Omega::DATABASE を tuplespace としてスクリプトに書き上げているソースをデータベースの下地とする。これをコンピュータに多様体として表現、実行、流れとして、動的に実行する。この実行した後に、スクリプト言語の動作を停止した場合は、ガベージコレクションとして破棄されるとする。この動作している状態のときに、同時に実行される関数、オブジェクト、文字出力は、このときに同時に起動している多様体の性質をウェブのネットワーク上で多様体の記述されている規則、ルールに則ってプログラミング言語でコンピュータに作用させている、最終的な産物のゼータ関数としてのガンマ関数の大域的微分多様体を熱エントロピー値として、この熱値の性質として分類、整列される TupleSpace 上の関数の群論として、なにがコンピュータ上だけでなく、存在論だけにとどまらない電気信号かが、数学と情報科学で研究されるべきと、この多様体を調べることが必要と、目下の課題になっている。

現実の世界として、この世界を架空化する空間が同型としてのフェルミオンとボソンが、この空想上での入れ物に電気信号としての文字列がバーチャルネットワークに出力されて、この出力される文字列と電気信号が架空の性質として、物体や生命に現実の世界としての相対的な実存を特徴、成分、性質、分類としてコンピュータに文字列として命を吹き込む機能をプログラミング言語で生成されたバーチャルコードによって生み出せる可能性を秘めている。

```

Omega::DATABASE[tuplespace]
{
  Z \supset C \bigoplus \nabla R^{+}, \nabla(R^{+}
  \cap E^{+}) \ni x, \Delta(C \subset R) \ni x
  M^{+}_{-}\bigoplus R^{+}, E^{+} \in
  \bigoplus \nabla R^{+}, S^{+}_{-} \subset R^{+}_{2},
  V^{+}_{-} \times R^{+}_{-} \cong \{V \over S\}
  C^{+} \cup V^{+}_{-} \ni M_{1} \bigoplus \nabla C^{+}_{-},
  Q \supseteq R^{+}_{-},
  Q \subset \bigoplus M^{+}_{-},
  \bigotimes Q \subset \zeta(x), \bigoplus \nabla C^{+}_{-} \cong M_3
  R \subset M_3,
  C^{+} \bigoplus M_n, E^{+} \cap R^{+},
  E_2 \bigoplus E_1, R^{-} \subset C^{+}, M^{+}_{-}
  C^{+}_{-}, M^{+}_{-} \nabla C^{+}_{-}, C^{+} \nabla H_m,
  E^{+} \nabla R^{+}_{-}, E_2 \nabla E_1,
  R^{-} \nabla C^{+}_{-}

  [- \Delta v + \nabla_{\{i\}} \nabla_{\{j\}} v_{\{ij\}} - R_{\{ij\}} v_{\{ij\}}
  - v_{\{ij\}} \nabla_{\{i\}} \nabla_{\{j\}} + 2 < \nabla f, \nabla h >
  + (R + \nabla f^2)(\{v \over 2\} - h)]

  S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1,
  H^1 \times S^1, H^1, S^2 \times E
}

```

クラウドにおけるデータベースを多様体が機能する仕組みからデータの相互関係と各データの処理対応が数学における多様体からソースコード化できる。

まず始めに、ソースコードを記述する人が定義したデータベースをライブラリーとして、動的にスクリプト言語に取り込む

```

import Omega::Tuplespace < DATABASE
{
  {\bigoplus M^{+}_{-} -> =: \nabla R^{+} \nabla C^{+}}-< [construct_emerge_equation.built]
  >> VIRTUALMACHINE[tuplespace]
  => {regextpt.pattern |w|
    w.scan(equal.value) [ > [\nabla \int \int \nabla_{\{i\}} \nabla_{\{j\}} f \circ g(x)]
    equal.value.shift => tuplespace.value
    w.emerged >> |value| value.equation_create
    w <- value
    w.pop => tuplespace.value
  }
}

```

多様体の式をバーチャルマシンに方程式としてと、データベースとして

```

{\bigoplus M^{+}_{-} -> =: \nabla R^{+} \nabla C^{+}}-< [construct_emerge_equation.built]

```

```
>> VIRTUALMACHINE[tuplespace]
```

多様体の式を分岐したリストで、配列に生成される方程式の再構築とバーチャルマシンに >> で入力する。

```
=> {regextpt.pattern |w|
  w.scan(equal.value) [ > [\nabla \int \int \nabla_{i}\nabla_{j} f \circ g(x)]]
  equal.value.shift => tuplespace.value
  w.emerged >> |value| value.equation_create
  w <- value
  w.pop => tuplespace.value
}

}
```

このバーチャルマシンに入力されたデータを正規表現で共通要素を抽出して、これも配列に入っている定義されている多様体へ、数値解析として>と入力する。この共通データをデータの端から取り除く値を tuplespace の値としてリスト化する。この抽出されてデータを、データベースに取り込んでいる多様体の規則からトリガーとして機能を発動させる。この多様体の値を再び正規表現として<-と入力する。このデータベースの全データを取り入れた段階で再構築して、生成し直す。

もとのデータ >> 対象物のデータ、>> は文字入力機能を表す。
もとのデータ >- 対象物のデータ、>- はデータの分岐の流れを作る。

```
{\vee (\int \nabla_{i}\nabla_{j} (R + \Delta f)^2)
 \over \exists (R + \Delta f)} -> =: variable array[]
>> VIRTUAL_MACHINE[tuplespace]
=> {regextpt.pattern |w|
  w.emerged => tuplespace[array]
  w <- value
  w.pop => tuplespace.value
}
```

多様体を入力する配列を -> =: 変数 array[] と表す。
>> は、データベースに配列として入力する。
このデータベースに機能しているデータを正規表現として扱うように{equation}=>{regextpt.pattern}へデータを流す。そのあとに、自動でデータを更新する。

```
Omega.DATABASE[tuplespace]->w.emerged >> |value| value.equation_create
{
  w.process <- Omega.space
  {=>
    cognitive_system :=> tuplespace[process.excluded].reload
    assembly_process <- w.file.reload.process
```

```

=> : [regext.pattern(file)=>text_included.w.process]
}
}

```

データベースから正規表現で生成された変数値から、それにポインタされた方程式を、データベースをもとで生成する。この生成された中で、ソースコードを正規表現にプロセス、マルチスレッド化して、外部のデータを後ろからポインタとして、連結する。w.process <- Omega.space として上の表現として表している。

これもデータベースとして扱い、そのコード実行の中で、作用素環機能として、だが、機能ではなく、機能をポインタで指されているアドレスに存在定義されている cognitive_system を機能を実現させる一種の合言葉として、tuplespace[process.excluded] ヘデータを :=> を使い、流す。これを reload する。assembly_process も変数のような定数で表せられて、この変数に w.file.reload.process としてポインタを当てる。この当てられた assembly_process を配列のデータベースへ正規表現をファイルに記述されているデータとして再取り込みを行う。

```

Omega.DATABASE[tuplespace]->w.emerged >> |list| list.equation_create
{
  w.process <- Omega.space
  {=>
    poly w.process.cognitive_system :=> tuplespace[process.excluded].reload
    homology w.process :=> tuplespace[process.excluded].reload
    mesh.volume_manifold :=> tuplespace[process.excluded].reload
    \nabla_{i}\nabla_{j} w.process.excluded :=> tuplespace[process.excluded].reload
    {\exp[\int \int (R + \Delta f)^2 e^{-x \log x}dV}.emerge_equation.reality{|repository|
      repository.regext.pattern => tuplespace[process.excluded].reload
      tuplespace[process.excluded].rebuild >> Omega.DATABASE[tuplespace]
      {\imaginary.equation => e^{\cos \theta + i \sin \theta}} <=> Omega.DATABASE[tuplespace]
      {{d \over df}F ==> {d \over df}{1 \over {(x \log x)^2 \circ (y \log y)
        ^{1 \over 2}}}}dm}.cognitive_system.reload
      :=> [repository.scan(regext.pattern) { <=> btree.scan |array| <-> ultranetwork.attachment}
      repository.saved
    }
  }
}
}

```

データベースから連想リスト構造の方程式を生成して、

このデータの tuple から、外部でのスペースに記述されているデータをポインタとして指して、取り込む。

この取り込んでいるデータを、作用素環の半静性宣言子としての、poly,homolgy,equation が記述されているソース

の式を使って、データを各ポインタを指しているデータ自体にリンクとして双対性をプログラミングしていく。

:=>, >> ==> ,<=> .emerge_equation.reality, .reload, .cognitive_system, .reload, .saved の各ポインタを指すための代入子、入力子、等号入力子、倒置入力子、記述されている方程式を生成する、それを実行する。再取り込み、連想配列生成、保存を各レシーバはオブジェクトから保持している機能呼び起こせられる。

```

import ultra_database.included
def < this.class::Omega.DATABASE[first,second,third.fourth] end
  def.first.iterator => array.emerge_equation
  def.second.iterator => array.emerge_equation
  def.third.iterator => array.emerge_equation
  def.fourth.iterator => array.emerge_equation
  _ struct_ {
    Omega.iterator => repository.reload
  }
end
  typedef _ struct_ :Omega.aspective
end

```

今までのデータベースをウルトラネットワークとくして、取り込む。この多様体がデータベースとして宣言されている情報空間へ関数のメソッドとして、データベースに記述されている機能としてのメソッドとして各正規表現を配列に入っている文字列から方程式として、多様体のデータベースの単体量へポインタを介して、関数のメソッドのハッシュを作り、このポインタへの各要素のデータベースをリポジトリとしての構造体へとアスペクト指向として、関数定義する。

```

Omega::DATABASE[reload]
{
  [category.repository <-> w.process] <=> catastrophe.category.selected[list]
  list.distributed => ultra_database.exist ->
  w.summurate_pattern[Omega.Database]
  btree.exclude -> this.klass
  list.scan(regexpt.pattern) <-> btree.included
  list.exclude -> [Omega.Database]
  all_of_equation.emerged <=> Omega.Database
  {
    list.summuate -> Omega.Database.excluded
  }
}

```

今までのデータをデータベースにリロードして、その中で、不変性を見つけて分類していく。この分類された連想配列によるリスト構造をウルトラネットワークへ双対性 $=_i$ をつかって、 $-_i$ と統合されるべきパターンへと流す。これを btree 構造体にポインタをつなげて、リスト化して、各リストを再びデータベースへとつなげる。今までの方程式をデータベースの中の多様体に入れて、相互に比較してリスト構造体を再編成する。

```

list.distributed => {
  {\bigoplus \nabla M^{+}_{-}}.constructed <-> Omega.Database[import]
  {=>
    each_selected :file.excluded
  }
}

```

この再編成されたリストを自分が導いた方程式が、どの範疇のデータで、何の方程式かを、多様体から意思

が生成された認知でもある場の理論として、判断させて、未知の理論を多様体からの人工知能で見つける。

これらのデータベース化されたリストから、レシーバでもある、前からの宣言と後ろからの、レシーバで

オブジェクトとして、リスト化したデータへ、以下の式たちを入力させる。equation_manifolds.scan(value) |value| と=|value| 以下で数式たちを代入=している。

```
Omega::DATABASE[tuplespace] >> list.cognitive_system |value|
= { x^{\frac{1}{2}} + iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h
x^{\frac{1}{2}} + iy = \mathrm{exp}[\int \nabla_i \nabla_j f(g(x)) g'(x) /
\partial f \partial g]

\mathcal{O}(x) = \{[f(x) \circ g(x), \bar{h}(x)], g^{-1}(x)\}

\exists [\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty}
\nabla \int \nabla_i \nabla_j f(x) dm

\vee (\nabla_i \nabla_j f) = \bigotimes \nabla E^{+}

g(x,y) = \mathcal{O}(x)[f(x) + \bar{h}(x)] + T^2 d^2 \phi

\mathcal{O}(x) = \left( \int [g(x)] e^{-f} dV \right)' - \sum \Delta(x)
\mathcal{O}(x) = [\nabla_i \nabla_j f(x)]' \cong {}_nC_r f(x)^n
f(y)^{n-r} \Delta(x,y),
V(\tau) = \int [f(x)] dm / \partial f_{xy}

\square \psi = 8 \pi G T^{\mu\nu}, (\square \psi)' = \nabla_i \nabla_j
(\Delta(x) \circ G(x))^{\mu\nu}
\left( \frac{p}{c^3} \circ \frac{V}{S} \right), x^{\frac{1}{2}} + iy = e^{x \log x}

\Delta(x) \phi = \{ \vee [\nabla_i \nabla_j f \circ g(x)] \over
\exists (R + \Delta f) \}

{}_nC_r = {}_{\frac{1}{i}H\psi} C_{\hbar \psi} + {}_{[H, \psi]} C_{n-r}
{}_nC_r = {}_nC_{n-r}

\int \int \frac{1}{(x \log x)^2} dx_m \to \mathcal{O}(x) =
[\nabla_i \nabla_j f]' / \partial f_{xy}

\bigcup_{x=0}^{\infty} f(x) = \nabla_i \nabla_j f(x) \oplus \sum f(x)
= \bigoplus \nabla f(x)
\nabla_i \nabla_j f \cong \partial x \partial y \int
\nabla_i \nabla_j f dm
\cong \int [f(x)] dm
```

$\int [f(x), g(x)] g^{-1}(x) dx$
 $\int \sqrt{\psi}$
 $\int \nabla \psi^2$
 $f(x \circ y) \leq f(x) \circ g(x)$
 $\int |f(x)| + |g(x)|$

$\Delta(x) \psi = \langle f, g \rangle \circ |h^{-1}(x)|$
 $\partial_x \cdot \Delta(x) \psi = x$
 $x \in \mathcal{H}_0(x)$
 $\mathcal{H}_0(x) = \{[f \circ g, h^{-1}(x)], g(x)\}$

$\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f = [\nabla \int$
 $\nabla_{i_1} \nabla_{j_1} f(x) dx_m, g^{-1}(x)] \rightarrow \bigoplus_{k=0}^{\infty}$
 ∇E^{+}_{-}
 $= M_3$
 $= \bigoplus_{k=0}^{\infty} E^{+}_{-}$
 $dx^2 = [g^2_{\mu\nu}, dx], g^{-1} = dx \int \Delta(x) f(x) dx$
 $f(x) = \exp[\nabla_{i_1} \nabla_{j_1} f(x), g^{-1}(x)]$
 $\pi(\chi, x) = [\pi(\chi, x), f(x)]$
 $\left(\frac{g(x)}{f(x)} \right)' =$
 $\lim_{n \rightarrow \infty} \left\{ \frac{g(x)}{f(x)} \right\}$
 $= \left\{ \frac{g'(x)}{f'(x)} \right\}$

$\nabla F = f \cdot \{1 \over 4\} |r|^2$
 $\nabla_i \nabla_j f = \{d \over dx_i\}$
 $\{d \over dx_j\} f(x) g(x)$
 $D^2 \psi = \nabla \int (\nabla_i \nabla_j f)^2 d\eta$
 $E = m c^2, E = \{1 \over 2\} m v^2 - \{1 \over 2\} k x^2, G^{\mu\nu} =$
 $\{1 \over 2\} \Lambda g_{ij},$
 $\square = \{1 \over 2\} k T^2$

$\ker f / \operatorname{im} f \cong S^{\mu\nu}_m,$
 $S^{\mu\nu}_m = \pi(\chi, x) \otimes h_{\mu\nu}$

$D^2 \psi = \mathcal{H}_0(x) \left(\frac{p}{c^3} + \right.$
 $\left. \frac{V}{S} \right), V(x) = D^2 \psi \otimes M^+_3$

$S^{\mu\nu}_m \otimes S^{\mu\nu}_n =$
 $- \{2R_{ij} \over V(\tau)\} [D^2 \psi]$

$\nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] =$
 $\int \{V(\tau) \over f(x)\} [D^2 \psi]$
 $\nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] =$
 $\int \{V(\tau) \over f(x)\} \mathcal{H}_0(x)$

$z(x) = \{g(cx + d) \over f(ax + b)\} h(ex + l)$
 $= \int \{V(\tau) \over f(x)\} \mathcal{H}_0(x)$

$\{V(x) \over f(x)\} = m(x), \mathcal{H}_0(x) = m(x) [D^2 \psi(x)]$
 $\{d \over df\} F = m(x), \int F dx_m = \sum_{k=0}^{\infty} m(x)$

$$\mathcal{O}(x) = \left([\nabla_i \nabla_j f(x)] \right)^{\prime}$$

$$\cong \{ {}_nC_r(x)^n (y)^{n-r} \Delta(x,y)$$

$$(\square \psi)' = \nabla_i \nabla_j (\Delta(x) \circ$$

$$G(x))^{\mu \nu} \left(p \over c^3 \right) \circ$$

$$\{ V \over S \} \right)$$

$$F^m_t = \{ 1 \over 4 \} g^2_{ij}, \ x^{\{ 1 \over 2 \} + iy} = e^{x \log x}$$

$$S^{\mu \nu}_m \otimes S^{\mu \nu}_n = G_{\mu \nu} \times T^{\mu \nu}$$

$$S^{\mu \nu}_m \otimes S^{\mu \nu}_n = -2 R_{ij} \over V(\tau) [D^2 \psi]$$

$$S^{\mu \nu}_m = \pi(\chi, x) \otimes h_{\mu \nu}$$

$$\pi(\chi, x) = \int \mathrm{exp}[L(p, q)] d\psi$$

$$ds^2 = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu \nu} dx^{\mu \nu} dx^{\mu \nu} +$$

$$T^2 d^2\psi$$

$$M_3 \bigotimes_{k=0}^{\infty} E^{+}_{-} = \mathrm{rot}$$

$$(\mathrm{div} \, E, E_1)$$

$$= m(x), \{ P^{2n} \over M_3 \} = H_3(M_1)$$

$$\exists [R + |\nabla f|^2]^{\{ 1 \over 2 \} + iy}$$

$$= \int \mathrm{exp}[L(p, q)] d\psi$$

$$= \exists [R + |\nabla f|^2]^{\{ 1 \over 2 \} + iy} \otimes$$

$$\int \mathrm{exp}[L(p, q)] d\psi +$$

$$N \bmod (e^{x \log x})$$

$$= \mathcal{O}(\psi)$$

$$\{ d \over dt \} g_{ij}(t) = -2 R_{ij}, \{ P^{2n} \over M_3 \}$$

$$= H_3(M_1), H_3(M_1) = \pi(\chi, x) \otimes h_{\mu \nu}$$

$$S^{\mu \nu}_m \times S^{\mu \nu}_n$$

$$= [D^2 \psi], S^{\mu \nu}_m \times S^{\mu \nu}_n$$

$$= \mathrm{ker} f / \mathrm{im} f, S^{\mu \nu}_m \otimes$$

$$S^{\mu \nu}_n = m(x) [D^2 \psi], \{ -2 R_{ij} \over V(\tau) \} = f^{-1} x f(x)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \circ$$

$$\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}_{-} \right] dx dy dz,$$

$$\to f_z^{\{ 1 \over 2 \}} \to (0,1) \cdot (0,1) = -1, i =$$

$$\sqrt{-1}$$

$$\{ \begin{pmatrix} x, y, z \\ \end{pmatrix}^2 = (x, y, z) \cdot (x, y, z) \to -1$$

$$\mathcal{O}(x) = \nabla_i \nabla_j \int e^{\{ 2 \over m \} \sin \theta}$$

$$\cos \theta \times \{ N \bmod$$

$$(e^{x \log x})$$

$$\over \mathcal{O}(x) (x + \Delta |f|^2)^{\{ 1 \over 2 \}}$$

$$x \Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta,$$

$$\mathcal{O}(x) = m(x) [D^2 \psi]$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$f^{-1}(x) \times f(x) = I_m, \quad I_m = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$i^2 = (0,1) \cdot (0,1), |a||b|\cos \theta = -1,$$

$$E = \operatorname{div}(E, E_1)$$

$$\left(\frac{f,g}{[f,g]} \right)' = i^2, \quad E = mc^2, \quad I' = i^2$$

$$\mathcal{C}_0(x) = || \nabla \int [\nabla_i \nabla_j f \circ g(x)]^{\frac{1}{2} + i y} ||, \quad \partial r^n$$

$$||\nabla||^2 \rightarrow \nabla_i \nabla_j ||\vec{v}||^2$$

$$\nabla^2 \phi$$

$$\nabla^2 \phi = 8 \pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$(\log x^{\frac{1}{2}})' = \frac{1}{2} \frac{1}{x \log x},$$

$$(\sin \theta)' = \cos \theta, \quad (f_z)' = i e^{i x \log x},$$

$$\frac{d}{df} F = m(x)$$

$$\frac{d}{df} \int \int \frac{1}{x \log x} dx_m$$

$$+ \frac{d}{df} \int \int \frac{1}{y \log y} dy_m$$

$$= \frac{d}{df} \int \int \left(\frac{1}{x \log x} \right)$$

$$+ \frac{1}{y \log y} dy_m$$

$$\geq \frac{d}{df} \int \int \left(\frac{1}{x \log x} \right)$$

$$(x \log x)^2 \circ (y \log y)^{\frac{1}{2}} dy_m$$

$$\geq 2h$$

$$\frac{d}{df} \int \int \left(\frac{1}{x \log x} \right)^2 \circ (y \log y)^{\frac{1}{2}} dy_m \geq \hbar$$

$$y = x, \quad xy = x^2, \quad (\square \psi)' = 8 \pi G$$

$$\left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

$$\square \psi = \int \int \operatorname{exp}[8 \pi G (\bar{h}_{\mu\nu})$$

$$\circ \eta_{\mu\nu}] d\mu d\psi,$$

$$\sum a_k x^k = \frac{d}{df} \sum \sum \frac{1}{a_k^2} f^k dx_k$$

$$\sum a_k f^k = \frac{d}{df} \sum \sum$$

$$\{ \zeta(s) \over a_k dx_{km} \},$$

$$a_k f^{\frac{1}{2}} \rightarrow \lim_{k \rightarrow 1} a_k f^k = \alpha$$

$$ds^2 = [g_{\mu\nu}^2, dx]$$

$$M_2$$

$$ds^2 = g_{\mu\nu}^{-1} (g_{\mu\nu}^2(x) - dx g_{\mu\nu}^2)$$

$$M_2$$

$$= h(x) \otimes g_{\mu\nu} d^2 x - h(x) \otimes dx g_{\mu\nu}(x),$$

$$h(x) = (f^2(\vec{x}) - \vec{E}^+)$$

$$G_{\mu\nu} = R_{\mu\nu} T^{\mu\nu},$$

$$\begin{aligned} \partial M_2 &= \bigoplus \nabla C^{+}_{-} \\ G_{\{\mu\nu\}} &= R_{\{\mu\nu\}} \{d \over dt\} g_{ij} = -2 R_{ij} \\ r &= 2 f^{\{1 \over 2\}}(x) \\ E^{+} &= f^{-1} x f(x), \\ h(x) \otimes g(\vec{x}) &\cong \{V \over S\}, \\ \{R \over M_2\} &= E^{+} - \{\phi\} \\ &= M_3 \supset R, \\ M^{+}_2 &= E^{+}_{\{1\}} \cup E^{+}_{\{2\}} \rightarrow E^{+}_1 \oplus E^{+}_2 \\ &= M_1 \oplus \nabla C^{+}_{-}, (E^{+}_{\{1\}} \oplus E^{+}_{\{2\}}) \\ &\quad \cdot (R^{-} \subset C^{+}) \\ \{R \over M_2\} &= E^{+} - \{\phi\} \\ &= M_3 \supset R \\ M^{+}_3 &\cong h(x) \cdot R^{+}_3 \\ &= \bigoplus \nabla C^{+}_{-}, \\ R &= E^{+} \oplus M_2 - (E^{+} \cap M_2) \\ E^{+} &= g_{\{\mu\nu\}} dx_{\{\mu\nu\}}, \\ M_2 &= g_{\{\mu\nu\}} d^2x, \\ F &= \rho \, g \, 1 \rightarrow \{V \over S\} \\ \mathcal{H}_0(x) &= \delta(x) [f(x) + g(\bar{x})] + \rho \, g \, 1, \\ F &= \{1 \over 2\} m v^2 - \{1 \over 2\} k x^2, \\ M_2 &= P^{2n} \\ r &= 2 f^{\{1 \over 2\}}(x), \\ f(x) &= \{1 \over 4\} |r|^2 \\ \\ V &= R^{+} \sum K_m, W = C^{+} \sum^{\infty}_{k=0} K_{n+2}, \\ V/W &= R^{+} \sum K_m / C^{+} \sum K_{n+2} \\ &= R^{+} / C^{+} \sum \{x^k \over a_k f^k(x)\} \\ &= M^{+}_{-}, \{d \over df\} F = m(x), \rightarrow M^{+}_{-}, \sum^{\infty}_{k=0} \\ \{x^k \over a_k f^k(x)\} &= \{a_k x^k \over \\ \zeta(x)\} \\ \\ \{f, g\} \over [f, g] &= \{fg + gf \over fg - gf\}, \\ \nabla f = 2, \partial H_3 = 2, \{1 + f \over 1 - f\} &= 1, \\ \{d \over df\} F = \bigoplus \nabla C^{+}_{-}, \vec{F} &= \\ \{1 \over 2\} \\ H_1 \cong H_3 &= M_3 \\ \\ H_3 \cong H_1 \rightarrow \pi(\chi, x), H_n, H_m &= \\ \mathrm{rank}(m, n), \mathrm{mesh}(\mathrm{rank}(m, n)) \lim \mathrm{mesh} \rightarrow 0 \\ (fg)' &= fg' + gf', (\{f \over g\})' = \{f'g - g'f \over g^2\}, \\ \{f, g\} \over [f, g] &= \{(fg)' \otimes dx_{fg} \over \\ (\{f \over g\})' \otimes g^{-2} dx_{fg}\} \\ &= \{(\{fg\}' \otimes dx_{fg}) \over (\{f \over g\})' \otimes g^{-2} dx_{fg}\} \\ &= \{d \over df\} F \\ \\ \hbar \psi = \{1 \over i\} H \Psi, i[H, \psi] = -H \Psi, \{f, g\} \over [f, g] &= (i)^2 \\ \\ [\nabla_i \nabla_j f(x), \delta(x)] &= \nabla_i \nabla_j \\ \int f(x, y) dm_{xy}, f(x, y) &= [f(x), h(x)] \times [g(x), h^{-1}(x)] \\ \delta(x) &= \{1 \over f'(x)\}, [H, \psi] = \Delta f(x), \\ \mathcal{H}_0(x) &= \nabla_i \nabla_j \int \delta(x) f(x) dx \\ \mathcal{H}_0(x) &= \int \delta(x) f(x) dx \end{aligned}$$

$R^{+} \cap E^{+}_{-} \ni x, M \times R^{+} \ni M_3, Q \supset C^{+}_{-},$
 $Z \in Q \nabla f, f \cong \bigoplus_{k=0}^n \nabla C^{+}_{-}$
 $\bigoplus_{k=0}^{\infty} \nabla C^{+}_{-} = M_1, \bigoplus_{k=0}^{\infty} \nabla M^{+}_{-} \cong E^{+}_{-},$
 $M_3 \cong M_1 \bigoplus_{k=0}^{\infty} \nabla \{V^{+}_{-} \over S\}$
 $\{P^{2n} \over M_2\} \cong \bigoplus_{k=0}^{\infty} \nabla C^{+}_{-}, E^{+}_{-} \times R^{+}_{-} \cong M_2$
 $\zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k,$
 $M_2 \cong P^{2n} / \ker f, \to \bigoplus \nabla C^{+}_{-}$
 $S^{+}_{-} \times V^{+}_{-} \cong \{V \over S\} \bigoplus_{k=0}^{\infty} \nabla C^{+}_{-},$
 $\nabla C^{+}_{-}, V^{+} \cong M^{+}_{-} \bigotimes S^{+}_{-},$
 $Q \times M_1 \subset \bigoplus \nabla C^{+}_{-}$
 $\sum_{k=0}^{\infty} Z \otimes Q^{+}_{-} = \bigotimes_{k=0}^{\infty} \nabla M_1$
 $= \bigotimes_{k=0}^{\infty} \nabla C^{+}_{-} \times \sum_{k=0}^{\infty} M_1, x \in R^{+} \times C^{+}_{-}$
 $\supset M_1, M_1 \subset M_2 \subset M_3$

$S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1,$
 $H^1, S^2 \times E^1.$
 $\bigoplus \nabla C^{+}_{-} \cong M_3, R \supset Q, R \cap Q,$
 $R \subset M_3, C^{+} \bigoplus M_n, E^{+} \cap R^{+}$
 $M^{+}_{-} \nabla C^{+}_{-}, C^{+} \nabla H_m, E^{+} \nabla R^{+}_{-}, E_2 \nabla E_1$
 $\$ R^{-} \nabla C^{+}_{-} \$.$ $\{\nabla \over \Delta \int x f(x) dx,$
 $\{\nabla R \over \Delta f\}, \square = 2 \{\int (R + \nabla_{\{i\}} \nabla_{\{j\}} f)^2$
 $\over -(R + \Delta f)\} e^{-f} dV$
 $\square = \{\nabla R \over \Delta f\}, \{d \over dt\} g_{\{ij\}}$
 $= \square \to \{\nabla f \over \Delta x\}, (R +$
 $|\nabla f|^2) dm \to -2(R + \nabla_{\{i\}} \nabla_{\{j\}} f)^2 e^{-f} dV$
 $x^n + y^n = z^n \to \nabla \psi^2 = 8 \pi G T^{\mu \nu},$
 $f(x+y) \geq f(x) \circ f(y)$
 $\mathrm{im} f / \ker f = \partial f, \ker f$
 $= \partial f, \ker f / \mathrm{im} f \cong$
 $\partial f, \ker f = f^{-1}(x) x f(x)$
 $f^{-1}(x) x f(x) = \int \partial f(x) d(\ker f) \to \nabla f = 2$
 $_{\{n\}} C_{\{r\}} = \{_{\{n\}} C_{\{n-r\}} \to \mathrm{im} f / \ker f$
 $\cong \ker f / \mathrm{im} f$

$\$ \sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi \$.$ this equation $\$ a_k \cong$
 $\sum_{r=0}^{\infty} \{_{\{n\}} C_{\{r\}} \$.$
 $V/W = R/C \sum_{k=0}^{\infty} \{x^k \over a_k f^k\}, W/V = C/R$
 $\sum_{k=0}^{\infty} \{a_k f^k \over x^k\}$
 $V/W \cong W/V \cong R/C (\sum_{r=0}^{\infty} \{_{\{n\}} C_{\{r\}})^{-1}$
 $\sum_{k=0}^{\infty} x^k$
 This equation is differential equation, then $\$ \sum_{k=0}^{\infty} a_k f^k \$$
 is included with $\$ a_k \cong \sum_{r=0}^{\infty} \{_{\{n\}} C_{\{r\}} \$$
 $W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{\{1-t\}} dx,$
 $\sum_{k=0}^{\infty} a_k f^k = (f^k)'$
 $\sum_{k=0}^{\infty} a_k f^k = \sum_{k=0}^{\infty} \{_{\{n\}} C_{\{r\}} f^k$
 $= (f^k)',$
 $\sum_{k=0}^{\infty} a_k f^k = [f(x)],$

```

\sum^{\infty}_{k=0} a_k f^k = \alpha, \sum^{\infty}_{k=0}
{1 \over a_k f^k}, \sum^{\infty}_{k=0} (a_k f^k)^{-1} = {1 \over 1 - z}

{\int \int {1 \over (x \log x)(y \log y)}dxy} =
{{}_nC_r^{xy} \over {({}_nC_{n-r})}}
(x \log x)(y \log y))^{-1}}
= ({}_nC_{n-r})^2 \sum_{k=0}^{\infty} ({1 \over x \log x}
- {1 \over y \log y})d{1 \over nxy} \times {xy}
= \sum_{k=0}^{\infty} a_k f^k
= \alpha
}

```

これまでのデータベース化された機能のもとでもある方程式たちを構造体として、
まとめて、=> [tuplespace] としてポインタを当てる。

```

_ struct_ :asperal equation.emerged => [tuplespace]
tuplespace.cognitive_system => development -> Omega.Database[import]
value.equation_emerged.exclude >- Omega.Database[tuplespace]

```

この連想ポインタは tuplespace 自体をオブジェクト化して、レシーバの.cognitive_system から
実装段階で、Omega データベース化する。
このデータベースの仮の方程式、未定義な式をデータベースから多様体の仕組みを利用して、
データベースから分岐して、value のオブジェクトとしてポインタを当てる。

以下のソースコードは、今まで扱っている多様体のデータを使って、アプリケーションプログラミングとし
て、即席スクリプト言語を DSL として書いている。

```

Omega::DataBase <-> virtual_connect(VIRTUALMACHINE)
{
  blidge_base.network => localmachine.attachment
  :=> {
    dhcp.etc_load_file(this.klass) {|list|
      list.connect[XWin.display _ <- xhost.in(regexpt.pattern)]
      {
        ultranetwork.def _struct {
          asperal_language :this.network_address.included[type.system_pattern]
          {|regexpt.pattern|
            <- w.scan
              |each_string| <= { ipv4.file :file.port
                                subnetmask :file.address
                                file.port <=> file.address
                                FILE *pointer

```



```

def < host_base.ethernet.connect
{
    host_name.connect => local_network
}
}
end

def < etc.load_file
{
    etc.include(inetd.rc)
    {
        virtual_connect(VIRTUAL_MACHINE){|list|
            list.attachment(etc.load_file)
        }
    }
}
end

mainloop{
    def.virtual_connect => xhost.localmachine
    {
        xhost.client <-> xhost.server
    }
    def.network.type <- [Omega.DATABASE] end
    def.etc.load_file.attachment(VIRTUAL_MACHINE) end
end
end

class UltraNetwork::DATABASE import OMEGA.TUPLESAPCE
def load_file >- VIRTUAL_MACHINE
{ in . => attachment_device |for|
for.load -> acceptance.hardware
virtual_machine.new
{
    tk.loop-> start
    XWin -multiwindow
    if dwm <-> new_xwin.start
    localhost :xhost :display -x
    xdisplay :-> [preset :XFree.demand>=needed
    for.set_up
    install_process >- tar -xvfz "#{load_file}" <-> install_attachment
    ]
    else if
    only :new_xwin.start
    localhost :xhost :multiwindow . { in
    display -x
    attachment :localhost -client
    from -client into

```

```

server.XWin -attachment}
end
condition :{ in .=>
check->[xdisplay.install_process]}
end

def < network_rout
  wireshark.start -> ethernet.device >- define rout
    rout.ipstate do |file|
      file.type <- encoding XWin -filesystem
      file.included >- make kernel_system.rebuild
      file.vmware.start do |rout|
        rout.blidgebase | rout.hostbase
      end
    end
    -> file.install
    file.address_ipstate
    => {"{file}" :> dwm.state_presense
    virtual_machine.included[file]
  }
end

def < launcher_application
  network_rout.new
  |file|
  file.attachment => { in .
  new_xwin.start :> file.included
  demand.file <- success_exit}
end

def < terminal_port
  network_rout.new
  launcher_application.new |rout|
  rout.acceptance {
    vmware.state.process |new_rout|
    new_rout : attachment.class <-> dwm.state_attachment
    new_rout -> condition.start_wmware.process}
end

def < kterm_port
  launcher_application.new
  def.included[DATABASE]
  |rout|
  rout.attachment <- |new_rout|
  new_rout.attachment do
    install.condition < rout.def.terminal_port.exclude[file]
  end
end

main_loop :file do
  kterm_port.excluded :=> VIRTUAL_MACHINE
  |new_rout| start do
    rout.process -> network_rout.rout [
    file,launcher_application, terminal_port, kterm_port].def < included
    |file|
  end
end

```

```

        file.all_attachment: file_type :=> encoding-utf8
    end
end

class < def {
    pholograph_data[] = [R,V,S,E,U,M_n,Z_n,Q,C,N,f,g]
    source_array <- pholograph_data[]
}

def > operator_data[] = {nabla,nabla_i nabla_j,Delta,partial,
                        d, int, cap,cup,ni,in,chi,oplus,otimes,bigoplus,bigotimes,d /over df,
end

def > manifold_emerge

    c = def.inject >- source_array times def.operator_data[]
    repository_data <=> c{

    c.scan(/tuplespace[]/)
    import |list| list{
        kerf = -2 \int (R + nabla_i nabla_j f)^2e^{-f}dV
        kerf / imf
        =< {d \over df}F}
    }

    equals_data =~ /list/
    list.match(/"#{c}"/) {|list|
        list.delete
        jisyo_data_mathmatics <=> list{
        list.emerge => {asperal function >- pholograph_data[] times repository_data
            =< list.update}
        }

        ln -s operator_named <= {list}
        define _struct |list|
            -> list.element -> manifold_emerge
            => list.reconstruct > def.inject /"#{pattern}"/}
    end

import Omega::Tuplespace < Database
{
    {\bigoplus \nabla M^{+}_{-}}.equation_create -> asperal :variable[array]
    :=> [cognitive_system <-> def < VIRTUALMACHINE.terminal
        {
            [ipv4.bloodcast.address :
                ipv4.network.address].subnetmask
            <-> file.port.transport_import :
                Omega[tuplespace]
        }
    }
}

```



```

_struct _ Omega[tuplespace] >> VIRTUALMACHINE.terminal.value

class < def.VIRTUALMACHINE.system_environment

    file.reload[hardware] => file.exclude >> file.attachment
    {=>
        |file|
        file.port(wireshark.rout <-> {file.port.transport_export
            :=> Omega[tuplespace]})
        assembly_process.file.included >- file.reloaded
            :- |file.environment| {=>
                file.type? :=> exist
                file.regexpt.pattern[scan.flex]
                    => |pattern|
                        <->
                            file.[scan.compiler]
            }
        end
    end
    file <<
}

Omega::Database[tuplespace]
{
    cognitive_system |: -> { DATABASE.create.regexpt_pattern >-
        cognitive_system[tuplespace].recreated >- : =< DATABASE.value
        >> system_require.application.reloaded[tuplespace]
        } : _struct _ def.VIRTUALMACHINE.terminal >> {
            ||machine.attachment|| <-> OBJECT.shift => system.reloaded
            . in {
                : _struct _ class.import :-> require mechanics.DATABASE
                {|regexpt_pattern| :|-> aspective _union _
                    def _union _}
            }
        }

    end
}

system.require <- import library.DATABASE
{
    Omega[tuplespace]
    {
        cognitive_system : VIRTUALMACHINE.equality_realized
    }
}

```

```

    {|regextpt_pattern| => value | key [ > cognitive_system.loop.stdout]
      value : display -bash :xhost -number XWin.terminal
      key   : registry.edit :=> {[cognitive_system.reloaded]}
    }
  }
}

_union _ => DATABASE[tuplespace].aspective_reloaded
_union _ :fx | -> |regextpt_pattern| => {
  VIRTUALMACHIE.recreated-> _union _ |
  _struct _ def.DATABASE.recreated <- fx
  >> DATABASE[tuplespace].rebuild
}

DATABASE[tuplespace] -< {[ > aimed.compiler | aimed.interpreter] | btree.def.distributed >-
  aimed[tuplespace]}
aimed[tuplespace] -< btree.class.hyperroot_ struct _ => Omega::Database[tuplespace].value
sheap_ union _ :aspective | -> Omega[tuplespace]: | aimed[tuplespace].differented_review
}

aimed[tuplespace].process => DATABASE[tuplespace].reloaded
aimed.different | aimed.stdout >> vale | key [ > cognitive_system.loop.stdin] {|pattern|
  pattern.scan(value : aimed[def.value]
    key       : aimed[def.key])
  } _ struct _ : flex | interpreter.system
  => expression.iterator[def.first,def.second,def.third,def.fourth]
  { def < Omega[tuplespace]
    def.cognitive_system |: -> DATABASE[tuplespace] | aimed[tuplespace]
  }
}

Omega::Tuplespace < DATABASE
{=>
  norm[Fx] -> . in for def.all_included < aimed[tuplespace].each_scan([regextpt_pattern]
    <->
      DATABASE[tuplespace]) << streem database.excluded
  >- more_pattern.scan(value : aimed[def.value]

  key :aimed[def.key])
    . in { _struct _ :flex | interpreter.system
      => expression.iterator[def.all.each -> |value, key|
        included >- norm[Fx] | [DATABASE[tuplespace]
,aimed[tuplespace]] |
        finality : aimed[tuplespace], DATABASE[tuplespace]

: -> def.included(in_all)
    {
      def.key | def,value => [DATABASE].recompile
      : in_all -> _struct _ :aspective :tuplespace
    : all_homology_created}
    }
}

```

```

def < Omega::Tuplespace[DATABASE]
  def.iterator -> |klass,define_method,constant,variable,infinity_data : -> finite_data|
  def.each_klass?{|value, key|
    _struct _ :aspective -> tuplespace :all_homology_recreated :make menuconfig
    {+=
      def.key -> aimed[def.key],def.value -> aimed[def.value] {|list|
        list.developed => <key,value> | <aimed[$',$']
        -> _union _ :value,key : _struct _
        <- (_union _ <-> _struct _ +)
      begin
        def.key <-> aimed[value]
        case :one_ exist :other :bug
        {
          result <-> def.key
          {
            differentiated :DATABASE[tuplespace]
          }
          return :tuplespace.value.shift -> included<tuplespace>
        else if
        :other :bug
        {
          success_exit <- bug[value]
          {
            cognitive_system.scan(bug[value])
            {
              {[e^{-f}][{2 \int (R + \nabla f^2) \over -(R + \Delta f)}e^{-f}dV}
            }
          }
        }
      }
    }
    .created_field
    {=>
      regxpt.pattern \native_function <-> euler-equation
      {
        $variable =< diff e^{-f} >- $'
        all_included <- def.key <-> aimed[value]
        $variable - all_included.diff
        \summate_manifold.recreated
      }
    }
    <- \native_function : euler-equation
    }
  }
  }
  } _union _ :cognitive_system.rebuild(one_ exist)
}
}
ensure
{
  return :success_exit
  => Tuplespace[DATABASE]
}
}
}
end

```

```

end
}

int
streem_style {
  :Endire <- [ADD,EVEN,MOD,DEL,MIX,INCLUDED,EXCLUDED,EBN,EXN,EOR,EXOR,
             SUM,INT,DIFF,PARTIAL,ROUND,HOMOLOGY,MESH]
  Endire.iterator -> {def < :Endire.element, -> def.means_each{x -> expression.define.included
    def.each{x -> case :x.each => :lex.include_ . in [ > [x.all_expire] ]}}
}

main_loop {
  FILE *fp :=> streem_style.address_objective_space
  fp.each{x -> domain_specific_language_style_included[array]}
  array << streem.DATABASE[tuplespace]
  array.each{[tuplespace] -> aimed[tuplespace] | OMEGA_DATABASE[tuplespace]}.excluded <-> array
  def.key <-> def.value => {x -> stdin | stdout | => streem_style <- def.each.klass.value}
}

```

リバイザを使うと、独自のタプルスペースで一時保存の書き換えの分派スクリプトが出来て、
 その応用で、例外処理機能として、そのスクリプトを使うと、独自に機能拡張できる。
 書き換え機能自体、書き換えたいとおもっている好きなところを書き換えられる。
 構文解析器も文字抽出器も全部書き換えられる。
 その書き換えたのをライブラリとして、データベースに取り込むと、この部分が例外処理機能の
 おかげで、タプルスペースが働き、今まで使ってきた機能と一線を描く。

```

@reviser : def < OmegaDatabase[tuplespace].mechanism
{
  aspective : _union _ {
    int streem_style : [ > [def.each{x -> stdin | stdout > display :xhost in XWin -multiwind
    {
      Endire <- [ADD,EVEN,ODE,EXOR,XOR,DEL,DIFF,PARTIAL,INT].included > struct _ :-> _union
      Endire.each{def.value -> def.key :hash.define}.included > _union}
    }
  }
}

@reviser : def.reconstructed.each{_union <-> _struct _ .recreated} : [def.del - def.before_deter

import perl.lib | python.lib <-> ruby.lib
{
  int @reviser : def.each{x -> x.klass |-> $variable in $stdin | $stdout}
  .developed >= {

```

```

        ping localhost -> blidgebase <-> hostbase.virtualmachine.attachment
        {
            xhost :display -> streem_style.value
            networkconnect.hostbase -> localarea.virtualmachine
        } :connected -> networkrout : flow_to :localhost.attachment
    }
}

_struct : def < hostbase.virtualmachine.attachment => : networkrout.area.build

@reviser <-> def.add [ < _struct]
@reviser : def.each{listmenu -> listlink | unlinklist > [developed -> {def.key , def.value}.current]
@reviser <-> def.rebuild [ < _struct]

@reviser.def.<value|key>networkrout-> def.present
def.present.flow_to -> hostbase.rout << networkrout.data.<value|key>

XWin -multiwindow <-> networkrout.data[$', $']
def < $'
@reviser <-> def.present.state
@reviser.def.each{x | -> key.rebuild | value.rebuild}.flow_to :redefined

def < OmegaDatabase[tuplespace]
{
    FILE *fp -> cmd.value : cmd.key {fp |-> synchronized.file[tuplespace] | aimed.file[tuplespace]}
    cmd.key => [ > fp.($':$')] <-> registry.excluded<fp.file[cmd.state]>
}

def.each{fp|-> def.first,def.second,def.third,def.fourth}


cmd _struct : {
[ ^C-O : ^C-X-F, exit.cmd : ^C-X-C, shift-up : ^C-P, shift-down : ^C-N]
}

cmd _union : def.restructured
keyhook.cmd <- : [_struct ]
{
    @reviser :def._struct <-> def. _union
}

```


素数分布論と素粒子、ゼータ関数 超対称性理論

Masaaki Yamaguchi




$$\begin{aligned}
 (u+d)+c &= e^{-f}-e^f, b+(w+s)=e^{-f}+e^f \\
 \int\int\frac{1}{x^s}\mathrm{dvol}-\int\log x\mathrm{dvol} &= \frac{d}{df}\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m-\int Cdx_m \\
 &= e^{-f}-e^f \\
 \frac{d}{df}F &= \frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m \\
 &= e^f+e^{-f}
 \end{aligned}$$

素粒子方程式は、2種類の素粒子と1種類の素粒子の組み合わせで、Jones 多項式をペアで構成されている。Rico level 理論をベースにして、強い力と弱い力、電磁気力が、誤差を D-brane 間を重力が graviton として伝わっているのが、洩れて、反重力をこの Weak electric theory に加えると、全部の力が合わさって、統一場理論が完成される。これが、ヒッグス場とオイラーの定数を多様体を次元として、加群すると、ゼータ関数になる。これが、微分幾何の量子化とガンマ関数でもある。



$$||ds^2||=e^{-2\pi T|\psi|}[\eta+\bar{h}(x)]dx^\mu dx^\nu+T^2d^2\psi$$



素数分布論が、ゼータ関数を開集合として同型でもあり、ノルム空間、経路積分として、位相差が原子間力にもなっている。このエネルギー分布が、素数と素粒子方程式、宇宙と異次元のフェルミオンとして、一致している。このエネルギー分布が、複素多様体である。

$$\square = 2(\cos(ix \log x) + i \sin(ix \log x))$$

宇宙と異次元が、双対性として、超対称性の素粒子を形成している。

[reference Lisa Randall, Toshihide Masuoka, Makoto Kobayashi]

整数論から代数、そして幾何学

Masaaki Yamaguchi

ある整数を a とおく。この a を $\log x$ で割ると

$$x = \frac{a}{\log x}$$

で求まり、ゼータ関数の x が導かれる。この x が、

$$a = x \log x$$

とシャノンのエントロピー値として、

$$\int a dx = \int x \log x dx$$

と、コルモゴロフ・シナイの値にもなる。これが、別の道でもある、求まるゼータ関数の式にもなる。この式から、

$$C = \int \left(\int \frac{1}{x^s} dx + \log x \right) d\text{vol}$$

このオイラーの定数を多様体積分すると

$$\int C dx_m = \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m + \int \log x d\text{vol}$$

となり、hartshorn conjecture の多世界理論となり、いろいろな式に分岐していく。

始めの x を見つけにくいので、オイラーの定数の多様体積分の分岐の式達がある。要するに、 x の式のヒントが幾多もある。

ガンマ関数を大域的微分多様体として共変微分すると、オイラーの定数を多様体積分した式と合流する。次元を多様体を基底群として、オイラーの定数とヒッグス場の式を加群すると、一般座標変換を大域的微分方程式として求めた結果と同型ともなる。

$$\int C dx_m + \frac{d}{df} F(x) = \Gamma^{\gamma'}$$
$$\frac{d}{d\gamma} \Gamma = \Gamma^{\gamma'}$$

これらの式が Jones 多項式となり、経済理論を Reco level theory としての一般周期関数ともなる。

$$2i \sin(ix \log x) = e^f + e^{-f}$$

これらから、素数分布論が $x = \frac{a}{\log x}$ が基底群を成して、数として、オイラーの定数から、幾多の式になり、Jones 多項式を最大数と最小数の間を一般周期関数として数学を形成して、整数論が群論となり、代数、解析、幾何学を成り立たせている。

Global Integral manifold equal with oneselves being devided with logment element

Masaaki Yamaguchi

$$\begin{aligned}
 \int \Gamma' dx &= \frac{\Gamma'}{\log x} = \Gamma \\
 \int a dx &= \frac{a}{\log x} = \Gamma \\
 \Gamma &= \int e^{-x} x^{1-t} dx \\
 a = 1, a &= \int e^{-x} x^{1-t} dx, 1 < a \\
 \frac{1}{\log x} &= \vee \int \wedge (R + \nabla_i \nabla_j f)^x \\
 &= \frac{\wedge (R + \nabla_i \nabla_j f)^n}{\exists (R + \nabla_i \nabla_j f \circ g)^n} \\
 \wedge (R + \nabla_i \nabla_j f)^x &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
 \vee (R + \nabla_i \nabla_j f)^n &= \int \frac{\wedge (R + \nabla_i \nabla_j)^2}{\exists (R + \Delta f)^n} \\
 \int \gamma dx_m &= \frac{\gamma}{\log x} = \Gamma \\
 \frac{d}{df_m} \gamma &= \frac{\gamma}{\log x} \\
 &= \frac{\Gamma'}{\log x} \\
 &= \int e^{-x} x^{1-t} dx \\
 \Gamma' &= \int e^{-x} x^{1-t} \log x dx
 \end{aligned}$$

All of essential number is constructed with primer of pair, this element have two of pair. Thus is, 1 and n are exceed with prime prisest of nature in world of mathmatics series. This set of number exceed with prime number, and this of one is devide with logment of this pair number. moreover this pair of number has with already of a in this conceel with being decided to resolve with a for this devition of x. After

all, this redevide of x is concerned for being Differential geometry of quantum level quality equale with Gamma function.

Prime number of value have with all of pair in count of $\log x$, and this count of number devide with one element, exceed with this value call $\log x$ to decide with different of value oneselved.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

$2(y \log y)^{\frac{1}{2}}$ have all of primer number and Higgs field + Euler product equal with three particle element, moreover this element also constructed with sixth particle of three of two in one pair. And this concerned of element in Farvat theorem have relativty with Euler equation of reverse in Imaginary of complex equation. Also this equation call one to equal with circle function of themselves. Resolved with all of Gloal differential manifold are jones manifod of equation. Paricle of two pair are constucted with three of one' element, this element are three of pair in two of paricles. Farvat theorem are $n \leq 3 x^n + y^n \leq z^n$, and real of result with conceel is priset of element, and imaginary value of resulted of conceel are particle with complex manifold, and this formation is super symmetry element. Dimension of number relate with Goaris equation of extension, and this extend of over resolved value are complex resulted of elements.

ブレインインターフェースとその応用、機知の仕組み

ブレインインターフェースは、機械において、音声を文字列に表示するには、まず、音声を音階の7階層に分解して、その上に、大脳基底核を経由する共振回路の共鳴を使って、唇の動きのデータを電気信号に変えて、この2種類のデータから、音声を文字列に表示する。頭頂葉から大脳基底核を経由する体感触覚によるデータを使って、そのデータをイメージと言語と文字列に変えて、画像処理を使って、動的に映像やページを処理する。例えば、Wordのページをめくったりする。ASを頭頂葉から大脳基底核を経由するデータを電気信号に変えて、機知の仕組みを担う第32, 33野を経由するドーパミンがそれぞれの役割をする共振回路の共鳴を使って、血液のリボ核酸のヘリコバクターがこの共鳴する共振回路に接続して、ASを動かす。

$$H_n = \sigma(E_n) \times \sigma(K_n), e_n = f^{-1}(x)xf(x)$$

$$e_n = \ker f \oplus \operatorname{im} f, H_n^2 = |(|ds^2|)|, = H_n \times K_n$$

$$E_n = \sigma(K_n) \times \sigma(H_n), \pi(\sigma_1, \sigma_2) = \frac{d}{df} F(x)$$

$$x\Gamma(x) = \pi(\sigma_1, \sigma_2) - (8\pi G(\frac{p}{c^3} + \frac{V}{S}))$$

$$= 2 \int ||\sin 2x||^2 d\tau$$

$$= 4 \int ||\sin x \cos x||^2 d\tau \rightarrow 4V = g_{ij}^2$$

固有エントロピー = 原子の固有エントロピー - 3次元多様体のエントロピー
ブレインインターフェースの大脳基底核の固有エントロピー値

三角関数と虚数、そしてオイラー方程式、 電弱相互理論と電強相互理論から重力と反重力により、 派生する時間のルーツ

Masaaki Yamaguchi

次元の対称性による回転は異次元の系列による要素に変化して、この真空エネルギーはヒッグス場を形成している素粒子から生成されていて、その上に零次元からのエネルギーがこの場から生成されることから宇宙と異次元がダークマターから表現論によりビッグバンとして始まっている。それ故に、この要素による物質場は原子の貯蔵庫が素粒子 1 2 種類に属している。

$$(dx, \partial x) \cdot (\epsilon x, \delta x) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$\frac{d}{df} F = m(x)$$

それ故に、この素粒子は超対称性理論により派生して、物理作用が化学変化を統合してもいる。これらの作用素は幾何学変化から空間により生物と宇宙を形つくるネットワークをも生み出ししている。この対称性による次元の輸送による転位は虚軸と実軸による回転をも空間のニュートンリングとして派生してもいる。そして、この対称性による転位からの次元による回転は時間の一方方向性にも属している。量子力学は時間の別側面による虚数による系列をも有している。この両側面による時間の流れは時間自体が止まり、重力と反重力へと変化してもいる。時間は虚時間をも有している。電弱相互理論は時間の一方方向性でもあり、このトポロジーは時間の流れの結果からの鎖体でもある。

$$\square(\frac{\sigma_1 + \sigma_2}{2}) = [3\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\square\psi = 8\pi GT^{\mu\nu}$$

$$\frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$\nabla\psi^2 = 4\pi G\rho$$

$$\square(\sigma_1 + \sigma_2) = [6\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$= [i\pi(\chi, x), f(x)]$$

$$\square\psi = [12\pi(\chi, x) \circ f(x), \sigma(x)]$$

$$\frac{d}{df} F = \int e^{-f} [-\Delta v + R_{ij} v_{ij} + \nabla_i \nabla_j f + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(\frac{v}{2} - h)]$$

$$= [i\pi(\chi, x), f(x)]$$

これらの方程式は時間の流れから一方向性をも感じて、未来と過去へと流れる時間の機構へと行き着く。それ故に、この時間のシステムの結果電弱相互理論は時間の流れにもなっている。そして、虚時間による反重力が時間の別側面にもなっている。ここでも注目すべきは、電磁気力が2方向の重力と弱い力が一方で、もう一方が反重力と強い力として結合されている故に、電弱相互理論と電強相互理論として分かれている力が統一場理論として量子力学として時間が止まっている仕組みをつくっている。

$$f^{-1}(x)xf(x) = 1, H_m = E_m \times K_m$$

この非可換代数方程式はローレンツアトラクターとして複素多様体になり、世界面を形成していて、この弦理論は宇宙を構成している6種類の素粒子と異次元を形成している6種類の素粒子をも示している。これらの素粒子は超対称性理論をも示している。

$$i = (1, 0) \cdot (1, 0), e^{i\theta} = \cos \theta + i \sin \theta$$

$$e^{-i\theta} = \cos \theta - i \sin \theta$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

$$\sin i\theta = \frac{e^{-\theta} - e^{\theta}}{2}$$

$$\pi(\chi, x) = \cos \theta + i \sin \theta$$

この方程式から2次元が3次元へと逆方向へと分解して、この崩壊から5次元が統合されている。この分解による逆方向へと統合される3次元多様体が同型としての5次元多様体を二重被覆として準同型写像となり、3次元多様体をも包み込む結果を生む機構にもなっている。3次元多様体と2次元射影空間がミンコフスキー空間として、2重被覆となり、アーベル多様体と同型と求まり、リッチ・フロー方程式となる。

$$R(-\alpha) = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$$

$$R(\alpha)MR(-\alpha) = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos^2 \alpha - \sin^2 \alpha & 2 \sin \alpha \cos \alpha \\ 2 \sin \alpha \cos \alpha & -\cos^2 \alpha + \sin^2 \alpha \end{pmatrix}$$

$$= \begin{pmatrix} \cos 2\alpha & \sin 2\alpha \\ \sin 2\alpha & -\cos 2\alpha \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & -1 \\ 1 & -\sin \alpha \end{pmatrix} \begin{pmatrix} \cos \alpha \\ \sin \alpha \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{aligned}
f^{-1}xf(x) &= 1 \\
(\log x)' &= \frac{1}{x}, x^n + y^n = z^n \\
x^n &= -y^n + c, nx^{n-1} = -ny^{n-1}y' \\
y' &= \frac{nx^{n-1}}{ny^{n-1}} \\
&= \frac{x^{n-1}}{y^{n-1}} = -\frac{y}{x} \cdot \left(\frac{x}{y}\right)^n \\
-\frac{\cos x}{(\cos x)'} (\sin x)' &= z_n \\
z^n &= -2e^{x \log x} \\
\lim_{x \rightarrow \infty} f(x) = a, \lim_{y \rightarrow \infty} f(y) = b, \lim_{x, y \rightarrow \infty} \{f(x) + f(y)\} &= a + b \\
\lim_{z \rightarrow 0} f(z) = c, \delta \int z^n &= \frac{d}{dV} x^3 \\
\lim_{x \rightarrow \infty} = c - \lim_{y \rightarrow \infty} f(y) \lim_{x \rightarrow \infty} f(x) + \lim_{y \rightarrow \infty} f(y) &= \lim_{z \rightarrow \infty} f(z) \\
z^n &= \cos n\theta + i \sin n\theta \\
&= -2e^{x \log x}
\end{aligned}$$

これらの方程式は重力と反重力方程式でもある。

$$\begin{aligned}
&\frac{d}{d\sigma} \left[\frac{(\sigma_1 + \sigma_2)}{2} \right] \\
&= \sigma(\downarrow\downarrow) + \sigma(\uparrow\uparrow) + \sigma(\downarrow\uparrow) + \sigma(\uparrow\downarrow) + \sigma(\rightleftharpoons) \\
&\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m = \sigma(\downarrow) \\
&\frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = \sigma(\downarrow) \\
\sigma(\leftarrow) + \sigma(\rightarrow) &= \int e^{-f} [-\Delta v + R_{ij} v_{ij} + \nabla_i \nabla_j v + v \nabla_i \nabla_j + 2 \langle f, h \rangle + (R + \nabla f)(v - \frac{h}{2})] \\
\sigma(\downarrow) &= \sigma(\downarrow\downarrow + \uparrow\uparrow + \rightleftharpoons) \\
\sigma(1) &= \sigma(\downarrow\downarrow + \downarrow\uparrow + \rightleftharpoons)
\end{aligned}$$

$$\text{weak electric theorem} = \sigma(1)$$

$$\text{strong electric theorem} = \sigma(\downarrow)$$

これらの方程式は、トポロジーとしての弦理論のモデルでもあり、電弱相互理論を構成しているマクスウェル方程式と弱い力と重力として、カルーツァ・クライン理論が部分群として強い力を反重力に分配して、この力もマクスウェル理論に結合して、3種類ずつの力へと統合されている。この2種類の結合力が重力と反重力へと結びつけられて、ゼータ関数となっている。

Report of Quantum level of differential structure, zeta function and Global differential equation

Masaaki Yamaguchi, with my son

Higgs field + Euler constant = zeta function

$$m(x) + C = 2 \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$\text{Euler constance} = \frac{d}{d\gamma} \Gamma^{-1} - \gamma^{(\gamma)'} = \left(\Gamma^{(\gamma)'} \right)^{-1} - \gamma^{\gamma'}$$

gravity + anti gravity = zeta function

Gamma function in Global differential equation developed with part integrate manifold cohomologite with helmader operator, and this subrace of differantial form start with first minimalized function, this operator emerged from quato algebra to fermion partial specture in Higgs fields.

$$\Gamma dx_m + \int \Gamma dx_m = \Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$\begin{aligned} & \frac{d}{d\gamma} \Gamma + \int \Gamma dx_m \\ &= \int f(g(x)) g'(x) dx \\ &= \left(\int f(g(x)) dx \right)' \\ & \nabla_i \nabla_j \int \nabla f(x) d\eta \end{aligned}$$

Global differential equation is partial integrate operator in resolved zone of frobenius theorem and this seminal concept with zone, Jones formula equation conqure with rico level theorem, this spectireum releafe in zeta function in CP symmetry broken, this broken from being integrate with blance into universe of zeta function, all of this renze equation into one class of Euler equation concerned with seminal concept in Jones formula equation, zeta function is proofed that this relativity level emerged from Quantum level of differential structure.

$$|f \circ g| \leq |f + g|$$

$$\frac{f}{\log x} = m(x)$$

$$\frac{\gamma}{\log x} = \frac{d}{d\gamma} \Gamma$$

$$\gamma = \Gamma^{(\gamma)'} \log x$$

$$\begin{aligned} \Gamma^{(\gamma)'} &= \gamma (\log x)^{-1} \\ &= r dx_m \end{aligned}$$

$$\begin{aligned} \Gamma^{(\gamma)'} &= \left(\int e^{-x} x^{1-t} dx \right)^{\left(\int e^{-x} x^{1-t} \log x dx \right)'} \\ &= e^f \\ &= r dx_m = e^f \end{aligned}$$

$$\int \Gamma^{(\gamma)'} dx_m$$

$$\nabla_i \nabla_j \int \nabla \Gamma(\gamma) dx$$

$$= \int \Gamma dx_m \cdot \frac{d}{d\gamma} \Gamma dx_m$$

$$= \int \Gamma \cdot \frac{d}{d\gamma} \Gamma dx_m$$

$$= \left(\int \Gamma dx_m \circ \frac{d}{d\gamma} \Gamma \leq \int \Gamma dx_m + \Gamma dx_m \right)' = e^{-f} \cdot e^f \leq e^{-f} + e^f$$

$$(1 \leq e^{-f} + e^f)'$$

$$0 = -(e^{-f} - e^f)$$

$$\Gamma^{(\gamma)'} + \Gamma^{(\gamma)}$$

$$(= e^f + e^{-f})'$$

$$= -e^{-f} + e^f$$

$$= -(e^{-f} - e^f)$$

$$(R_{ij})' = -(R_{ij})$$

$$\int C dx_m = \int \left(\int \frac{1}{x^s} dx - \log x \right) dx$$

$$= \int \int \frac{1}{x^s} \mathrm{dvol} - \int \log x \mathrm{dvol}$$

$$\Gamma dx_m = \gamma dx_m$$

$$= \frac{d}{d\gamma} \Gamma^1 - (\gamma)^\gamma'$$

$$= e^{-f} - e^f$$

Quantum level of differential structure equal with Gamma function and related with Beta function, escorted into Euler constant and symmetry build with Higgs field, and proof with zeta function summarized with Pólya and Riemann conjecture. Mathematicians of Russia, America and Japan resolved with this conjecture. My son resolved with this conjecture build in concepted with quantum level of differential structure into zeta function. All of equation is integrated with manifold into Hilbert space, this manifold emerged from general relativity and quantum physics in manifold of Hilbert space, this manifold resolved with D-brane and Gauss surface into gravity and atom of element with this theory, this theory is integrated theorem released with one class of universe equation.

AdS₅ 多様体と別次元の AdS₅ 素数と素粒子方程式

Masaaki Yamaguchi

$$e^{-2\pi||psi||}[\eta_{\mu\nu} + \hbar x]dx^\mu dx^\nu + T^2 d^2\psi \\ = e^{-f} + e^f = (u + d) + c, (w + s) + b = -e^{-f} + e^f$$

この式は、時空にどれだけの原子が凝縮しているかを表していて、その上に、この逆数は、密度エネルギーをも示している。空間の体積に原子を商代数にすると、濃度にもなる。逆数は、a=原子のエネルギー が全空間を 1 とすると $\frac{1}{x}$ すると、 $\frac{x}{y}$ は密度になる。 x=原子,y=全空間、個数は $\frac{y}{x}$

$$\square = \text{原子} \rightarrow \frac{[\eta_{\mu\nu} + \hbar(x)]dx^\mu dx^\nu}{e^{2\pi T||\psi||}}$$

$$||ds^2|| = e^{2\pi T||\psi||} ([\eta_{\mu\nu} + \hbar(x)]d^\mu dx^\nu)^{-1} + (T^2 d^2\psi)^{-1}$$

$$||ds^2|| = e^{-2\pi T||\psi||} [\eta_{\mu\nu} + \hbar(x)]d^\mu dx^\nu + T^2 d^2\psi$$

$$(u + d) + c = e^{-f} + e^f, (w + s) + b = e^{-f} + e^f$$

$$R'_{ij} = -R_{ij}$$

この AdS₅ 多様体の $e^{-2\pi T||\psi||}$ は原子間距離を表していて、 $[\eta + \bar{h}(x)]d^\mu dx^\nu$ で宇宙の D-brane の構造を表している。この数式が $T^2 d^2\psi$ とアーベル多様体が包み込んでいる。原子が回転するのと、ニュートンリングが回転する宇宙は同じ速さで回転する。

$$||ds^2|| = \square + \rho, ||ds^2|| = \nabla\Psi^2 + \psi$$

$$\eta_{\mu\nu} + \bar{h}(x)$$

$$\text{proximity} = \Delta x \Delta p - \Delta p \Delta x + \delta(p, x)$$

$$||ds^2|| = e^{-2\pi T||\psi||} + [\eta_{\mu\nu} + \bar{h}(x)]dx^{\mu\nu} + T^2 d^2\psi$$

$$\frac{d}{df}F = e^f + e^{-f}, \frac{d}{df} \int C dx_m = e^f - e^{-f}$$

$$R'_{ij} = -R_{ij}$$

宇宙と異次元では0だが宇宙だけだと誤差が生じる。これが不確定性原理であり、異次元が誤差になり、宇宙と加群すると0になる。 $\delta(p, x)$ が隠れた変数となり、異次元で誤差をこの不確定性原理と表している。全ては素粒子方程式から生成される式達である。

オイラーの定数とガンマ関数、 そして微分幾何の量子化と虚数による逆三角関数、 ベータ関数による場の理論

Masaaki Yamaguchi

オイラーの定数は、次の式で成り立っている。

$$C = \int \frac{1}{x^s} dx - \log x$$

その式を単体積分をすると、

$$= \int \left(\int \frac{1}{x^s} dx - \log x \right) d\text{vol}$$

ゼータ関数を多重積分すると、

$$= \int \int \frac{1}{x^s} dx - \int \log x d\text{vol}$$

ここで、対数方程式は、多様体積分がガンマ関数を微分した方程式と同値より、

$$\begin{aligned} F = \Gamma &= \int e^{-x} x^{1-t} dx \\ &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dx_m - \int e^{-x} x^{1-t} \log x dx_m \\ &\quad \int x^{1-t} e^{-x} dV = \int x^{1-t} dm \\ &\quad \int x^{1-t} e^{-x} dV = \int x^{1-t} d\text{vol} \\ f = \gamma = \Gamma' &= \int e^{-x} x^{1-t} \log x dx \\ &= \frac{d}{d\gamma} \Gamma^{-1} - \int e^{-x} x^{1-t} \log x dx_m \\ &= \frac{d}{d\gamma} \Gamma^{-1} - (\gamma)^{\gamma'} \end{aligned}$$

これらより、大域的微分方程式のヒッグス場方程式の逆三角関数の双曲多様体に対極する、双曲多様体の余弦定理と同値になる。

$$\begin{aligned} &= e^{-f} - e^f \\ &= 2 \cos(ix \log x) \end{aligned}$$

結局は、微分幾何の量子化がガンマ関数でもあり、その逆関数はゼータ関数でもあり、そして、オイラーの定数を多様体微分をすると、ヒッグス場の方程式は正弦定理の逆三角関数であり、このヒッグス場の方程式の逆三角関数が、余弦定理の双曲多様体として、オイラーの定数になる。オイラーの定数は、このガンマ関数から、無理数と証明される。

六星占術の運命星と宿命星の求め方

細木数子、山口雅旭、苔米地英人

$$wxyz \cong wx \oplus yz, wx \oplus yz = w'x' \otimes y'z'$$

$$x|_{1979} \equiv 19 \oplus 16, 19 \oplus 16 = 35$$

$$19 \oplus 16 = 36 - 1, 36 + 2 = 38, 1979 \mod n = 2$$

$$38 \otimes 2 = 76, x|_{1940, 1979} = 24, x| = \frac{76 - 12}{2}, x|_{1979} = 32 \oplus 6$$

易学の64卦を38周期で求めると、同型写像から32となり、この真逆が六星占術の暦となる。日にちの干支と十干が月の干支と十干となり、真逆が六星占術となる。1940と1979の38周期から干支の素因数分解の剰余項と1979の準同形写像からも運命星がもとまる。この運命星は月運の性質でもある。 $x = 2n - 2$ は、5行の月運の十干を表している。1979年周期で、西暦の加群分解がある。この1979年前は、-1979年で0年と おけて、旧約聖書の12月が、0年の2月が36より、1月が5から5-31で-26から60-26より12月は34となり、 $34+31=65$ によって、0年の1月の生命エネルギーは5となり、キリストが生誕した、B.C.4年から、1年ごとに5と干支と十干が動くこのから、キリストを1とすると、5年前は0年となり、-1979年と同型から-1979年の0年は2月が38となる。1979年周期でこの2倍の2958年の加群分解は、 $29+58=97-60=37$ となり、この2958年からの2倍の1979年の3倍は、 $5937=59+37=96-60=39$ となり、神様と人類の即身成仏の数秘術となる。加群分解が成り立つのは、この1979を起点とする倍数の±2前後の西暦の数値となっている。六星占術は、生命が誕生したら-1加群となる。これが運命数から運命星となる。運命から宿命そして立命となる。五行で運気が悪いのが六星占術ではよい運気になっている。六星占術で運気が悪いのが五行では運気がよい。易学を細分化すると六星占術となる。相性殺界となる運気が六星占術と五行の相殺となる。宇宙が始まって1940と1979の38周期が易学の同型写像となり、六星占術となる。六星占術の0から9までは、土星の陽における種子から陰における安定10から19までは、金星の陽の種子から陰の安定20から29までは、火星の陽の種子から陰の安定30から39までは、天王星の陽の種子から陰の安定40から49までは、木星の陽の種子から陰の安定50から59までは、水星の陽の種子から陰の安定

5行では、金星から木星、火星、土星の申から未までを通っている。六星占術の大殺界は、本来真逆の財になっている。九紫、大黒、緑水星、大善星、妙雅、火竹、白水、静雲、白鷗、光美と宿命星が廻っている。各宿命星の各星人の周りは別にこの通りではない。

$$x|_{1979} = 35, x = 36 + \sum_{k=1}^{30} x_1 + \sum_{k=1}^{31} x_2 + \sum_{k=1}^{28} x_3 + x_4$$

$$x = 36 + \frac{n(n-1)}{2} x_9 + x_4$$

$$x_5 = \frac{x}{60}, x_6 = \frac{x_5}{12}, x_7 = \frac{x_6}{10}$$

$$x_8 = \frac{2x + x' - 1}{2}$$

西暦 1979 年を西暦 0 年と生命エネルギー 36 を閾値としての、各生命エネルギーの上下の平行線となっていて、各星人の生命エネルギーは、この線の上と下の行き来となる周期で表されている。

六星占術の始まりの年の月における日の求め方は、

$$-1979 \cong 1979, 1979 = 19 \oplus 79 \cong 19 \oplus 16 = 35 \cong 38$$

1979 年の日の五行の月は、

$$x = 2n - 2$$

と表せられて、2 月から宿命星の年が始まることにより、求める星人の運命星は、運命数-1 加群より、求める生命エネルギーを x とすると、

$$x = 35(2 \text{ 月の生命エネルギーから}) + \sum(1, 3, 5, 7, 8, 10, 12) \sum(4, 6, 9, 11) + (\text{うるう年の 4 年に一回} + 1)$$

ここで、うるう年が 4 年に一回なので、 x が閏年があるとしたと、2 月 29 日として、4 年に 1 回、2 月に +1 する。キリスト誕生の年が B.C.4 年で、キリストが旧約聖書のときの西暦 0 年の前の 12 月 25 日が -1979 年とすると、西暦 0 年の 2 月も生命エネルギーが 35 前後になる。神様と人類の即身成仏の数秘術が六星占術となる。

天運、本人の月運、地運は、六星占術は、地運の誕生のときが生命エネルギーとしての本人の月運となり、この運命星の十二支が先祖の十二支によって、天運となり、この基質によって人生の天運のときの相性、月運、地運が動き、六星占術の各星人の適職、才能、特技、性格によって、運命星の運勢として運命が流れる。大殺界のときは、得手不得手として、流れるが、5 行を見ると大丈夫になっている。各星人の得手、不得手によって、方位が凶か吉として巡る。九星は六星占術の相性殺界を使って、凶を吉にしている大人の占いでもある。宿命星は、運命数の単体量をだして、この数値に +10 として十の宿命星の流れを占っている。運命が固定されている。要するに、日の各運命数が十干の月運 $2n-2$ と同じく、 ± 2 の範囲で動き、十干と十二支によって決まっているのとは、違う。その上に、常占術としての月運が 1 年に 12 あるのとは違い、運命星の日の運命数の星人と十二支を月に移動させて、先祖の天運の十二支をこの日の星人と干支を月とみなしての、各星人が同じ月なのに違う占いとして成り立っている。

Instance of acknowledged from refrent existed of humanity theorem and reality of thought in modern technology 実存主義と現代文明、双極性による精神機能からの未知の智

Masaaki Yamaguchi

Refrent existed of humanity theorem is built with acasic record from human being constructed with world is setting from view of reality human being created from. This theorem mechanical from this type afford into world viewed of this human being, and this thought of history record having with the type recieved with acknowlege of acasic record and reality of human being seen world to accept with native reality, this type of human being become with enginner and human brain interface created with modern history existed. These type of being recieved from the world of acknowlege is existed in reality of non dreamly world of being seen. Whisperd of this type called with world of being existed, estling of this type called and this type of being selected with is recieved with acasic record. Photo Reader is reffer to being selected with estling of acknowlege in world of tounge. Refrent existed of humanity theorem is inspired with Philosophy of being called human being to recieve in this record. Refrent means that jugement in person and creature are accepted with acasic record from recognized of system how registance of recieve with nature and universe of life.

実存主義は、サルトル・アッカーシュマンが去った後も、現実主義者がこの実存主義に分類されている上に、ブレインインターフェースやコンピュータを作っている人がこの宗派に分類されている。この実存主義は、無の状態から零次元がエターナルな場に同型とされている瞑想を対象として、アカシックレコードから未知の情報を取得する。この無は禅とも言われていて、実存と虚存としての物理学を対象とすると、閉3次元多様体として、ザイフェルト多様体から未知の智を感知する。この智は、他者との関係から関与して、この世は天啓のごとく、天からの啓示を本として接点として、未知の智を感知する。現実主義者は、うつの人に傾向として言われているが、実存主義にこのうつの人を対象として、夢を誇大妄想とは別として、研究対象にされている。なぜ、うつの人が現実主義者に多いのに実存主義の能力があるかは、瞑想から無の状態を使い、アカシックレコードにアクセスするのと、本当に現代文明を作り上げることができる能力をもっているからである。無からの知識はエターナルな場から零次元が世界に存在していて、感知していない次元が零次元と無としての禅による相対性としての他者との関与から同型として、禅の無からアカシックレコードとこの存在している次元から実存と虚存から未知の智をこの世界に降臨させてもいる。それゆえに、このタイプの人、ウィスパードと言われている、臥龍をこのエターナルな場を生成元として、エストリングをこのウィスパードを育成する場と言われている。カロリング研究所もこの分類に入る。このエストリングな場は、若返りの場にもなっている。アカシックレコードは絶大な治癒力を有している。宇宙は、始まりと終わりが真逆な時間の矢にもなっている。過去の宇宙が未来の宇宙になってもいて、時間と空間がシンメトリックな性質になってもいる。年を取る事が勉強との学問の教訓から身につけられていると、若返りの効果が生成される。このタイプには、テクノ

ロジーとマネジメント、ストラテジストが分類として研究されている。エストリングな場から、今まで誰も思いつかなかった数式や理論をアカシックレコードから投射として得ることが出来る人もいる。実存主義とは、判断と認識、そして無による愛でもある。


Infinity of acknowledge in world is existed zero dimension and rebuiled with eternal space, non-existed dimension equaled with zero dimension and zen of skill relativity, and zen of infinity of atomasphere required with acasic record from this space of dimension. Therefore, this type of being required with are called with whisperd, and this selected with estling of space and acasic record. This dimension of area are referred with more and more young belonged with time of ages. In this referred with time of ages are cured from any built of system. Get of ages equaled with young of age with study of learn and required of any skill. Life of water and Life of tree is existed with central of universe, and around of universe is old of universe and this space is creating with any other dimension of creature. Time followed is reverse of system. Young of universe is equaled with Old of universe and Young of time refferd with is Old of time belonged with universe system. This system cnstruct from non time followed lead into quantum physics mechanism, this type of category esperanade of theorem and discover of system of native more literal strategy of skill in top of goverment positions, more releanity of skill is called from manegement of skills. Any type of this required skills lead from every acknowleged of technical and potential possibility, more spectrum influence of being inspired into a respired of reluctant with acquiring lives of flowing days on stress pressiers. Life of insurance need every moment to avoid from varnished any others. Incoming of value space revealed from any intransit of memories, this acceptance of being invited from accerity of geotics.

無の状態を維持すると空間に同化する。それと同時に現実と隔離できる。この作用で、一念三千を感じることに無による瞑想はアカシックレコードに接することの必要十分条件にもなっている。無とはそれほどにも大切な技能に言えることでもある。速読法がこの技術を使っている。無の音読が確の目によりこの技術を使っている。

Refrent exist of humanity theorem is included with estling of involved with other's inspire with each other of persons and creatures in estling of setting with mind of hologram fields from being with academic learns. This estling of fields is also catastrophe of fields how converted from being with harmony and nestling of mind inspect with each persons and each creatures in other's of minds. Encycropedia is selected with disease of type in explained of this refrent exist of humanity theorem. Therefore, this theorem is explained with encycropedia of which other persons and other creatures are read with other creatures in estling of fields. This psychological of mind in refrent exist of humanity theorem is explained with indivisual of mind in zero dimension and zen of mind recieved from future and past of accident what any matter are incident of other's distant with involved of counsels.

実存主義は、他者と自己においての精神作用の関与による、精神の共鳴と共振の機構を述べてもいる。自己のどういう思考の仕方、他者との心理作用がカタストロフエネルギーの位置関係に在るかを、他者がどういう態度で他者同士の心理がエストリングの場でどのように変化するかをこの実存主義は、弁証論として述べてもいる。心因反応は、この実存主義で分析されてもいる。自己の心理を他者が読むのは、この実存主義の関与からも分かる。この実存主義には、自己による無の状態から未来と過去に何が起きるかも他者との相対関係から分かることの心理をも述べてもいる。プリンストン高等研究所では、この無からの超高速コミュニケーションが研究者同士でされてもいる。

孝君のお嫁さんが、まさか持田さんと同じで、母さんの若かりし頃と同じとは、知らなかった上に、美鈴ヶ丘に行かれたのは、本当に申し訳ございませんでした。投射を改良していますが、情報空間の存在は、苫米地博士で知りました。その投射とは、違うが、経路は同じ感覚で、アカシックレコードに接続しますが、ロールシャッハテストと点字を探る要領で接触します。痴漢ではありません。ドアは使います。



ユウの直立と私の座禅の上のユウ

ユウの誇大妄想は、私はここまではさすがに思ってもいなかったが、同レベルとしか言いようがない。ゼータ関数をアカシックレコードから投射で得たときに、前のユウちゃんが目撃していたのが、事の発端だった。それが、このあっかんべーらしい。

〒721-0942 広島県福山市引野町 458

山口道弘 ・ 山口雅旭