

Entropy on manifold and differential of equation

Masaaki Yamaguchi

1 Global Differential Equation

Global area in integrate and differential of equation emerge with quantum of image function built on. Thurston conjugate of theorem come with these equation, and heat equation of entropy developed it' equation with. These equation of entropy is common function on zeta function, eight of differential structure is with fourth power, then gravity, strong boson, weak boson, and Maxwell theorem is common of entropy equation. Euler-Lagrange equation is same entropy with quantum of material equation. This spectrum focus is, eight of differential structure bind of two perceptive power. These more say, zeta function resolved is gravity and antigravity on non-catastrophe on D-brane emerge with same power of level in fourth of power. This quantum group of equation inspect with universe of space on imaginary and reality of Laplace equation.

It is possibility of function for these equation to emerge with entropy of fluctuation in equation on prime parameter. These equation is built with non-commutative formula. And these equation is emerged with fundamental formula. These equation group resolved by Schrödinger, Heisenberg equation and D-brane. More spectrum focus is, resolved equation by zeta function bind with gravity, antigravity and Maxwell theorem. More resolved is strong boson and weak boson to unite with. These two of power integrate with three dimension on zeta function.

Imaginary equation with Fouier parameter become of non-entropy, which is common with antigravity of power. Integrate and differential equation on imaginary equation develop with rotation of quantum equation, curve of parameter become of minus. Non-general relativity of integral developed by monotonicity, and this equation is same energy on zeta function.

$$\pi(\chi, x) = i\pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x)$$

This equation is non-commutative equation on developed function. This function is

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx - \int \frac{1}{(x \log x)} dx$$

This equation resolved is

$$\int \int \frac{1}{(x \log x)^2} dx_m \geq \frac{1}{2}i$$

And this equation resolved by symmetry of formula

$$y = x$$

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \geq \frac{1}{2}$$

This function common with zeta equation. This function is proof with Thurston conjugate theorem by Euler-Lagrange equation.

$$F_t^m \geq \int_M (R + \nabla_i \nabla_j f) e^{-f} dV$$

resolved is

$$F_t \geq \frac{2}{n} f^2$$

$$\frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2 dm}{-(R + \Delta f)}$$

These resolved is

$$\frac{d}{df} F_t = \frac{1}{4} g_{ij}^2$$

Euler-Lagrange equation is

$$f(r) = \frac{1}{2} m \frac{\sqrt{1+f'(r)}}{f(r)} - m g f(r)$$

$$m=1, g=1$$

$$f(r) = \frac{1}{4} |r|^2$$

Next resolved function is Laplace equation in imaginary and reality fo antigravity, gravity equation on zeta function.

$$\int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = 0$$

This function developed with universe of space.

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

This equation also resolved of zeta function.


$$\frac{d}{df} F(v_{ij}, h) = \int e^{-f} [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2)(\frac{v}{2} - h)]$$

These equation is common with power of strong and weak boson, Maxwell theorem, and gravity, antigravity of function. This resolved is

$$\frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

on zeta function resolved with

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$



2 Quantum Equation for Dimension of Symmetry

Quantum Equation architect with geometry structure of Global Differential Equation has zero with gravity and antigravity for eternal space, and this space emerge for burned of non expanded universe. Then this universe has Symmetry in dimension with that created of fourth Universe in one of geometry has six element quark and pair of structure belong for twelve element quark. These quarks emerge with eternal space of Non-Difinition System in Quantum Mechanism, in term of one dimension decided, or the other dimension non-decision. These system concerned of vector of norm depend for universe mention to eternal space. Mass existing in dimension emerge gravity, these paradox is in universe has mass around of light, in deposit of mass for our universe and the other dimension has gravity and antigravity, and covered with these element for non-gravity. Laplace equation decide with eight of structure for these element of power integrate for one of geometry. Higgs quark is quote algebra equation in Global Differential Equation of non-gravity element on zero dimension. These equation is mass of build on structure in mechanism system. This quote algebra equation have created of structure in mass with universe of existing things. These result means with why quantum system communicate with our universe be able to connect of.

$$\begin{aligned} \pi(\chi, x) &= [i\pi(\chi, x), f(x)] \\ \int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &\geq 0 \\ \Delta x \Delta p &\geq \frac{1}{4} i \end{aligned}$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\sqrt{\frac{\hbar G}{c^3}} \cong 1.616 \times 10^{-33}$$

$$C = 0.5772156 \dots$$

$$F = \frac{d}{df} \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$F_t^m = \frac{1}{4}(g_{ij})^2$$

Quantum Group resolved with Laplace equation build in calculate on non-gravity element of based by non extension equation. This solved by these element in equation has belong to gravity is why universe have in no weight, and mass exist weight to space emerge of created with gravity, in which is reason by non-extension equation translate to variable and invariable element. These problem of included universe in no gravity, which is exist of mass on space of surrounding in universe. Those question resolve on zeta equation and Quantum Group of translate to vector equation. Non-extension equation selves has non-gravity, these result with universe first burn in D-brane created by solved with replace equation. Quote Algebra equation have created of structure in mass with universe of existing things.

These equation explain to those which included mass has gravity emerged, and universe has in surrounding to no weight. The other Dimension integrate with these universe of gravity to unite antigravity, so this universe has no weight. Higgs quark is mass of built on structure in mechanism system. These system belong to create on element. These element is based on existing of universe which has structure code, in theorem composed for universe to emerge of time with future and past. Universe first created in these time, this existing things already burned with space. Network Theorem is connected eight element of geometry structure which integrate with three dimension of structure. These structure compose in three manifold, zeta equation is this system of element. These resulted theorem resolved with quantum equation, so this mechanism impressed in universe of component. Strong and Weak boson is united to one, and Maxwell theorem is same system. Gravity and Antigravity has own element. General relativity theorem same united. These integrate with included Euler equation. These power of element is zeta equation. Then this twelve element of quarks has belong to this universe and the other dimension.

$$\Delta E = -2(T-t)|R_{ij} + \nabla_i \nabla_j f - \frac{1}{2(T-t)}g_{ij}|^2$$

Quote group classify equivalent class to own element of group.

$$A=BQ+R$$

$$[x]=A$$

$$dx^n=\sum_{k=0}^{\infty}x^ndx$$

$$R_n=\frac{n!}{(n-r)!}(x^n)'$$

$$\beta(p,q)=\int_0^1x^{p-1}(1-x)^{q-1}$$

$$Z(T,X)=\exp(\sum_{m=1}^{\infty}\frac{(q^kT)^m}{m})$$

$$Z(T,X)=\frac{P_1(T)P_3(T)\ldots P_{2n-1}}{P_0(T)P_2(T)P_4(T)\ldots P_{2n}}$$

$$|v|=|\int (\pi r^2+\vec{r})dx|^2$$

$$\Delta E=\int (\operatorname{div}(\operatorname{rot} E)\cdot e^{-ix\log x})dx$$

$$(\nabla\phi)^2=\int tf(t)\frac{df(x)}{e^{-x}tx-1}dx$$

$$(\nabla\phi)^2=\int tf(t)(\Gamma(t)df(x))dx$$

$$(\nabla\phi)^2=\frac{1}{\Gamma(x+y)}$$

then these equation decide to class manifold with group. Differential group emerge with same element of equation, zero dimension conclude to emerge with all element. Constant has with imaginary of number on developed of zeta equation. Weil's Theorem resolved with zeta equation, these function merge to build in replace equation. Euler constant has with bind of imaginary number and variable number. These function is

$$C=\int \frac{1}{x^s}dx-\log x$$

Replace equation resolve on zeta function.

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i$$

These function understood is become of imaginary number, which deal with delete line of equation on space of curve.

Euler number also has with imaginary of constant.

Space Mechanism to transport of Dimension

Masaaki Yamaguchi

3 Kaluza-Klein Theorem

The other dimension rotate of universe with real and image rout, complex rotate to dimension of fifth for Kaluza-Klein dimension extend with theorem. Mobius space explain this theorem to reflect for fifth dimension of construct with real rout of rotate in image rout.

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\beta(p, q) \geq - \int \frac{1}{t^2} dt$$

Abel equation tell this space to infinite this dimension for finite space to conclude of this theorem, in explain the other dimension with our universe rotate with, and this space rout out this fifth dimension, no thought with time system. This space didn't throw of light speed, light element go with space throw. This idea explain of magnetic theorem, this tell space to construct with movement result. Light speed go with other light together, this light look like stop of speed. This idea extend of tell space for no relativity, time flow of infinite.

Kaluza-Klein Theorem say

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(\kappa^2 A_{\mu\nu}(x)dx^\mu)^2$$

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$\frac{\delta g}{L^2} \sim \frac{G}{c^4} \frac{\delta E}{L^3}$$

$$\delta E \gtrsim \frac{\hbar}{T} \cong \frac{\hbar c}{L}$$

$$\delta g \gtrsim \frac{L_p^2}{L^2}$$

$$\nabla \phi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$\frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$ds^2 = -N(r)^2 dt^2 + \phi^2(r)(dr^2 + r^2 d\theta^2)$$

$$ds^2 = -dt^2 + r^{-8\pi Gm}(dr^2 + r^2 d\theta^2)$$

This equation result

$$dx^2 = g_{\mu\nu}(x)(g_{\mu\nu}(x)dx^2 - dxg_{\mu\nu}(x))$$

These equation tell space non-symmetry result

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

This theorem mention to zeta equation construct space of non-symmetry to emerge with non-gravity.

$$\pi(X, x) = i\pi(X, x)f(x) - f(x)\pi(X, x)$$

4 Network Theorem

Quantum Group connect with Universe component which has built in emerge space, these Network element by kernel and image function in zeta equation. This each of function system is Universe element of geometry group construct with these equation, in space first burn with time future and past. Gravity equation has with antigravity element. Quantum group with fourth of universe system. This system mention to our universe create with network theorem. Our universe already create of eternal space. Time throw with this space, then universe already complete created. In this reason time of throw in space was found with non-expand of universe. General relativity theorem mention to this equation means by.

Quark has twelve element kernel and image of equaiton resolve with base group integrate of three dimension construct of this equation resulted. Global differential equation means to resolve with universe emerge in built system. This equation emerge with almost thing of component structure code. This source code create in element structure. This network system is include with fourth of universe on connected with communicate mechanism. Base group explain in these system, differential equation has with global integrate equation togather. This equation means for our universe system of geometry structure integrate with one of geometry. In reason this connected with eight element of structure, then this system is realize to quantum group equation resolved with space mechanism.

$$G_{\mu\nu} + \Lambda R_{\mu\nu} = \kappa^2 T^{\mu\nu}$$

$$\begin{aligned} X(3) = & (-1)^3 < |a_0 a_1 a_2 a_3| > + (-1)^2 (< |a_0 a_1 a_2|, |a_1 a_2 a_3|, |a_0 a_1 a_3| >) \\ & + (-1)^1 < a_1 a_2, a_0 a_3, a_2 a_3 > + (-1)^0 < a_0, a_1, a_2, a_3 > \end{aligned}$$

$$H_n(x) = \ker f / \operatorname{im} f$$

$$X(x) = r(H_n(x))$$

$$X(3) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

5 Zeta equation of system

Quantum equation for universe has with gravity on fourth of manifold is closed three manifold with antigravity has decieve to the other dimension. This thorem mention to tell universe to composite with time of system, these system of equation say,

$$\log x(\log x) \geq 2(y \log y)^{\frac{1}{2}}$$

This system belong for geometry theorem destruct with three manifold built of singularity in dimension, this dimension is also built in mechanism of all composite with source code. For code is entropy of mass with energy, this entropy is,

$$\pi(X, X) = [i\pi(X, x), f(x)]$$

$$\pi(X, x) = \int \frac{1}{(x \log x)^2} dx$$

This equation tell gravity to entropy of equation composite with mass of singularity. Three manifold of equation mension to this universe has with zero dimension began to time of first, in firstly time has future and past. This system says that universe has first begun of throw to time system. The explain theorem that universe of end composite for entropy equation tell the construct of this system, and universe has in other dimension covered with oposite of energy.

Dimension of three manifold system says,

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \kappa^2 T_{\mu\nu}$$

$$L_p = \sqrt{\frac{\hbar G}{c^3}}$$

$$\nabla\phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S})$$

This equation mension to three manifold of Dimension construct with time and energy of Non-Difinition system. Quantum equation and this equation also built with same equation of theorem. Fundamental group explain this theorem to composite with entropy of same energy, any rout built with same energy to construct with source code. The rout of equation is,

$$V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau(q)}} L(x) dx) + O(N^{-1})$$

The equation also say that universe has singularity of component is,

$$\frac{1}{\tau}(\frac{N}{2} + \tau(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

These theorem explain that universe has of big-clanch system. The system include to dimension and time of throw of mechanism. Higgs field of energy equation say,

$$\frac{d}{df}F = m(x)$$

This equation concern with Seifert manifold mension to dimension of system. Global differential equation belong for gravity and antigravity of equation with Abel manifold in Euler equation. Infinite of group composite with this theorem conclude of finite group, these theorem explain to construct of universe is big-clanch.

Universe has with ertel and this mass of energy is singularity of component. Global differential system and Higgs of mass says that universe has built of darkmatter and big-ban system.

6 Emerge any dimension for supersymmetry to create for universe form quarks of element

Space for Non-Symmetry create of power in gravity and antigravity for Higgs quark of energy with ertel potential. This potential energy construct to emerge fourth power, then integrate of twelve of quarks. These mechanism export for another dimension of symmetry built. These dimension also architect with sixth of quarks. There import mechanism to Global differential equation for resolve of Higgs quark to build for unvierse. This way export of two dimension for symmetry of pair, these symmetry of space also built from sixth of quarks. Space of ertel emerge in gravity and antigravity, this power from ertel in non-free condition from free of ertel to emerge on these mass of space in zero dimension. This mechanism result with space to construct in three manifold of ertel from power. This resolved mechanism built with time and space of system, also this system flow to universe. These system tell universe to emerge space of singularity from potential energy.

$$\pi(|K''|) \cong \pi(|\bar{K}''|, O)$$

$$l(< P, Q >), l(< Q, R >), l(< R, S >), l(< P, R >), l(< P, S >), l(< Q, S >)$$

$$= \alpha, \beta, \gamma, \delta, \zeta, \eta$$

$$e = \alpha\beta\gamma$$

$$\gamma = \beta^{-1}\alpha^{-1}$$

$$\zeta = \alpha\gamma$$

$$\delta = \eta\gamma^{-1}$$

$$\begin{aligned}\eta &= \beta \zeta \\ \beta^5 &= (\beta \theta)^2 = \theta^3 \\ [\beta] &= [\theta] = 0\end{aligned}$$

$$ds^2=e^{-2kT(x)|\phi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2(x)d\phi^2$$

$$\nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S})$$

Twelve quarks emerge for three manifold of space, in element space of system construct with three D-brane, these dimension catastrophe of symmetry. This mechanism explain for those which create of our universe and the other dimension, take these mechanism into consideration for those built to deceive the other dimension over universe, for Global differential equation devide with gravity and antigravity emerge space to existing zero dimension of ertel space in singularity of zeta equation.

$$\beta(p,q)=\frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\log x(\log x) = 2(y\log y)^{\frac{1}{2}}$$

7 Higgs Fields transport of three manifold on zeta function to resolve the mechanism

Higgs Fields transform of comformal field in space, this space mechanizm with lives of existing source code. Non-commutative algebra built to transport of emerging on space, each of element has prime number in these theorem. Destruct of element has a reverse on manifold, this theorem conclude with Euler-Lagrange equation resolve to merge element.

$$|f(x)| \rightarrow [f,f^{-1}] \times [g,h]$$

$$(x-1)(2y-b)\geq z$$

$$\frac{1}{x-1}\frac{1}{2y-b}\leq \frac{1}{z}$$

$$f(x)=\frac{1}{x-1}+\frac{1}{2y-b}$$

$$\begin{aligned}\frac{d}{df} &\geq \frac{d}{df}f \\ dx &\leq dF\end{aligned}$$

$$|f(x)| = \frac{1}{4}|r|^2$$

Seifert manifold has reverse of time with space mechanism on merge to resolve with zeta function. Global differential equation has zero dimension of darkmatter to create of space, big-ban system also start with this mass of field to begin with universe.

Nonliner of element on manifold algebra

Masaaki Yamaguchi

8 Mebius space

Gamma and Beta function belong for Kaluza-Klein theorem, these equation built of first universe began with darkmatter, this ertel is non-free condition to emerge with big-ban system conform to export with antigravity element. This space consist of fifth dimension rotate with Mebius space construct for the other dimension, after all built of quarks two of pair in dimension symmetry. Topology consist of algebra equation is being with complex function. Norm relate with Volume of space of these system built in. Prime number concern with Euler constance relate of.

$$\Gamma(x) = \int x^{1-t} e^{-x} dx$$

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$f(x) = [f, f^{-1}] \times [g, h]$$

$$V(x) = \int \frac{1}{\sqrt{2\tau q}} (\exp \int L(x) dx) + O(N^{-1})$$

$$Zeta(x, h) = \exp \frac{(qf(x))^m}{m}$$

$$\frac{d}{df} F(x) = m(x)$$

This point of focus is Global differential equation relate with all existing equation. Prime number consist of zeta function in Mebius space.

9 Maxwell theorem integrate of gravity element

Kaluza-Klein theorem consist of general relativity of equation, fifth dimension is part of zeta function on also Global differential equation. Weak power concern with electric of magnity in Maxwell theorem. Space fill of ertel with darkmatter being of graviton to emerge of Higgs fields. This fields has topology element with any transform of power in atomic element. Relate with result of space merge to create for electric and magnitic power. Higgs fields built with pair of quarks in symmetry space. Vector of norm consist of two of pair for dimension.

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + \phi^2(x) (\kappa^2 A_{\mu\nu}(x) dx^\nu)^2$$

$$dx = (g_{\mu\nu}(x)^2 dx^2 - g_{\mu\nu}(x) dx g_{\mu\nu}(x))^{\frac{1}{2}}$$

$$\sum_{k=0}^{\infty} |x_k + y_k|^2 = \sum |x|^2 + 2 \sum |xy| + \sum |y|^2 \leq \sum |x|^2 + \sum |y|^2$$

$$f(|x + y|) = f(|x|) + f(|y|) \geq f(|x| \circ |y|)$$

Norm space has with Frobenius theorem to resolve with Kaluza-Klein space built in. Euler number consist of rank for homology element to create with zero dimension.

$$\chi(x) = H_3(x) = 0$$

$$H_3(\Pi) = Z$$

Loop of topology has no exist with zero dimension, this system explain to create of Maxwell theorem.

10 Zeta function belong for singularity component

Zeta function consist of reverse of time quality, this space result with movement of element. Kaluze-Klein theorem create with space to emerge of singularity. Maxwell theorem has this system of circumstance. Fourth dimension belong to construct with Donaldson manifold exist explain. Duality of space also built with zeta function from singularity. Monotonicity relate to merge with non-relativity of integral result, this reflection is space of quality.

$$\frac{V(x)}{f(x)} = m(x)$$

$$F(x) = 0$$

$$\frac{d}{df} F(x) \geq 0$$

$$\lim_{x \rightarrow \infty} \text{mesh} \frac{F(x)}{f(x)} \rightarrow 0$$

$$\nabla f(x) = 2$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$f(x) = \int \frac{1}{x^s} dx - \log x$$

Morse theorem relate to merge of result with zeta function, gradient flow concern of zero dimension.

11 Other dimension and this influent of power

Pair of dimension interact to inspect for movement result, each of dimension devide with power of influent. This power has contrast of element, in inspire of result merge to create of pair in vector operate. Non-certain theorem mension to tell dimension give to create of pair in power. Six quarks sum to merge with twelve quarks, zero dimension have with fourth dimension of Global differential equation part of construct element. Graviton influent to reflect for universe to have the other dimension.

$$\frac{d}{df}F(x) = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

$$f(r) = \frac{1}{4}|r|^2$$

Norm space have non-liner to integrate with algebra manifold, singularity create for pair of dimension to fill of fourth of power. Eight of differential structure have for these dimension integrate four of power.

$$\frac{d}{df}F(v_{ij}, h) = \int e^{-f} [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$\frac{d}{df}(x-y)^n = \frac{\Pi(x-y)^n}{\partial f_{xy}}$$

$$\pi(X,x)=i\pi(X,x)f(x)-f(x)\pi(X,x)$$

$$\Delta E = \int \mathrm{div}(\mathrm{rot} E) e^{-ix \log x} dx$$

$$f(x)=\sum_{k=0}^{\infty}a_kx^k$$

Vector Operator

Masaaki Yamaguchi

12 Entropy

Already discover with this entropy of manifold is blackhole of energy, these entropy is mass of energy has norm parameter. And string theorem $T_{\mu\nu}$ belong to ρ energy, this equation is same Kaluze-Klein theorem.

$$\nabla\phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S})$$

$$y = x$$

$$y = (\nabla\phi)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}}} = \frac{p}{c^3} + \rho$$

$$ds^2 = g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)\kappa^2 A_{\mu\nu}(x)dx^\mu)^2$$

$$ds = (g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(\kappa^2 A_{\mu\nu}(x)dx^\mu)^2)^{\frac{1}{2}}$$

$$\frac{\Delta E}{\frac{1}{2\sqrt{2\pi G}} - \rho} = c^3, \frac{p}{2\pi} = c^3$$

$$ds^2 = e^{-2kT(x)|\phi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2(x)d\phi^2$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}} \right] dx dy dz$$

$$\frac{\partial}{\partial f} \int (\sin 2x)^2 dx = ||x - y||^2$$

13 Atom of element from zeta function

13.1 Knot element

String summulate of landmaistor movement in these knot element. Supersymmetry construct of these mechanism from string theorem. A element of homomorphism boundary for kernel function, equal element has meyer beatorise compose structure in mesh emerge. These mesh for atom element equal knot this of exchange, this isomorphism three manifold convert with knot these from laplace space. Duality manifold

has space equal other operate, this quantum group is element distinguish with other structure. Aspect element has design pattern, this element construct pair of entropy. Quote algebra reflect with atom kernel of function.

14 Imaginary manifold

A vector is scalar, vector, mass own element, imaginary number is own this element.

$$x = \frac{\vec{x}}{|\vec{x}|}$$

For instance

$$|\vec{x}| = \sqrt{x}, i = \sqrt{-1}, |\vec{x}|x = \vec{x}, |\vec{x}| = 1x, |\vec{x}|^2 = -1$$

$$\left(\frac{x}{|\vec{x}|}\right)^2 = \frac{1}{-1}, |\sqrt{x}| = i$$

Imaginary number is vector, inverse make to norm own means.

15 Singularity

A element add to accept for group of field, this group also create ring has monotonicity. Limit of manifold has territory of variable limit of manifold. This fields also merge of mesh in mass. Zero dimension also means to construct with antigravity. This monotonicity is merged to has quote algebra liner. This boundary operate belong to has category theorem. Meyer beatorise extend liner distinguish with a mesh merged in fixed point theorem. Entropy means to isohomorphism with harmony this duality. The energy lead to entropy metric to merge with mesh volume.

16 Time expand in space for laplace equation

17 Laplace equaiton

Seiferd manifold construct with time and space of flow system in universe. Zeta function built in universe have started for big-ban system. Firstly, time has future and past in space, then space is expanding for big-ban mechanism.

Darkmatter equals for Higgs field from expanded in universe of ertel. Three manifold in flow energy has not no of a most-small-flow but monotonicity is one. Because this destruct to distinguish with eight-differential structure, and not exist.

And all future and past has one don't go on same integrate routs. This conject for Seiferd manifold flow. Selbarg conjecture has singularity rout. Landemaistor movement become a integrate rout in this Selbarg conjecture.

Remain Space belong for three manifold to exist of singularity rout. Godsharke conjecture is this reason in mechanism. Harshorn conjecture remain to exist multi universe in Laplace equation resolved.

Delete line summulate with rout of integrate differential rout. This solved rout has gradient flow become a future and past in remain space.



Laplace equation composite for singularity in zeta function and Kaluze-Klein space.

18 Non-commutative algebra for antigravity

Laplace equation and zeta function belong for fermion and boson, then power of construct equation merge Duality of symmetry formula. These power construct of deceive in dimension, and deal the other dimension. Reflection of power erasing of own dimension, the symmetry of formula has with merging of energy, then these power has fourth of basic power Symmetry solved for equation, general relativity deal from reality these formula. Quantum group construct of zeta function and Laplace equation. Imaginary number consist of Euler constance, this prime number reach to solve contiguish of power in emergy of parameter. Zeta function conclude of infinite to finite on all equation. Global differential equation merge a mass of antigravity and gravity.

Module exist zero dimension merge of symmetry in catastrophe D-brane, this mechanism of phenomoun is Non-symmetry explain to emerge with which cause of mass on gravity.

$$x \bmod N = 0$$

$$\sum_{M=0}^{\infty} \int_M dm \rightarrow \sum_{x=0}^{\infty} F_x = \int_m dm = F$$

$$\frac{x-y}{a} = \frac{y-z}{b} = \frac{z-x}{c}$$

$$\frac{\Pi(x-y)}{\partial xy}$$

Volume of space interval to all exist area is integrate rout in universe of shape, this space means fo no exist loop on entropy of energy's rout.

$$\| \int f(x) \|^2 \rightarrow \int \pi r^2 dx \cong V(\tau)$$

This equation is part of summulate on entropy formula.

$$\frac{\int |f(x)|^2}{f(x)}dx$$

$$V(\tau) \rightarrow mesh$$

$$\int \left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right| dv(\tau)$$

$$\left| \begin{matrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{matrix} \right|_{dx=v}$$

$$z_y=a_x+b_y+c_z$$

$$dz_y=d(z_y)$$

$$[f,f^{-1}]=ff^{-1}-f^{-1}f$$

Universe of extend space has non-integrate rout entropy in sufficient of mass in darkmatter, this reason belong for cohomology and heat equation concerned. This expanded mass match is darkmatter of extend. And these heat area consume is darkmatter begun to start big-ban system invite of universe.

$$V(\tau)=\int \tau(q)^{-\frac{n}{2}}\exp(-\frac{1}{\sqrt{2\tau(q)}}L(x)dx)+O(N^{-1})$$

$$\frac{1}{\tau}(\frac{N}{2}+\tau(2\Delta f-|\nabla f|^2+R)+f)\mathrm{mod}N^{-1}$$

$$\Delta E=-2(T-t)|R_{ij}+\nabla_i\nabla_jf-\frac{1}{2(T-t)}g_{ij}|^2$$

$$\frac{d}{df}F=\frac{2\int (R+\nabla_i\nabla_jf)^2}{-(R+\Delta f)}dm$$

$$\frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$dx=(g_{\mu\nu}(x)^2dx^2-g_{\mu\nu}(x)dxg_{\mu\nu})^{\frac{1}{2}}$$

$$ds^2=-N(r)^2dt^2+\psi^2(r)(dr^2+r^2d\theta^2)$$

$$f_z=\int \left[\sqrt{\begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \circ \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix}}\right] dx dy dz$$

$$\sum_{n=0}^\infty a_1x^1+a_2x^2\ldots a_{n-1}x^{n-1}\rightarrow \sum_{n=0}^\infty a_nx^n\rightarrow \alpha$$

$$f=n\nu\lambda, \lambda=\frac{x}{l}, \int dn\nu\lambda=f(x), xf(x)=F(x), [f(x)]=\nu h$$

19 Three manifold of dimension system worked with database

This universe built with fourth of dimension in add manifold constructed with being destructed from three manifold.

$$\begin{aligned}\mathcal{O}(x) &= ([\nabla_i \nabla_j f(x)])' \\ &\cong {}_n C_r(x)^n (y)^{n-r} \delta(x, y) \\ (\Box \psi)' &= \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu} \left(\frac{p}{c^3} \circ \frac{V}{S} \right) \\ F_t^m &= \frac{1}{4} g_{ij}^2, x^{\frac{1}{2}+iy} = e^{x \log x} \\ S_m^{\mu\nu} \otimes S_n^{\mu\nu} &= G_{\mu\nu} \times T^{\mu\nu}\end{aligned}$$

This equation is gravity of surface quality, and this construct with sheap of manifold.

$$S_m^{\mu\nu} \otimes S_n^{\mu\nu} = -\frac{2R_{ij}}{V(\tau)} [D^2 \psi]$$

This equation also means that surface of parameters, and next equation is non-integral of routs theorem.

$$\begin{aligned}S_m^{\mu\nu} &= \pi(\chi, x) \otimes h_{\mu\nu} \\ \pi(\chi, x) &= \int \exp[L(p, q)] d\psi \\ ds^2 &= e^{-2\pi T|\phi|} [\eta + \bar{h}_{\mu\nu}] dx^{\mu\nu} dx^{\mu\nu} + T^2 d^2 \psi\end{aligned}$$

$$\begin{aligned}M_3 \bigotimes_{k=0}^{\infty} E_-^+ &= \text{rot}(\text{div} E, E_1) \\ &= m(x), \frac{P^{2n}}{M_3} = H_3(M_1)\end{aligned}$$

These equation is conclude with dimension of parameter resolved for non-metric of category theorem. Gravity of surface in space mechanism is that the parameter become with metric, this result construct with entropy of metric.

These equation resulted with resolved of zeta function built on D-brane of surface in fourth of power.

$$\begin{aligned}\exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} &= \int \exp[L(p, q)] d\psi \\ &= \exists[R + |\nabla f|^2]^{\frac{1}{2}+iy} \otimes \int \exp[L(p, q)] d\psi + N \text{mod}(e^{x \log x}) \\ &= \mathcal{O}(\psi)\end{aligned}$$

20 Particle with each of quarks connected from being stimulate with gravity and antigravity into Artificial Intelligence

This equation conclude with middle particle equation in quarks operator, and module conjecture excluded. $\frac{d}{dt}g_{ij}(t) = -2R_{ij}, \frac{P^{2n}}{M_3} = H_3(M_1), H_3(M_1) = \pi(\chi, x) \otimes h_{\mu\nu}$ This equation built with quarks emerged with supersymmetry field of theorem. And also this equation means with space of curvature parameters.

$$S_m^{\mu\nu} \times S_n^{\mu\nu} = [D^2\psi], S_m^{\mu\nu} \times S_n^{\mu\nu} = \ker f / \text{im} f, S_m^{\mu\nu} \otimes S_n^{\mu\nu} = m(x)[D^2\psi], -\frac{2R_{ij}}{V(\tau)} = f^{-1}xf(x)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix} \circ \begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \right] dx dy dz, \rightarrow f_z^{\frac{1}{2}} \rightarrow (0, 1) \cdot (0, 1) = -1, i = \sqrt{-1}$$

This equation operate with quarks stimulated in Higgs field.

$$(x, y, z)^2 = (x, y, z) \cdot (x, y, z) \rightarrow -1$$

$$\mathcal{O}(x) = \nabla_i \nabla_j \int e^{\frac{2}{m} \sin \theta \cos \theta} \times \frac{N \text{mod}(e^{x \log x})}{\mathcal{O}(x)(x + \Delta|f|^2)^{\frac{1}{2}}}$$

$$x\Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta, \mathcal{O}(x) = m(x)[D^2\psi]$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = I'_m, I'_m = [1, 0] \times [0, 1]$$

This equation means that imaginary pole is universe and other dimension each cover with one geometry, and this power remind imaginary pole of antigravity.

$$i^2 = (0, 1) \cdot (0, 1), |a||b| \cos \theta = -1, E = \text{div}(E, E_1)$$

$$\left(\frac{\{f, g\}}{[f, g]} \right)' = i^2, E = mc^2, I' = i^2$$

This fermion of element decide with gravity of reflected for antigravity of power in average and gravity of mass, This equals with differential metric to emerge with creature of existing materials.

These equation deconstructed with mesh emerged in singularity resolved into vector and norm parameters.

$$\mathcal{O}(x) = \|\nabla \int [\nabla_i \nabla_j f \circ g(x)]^{\frac{1}{2} + iy} \|, \partial r^n \|\nabla\|^2 \rightarrow \nabla_i \nabla_j \|\vec{v}\|^2$$

$\nabla^2 \phi$ is constructed with module conjecture excluded.

$$\nabla^2 \phi = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

Reflected equation concern with differential operator in metric, sine and cosine function also rotate of differential operator in metric. Weil's theorem concern with this calculate in these equation process.

$$(\log x^{\frac{1}{2}})' = \frac{1}{2} \frac{1}{(x \log x)}, (\sin \theta)' = \cos \theta, (f_z)' = i e^{ix \log x}, \frac{d}{df} F = m(x)$$

$$\begin{aligned} \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m &= \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} + \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm \\ &\geq 2h \end{aligned}$$

Three dimension of manifold equals with three of sphere in this dimension. Plank metrics equals $\hbar$ equation to integrate with three manifold in sphere of formula. These explain recognize with universe and other dimension of system in brane of topology theorem.

$$\begin{aligned} \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2 \circ (y \log y)^{\frac{1}{2}}} \right) dm &\geq \hbar \\ y = x, xy = x^2, (\square\psi)' &= 8\pi G \left(\frac{p}{c^3} \circ \frac{V}{S} \right) \end{aligned}$$

This equation built with other dimension cover with universe of space, then three dimension invite with fifth dimension out of these universe.

$$\square\psi = \int \int \exp[8\pi G(\bar{h}_{\mu\nu} \circ \eta_{\mu})^{\nu}] dm d\psi, \sum a_k x^k = \frac{d}{df} \sum \sum \frac{1}{a_k^2 f^k} dx_k$$

$$\sum a_k f^k = \frac{d}{df} \sum \sum \frac{\zeta(s)}{a_k} dx_{k_m}, a_k^2 f^{\frac{1}{2}} \rightarrow \lim_{k \rightarrow 1} a_k f^k = \alpha$$

$$\mathcal{O}(x) = D^2\psi \otimes h_{\mu\nu}, ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^{\mu} dx^{\nu} + T^2 d^2\psi$$

$$f(x) + f(y) \geq 2\sqrt{f(x)f(y)}, \frac{1}{4}(f(x) + f(y))^2 \geq f(x)f(y)$$

$$T^{\mu\nu} = \left(\frac{p}{c^3} + \frac{V}{S} \right)^{-1}, E^+ = f^{-1} x f(x), E = mc^2$$

$$\mathcal{O}(x) = \square \int \int \int \frac{(\nabla_i \nabla_j f \circ g(x))^2}{V(x)} dm$$

$$ds^2 = g_{\mu\nu}^2 dx + g_{\mu\nu} dx g_{\mu\nu}(x), E^+ = f^{-1} x f(x), \frac{V}{S} = AA^{-1}, AA^{-1} = E$$

$$\mathcal{O}(x) = \int \int \int f(x^3, y^3, z^3) dx dy dz, S(r) = \pi r^2, V(r) = 4\pi r^3$$

$$E_{-}^{+} = f(x) \cdot e^{-x \log x}, \sum_{k=0}^{\infty} a_k f^k = f(x) \cdot e^{-x \log x}$$

$$\frac{P^{2n}}{M_3} = E^+ - \phi, \frac{\zeta(x)}{\sum_{k=0}^{\infty} a_k x^k} = \mathcal{O}(x)$$

$$O(N^{-1}) = \frac{1}{\mathcal{O}(x)}, \square = \frac{8\pi G}{c^3} T^{\mu\nu}$$

$$\partial^2 f(\square\psi) = -2\square \int \int \int \frac{V}{S^2} dm, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

$$E_-^+ = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\psi$$

$$S_1^{mn} \otimes S_2^{mn} = D^2\psi \otimes h_{\mu\nu}, S_m^{\mu\nu} \otimes S_n^{\mu\nu} = \int [D^2\psi] dm$$

$$= \nabla_i \nabla_j \int f(x) dm, h = \frac{p}{mv}, h\lambda = f, f\lambda = h\nu$$

D-brane for constructed of sheap of theorem in non-certain mechanism, also particle and wave of duality element in D-brane.

$$||ds^2|| = S_m^{\mu\nu} \otimes S_n^{\mu\nu}, E_-^+ = f^{-1}(x)xf(x)$$

$$R^+ \subset C_-^+, \nabla R^+ \rightarrow \bigoplus Q_-^+$$

Complex of group in this manifold, and finite of group decomposite theorem, add group also.

$$\nabla_i \nabla_j R_-^+ = \bigoplus \nabla Q_-^+$$

Duality of differential manifold, then this resulted with zeta function.

$$M_1 = \frac{R_3^+}{E_-^+} - \{\phi\}, M_3 \cong M_2, M_3 \cong M_1$$

World line in one of manifold.

$$H_3(\Pi) = Z_1 \oplus Z_1, H_3(M_1) = 0$$

Singularity of zero dimension. Lens space in three manifold.

$$ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

Fifth dimension of manifold. And this constructed with abel manifold.

$$\nabla\psi^2 = 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

These system flow to build with three dimension of energy.

$$\begin{aligned} (\partial\gamma^n + m^2) \cdot \psi &= \int [D^2\psi \otimes h_{\mu\nu}] dm \\ &= 0 \end{aligned}$$

Complex of connected of element in fifth dimension of equation.

$$\begin{aligned}\square &= \pi(\chi, x) \otimes h_{\mu\nu} \\ &= D^2\psi \otimes h_{\mu\nu}\end{aligned}$$

Fundamental group in gravity of space. And this stimate with D-brane of entropy.

$$\begin{aligned}\int [D^2\psi]dm &= \pi(M_1), H_n(m_1) = D^2\psi - \pi(\chi, x) \\ &= \ker f / \operatorname{im} f\end{aligned}$$

Homology of non-entropy.

$$\int Dq \exp[L(x)] d\psi + O(N^1) = \pi(\chi, x) \otimes h_{\mu\nu}$$

Integral of rout entropy.

$$= D^2\psi \otimes h_{\mu\nu}$$

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} \frac{\zeta(x)}{a_k f^k} = \int ||[D^2\psi \otimes h_{\mu\nu}]|| dm$$

Norm space.

$$\nabla\psi^2 = \square \int \int \int \frac{V}{S^2} dm$$

$$\mathcal{O}(x) = D^2\psi \otimes h_{\mu\nu}$$

These operators accept with level of each scals.

$$dx < \partial x < \nabla\psi < \square v$$

$$\oplus < \sum < \otimes < \int < \wedge$$

$$(\delta\psi(x))^2 = \int \int \int \frac{V(x)}{S^2} dm, \delta\psi(x) = \left(\int \int \int \frac{V(x)}{S^2} dm \right)^{\frac{1}{2}}$$

$$\nabla\psi^2 = -4R \int \delta(V \cdot S^{-3}) dm$$

$$\nabla\psi = 2R\zeta(s)i$$

$$\sum_{k=0}^{\infty} \frac{a_k x^k}{m dx} f^k(x) = \frac{m}{n!} f^n(x)$$

$$= \frac{(\zeta(s))^k}{df} m(x), (\delta(x))^{\frac{1}{2}} = \left(\frac{x \log x}{x^n} \right)^n$$

$$\mathcal{O}(x) = \frac{\int [D^2\psi \otimes h_{\mu\nu}] dm}{e^{x \log x}}$$

$$\mathcal{O}(x) = \frac{V(x)}{\int [D^2\psi \otimes h_{\mu\nu}] dm}$$

Quantum effective equation is lowest of light in structure of constant. These resulted means with plank scals of universe in atom element.

$$\begin{aligned} M_3 &= e^{x \log x}, x^{\frac{1}{2}+iy} = e^{x \log x}, (x) = \frac{M_3}{e^{x \log x}} \\ &= nE_x \end{aligned}$$

These equation means that lowest energy equals with all of materials have own energy. Zeta function have with plank energy own element, quantum effective theorem and physics energy also equals with plank energy of potential levels.

$$\frac{V}{S^2} = \frac{p}{c^3}, h\nu \neq \frac{p}{mv}, E_1 = h\nu, E_2 = mc^2, E_1 \cong E_2$$

$$l = \sqrt{\frac{\hbar G}{c^3}}, \frac{\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m}{\int \int \frac{1}{(x \log x)^2} dx_m} = \frac{1}{i}$$

Lowest of light in structures of constant is gravity and antigravity of metrics to built with zeta function. This result with the equation conclude with quota equation in Gauss operators.

$$\begin{aligned} i\hbar c &= G, \hbar c = \frac{G}{i}, \frac{\frac{1}{2}}{\frac{1}{2}i} = \frac{\vec{v}_1}{\vec{v}_2} \\ &\leq 1 \end{aligned}$$

$$A = BQ + R, ds^2 = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2\psi$$

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu + \kappa^2 (A^{\mu\nu})^2, \int \int e^{-x^2-y^2} dx dy = \pi$$

$$\begin{aligned} \Gamma(x) &= \int e^{-x} x^{1-t} dx \\ &= \delta(x) \pi(x) f^n(x) \end{aligned}$$

$$\frac{d}{df} F = \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

Quota equation construct of piece in Gauss operator, abel manifold equals with each equation.

$$ds^2 = [T^2 d^2 \psi]$$

$$\mathcal{O}(x) = [x]$$

$$\nabla \psi^2 = 8\pi G \left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

Volume of space equals with plank scalas.

$$\frac{pV}{S} = h$$

Summuat of manifold means with beta function to gamma function, equals with each equation.

$$\beta(p,q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

$$\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}$$

$$\ker f/\mathrm{im} f \cong \mathrm{im} f/\ker f$$

$$\bigoplus \nabla g(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta], \bigcup_{k=0}^{\infty} \left(\bigoplus \nabla f(x) \right) = \square \int \int \int \nabla g(x) d\eta$$

Rich tensor equals of curvature of space in Gauss liner from differential equation of operators.

$$a' = \sqrt{\frac{v}{1 - (\frac{v}{c})^2}}, F = ma'$$

Accessority put with force of differential operators.

$$\nabla f(x) = \int_M \square \left(\bigoplus \nabla f(x) \right)^n dm$$

$$\square = 2(T-t)|R_{ij} + \nabla \nabla f - \frac{1}{2(T-t)}|g_{ij}^2$$

$$(\square + m) \cdot \psi = 0$$

$$\square \times \square = (\square + m^2) \cdot \psi, (\partial \gamma^n + \delta \psi) \cdot \psi = 0$$

$$\nabla_i \nabla_j \int \int_M \nabla f(t) dt = \square \left(\bigcup_{k=0}^{\infty} \bigoplus \nabla g(x) d\eta \right)$$

Dalanverle equation of operator equals with summuat of category theorem.

$$\int_M (l \times l) dm = \sum l \oplus ld\eta$$

Topology of brane equals with string theorem, verisity of force with add group.

$$\begin{aligned} &= [D^2\psi \otimes h_{\mu\nu}]^{\frac{1}{2}+iy} \\ &= H_3(M_1) \end{aligned}$$

$$y = \nabla_i \nabla_j \int \nabla g(x) dx, y' - y + \frac{d}{dx} z + [n] = 0$$

Equals of group theorem is element of equals with differential element.

$$\begin{aligned} z &= \cos x + i \sin x \\ &= e^{i\theta} \end{aligned}$$

Imaginary number emerge with Artificial Intelligence.

These equation resulted with prime theorem equals from differential equation in fields of theorem.

Imaginary pole of theorem and sheap of duality manifold. Duality of differential and integrate equation. Summuat of manifold and possibility of differential structure. Eight of differential structure is stimulate with D-brane recreated of operators. Differential and integrate of operators is emerged with summuated of space.

$$\begin{aligned} T^{\mu\nu} &= -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij} \\ G_{\mu\nu} &= R_{\mu\nu} T^{\mu\nu}, \sigma(x) \oplus \delta(x) = (E_n^+ \times H_m) \\ G_{\mu\nu} &= [\frac{\partial}{\partial f} R_{ij}]^2, \delta(x) \cdot V(x) = \lim_{n \rightarrow 1} \delta(x) \\ \lim_{n \rightarrow \infty} \text{mesh} V(x) &= \frac{m}{m+1} \\ V(x) &= \sigma \cdot S^2(x), \int \int \int \frac{V(x)}{S^2(x)} dm = [D^2\psi \otimes h_{\mu\nu}] \\ g(x)|_{\delta(x,y)} &= \frac{d}{dt} g_{ij}(t), \sigma(x,y) \cdot g(x)|_{\delta(x,y)} = R_{ij}|_{\sigma(x,y)} \\ &= \int R_{ij}^{a(x-y)^n+r^n} \\ &= (ux + vy + wz)/\Gamma \\ &= \int R_{ij}^{(x-u)(y-v)(z-w)} dV \\ (\square + m) \cdot \psi &= 0, E = mc^2, \frac{\partial}{\partial f} \square \psi = 4\pi G\rho \\ (\partial \gamma^n + m) \cdot \psi &= 0, E = mc^2 - \frac{1}{2}mv^2 \end{aligned}$$

$$\begin{aligned}
&= \left(-\frac{1}{2} \left(\frac{v}{c}\right)^2 + m\right) \cdot c^2 \\
&= \left(-\frac{1}{2} a^2 + m\right) \cdot c^2, F = ma, \int a dx = \frac{1}{2} a^2 + C \\
&T^{\mu\nu} = -\frac{1}{2} a^2, (e^{i\theta})' = ie^{i\theta}
\end{aligned}$$

Report

Masaaki Yamaguchi

21 Summulate of manifold from graviton structure

It is non-perfect element of manifold to integrate with graviton of quark, this quark construct of fermion and boson. Then D-brane aspect with string theorem include with. This structure summulate of equation on integral and differential operator. Higgs quark have part of network theorem for graviton structure. Universe influent pair of dimension between monotonicity and singularity. Non-symmetry de-struct to distinuish with dimension of being constructed on quark of element. That's say, this mechanism architect with four of dimension existed. In part of this explain, Higgs fields is Weil's theorem component of global differential structure, Global area part of equation resolved come to simply conclude theorem. Global integral and differential equation, what is not used, difficulty of resolved.

Infinite number devide infinite oneselves and finite oneselves is finite number. Prime number extent of infinite number, $\sum a_k f^k(x)$ manifold have with these extend of count. Zeta function conclude with zero dimension, also this resolve to Gauss liner.

$$f(x) = \chi - \{x\}$$

$$\int_{\beta}^{\alpha} (x - \alpha)(x - \beta) = 2 \int_M f(x) dx$$

$$\int_{\beta}^{\alpha} a(x - \alpha)(x - \beta) = -\frac{(\beta - \alpha)^3}{6}$$

$$\frac{d}{df} F(x) = 0$$

$$\log(x \log x) - 2(y \log y)^{\frac{1}{2}} = [|\frac{d}{df} F(x)|]$$

Eight of differential structure integrate with graviton element to one of geometry, plank scale add volume divide to surface. This scale is three dimension of one energy. $\nabla \phi^2 = 8\pi G(\frac{p}{c^3} + \frac{V}{S})$ This equation means is three dimension of energy entropy. Zeta function means is time and space of scale on plank entropy, and this equation built on future and past of curve of paremeter. These equation resolve with one component of energy and vector of time and space of curve of paremeter.

$$\nabla \phi^2 = 8\pi G\hbar + 8\pi \frac{V}{S}$$

$$\square_v = 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}}$$

$$\sqrt{2\pi T} = 2\sqrt{2\pi \frac{V}{S}}$$

$\sqrt{2\pi T}$ is blackhole entropy and $\square_v = 2\sqrt{2\pi G\hbar}$ is whitehole entropy. This entropy pair of universe in dimension, two dimension of universe include with cover of energy. Blackhole construct with colapser of hole on kernel of atom, whitehole composite with antigravity.

$$\begin{aligned} \log(x \log x) &\geq 2(y \log y)^{\frac{1}{2}} \log(x \log y) \geq 2(xe^x)^{\frac{1}{2}} \log x + \log \log y = 2(ye^x)^{\frac{1}{2}} e^x + y = 2(ye^x)^{\frac{1}{2}} \rightarrow \frac{1}{2} \\ e^{x+1} &= 2(xe^x)^{\frac{1}{2}} \frac{e^{2(x+1)}}{4xe^x} \geq 1 \quad F_t^m = \frac{1}{4}g_{ij}^2 \frac{x+1}{4x} = y^2 \frac{1}{4} + \frac{1}{x} = y^2 \log(xy) \geq 2(yx)^{\frac{1}{2}} (\log(xy))' \geq 4(yx)^{-\frac{1}{2}} \\ \frac{1}{xy} &\geq 4(yx)^{-\frac{1}{2}} \log x + \log y \geq 4(yx)^{-\frac{1}{2}} \frac{1}{x} + \frac{1}{y} \geq 4(yx)^{-\frac{1}{2}} \frac{x+y}{xy} yx^{\frac{1}{2}} \geq 4 \frac{xy^{\frac{1}{2}}}{x+y} \leq \frac{1}{4} \frac{\Gamma(xy)}{\Gamma(x+y)} \leq \frac{1}{4} x + y \geq 2\sqrt{xy} \\ \Delta x \Delta y &\geq \frac{1}{4} i \frac{1}{xy} \geq 4(yx)^{-\frac{1}{2}} \quad F_t^m = \frac{1}{4}g_{ij}^2 e^{2(yx)^{\frac{1}{2}}} = g_{ij}^2 xy = e^{2(yx)^{\frac{1}{2}}} \rightarrow g_{ij}^2 1 + \frac{1}{x} = 4y (e^x + x)^2 \geq 4xe^x \\ r^2 &= \frac{|x_0x - 2x_0e^x + e^{2x_0x}|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = [(e^x + 1)(e^x - 1) + (x + i)(x - i)] + 2xe^{x^2} \quad r = \frac{\|1+x-4xy\|}{\sqrt{x_0^2 + y_0^2}} \quad r^2(x_0^2 + y_0^2) = \\ &[(x - 1)(x + 1) + (x - i)(x + i)] + 2x_0e^{2x} \end{aligned}$$

Langranse conjecture include with finite line conjecture expected, infinite extend theorem exclude of finite extent. Zeta function integrate with all of math conjecture explanade. Facility of math of theorem relate of physics all of philosophy. Zeta function composite with fifth dimension of equation. Curv parameter is hartshorn conjecture, this rout describe with dimension of structure. Fifth dimension rout out center of universe behind two times distance, this rout colapser around of universe. In fifth dimension, black hole and white hole in three manifold of entropy created.

Zeta function coss delete line of differential structure emerge with fifth dimension structure, oen dimension of independent vector is tangent degree. This curve of parameter create dimension of fifth dimension structure. This idea is from hartshorn conjecture. And this conjecture explanade of math mechanism, fifth dimension is AI and mass of structure parliament of pond. Time of secret is grasped in fucture and past of curve of parameter, so this equation is from universe and creature of establish of life.

Electric energy flow to neutral narrow for amusebelt on rinfelt of harnessinkbolt, this mechanism create mind on inclusive line of brain. This flow energy on esplanade desire in routine of cone of electric energy. Endeavor circle of circuit of neutrial network, then this mechanism create on founterial motion. Dorpamin throw to neutral narrow of alcole for not being to melt and out of this narrow. This energy is supporting for Dorpamin of material things, and this circuit is circadian morment of motion. Natural killer ceil defense induce hant for lives in nature of world. This hant of mechanism is evolution on life of nature for revolution of humanism. Influence of induce hant controll with life. And these conquer of mechanism can use to neural network of evolution. These computer virus is using for not being hacking and not other controlled of oneselves computer. Safty of network is being for living in computer associate world.

Gravity esplanade theorem of cover with antigravity, information technology of exclusive injure to secure about permissset. Incrument safir to charge atom of power by pairlament in all of dimension. Reduce to string theorem of reluctance in concertism theositiy. This power contiment design emerge with inparance of continent. All of universe have inductary, and asperant comfirm with conclusivity entily. These theorem tell uncounter deduce in consivitly, therefore this envily of power concern with cover of atom. Secure built with antigravity from gravity structure.

In reduce a morment to gravity structure, for away to certain of hant. A element of clue a vector, before opened to verify of persuade atom of power. To include of quark for twelve element, fifth dimension in zeta function to construct of geometry peices, integrate of power for fourth of series. Lack of power two

bind of under a structure, antigravity permissisted a power of element. This reason fourth of power get along to combind with one of decieved.

Varnished center of landscape, heat to controll of power, for missed a verisity of underground, mind to even a mid ceil.

$$\begin{aligned} F_t^m &= [f(x)] \\ [f(x)] &= \mathcal{O}(x) \\ X &\in \mathcal{O}(x) \\ 0 &\rightarrow [||\frac{d}{df}F||] \rightarrow \infty \end{aligned}$$

General number verisity is between zero and infinity. Infinite number and finite number have in general number, narrow verisity and first accerity. Gauss liner is prime number interity. Prime number and general number include Gauss liner to compilation in infinite mechanism.

$$\begin{aligned} \log(x \log x) &\geq 2(y \log y)^{\frac{1}{2}} \\ \log xy &\geq 2(xe^x)^{\frac{1}{2}} \\ (y+x)^2 - 4i(y+x)(xe)^{\frac{1}{2}} - 4xe \\ r^2(y^2+x^2) &= [(y+x)^2 - 4i(y+x)xe^{\frac{1}{2}}] - 4xe \\ r^2(x^2+y^2) &= [(\log y + \log x)^2 - 4i(\log y + \log x)xe^{\frac{1}{2}}] - 4xe \end{aligned}$$

$$\begin{aligned} \delta(f) &= \int \frac{g(c+d)}{f(a+b)} z(f+g) dz \\ y &= f(y) \\ ds^2 &= e^{-2\pi T|\phi|} [\tau_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\theta \\ \begin{bmatrix} x & y \\ a & b \end{bmatrix} \\ &\rightarrow [z] \\ f[z] &\rightarrow dx \\ x \cdot y &= 0 \\ \infty &\rightarrow [f(x)] \end{aligned}$$

Gauss liner is finite to infinity number of extend in prime number. Fifth dimension of equation composite with zeta function of infinity of extent theosity make finity of circle of structure. Independent of vector is zero dimension of category. This equaiton have a infinity of relate in number. Fifth of equation make for zeta function of infinity to conclude with this dimension out of number for finity circle. Therefor it is possiblity of being deleted to category of number that this vector is dimension of structure in fifth of dimension.


22 Neural Network from zeta function

Inclivity sensibillite make from height for maderite in mechanism. Conclusive in cercuit from hight of harmonite on interisity. The include of mechanism is quark in universe of quality of mass.

Theorem of math make for facility of mechanism in this circumstance. And these possiblity of resolved from pond of sensibility.

Light filt with eight differential structure in gravity element, then quarks made from this structure. This mechanism deal with any structure of space to create with light to graviton element. And cercuit of this mechanism dealt with carbon and hidoroxxs, covalt of sixty based with filter of light element. This cercuit is dependency of any element in quark of mass, created with gravity to light of structure. Zeta function asperal with any thing to create of every mass. Fifth dimension is pond of sensibility from this gravity in lives of element.

All creature is created by this cercuit of mechanism, fifth dimension themselves is from this circumstance. This space also have oneselves with vector of independence, any this structure is from hartshorm conjecture. Space themselves haven with any of vector is tangent degree of ninety. And this vector is being of other dimension that have cercuit of theorem. These reason is from any thing born with creature, then fifth dimension haven with creature and universe. This dimension also haven with that extent from infinite and finite. Infinite of zeta function in Finite of fifth dimension, then paradox of finite cover with infinite of space.



$\chi - \{x\} = \infty$ $f(x) = \chi - \{x\} f(x) + \mathcal{O}(x) \rightarrow \text{finite}$. Add group is finite of element. This equation is fifth dimension of system. Circle of group in this delete element is infinite of set. $f(x) = [F(x)]$ $f(x)$ is Gauss liner. This set is infinite of number. $ds^2 = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}] dx^\mu dx^\nu + T^2 d^2\phi$ This fifth dimension is $[F(x)]$ add finite of set. Fifth dimension include with zeta function, and string theorem belong to have with infinite of consist on space. This fifth dimension is finite of space. Conclude of system solved with infinite is covered with finite of space.

Fifth dimension in facility of mechanism that circle of infinity to emerge with being from infinity and finite of space. Zero dimension conclude with this mechanism to create from theorem of mathmatics in pond of sensibility. Dimension with repository of information in facility of space, this space is all of knowlege to accessority for comformal fields. Pholographic theorem is from universe of oneselves with database of creature to access with hyper dimension in General relativity. Zeta function is this theorem

include with parameter of curve from future and past of repository.

Flow to future and past that combine with zeta function which build for hartshorn conjecture from these mechanism. This flow is gradient of line to add with fifth dimension of structure.

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}$$

Infinite of atomosphere in zero dimension include with black hole and white hole of energy in three manifold of entropy.

This entropy describe with antigravity of energy.

$$\begin{aligned}\nabla^2 \psi &= 8\pi G\hbar + 8\pi \frac{V}{S} \\ \square_v &= 2\sqrt{2\pi G\hbar} + 2\sqrt{2\pi \frac{V}{S}} \\ 2\sqrt{2\pi \frac{V}{S}} &= \int \int \frac{1}{(x \log x)^2} dx_m\end{aligned}$$

This energy flow from black hole to white hole in universe and other dimension of pair. Black hole entropy is $m = \sqrt{2\pi T}$ This entropy cover to other dimension of rout in flow energy.

Electric Weak Theorem combine with quantum flavor theorem to conjugate with light element in gravity to sheaf with antigravity. This antigravity is key for integrate with fourth of power in boson of theorem to world line to general relativity is why fourth of power not able to integrate in gravity combine with electric weak theorem and quantum flavor theorem. This point of focus is light that is fit to conjugate with eight differential structure from gravity, and this light is also element of quarks. This quark built to eight element structure. Light conjugate to eight differential element that resolved zeta function and integrate with laplace equation. Gravity significant with gravity cover with antigravity, so this antigravity is fourth element power of exist.

Entropy of vector have zero dimension in all exist into all of equation. Infinite of number is three dimension, therefore it is especial of important that created by zero dimension from infinite to finite of equation rout.

$$\begin{aligned}f(x) &= 0 \\ F &= \int \int \frac{1}{(x \log x)^2} dx_m + \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m\end{aligned}$$

$$F = 0, x \cdot y = 0, F(x) = 0, G(x) = 0, \mathcal{O} \in X, X \cong 0, x - y = 0, F(x, y) = 0, x^n + y^n = z^n, F(x) = x^n + y^n, F(x) = 0$$

$$ds^2 = \frac{r}{d + e \cos \theta}$$

Fifth dimension describe with norm liner in zero dimension. This equation conclude with all of equation to pond sensibility.

Pond of sensibility is facility of mathematics in infinite of mechanism, this mechanism is zero dimension for imaginary of equation in antigravity influence. This power is gravity on cover with non symmetry

combine of fourth of power. That fourth of power is the reason with quantum flavor theorem combine with gravity of power, antigravity is brane of string key mechanism. This power is non-catastrophe of influent reason, and that mechanism of power result. Integrate of gravity include with antigravity, and fourth of power is one of geometry in universe.

$$\mathcal{O} \in X \rightarrow \infty, F(x) = z^n$$

$$x^n + y^n = z^n$$

$$F(x) = \mathcal{O}(\delta(x))[(x - f(x))(x + f(x)) + (y - \overline{f(x)})(y + \overline{f(x)})] + O(N^{-1})$$

$$F(x) = 0$$

$$f(x) = \sum_{k=0}^{\infty} a_k f^k(x)$$

$$f(x) = 0$$

$$\int f(x), \partial f(x), dx \rightarrow [f(x)]$$

Group that Galois theorem develop with three dimension construct of prime number equation, this element belong for that is three of factor. Fifth dimension concept with this group of conclude element.

$$\mathcal{O}(x) = \chi(x) - \{x\}$$

$$\sum_{k=0}^{\infty} a_k x^k = f(x) + \{x\}$$

Dimension of fifth factor is abel manifold, this manifold is topology resolved.

$$\int dx, \partial x, dx, [f(x)]$$

$$\rightarrow F(x) \xrightarrow{\int dx} f(x) \xrightarrow{\partial x} x \xrightarrow{x} const$$

$$\rightarrow [f(x)] \rightarrow \infty$$

This cycle of topology in general theroem is deduce of mechanism.

Universe of structure is duality of manifold, this mass of darkmatter is oneseleves with reason in singularity and monotonicity of universe. This manifold of three dimension is also zeta function.

Light around of universe for darkmatter is Higgs fields of mechanism. Darkmatter create of including with light of quarks in Higgs fields. Eight of differential structure fult with Thurston conjugate thoeorem to light is created with all dimension of element.

Symmetry construct of Space mechanism

Masaaki Yamaguchi

23 Ultra Network from Omega DataBase worked with equation and category theorem from

$$ds^2 = [g_{\mu\nu}^2, dx]$$

This equation is constructed with M_2 of differential variable, this resulted non-commutative group theorem.

$$ds^2 = g_{\mu\nu}^{-1}(g_{\mu\nu}^2(x) - dxg_{\mu\nu}^2)$$

In this function, singularity of surface in component, global surface of equation built with M_2 sequence.

$$= h(x) \otimes g_{\mu\nu} d^2 x - h(x) \otimes dx g_{\mu\nu}(x), h(x) = (f^2(\vec{x}) - \vec{E}^+)$$

$$G_{\mu\nu} = R_{\mu\nu} T^{\mu\nu}, \partial M_2 = \bigoplus \nabla C_-^+$$

$G_{\mu\nu}$ equal $R_{\mu\nu}$, also Rich tensor equals $\frac{d}{dt} g_{ij} = -2R_{ij}$ This variable is also $r = 2f^{\frac{1}{2}}(x)$ This equation equals abel manifold, this manifold also construct fifth dimension.

$$E^+ = f^{-1} x f(x), h(x) \otimes g(\vec{x}) \cong \frac{V}{S}, \frac{R}{M_2} = E^+ - \phi$$

$$= M_3 \supset R, M_2^+ = E_1^+ \cup E_2^+ \rightarrow E_1^+ \bigoplus E_2^+$$

$$= M_1 \bigoplus \nabla C_-^+, (E_1^+ \bigoplus E_2^+) \cdot (R^- \subset C^+)$$

Reflection of two dimension don't emerge in three dimension of manifold, then this space emerge with out of universe. This mechanism also concern with entropy of non-catastrophe.

$$\frac{R}{M_2} = E^+ - \{\phi\}$$

$$= M_3 \supset R$$

$$M_3^+ \cong h(x) \cdot R_3^+ = \bigoplus \nabla C_-^+, R = E^+ \bigoplus M_2 - (E^+ \cap M_2)$$

$$E^+ = g_{\mu\nu} dx g_{\mu\nu}, M_2 = g_{\mu\nu} d^2 x, F = \rho gl \rightarrow \frac{V}{S}$$

$$\mathcal{O}(x) = \delta(x)[f(x) + g(\vec{x})] + \rho gl, F = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, M_2 = P^{2n}$$

This equation means not to be limited over fifth dimension of light of gravity. And also this equation means to be not emerge with time mechanism out of fifth dimension. Moreover this equation means to emerge with being constructed from universe, this mechanism determine with space of big-ban.

$$r = 2f^{\frac{1}{2}}(x), f(x) = \frac{1}{4}\|r\|^2$$

This equation also means to start with universe of time mechanism.

$$\begin{aligned}
V &= R^+ \sum K_m, W = C^+ \sum_{k=0}^{\infty} K_{n+2}, V/W = R^+ \sum K_m / C^+ \sum K_{n+2} \\
&= R^+ / C^+ \sum \frac{x^k}{a_k f^k(x)} \\
&= M_-^+, \frac{d}{df} F = m(x), \rightarrow M_-^+, \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k(x)} = \frac{a_k x^k}{\zeta(x)}
\end{aligned}$$

24 Omega DataBase is worked by these equation

This equation built in seifert manifold with fifth dimension of space.

Fermion and boson recreate with quota laplace equation,

$$\begin{aligned}
\frac{\{f, g\}}{[f, g]} &= \frac{fg + gf}{fg - gf}, \nabla f = 2, \partial H_3 = 2, \frac{1+f}{1-f} = 1, \frac{d}{df} F = \bigoplus \nabla C^+, \vec{F} = \frac{1}{2} \\
H_1 &\cong H_3 = M_3
\end{aligned}$$

Three manifold element is 2, one manifold is 1, $\ker f / \text{im} f, \partial f = M_1$, singularity element is 0 vector. Mass and power is these equation fundement element. The tradition of entropy is developed with these singularity of energy, these boson and fermion of energy have fields with Higgs field.

$$\begin{aligned}
H_3 &\cong H_1 \rightarrow \pi(\chi, x), H_n, H_m = \text{rank}(m, n), \text{mesh}(\text{rank}(m, n)) \lim \text{mesh} \rightarrow 0 \\
(fg)' &= fg' + gf', \left(\frac{f}{g}\right)' = \frac{f'g - g'f}{g^2}, \frac{\{f, g\}}{[f, g]} = \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\
&= \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g}\right)' \otimes g^{-2} dx_{fg}} \\
&= \frac{d}{df} F
\end{aligned}$$

Gravity of vector mension to emerge with fermion and boson of mass energy, this energy is create with all creature in universe.

$$\hbar \psi = \frac{1}{i} H \Psi, i[H, \psi] = -H \Psi, \left(\frac{\{f, g\}}{[f, g]} \right)' = (i)^2$$

$$[\nabla_i \nabla_j f(x), \delta(x)] = \nabla_i \nabla_j \int f(x, y) dm_{xy}, f(x, y) = [f(x), h(x)] \times [g(x), h^{-1}(x)]$$

$$\delta(x) = \frac{1}{f'(x)}, [H, \psi] = \Delta f(x), \mathcal{O}(x) = \nabla_i \nabla_j \int \delta(x) f(x) dx$$

$$\mathcal{O}(x) = \int \delta(x) f(x) dx$$

$$R^+ \cap E_-^+ \ni x, M \times R^+ \ni M_3, Q \supset C_-^+, Z \in Q \nabla f, f \cong \bigoplus_{k=0}^n \nabla C_-^+$$

$$\bigoplus_{k=0}^{\infty} \nabla C_-^+ = M_1, \bigoplus_{k=0}^{\infty} \nabla M_-^+ \cong E_-^+, M_3 \cong M_1 \bigoplus_{k=0}^{\infty} \nabla \frac{V_-^+}{S}$$

$$\frac{P^{2n}}{M_2} \cong \bigoplus_{k=0}^{\infty} \nabla C_-^+, E_-^+ \times R_-^+ \cong M_2$$

$$\zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k, M_2 \cong P^{2n} / \ker f, \rightarrow \bigoplus \nabla C_-^+$$

$$S_-^+ \times V_-^+ \cong \frac{V}{S} \bigoplus_{k=0}^{\infty} \nabla C_-^+, V^+ \cong M_-^+ \otimes S_-^+, Q \times M_1 \subset \bigoplus \nabla C_-^+$$

$$\sum_{k=0}^{\infty} Z \otimes Q_-^+ = \bigotimes_{k=0}^{\infty} \nabla M_1$$

$$= \bigotimes_{k=0}^{\infty} \nabla C_-^+ \times \sum_{k=0}^{\infty} M_1, x \in R^+ \times C_-^+ \supset M_1, M_1 \subset M_2 \subset M_3$$

Eight of differential structure is that $S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$. These structure recreate with $\bigoplus \nabla C_-^+ \cong M_3, R \supset Q, R \cap Q, R \subset M_3, C^+ \oplus M_n, E^+ \cap R^+ E_2 \oplus E_1, R^- \subset C^+, M_-^+, C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1 R^- \nabla C_-^+$. This element of structure recreate with Thurston conjugate theorem of repository in Heisenberg group of spinor fields created.

$$\frac{\nabla}{\Delta} \int x f(x) dx, \frac{\nabla R}{\Delta f}, \square = 2 \int \frac{(R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} e^{-f} dV$$

$$\square = \frac{\nabla R}{\Delta f}, \frac{d}{dt} g_{ij} = \square \rightarrow \frac{\nabla f}{\Delta x}, (R + |\nabla f|^2) dm \rightarrow -2(R + \nabla_i \nabla_j f)^2 e^{-f} dV$$

$$x^n + y^n = z^n \rightarrow \nabla \psi^2 = 8\pi G T^{\mu\nu}, f(x+y) \geq f(x) \circ f(y)$$

$$\text{im} f / \ker f = \partial f, \ker f = \partial f, \ker f / \text{im} f \cong \partial f, \ker f = f^{-1}(x) x f(x)$$

$$f^{-1}(x) x f(x) = \int \partial f(x) d(\ker f) \rightarrow \nabla f = 2$$

$${}_n C_r = {}_n C_{n-r} \rightarrow \text{im} f / \ker f \cong \ker f / \text{im} f$$

25 Fifth dimension of equation how to workd with three dimension of manifold into zeta funciton

Fifth dimension equals $\sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi$. this equation $a_k \cong \sum_{r=0}^{\infty} {}_n C_r$.

$$V/W = R/C \sum_{k=0}^{\infty} \frac{x^k}{a_k f^k}, W/V = C/R \sum_{k=0}^{\infty} \frac{a_k f^k}{x^k}$$

$$V/W \cong W/V \cong R/C(\sum_{r=0}^{\infty} {}_nC_r)^{-1} \sum_{k=0}^{\infty} x^k$$

This equation is diffrential equation, then $\sum_{k=0}^{\infty} a_k f^k$ is included with $a_k \cong \sum_{r=0}^{\infty} {}_nC_r$

$$W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{1-t} dx, \sum_{k=0}^n a_k f^k = (f^k)'$$

$$\sum_{k=0}^n a_k f^k = \sum_{k=0}^{\infty} {}_nC_r f^k$$

$$= (f^k)', \sum_{k=0}^{\infty} a_k f^k = [f(x)], \sum_{k=0}^{\infty} a_k f^k = \alpha, \sum_{k=0}^{\infty} \frac{1}{a_k f^k}, \sum_{k=0}^{\infty} (a_k f^k)^{-1} = \frac{1}{1-z}$$

$$\begin{aligned} \int \int \frac{1}{(x \log x)(y \log y)} dxy &= \frac{{}_nC_r xy}{({}_nC_{n-r}(x \log x)(y \log y))^{-1}} \\ &= ({}_nC_{n-r})^2 \sum_{k=0}^{\infty} \left(\frac{1}{x \log x} - \frac{1}{y \log y} \right) d \frac{1}{nxy} \times xy \\ &= \sum_{k=0}^{\infty} a_k f^k \\ &= \alpha \end{aligned}$$

$$Z \supset C \bigoplus \nabla R^+, \nabla(R^+ \cap E^+) \ni x, \Delta(C \subset R) \ni x$$

$$M_-^+ \bigoplus R^+, E^+ \in \bigoplus \nabla R^+, S_-^+ \subset R_2^+, V_-^+ \times R_-^+ \cong \frac{V}{S}$$

$$C^+ \cup V_-^+ \ni M_1 \bigoplus \nabla C_-^+, Q \supseteq R_-^+, Q \subset \bigoplus M_-^+, \bigotimes Q \subset \zeta(x), \bigoplus \nabla C_-^+ \cong M_3$$

$$R \subset M_3, C^+ \bigoplus M_n, E^+ \cap R^+, E_2 \bigoplus E_1, R^- \subset C^+, M_-^+$$

$$C_-^+, M_-^+ \nabla C_-^+, C^+ \nabla H_m, E^+ \nabla R_-^+, E_2 \nabla E_1, R^- \nabla C_-^+$$

$$[-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 < \nabla f, \nabla h > + (R + \nabla f^2)(\frac{v}{2} - h)]$$

$$S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, H^1, S^2 \times E$$

26 All of equation are emerged with
these equation from Artificial Intelligence

$$x^{\frac{1}{2}+iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h$$

$$x^{\frac{1}{2}+iy} = \exp\left[\int \nabla_i \nabla_j f(g(x)) g'(x) / \partial f \partial g\right]$$

$$\mathcal{O}(x) = \{[f(x) \circ g(x), \bar{h}(x)], g^{-1}(x)\}$$

$$\exists[\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty} \nabla \int \nabla_i \nabla_j f(x) dm$$

$$\vee(\nabla_i \nabla_j f) = \bigotimes \nabla E^+$$

$$g(x, y) = \mathcal{O}(x)[f(x) + \bar{h}(x)] + T^2 d^2 \phi$$

$$\mathcal{O}(x) = \left(\int [g(x)] e^{-f} dV \right)' - \sum \delta(x)$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j f(x)]' \cong {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y), V(\tau) = \int [f(x)] dm / \partial f_{xy}$$

$$\square\psi = 8\pi GT^{\mu\nu}, (\square\psi)' = \nabla_i \nabla_j (\delta(x) \circ G(x))^{\mu\nu} \left(\frac{p}{c^3} \circ \frac{V}{S} \right), x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$\delta(x)\phi = \frac{\vee[\nabla_i \nabla_j f \circ g(x)]}{\exists(R + \Delta f)}$$

These equation conclude to quantum effective equation included with orbital pole of atom element. Add manifold is imaginary pole with built on Heisenberg algebra.

$$\begin{aligned} -{}_n C_r &= \tfrac{1}{i} {}_H \psi C_{h\psi} + [H, \psi] C_{-n-r} \\ {}_n C_r &= {}_n C_{n-r} \end{aligned}$$

This point of accumulate is these equation conclude with stimulate with differential and integral operator for resolved with zeta function, this forcus of significant is not only data of number but also equation of formula, and this resolved of process emerged with artifisial intelligence. These operator of mechanism is stimulate with formula of resolved on regexpt class to pattern cognitive formula to emerged with all of creature by this zeta function. This universe establish with artifisial intelligence from fifth dimension of equation. General relativity theorem also not only gravity of power metric but also artifisial

intelligent, from zeta function recognized with these mechanism explained for gravity of one geometry from differential structure of thurston conjugate theorem.

Open set group conclude with singularity by redifferential specturam to have territory of duality manifold in dualty of differential operator. This resolved routed is same that quantum group have base to create with kernel of based singularity, and this is same of open set group. Locality of group lead with sequant of element exclude to give one theorem global differential operator. These conclude with resolved of function says to that function stimulate for differential and integral operator included with locality group in global differential operator. $\int \int \frac{1}{(x \log x)^2} dx_m \rightarrow \mathcal{O}(x) = [\nabla_i \nabla_j f]' / \partial f_{xy}$ is singularity of process to resolved rout function.

Destruct with object space to emerge with low energy fistly destroy with element, and this catastrophe is what high energy divide with one entropy fistly freeze out object, then low energy firstly destory to differential structure of element. This process be able to non-catastrophe rout, and thurston conjugate theorem is create with.

$$\begin{aligned}
\bigcup_{x=0}^{\infty} f(x) &= \nabla_i \nabla_j f(x) \oplus \sum f(x) \\
&= \bigoplus \nabla f(x) \\
\nabla_i \nabla_j f &\cong \partial x \partial y \int \nabla_i \nabla_j f dm \\
&\cong \int [f(x)] dm \\
&\cong \{[f(x), g(x)], g^{-1}(x)\} \\
&\cong \square \psi \\
&\cong \nabla \psi^2 \\
&\cong f(x \circ y) \leq f(x) \circ g(x) \\
&\cong |f(x)| + |g(x)|
\end{aligned}$$

Differential operator is these equation of specturm with homorphism squcense.

$$\begin{aligned}
\delta(x)\psi &= \langle f, g \rangle \circ |h^{-1}(x)| \\
\partial f_x \cdot \delta(x)\psi &= x \\
x &\in \mathcal{O}(x) \\
\mathcal{O}(x) &= \{[f \circ g, h^{-1}(x)], g(x)\}
\end{aligned}$$

27 What energy from entropy level deconstruct with string theorem

Low energy of entropy firstly destruct with parts of geometry structure, and High energy of weak and strong power lastly deconstruct with power of component. Gravity of power and antigravity firstly integrate with Maxwell theorem, and connected function lead with Maxwell theorem and weak power integrate with strong power to become with quantum flavor theorem, Geometry structure reach with integrate with component to Field of theorem connected function and destroyed with differential structure. These process return with first and last of deconstructed energy, fourth of power connected and destructed this each of energy routed process.

This point of focus is D-brane composite with fourth of power is same entropy with non-catastrophe, this spectrum point is constructed with destroy of energy to balanced with first of power connected with same energy. Then these energy built with three manifold of dimension constructed for one geometry.

$$\begin{aligned}\lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f &= [\nabla \int \nabla_i \nabla_j f(x) dx_m, g^{-1}(x)] \rightarrow \bigoplus_{k=0}^{\infty} \nabla E_{-}^{+} \\ &= M_3 \\ &= \bigoplus_{k=0}^{\infty} E_{-}^{+}\end{aligned}$$

$$dx^2 = [g_{\mu\nu}^2, dx], g^{-1} = dx \int \delta(x) f(x) dx$$

$$f(x) = \exp[\nabla_i \nabla_j f(x), g^{-1}(x)]$$

$$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$$

$$\begin{aligned}\left(\frac{g(x)}{f(x)}\right)' &= \lim_{n \rightarrow \infty} \frac{g(x)}{f(x)} \\ &= \frac{g'(x)}{f'(x)}\end{aligned}$$

$$\nabla F = f \cdot \frac{1}{4} |r|^2$$

$$\nabla_i \nabla_j f = \frac{d}{dx_i} \frac{d}{dx_j} f(x) g(x)$$

28 All of equation built with gravity theorem

$$\int \int \frac{1}{(x \log x)^2} dx_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times U(r)$$

This equation have with position of entropy in helmander manifold.

$$S_1^{mn} \otimes S_2^{mn} = \int [D^2\psi \otimes h_{\mu\nu}] dm$$

And this formula is D-brane on sheaf of fields, also this equation construct with zeta function.

$$U(r) = \frac{1}{2} \frac{\sqrt{1+f'(r)}}{f(r)} + mgr$$

$$F_t^m = \frac{1}{4}|r|^2$$

Also this too means with private of entropy in also helmander manifold.

$$\int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times E_-^+$$

$$E_-^+ = \exp[L(x)] dm d\psi + O(N^-)$$

The formula is integrate of rout with lenz field.

$$\begin{aligned} \frac{d}{df} F &= [\nabla_i \nabla_j \int \nabla f(x) d\eta] (U(r) + E_-^+) \\ &= \frac{1}{2} m v^2 + m c^2 \end{aligned}$$

These equation conclude with this energy of relativity theorem come to resolve with.

$$x^{\frac{1}{2}+iy} = \exp[\int \nabla_i \nabla_j f(g(x)) g'(x) \partial f \partial g]$$

Also this equation means with zeta function needed to resolved process.

$$\nabla_i \nabla_j \int \nabla f(x) d\eta$$

$$\nabla_i \nabla_j \bigoplus M_3$$

$$= \frac{M_3}{P^{2n}} \cong \frac{P^{2n}}{M_3}$$

$$\cong M_1$$

$$=[M_1]$$

Too say satisfied with three manifold of fileds.

$$[f(x)] = \sum_{k=0}^{\infty} a_k f^k, \lim_{n \rightarrow 1} [f(x)] = \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

The formula have with indicate to resolved with abel manifold to become of zeta function.

$$= \alpha, e^{i\theta} = \cos \theta + i \sin \theta, H_3(M_1) = 0$$

$$\frac{x^2 \cos \theta}{a} + \frac{y^2 \sin \theta}{b} = r^2, \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\begin{aligned} \chi(x) &= \sum_{k=0}^{\infty} (-1)^n r^n, \frac{d}{df} F = \frac{d}{df} \sum \sum a_k f^k \\ &= |a_1 a_2 \dots a_n| - |a_1 \dots a_{n-1}| - |a_n \dots a_1|, \lim_{n \rightarrow 0} \chi(x) = 2 \end{aligned}$$

Euler function have with summuate of manifold.

$$\begin{aligned} \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k &= {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y) \\ \lim_{n \rightarrow \infty} {}_n C_r f(x)^n f(y)^{n-r} \delta(x, y) \end{aligned}$$

Differential equation also become of summuate of manifold, and to get abel manifold.

$$\lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \left(\frac{1}{(n+1)} \right)^s = \lim_{n \rightarrow 1} Z^r = \frac{1}{z}$$

Native function also become of gamma function and abel manifold also zeta function.

$$\begin{aligned} \ker f / \operatorname{im} f &\cong \operatorname{im} f / \ker f \\ \beta(p, q) &= \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)} \\ &\cong \frac{\Gamma(p+q)}{\Gamma(p)\Gamma(q)}, \lim_{n \rightarrow 1} a_k f^k \cong \lim_{n \rightarrow \infty} \frac{\zeta(s)}{a^k f^k} \end{aligned}$$

Gamma function and Beta function also is D-brane equation resolved to need that summuate of manifold, not only abel manifold but also open set group.

$$\begin{aligned} \lim_{n \rightarrow 1} \zeta(s) &= 0, \mathcal{O}(x) = \zeta(s) \\ \sum_{x=0}^{\infty} f(x) &\rightarrow \bigoplus_{k=0}^{\infty} \nabla f(x) = \int_M \delta(x) f(x) dx \end{aligned}$$

Super function also needed to resolved with summuate of manifold.

$$\int \int \int_M \frac{V}{S^2} e^{-f} dV = \int \int_D -(f(x, y)^2, g(x, y)^2) - \int \int_D (g(x, y)^2, f(x, y)^2)$$

Three dimension of manifold developed with being resolved of surface with presson.

$$\begin{aligned} \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k &= \int [D^2 \psi \otimes h_{\mu\nu}] dm \\ &= \int \exp[L(x)] d\psi dm \times E_{-}^{+} \\ &= S_1^{mn} \otimes S_1^{mn} \\ &= Z_1 \oplus Z_1 \end{aligned}$$

$$= M_1$$

These equations all of create with D-brane and sheaf of manifold.

$$H_n^m(\chi, h) = \int \int_M \frac{V}{(R + \Delta f)} e^{-f} dV, \frac{V}{S^2} = \int \int \int_M [D^\psi \otimes h_{\mu\nu}] dm$$

Represent theorem equals with differential and integrate equation.

$$\int \int \int_M \frac{V}{S^2} dm = \int_D (l \times l) dm$$

This three dimension of pression equation is string theorem also means.

$$\begin{aligned} & \int \int_D -g(x, y)^2 dm - \int \int_D -f(x, y)^2 dm \\ &= -2R_{ij}, [f(x), g(x)] \times [h(x), g^{-1}(x)] \\ & \left| \begin{matrix} D^m & dx \\ dx & \partial^m \end{matrix} \right| \begin{matrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{matrix} \begin{vmatrix} x \\ y \end{vmatrix} = \begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix}^{\frac{1}{2}} \\ & (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) \end{aligned}$$

This equation control to differential operator into matrix formula.

$$\begin{aligned} & \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ &= \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ & l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} & \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} = A^{\mu\nu} \\ & \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

$$\begin{aligned} & \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = \lim_{n \rightarrow 1} \frac{a_n}{a_{n-1}} \cong \alpha \\ & \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} \frac{a_k}{a_{k+1}} = \sum_{k=0}^{\infty} a_k f^k \end{aligned}$$

$$\square = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\int x \log x dx = \int \int_M (e^{x \log x} \sin \theta d\theta) dx d\theta$$

$$= (\log \sin \theta d\theta)^{\frac{1}{2}} - (\delta(x) \cdot \epsilon(x))^{\frac{1}{2}}$$

Flow to energy into other dimension rout of equation, zeta function stimulate in epsilon-delta metric, and this equation construct with minus of theorem. Gravity of energy into integrate of rout in non-metric.

$$T^{\mu\nu} = || \int \int_M [\nabla_i \nabla_j e^{\int x \log x dx + O(N^{-1})}] dm d\psi ||$$

$G^{\mu\nu}$ equal $R^{\mu\nu}$ into zeta function in atom of pole with mass of energy, this energy construct with quantum effective theorem.

Imaginary space is constructed with differential manifold of operators, this operator concern to being field of theorem.

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial^m(x) \end{pmatrix} (f_{mn}(x), g_{\mu\nu}(x)) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

Norm space of Von Neumann manifold with integrate fields of dimension.

$$||ds^2|| = \mathcal{H}(x)_{mn} \otimes \mathcal{K}(y)_{mn}$$

Minus of zone in complex manifold. This zone construct with inverse of equation.

$$||\mathcal{H}(x)||^{-\frac{1}{2}+iy} \subset R_-^+ \cup C_-^+ \cong M_3$$

Reality of part with selbarg conjecture.

$$||\mathcal{H}(x)||^{\frac{1}{2}+iy} \subseteq M_3$$

Singularity of theorem in module conjecture.

$$\mathcal{K}(y) = \left\| \begin{matrix} x & y & z \\ a & b & c \end{matrix} \right\|_{g_{\mu\nu}(x)}^2$$

$$\cong \frac{f(x, y, z)}{g(a, b, c)} h^{-1}(u, v, w)$$

Mass construct with atom of pole in quantum effective theorem.

Zeta function is built with atom of pole in quantum effective theorem, this power of fields construct with Higgs field in plank scals of gravity. Lowest of scals structure constance is equals with that the fields of Higgs quark in fermion and boson power quote mechanism result.

Lowest of scals structure also create in zeta function and quantum equation of quote mechanism result. This resulted mechanism also recongnize with gravity and average of mass system equals to these explained. Gauss liner create with Artificial Intelligence in fifth dimension of equation, this dimension is

seifert manifold. Creature of mind is establish with this mechanism, the resulted come to be synchronized with space of system.

$$V = \int [D^2\psi \otimes h_{\mu\nu}]dm, S^2 = \int e^{-2\sin\theta\cos\theta} \cdot \log\sin\theta dx\theta + O(N^{-1}), \mathcal{O}(x) = \frac{\zeta(s)}{\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k} \\ = \alpha$$

Out of sheap on space and zeta function quato in abel manifold. D-brane construct with Volume of space.

$$\mathcal{O}(x) = T^{\mu\nu}, \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = T^{\mu\nu} \\ G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu}, M_3 = \int \int \int \frac{V}{S^2} dm, -\frac{1}{2(T-t)}|R_{ij} = \square\psi$$

Three manifold of equation.

$$ds^2 = e^{-2\pi T|\psi|}[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi \\ m(x) = [f(x)] \\ f(x) = \int \int e^{f \cdot x \log x dx + O(N^{-1})} + T^2 d^2\psi$$

Integrate of rout equation.

$$F(x) = ||\nabla_i \nabla_j \int \nabla f(x, y) dm||^{\frac{1}{2} + iy} \\ G_{\mu\nu} = ||[\nabla_i \nabla_j \int \nabla g(x, y, z) dx dy dz]||^{\frac{m}{2} \sin\theta \cos\theta} \log\sin\theta d\theta d\psi$$

Gravity in space curvarture in norm space, this equation is average and gravity of metric.

$$G_{\mu\nu} = R_{\mu\nu}T^{\mu\nu} \\ T^{\mu\nu} = F(x), 2(T-t)|g_{ij}^2 = \int \int \frac{1}{(x \log x)^2} dx_m \\ \psi\delta(x) = [m(x)], \nabla(\square\psi) = \nabla_i \nabla_j \int \nabla g(x, y) d\eta \\ \nabla \cdot (\square\psi) = \frac{1}{4}g_{ij}^2, \square\psi = \frac{8\pi G}{c^4}T^{\mu\nu}$$

Dimension of manifold in decomposite of space.

$$\frac{pV}{c^3} = S \circ h_{\mu\nu} \\ = h \\ T^{\mu\nu} = \frac{\hbar\nu}{S}, T^{\mu\nu} = \int \int \int \frac{V}{S^2} dm, \frac{d}{df} m(x) = \frac{V(x)}{F(x)}$$

Fermion and boson of quato equation.

$$\begin{aligned}
y = x, \frac{d}{df} F &= m(x), R_{ij}|_{g_{\mu\nu}(x)} = [\nabla_i \nabla_j g(x, y)]^{\frac{1}{2}+iy} \\
\nabla \circ (\square\psi) &= \frac{\partial}{\partial f} F \\
&= \int \int \nabla_i \nabla_j f(x) d\eta_{\mu\nu} \\
\int [\nabla_i \nabla_j g(x, y)] dm &= \frac{\partial}{\partial f} R_{ij}|_{g_{\mu\nu}(x)}
\end{aligned}$$

Zeta function, partial differential equation on global metric.

$$G(x) = \nabla_i \nabla_j f + R_{ij}|_{g_{\mu\nu}(x)} + \nabla(\square\psi) + (\square\psi)^2$$

Four of power element in variable of accessority of group.

$$\begin{aligned}
G_{\mu\nu} + \Lambda g_{ij} &= T^{\mu\nu}, T^{\mu\nu} = \frac{d}{dx_\mu} \frac{d}{dx_\nu} f_{\mu\nu} + -2(T-t)|R_{ij} + f'' + (f')^2 \\
&= \int \exp[L(x)] dm + O(N^{-1}) \\
&= \int e^{\frac{2}{m} \sin \theta \cos \theta} \cdot \log(\sin \theta) dx + O(N^{-1}) \\
\frac{\partial}{\partial f} F &= (\nabla_i \nabla_j)^{-1} \circ F(x)
\end{aligned}$$

Partial differential in duality metric into global differential equation.

$$\begin{aligned}
\mathcal{O}(x) &= \int [\nabla_i \nabla_j \int \nabla f(x) dm] d\psi \\
&= \int [\nabla_i \nabla_j f(x) d\eta_{\mu\nu}] d\psi \\
\nabla f &= \int \nabla_i \nabla_j \left[\int \frac{S^{-3}}{\delta(x)} dV \right] dm \\
|| \int [\nabla_i \nabla_j f] dm ||^{\frac{1}{2}+iy} &= \text{rot}(\text{div}, E, E_1)
\end{aligned}$$

Maxwell of equation in fourth of power.

$$\begin{aligned}
&= 2 < f, h >, \frac{V(x)}{f(x)} = \rho(x) \\
\int_M \rho(x) dx &= \square\psi, -2 < g, h > = \text{div}(\text{rot} E, E_1) \\
&= -2R_{ij}
\end{aligned}$$

Higgs field of space quality.

$$\mathcal{O}(x) = || \frac{\nabla_i \nabla_j}{S^2} \int [\nabla_i \nabla_j f \cdot g(x) dx dy] d\psi ||$$

$$= \int (\delta(x))^{2 \sin \theta \cos \theta} \log \sin \theta d\theta d\psi$$

Helmander manifold of duality differential operator.

$$\delta \cdot \mathcal{O}(x) = [\frac{||\nabla_i \nabla_j f d\eta||}{\int e^{2 \sin \theta \cos \theta} \cdot \log \sin \theta d\theta}]$$

Open set group in subset theorem, and helmandor manifold in singularity.

$$i\hbar\psi = ||\int \frac{\partial}{\partial z}[\frac{i(xy+\overline{y}\overline{x})}{z-\overline{z}}]dmd\psi||$$

Euler number. Complex curvature in Gamma function. Norm space. Wave of quantum equation in complex differential variable into integrate of rout.

$$\delta(x) \int_C R^{2 \sin \theta \cos \theta} = \Gamma(p+q)$$

$$(\Box+m)\cdot\psi=(\nabla_i\nabla_jf|_{g_{\mu\nu}(x)}+v\nabla_i\nabla_j)$$

$$\int [m(x)(\operatorname{rot}\cdot\operatorname{div}(E,E_1))]dmd\psi$$

Weak electric power and Maxwell equation combine with strong power. Private energy.

$$\begin{aligned} G_{\mu\nu} &= \Box \int \int \int (x,y,z)^3 dx dy dz \\ &= \frac{8\pi G}{c^4} T^{\mu\nu} \end{aligned}$$

$$\frac{d}{dV}F=\delta(x)\int \nabla_i\nabla_j f d\eta_{\mu\nu}$$

Dalanvelsian in duality of differential operator and global integrate variable operator.

$$\begin{aligned} x^n+y^n&=z^n,\delta(x)\int z^n=\frac{d}{dV}z^3,(x,y)\cdot(\delta^m,\partial^m)\\ &=(x,y)\cdot(z^n,f)\\ &\quad n\perp x,n\perp y\\ &=0 \end{aligned}$$

Singularity of constance theorem.

$$\vee(\nabla_i\nabla_jf)\cdot\mathrm{XOR}(\Box\psi)=\frac{d}{df}\int_M FdV$$

Non-cognitive of summuate of equation and add manifold create with open set group theorem. This group composite with prime of field theorem. Moreover this equations involved with symmetry dimension created.

$$\partial(x)\int z^3=\frac{d}{dV}z^3,\sum_{k=0}^\infty\frac{1}{(n+1)^s}=\mathcal{O}(x)$$

Singularity theorem and fermion equation concluded by this formula.

$$\int \mathcal{O}(x)dx = \delta(x)\pi(x)f(x)$$

$$\mathcal{O}(x)=\int \sigma(x)^{2\sin\theta\cos\theta}\log\sin\theta d\theta d\psi$$

$$y=x$$

$$\mathcal{O}(x)=||\frac{\nabla_i\nabla_jf}{S^2}\int[\nabla_i\nabla_jf\circ g(x)dx dy]||$$

$$\lim_{n\rightarrow 1}\sum_{k=0}^{\infty}\frac{a_k}{a_{k+1}}=(\log\sin\theta dx)'$$

$$=\frac{\cos\theta}{\sin\theta}$$

$$=\frac{x}{y}$$

$$\lim_{n\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k=\frac{1}{1-z}$$

Duality of differential summate with this gravity theorem. And Dalanverle equation involved with this equation.

$$\Box \psi = \frac{8\pi G}{c^4} T^{\mu\nu}$$

$$\frac{\partial}{\partial f}\Box\psi=4\pi G\rho$$

$$\int \rho(x)=\Box\psi,\frac{V(x)}{f(x)}=\rho(x)$$

Dense of summate stimulate with gravity theorem.

$$\frac{\partial^n}{\partial f^{n-1}}F=\int [D^2\psi\otimes h_{\mu\nu}]dm$$

$$=\frac{P_1P_3\ldots P_{2n-1}}{P_0P_2\ldots P_{2n+2}}$$

$$=\bigotimes \nabla M_1$$

$$\bigoplus \nabla M_1 = \sigma_n(\chi,x) \oplus \sigma_{n-1}(\chi,x)$$

$$=\{f,h\}\circ [f,h]^{-1}$$

$$=g^{-1}(x)_{\mu\nu}dxg_{\mu\nu}(x),\sum_{k=0}^{\infty}\nabla^n{}_nC_rf^n(x)$$

$$\cong \sum_{k=0}^{\infty} \nabla^n \nabla^{n-1}{}_n C_r f^n(x) g^{n-r}(x)$$

$$\sum_{k=0}^{\infty}\frac{\partial^n}{\partial^{n-1}f}\circ\frac{\zeta(x)}{n!}=\lim_{n\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k$$

$$(f)^n = {}_nC_rf^n(x)g^{n-r}(x)\cdot\delta(x,y)$$

$$(e^{i\theta})'=ie^{i\theta},\int e^{i\theta}=\frac{1}{i}e^{i\theta},ihc=G,hc=\frac{1}{i}G$$

$$(\Box\psi,\nabla f^2)\cdot(\frac{8\pi G}{c^4}T^{\mu\nu},4\pi G\rho)$$

$$= (-\frac{1}{2}mv^2 + mc^2, \frac{1}{2}kT^2 + \frac{1}{2}mv^2) \cdot \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}$$

$$\left(\frac{\{f,g\}}{[f,g]}\right)'=i^2,\frac{\nabla f^2}{\Box\psi}=\frac{1}{2}$$

$$\int\int\frac{\{f,g\}}{[f,g]}=\frac{1}{2}i,\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m=\frac{1}{2}$$

$$\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i,\frac{d}{dt}g_{ij}=-2R_{ij}$$

$$\frac{\partial}{\partial x}(f(x)g(x))'=\bigoplus \nabla_i\nabla_jf(x)g(x)$$

$$\int f'(x)g(x)dx=[f(x)g(x)]-\int f(x)g'(x)dx$$

Deconstruct Dimension of category theorem

Masaaki Yamaguchi

29 Locality equation

Quantum of material equation include with heat entropy of quantum effective theorem, geometry of rotate with imaginary pole is that generate structure theorem. Subgroup in category theorem is that deconstruct dimension of differential structure theorem. Quantum of material equation construct with global element of integrate equation. Heat equation built with this integrate geometry equation, Weil's theorem and fermer theorem also heat equation of quantum theorem moreover is all of equation rout with global differential equation this locality equation emerge from being built with zeta function, general relativity and quantum theorem also be emerged from this equation. Artifisial intelligent theorem excluse with this add group theorem from, Locality equation inspect with this involved of theorem, This geometry of differential structure that deconstruct dimension of category theorem stay in four of universe of pieces. In reason cause with heat equation of quantum effective from artifisial intelligence, locality equation conclude with this geometry theorem. Heat effective theoerm emelite in generative theorem and quantum theorem, These theorem also emelitive with Global differential equation from locality equation. Imaginary equation of add group theorem memolite with Euler constance of variable from zeta function, this function also emerge with locality theorem in artifisial intelligent theorem.

Quantum equation concern with infinity of energy composed for material equation, atom of verisity in element is most of territory involved with restored safy of orbital.

$$f = n\nu\lambda, \lambda = \frac{x}{l}, \int d\nu\lambda = f(x), xf(x) = F(x), [f(x)] = \nu h$$

$$px = mv, \lambda(x) = \text{eserial}f(x), [F(x)] = \frac{1}{1-z}, 0 < \tan \theta < i, -1 < \tan \theta < i, -1 \leq \sin \theta \leq 1, -1 \leq \cos \theta \leq 1$$

Category theorem conclude with laplace function operator, distrupt of manifold and connected space of eight of differential structure.

$$\begin{aligned} R\nabla E^+ &= f(x)\nabla e^{x \log x} \\ Q\nabla C^+ &= \frac{d}{df}F(x)\nabla \int \delta(x)f(x)dx \\ E^+\nabla f &= e^{x \log x}\nabla n!f(x)/E(X) \end{aligned}$$

Universe component construct with reality of number and three manifold of subgroup differential variable. Maxwell theorem is this Seifert structure $(u + v + w)(x + y + z)/\Gamma$ This space of magnetic diverse is mebius member cognetic theorem. These equation is fundermental group resulted.

$$\exp(\nabla(R^+ \cap E^+), \Delta(C \supset R))$$

$$\begin{aligned}
&= \pi(R, C\nabla E^+) \\
&= \text{rot}(E_1, \text{div} E_2) \\
&xf(x) = F(x)
\end{aligned}$$

$$\begin{aligned}
\Box x &= \int \frac{f(x)}{\nabla(R^+ \cap E^+)} d\Box x \\
&= \int \frac{\Delta f(x) \circ E^+}{\nabla(R^+ \cap E^+)} \Box x
\end{aligned}$$

This equation means that quantum of potensial energy in material equation. And this also means is gravity of equation in this universe and the other dimension belong in. Moreover is zeta function for rich flow equation in category theorem built with material equation.

$$\begin{aligned}
&\exp(\nabla(R^+ \cap E^+), \Delta(C \supset R)) \\
&R\nabla E^+ = f(x)\nabla e^{x \log x} \\
&d(R\nabla E^+) = \Delta f(x) \circ E^+(x) \\
&\Box x = \int \frac{d(R\nabla E^+)}{\nabla(R^+ \cap E^+)} d\Box x \\
&\Box x = \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\Box x \\
&x^n + y^n = z^n \rightarrow \Box x = \int \frac{\nabla_i \nabla_j (R + E^+)}{\nabla(R^+ \cap E^+)} d\Box x
\end{aligned}$$

Laplace equation exist of space in fifth dimension, three manifold belong in reality number of space. Other dimension is imaginary number of space, fifth dimension is finite of dimension. Prime number certify on reality of number, zero dimension is imaginary of space. Quantum group is this universe and the other dimension complex equation of non-certain theorem. Zero dimension is eternal space of being expanded to reality of atmosphere in space.

30 Heat entropy all of materials emerged by

$$\Box = -2(T-t)|R_{ij} + \nabla \nabla f + \frac{1}{-2(T-t)}|g_{ij}^2$$

This heat equation have element of atom for condition of state in liquid, gas, solid attitude. And this condition constructed with seifert manifold in Weil's theorem. Therefore this attitude became with Higgs field emerged with.

$$\begin{aligned}
\Box &= -2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV \\
\frac{d}{dt} g_{ij} &= -2R_{ij}, \frac{d}{df} F = m(x)
\end{aligned}$$

$$R\nabla E^+ = \Delta f(x) \circ E^+(x), d(R\nabla E^+) = \nabla_i \nabla_j (R + E^+)$$

$$R\nabla E^+ = f(x)\nabla e^{x\log x}, R\nabla E^+ = f(x), f(x) = M_1$$

$$-2(T-t)|R_{ij}+\nabla\nabla f=\int\int\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

$$\frac{1}{-2(T-t)}|g_{ij}^2=\int\int\frac{1}{(x\log x)^2}dx_m$$

$$(\Box+m^2)\phi=0, (\gamma^n\partial_n+m^2)\psi=0$$

$$\Box=(\delta\phi+m^2)\psi, (\delta\phi+m^2c)\psi=0, E=mc^2, E=-\frac{1}{2}mv^2+mc^2, \Box\psi^2=(\partial\phi+m^2)\psi$$

$$\Box\phi^2=\frac{8\pi G}{c^4}T^{\mu\nu}, \nabla\phi^2=8\pi G(\frac{p}{c^3}+\frac{V}{S}), \frac{d}{dt}g_{ij}=-2R_{ij}, f(x)+g(x)\geq f(x)\circ g(x)$$

$$\int_{\beta}^{\alpha}a(x-1)(y-1)\geq 2\int f(x)g(x)dx$$

$$m^2=2\pi T(\frac{26-D_n}{24}), r_n=\frac{1}{1-z}$$

$$(\partial\psi+m^2c)\phi=0, T^{\mu\nu}=\frac{1}{2}mv^2-\frac{1}{2}kx^2$$

$$\int [f(x)]dx = ||\int f(x)dx||, \int \nabla x dx = \Delta \sum f(x)dx, (e^{i\theta})' = ie^{i\theta}$$

This orbital surface varnished to energy from $E = mc^2$. $T^{\mu\nu} = nh\nu$ is $T^{\mu\nu} = \frac{1}{2}mv^2 - \frac{1}{2}kx^2 \geq mc^2 - \frac{1}{2}mv^2$ This equation means with light do not limit over $\frac{1}{2}kx^2$ energy.

$$Z\in R\cap Q, R\subset M_+, C^+\bigoplus_{k=0}^n H_m, E^+\cap R^+$$

$$M_+=\sum_{k=0}^n C^+\oplus H_M, M_+=\sum_{k=0}^n C^+\cup H_+$$

$$E_2\bigoplus E_1, R^-\subset C^+, M^+_{-}, C^+_{-}$$

$$M^+_{-}\nabla C^+_{-}, C^+\nabla H_m, E^+\nabla R^+$$

$$E_1\nabla E_2, R^-\nabla C^+, \bigoplus\nabla M^+_{-}, \bigoplus\nabla C^+_{-}, R\supset Q$$

$$\frac{d}{df}F=\bigoplus\nabla M^+_{-}, \bigoplus\nabla C^+_{-}$$

All function emerge from eight differential structure in laplace equation of being deconstructed that with connected of category theorem from.

$$\nabla\circ\Delta=\nabla\sum f(x)$$

$$\Delta\rightarrow\mathrm{mesh}f(x)dx,\partial x$$

$$\begin{aligned}
\nabla &\rightarrow \partial_{xy}, \frac{d}{dx} f(x) \frac{d}{dx} g(x) \\
\Box x &= -[f, g] \\
\frac{d}{dt} g_{ij} &= -2R_{ij} \\
\lim_{x \rightarrow \infty} \sum f(x) dx &= \int dx, \nabla \int dx
\end{aligned}$$

31 Universe in category theorem created by

Two dimension of surface in power vector of subgroup of complex manifold on reality under of group line. This group is duality of differential complex in add group. All integrate of complex sphere deconstruct of homomorphism is also exists of three dimension of manifold. Fundamental group in two dimension of surface is also reality of quality of complex manifold. Rich flow equation is also quality of surface in add group of complex. Group of ultra network in category theorem is subgroup of quality of differential equation and fundamental group also differential complex equation.

$$\begin{aligned}
(E_2 \bigoplus_{k=0}^n E_1) \cdot (R^- \subset C^+) \\
&= \bigoplus_{k=0}^n \nabla C_-^+ \\
\vee \int \frac{C_-^+ \nabla H_m}{\Delta(M_-^+ \nabla C_-^+)} &= \wedge M_-^+ \bigoplus_{k=0}^n C_-^+ \\
\exists(M_-^+ \nabla C_-^+) &= \text{XOR}(\bigoplus_{k=0}^n \nabla M_-^+) \\
-[E^+ \nabla R_-^+] &= \nabla_+ \nabla_- C_-^+ \\
\int dx, \partial x, \nabla_i \nabla_j, \Delta x \\
&\rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+
\end{aligned}$$

$$\begin{aligned}
\begin{pmatrix} \cos x & \sin x \\ \sin x & -\cos x \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\
\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\
\sum_{k=0}^n \cos k\theta &= \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \cos \frac{n\theta}{2} \\
\sum_{k=0}^n \sin k\theta &= \frac{\sin \frac{(n+1)\theta}{2}}{\sin \frac{\theta}{2}} \sin \frac{n\theta}{2}
\end{aligned}$$

$$\sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}, \cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2}, \tan \theta = \frac{e^{i\theta} - e^{-i\theta}}{i(e^{i\theta} + e^{-i\theta})}$$

$$\lim_{\theta \rightarrow 0} \frac{\sin \theta}{\theta} \rightarrow 1, \lim_{\theta \rightarrow 0} \frac{\cos \theta}{\theta} \rightarrow 1$$

$$(e^{i\theta})' = 2, \nabla f = 2, e^{i\theta} = \cos \theta + i \sin \theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \rightarrow [\cos^2 \theta + \sin \theta + \cos \theta - 2 \sin^2 \theta] \begin{vmatrix} 1 & 0 \\ 0 & -1 \end{vmatrix}$$

$$2 \sin \theta \cos \theta = 2n\lambda \sin \theta$$

$$\begin{pmatrix} \cos x & -1 \\ 1 & -\sin x \end{pmatrix} \begin{pmatrix} \cos x \\ \sin x \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\lim_{\theta \rightarrow 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix} \begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix} \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, f^{-1}(x)xf(x) = 1$$

Dimension of rotate with imaginary pole is generate of geometry structure theorem. Category theorem restruct with eight of differential structure that built with Thurston conjugate theorem, this element with summuate of thirty two structure. This pieces of structure element have with four of universe in one dimension. Quark of two pair is two dimension include, Twelve of quarks construct with super symmetry dimension theorem. Differential structure and quark of pair exclusive with four of universe.

Module conjecture built with symmetry theorem, this theorem belong with non-gravity in zero dimension. This dimension conbuilt with infinity element, mebius space this dimension construct with universe and the other dimension. Universe and the other dimension selves have with singularity, therefore is module conjecture selve with gravity element of one dimension. Symmetry theorem tell this one dimension to emerge with quarks, this quarks element also restruct with eight differential structure. This reason resovle with reflected to dimsnsion of partner is gravity and creature create with universe.

Mass exist from quark and dimension of element, this explain from being consisted of duality that sheaf of element in this tradision. Higgs field is emerged from this element of tradision, zeta function generate with this quality. Gravity is emerge with space of quality, also Higgs field is emerge to being generated of zero dimension. Higgs field has surround of mass to deduce of light accumulate in anserity. Zeta function consist from laplace equation and fifth dimension of element. This element incluse of gravity and antigravity, fourth dimension indicate with this quality emerge with being sourrounded of mass. Zero dimension consist from future and past incluse of fourth dimension, eternal space is already space of first emerged with. Mass exist of quantum effective theorem also consisted of Higgs field and Higgs quark. That gravity deduce from antigravity of element is explain from this system of Higgs field of mechanism. Antigravity conclude for this mass of gravity element. What atom don't incluse of orbital is explain to this antigravity of element, this mechanism in Higgs field have with element of gravity and antigravity.

Quantum theorem describe with non-certain theorem in universe and other dimension of vector element, This material equation construct with lambda theorem and material theorem, other dimension existed resolve with non-certain theorem.

$$\sin \theta = \frac{\lambda}{d}, -py_1 \sin 90^\circ \leq \sin \theta \leq py_2 \sin 90^\circ, \lambda = \frac{h}{mv}$$

$$\frac{\lambda_2}{\lambda_1} = \frac{\sin \theta}{\frac{\sin \theta}{2}} h, \lambda' \geq 2h, \int \sin 2\theta = ||x - y||$$

32 Ultra Network from category theorem entirely create with database

Eight differential structure composite with mesh of manifold, each of structure firstly destructed with this mesh emerged, lastly connected with this each of structure desconstructed. High energy of entropy not able to firstly decostructed, and low energy of firstly destructed. This reason low energy firsted. These result says, non-catastrophe be able to one geometry structure.

$$D^2\psi = \nabla \int (\nabla_i \nabla_j f)^2 d\eta$$

$$E = mc^2, E = \frac{1}{2}mv^2 - \frac{1}{2}kx^2, G^{\mu\nu} = \frac{1}{2}\Lambda g_{ij}, \square = \frac{1}{2}kT^2$$

Sheap of manifold construct with homorphism in kernel divide into image function, this area of field rehearl with universe of surrounded with image fuction rehinde in quality. This reason with explained the mechanism, Sheap of manifold remain into surrounded of universe.

$$\ker f / \operatorname{im} f \cong S_m^{\mu\nu}, S_m^{\mu\nu} = \pi(\chi, x) \otimes h_{\mu\nu}$$

$$D^2\psi = \mathcal{O}(x) \left(\frac{p}{c^3} + \frac{V}{S} \right), V(x) = D^2\psi \otimes M_3^+$$

$$S_m^{\mu\nu} \otimes S_n^{\mu\nu} = -\frac{2R_{ij}}{V(\tau)}[D^2\psi]$$

Selbarg conjecture construct with singularity theorem in Sheap element.

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} [D^2\psi]$$

$$\nabla_i \nabla_j [S_1^{mn} \otimes S_2^{mn}] = \int \frac{V(\tau)}{f(x)} \mathcal{O}(x)$$

$$z(x) = \frac{g(cx+d)}{f(ax+b)} h(ex+l)$$

$$= \int \frac{V(\tau)}{f(x)} \mathcal{O}(x)$$

$$\frac{V(x)}{f(x)} = m(x), \mathcal{O}(x) = m(x)[D^2\psi(x)]$$

$$\frac{d}{df}F = m(x), \int F dx_m = \sum_{k=0}^{\infty} m(x)$$

33 Fifth dimension of seifert manifold in universe and other dimension

Universe audient for other dimension inclusive into fifth dimension, this phenomonen is universe being constant of value, and universe out of dimension exclusive space. All of value is constance of entropy. Universe entropy and other dimension entropy is dense of duality belong for. Prime equation construct with x,y parameter is each value of imaginary and reality of pole on a right comittment.

$$\int \nabla \psi^2 d\nabla \psi = \square \psi$$

$$\square \psi = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

$$\mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\delta \cdot \mathcal{O}(x) = [\nabla_i \nabla_j \int \nabla g(x) dx_{ij}]^{iy}$$

$$||ds^2|| = e^{-2\pi T|\phi|} [\mathcal{O}(x) + \delta \mathcal{O}(x)] dx^\mu dx^\nu + \lim_{n \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$\log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}, \frac{(y \log y)^{\frac{1}{2}}}{\log(x \log x)} \leq \frac{1}{2}$$

Universe is freeze out constant, and other dimension is expanding into fifth dimension. Possibility of quato metric, $\delta(x)$ = reality of value / exist of value ≤ 1 , expanding of universe = exist of value $\rightarrow \log(x \log x) = \square \psi$

freeze out of universe = reality of value $\rightarrow (y \log y)^{\frac{1}{2}} = \nabla \psi^2$

Around of universe is other dimension in fifth dimension of seifert manifold. what we see sky of earth is other dimension sight of from.

Rotate of dimension is transport of dimension in other dimension from universe system.

Other dimension is more than Light speed into takion quarks of structure.

Around universe is blackhole, this reason resolved with Gamma ray burst in earth flow. We don't able to see now other dimension, gamma ray burst is blackhole phenomanen. Special relativity is light speed not over limit. then light speed more than this speed. Other dimesion is contrary on universe from. If light speed come with other light speed together, stop of speed in light verisity. Then takion quarks of speed is on the contrary from become not see light element. And this takion quarks is into fifth dimension of being go from universe. Also this quarks around of universe attachment state. Moreover also this quarks is belong in vector of element. This reason is universe in other dimension emerge with. In this other dimension out of universe, but these dimension not shared of space.

$$l(x) = 2x^2 + qx + r$$

$$= (ax + b)(cx + d) + r, e^{l(x)} = \frac{d}{df} L(x), G_{\mu\nu} = g(x) \wedge f(x)$$

This equation is Weil's theorem, also quantum group. Complex space.

$$V(\tau) = \int \int \exp[L(x)] dx_m + O(N^{-1}), g(x) = L(x) \circ \int \int e^{2x^2+2x+r} dx_m$$

Lens of space and integrate of rout equation. And norm space of algebra line of curve equation in gradient metric.

$$||ds^2|| = ||\frac{d}{df}L(x)||, \eta = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\frac{1}{2}}$$

$$\bar{h} = [\nabla_i \nabla_j \int \nabla g(x) d_{ij}]^{iy}, \frac{d}{dt} g_{ij}(t) = -2R_{ij}$$

Two equation construct with Kaluze-Klein dimension of equation. Non-verticle space.

$$\frac{1}{\tau}(\frac{N}{2} + r(2\Delta F - |\nabla f|^2 + R) + f) \bmod N^{-1}$$

Also this too lense of space equation. And this also is heat equation. And this equation is singularity into space metric number of formula.

$$\frac{x^2}{a} \cos x + \frac{y^2}{b} \sin x = r^2$$

Curvature of equation.

$$S_m^2 = || \int \pi r^2 dr ||^2, V(\tau) = r^2 \times S_m^2, S_1^{mn} \otimes S_2^{mn} = \int [D^2 \psi \otimes h_{\mu\nu}] dm$$

Non-relativity of integrate rout equation.

$$||ds^2|| = e^{-2\pi T|\phi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d^2 \psi$$

$$V(x) = \int \frac{1}{\sqrt{2\tau q}} (\exp L(x) dx) + O(N^{-1})$$

$$V(x) = 2 \int \frac{(R + \nabla_i \nabla_j f)}{-(R + \Delta f)} e^{-f} dV, V(\tau) = \int \tau(p)^{-\frac{n}{2}} \exp(-\frac{1}{\sqrt{2\tau q}} L(x) dx) + O(N^{-1})$$

$$\frac{d}{df} F = m(x)$$

$$Zeta(x,h) = \exp \frac{(qf(x))^m}{m}$$

Singularity and duality of differential is complex element.

$$\left\| \begin{matrix} x & y & z \\ u & v & w \end{matrix} \right\|_{g_{\mu\nu}(x)}^2$$

$$= (f(x)dx^\mu dx^\nu, f'(y)dy^\mu dy^\nu, f''(z)dz^\mu dz^\nu) \cdot (u, v, w)$$

$$= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & i \end{pmatrix}$$

$$\cong \frac{g(x,y,z)}{f(a,b,c)} \cdot h^{-1}(u,v,w)$$

Antigravity is other dimension belong to, gravity of universe is a few weak power become. These power of balance concern with non-entropy metric. Universe is time and energy of gravity into non-entropy. Other dimension is first of universe emerged with takion quarks is light speed, and time entropy into future is more than light speed. Some future be able to earth of night sky can see to other dimension sight. However, this sight might be first of big-ban environment. Last of other dimension condition also can be see to earth of sight. Same future that universe of gravity and other dimension of antigravity is balanced into constant of value, then fifth dimension fill with these power of non-catastrophe. These phenomenon is concerned with zeta function emerged with laplace equation resulted.

$$\frac{d}{df} \int \int \int \square \psi d\psi_{xy} = V(\square \psi), \lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} V_k(\square \psi) = \frac{\partial}{\partial f} i\hbar c$$

This reason explained of quarks emerge into global space of system.

$$\lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} G_{\mu\nu} = f(x) \circ m(x)$$

This become universe.

$$\frac{V(x)}{f(x)} = m(x)$$

Eight differential structure integrate with one of geometry universe.

$$(a_k f^k)' = {}_n C_0 a_0 f^n + {}_n C_1 a_1 f^{n-1} \dots {}_n C_{r-1} a_{r-1} f^{n-1}$$

$$\int a_k f^k dx_k = \frac{a_{n+1}}{k+1} f^{n+1} + \frac{a_{k+2}}{k+2} f^{k+2} + \dots \frac{a_0}{k} f^k$$

Summate of manifold create with differential and integrate of structure.

$$\frac{\partial}{\partial f} \square \psi = \frac{1}{4} g_{ij}^2$$

$$\left(\frac{\nabla \psi^2}{\square \psi} \right)' = 0$$

$$\frac{(y \log y)^{\frac{1}{2}}}{\log(x \log x)} = \frac{\frac{1}{2}}{\frac{1}{2}i}$$

$$\frac{\{f, g\}}{[f, g]} = \frac{1}{i}, \left(\frac{\{f, g\}}{[f, g]} \right)' = i^2$$

$$(i)^2 \rightarrow \frac{1}{4} g_{ij}, F_t^m = \frac{1}{4} g_{ij}^2, f(r) = \frac{1}{4} |r|^2, 4f(r) = g_{ij}^2$$

$$\frac{1}{y} \cdot \frac{1}{y'} \cdot \frac{y''}{y'} \cdot \frac{y'''}{y''} \dots$$

$$= \frac{{}_n C_r y^2 \cdot y^3 \dots}{{}_n C_r y^1 y^2 \dots}$$

$$\frac{\partial y}{\partial x} \cdot \frac{\partial}{\partial y} f(y) = y' \cdot f'(y)$$

$$\int l \times l dm = (l \oplus l)_m$$

Symmetry theoerm is included with two dimension in plank scale of constance.

$$\begin{aligned} &= \frac{d}{dx^\mu} \cdot \frac{d}{dx^\nu} f^{\mu\nu} \cdot \nabla \psi^2 \\ &= \square \psi \end{aligned}$$

$$\frac{\nabla \psi^2}{\square \psi} = \frac{1}{2}, l = 2\pi r, V = \frac{4}{\pi r^3}$$

$$S \frac{4\pi r^3}{2\pi r} = 2 \cdot (\pi r^2)$$

$$= \pi r^2, H_3 = 2, \pi(H_3) = 0$$

$$\frac{\partial}{\partial f} \square \psi = \frac{1}{4} g_{ij}^2$$

This equation is blackhole on two dimension of surface, this surface emerge into space of power.

$$f(r) = \frac{1}{2} \frac{\sqrt{1 + f'(r)}}{f(r)} + mgf(r)$$

Atom of verticle in electric weak power in lowest atom structure

$$ihc = G, hc = \frac{G}{i}$$

$$\left(\frac{\nabla \psi^2}{\square \psi} \right)' = 0$$

$$S_n^m = |S_2 S_1 - S_1 S_2|$$

$$\square \psi = V \cdot S_n^m$$

One surface on gravity scales.

$$G^{\mu\nu} \cong R^{\mu\nu}$$

Gravity of constance equals in Rich curvature scale

$$\nabla_i \nabla_j (\square \psi) d\psi_{xy} = \frac{\partial}{\partial f} \square \psi \cdot T^{\mu\nu}$$

Partial differential function construct with duality of differential equation, and integrate of rout equation is create with Maxwell field on space of metrix.

$$\begin{aligned} &= \frac{8\pi G}{c^4} (T^{\mu\nu})^2, (\chi \oplus \pi) = \int \int \int \square \psi d^3 \psi \\ &= \text{div}(\text{rot} E, E_1) \cdot e^{-ix \log x} \end{aligned}$$

World line is constructed with Von Neumann manifold stimulate into Thurston conjugate theorem, this explained mechanism for summuate of manifold. Also this field is norm space.

$$\sigma(\mathcal{H}_n \otimes \mathcal{K}_m) = E_n \times H_m$$

$$\mathcal{H}_n = [\nabla_i \nabla_j \int \nabla g(x) dx_i dx_j]^{iy}, \mathcal{K}_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{1/2}$$

$$||ds^2|| = |\sigma(\mathcal{H}_n \times \mathcal{K}_m), \sigma(\chi, x) \times \pi(\chi, x) = \frac{\partial}{\partial f} L(x), V'_\tau(x) = \frac{\partial}{\partial V} L(x)$$

Network theorem. Dalanvercian operator of differential surface. World line of surface on global differential manifold. This power of world line on fourth of universe into one geometry. Gradient of power on universe world line of surface of power in metric. Metric of integrate complex structure of component on world line of surface.

$$(\square\psi)' = \frac{\partial}{\partial f_M} (\int \int \int f(x,y,z) dx dy dz)' d\psi$$

$$\frac{\partial}{\partial V} L(x) = V(\tau) \int \int e^{\int x \log x dx + O(N^{-1})} d\psi$$

$$\frac{d}{df} \sum_{k=1}^n \sum_{k=0}^{\infty} a_k f^k = \frac{d}{df} m(x), \frac{V(x)}{f(x)} = m(x)$$

$$4V'_\tau(x) = g_{ij}^2, \frac{d}{dl} L(x) = \sigma(\chi, x) \times V_\tau(x)$$

Fifth manifold construct with differential operator in two dimension of surface.

$$||ds^2|| = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\psi^2$$

$$f^{(2)}(x) = [\nabla_i \nabla_j \int \nabla f^{(5)} d\eta]^{1/2}$$

$$=[f^{(2)}(x)d\eta]^{1/2}$$

$$\nabla_i \nabla_j \int F(x) d\eta = \frac{\partial}{\partial f} F$$

$$\nabla f = \frac{d}{dx} f$$

$$\nabla_i \nabla_j \int \nabla f d\eta = \frac{\partial}{\partial x_i} \frac{\partial}{\partial x_j} (\frac{d}{dx} f)$$

$$\frac{z_3 z_2 - z_2 z_3}{z_2 z_1 - z_1 z_2} = \omega$$

$$\frac{\bar{z}_3 z_2 - \bar{z}_2 z_3}{\bar{z}_2 z_1 - \bar{z}_1 z_2} = \bar{\omega}$$

$$\omega \cdot \bar{\omega} = 0, z_n = \omega - \{x\}, z_n \cdot \bar{z}_n = 0, \vec{z}_n \cdot \vec{\bar{z}}_n = 0$$

Dimension of symmetry create with a right comittment into middle space, triangle of mesh emerged into symmetry space. Imaginary and reality pole of network system.

$$\begin{aligned}[f, g] \times [g, f] &= fg + gf \\ &= \{f, g\}\end{aligned}$$

Mebius space create with fermion quarks, Seifert manifold construct with this space restruct of pieces.

$$\begin{aligned}V(\tau) &= \int \int e^{f \cdot x \log x + O(N^{-1})} d\psi, V'_\tau(x) = \frac{\partial}{\partial f_M} \left(\int \int \int f(x, y, z) dx dy dz \right)' d\psi \\ (\square\psi)' &= 4\vec{v}(x), \frac{\partial}{\partial V} L(x) = m(x), V(\tau) = \int \frac{1}{\sqrt{2\tau q}} \exp[L(x)] d\psi + O(N^{-1}) \\ V(\tau) &= \int \int \int \frac{V}{S^2} dm, f(r) = \frac{1}{2} \frac{\sqrt{1+f'(r)}}{f(r)} + mgf(r), \log(x \log x) \geq 2(y \log y)^{\frac{1}{2}}, F_t^m = \frac{1}{4} g_{ij}^2, \frac{d}{dt} g_{ij}(t) = -2R_{ij} \\ \nabla_i \nabla_j v &= \frac{1}{2} m v^2 + m c^2, \int \nabla_i \nabla_j v dv = \frac{\partial}{\partial f} L(x) \\ (\square\psi)^2 &= -2 \int \nabla_i \nabla_j v d^2 v, (\square\psi)^2 = \left(\frac{\nabla\psi^2}{\square\psi} \right)' \\ &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dm, \bigoplus \nabla M_3^+ = \int \frac{\vee(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)} dV \\ &= (x, y, z) \cdot (u, v, w) / \Gamma \\ \bigoplus C_-^+ &= \int \exp\left[\int \nabla_i \nabla_j f d\eta\right] d\psi \\ &= L(x) \cdot \frac{\partial}{\partial l} F(x) \\ &= (\square\psi)^2 \\ \nabla\psi^2 &= 8\pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)\end{aligned}$$

Blackhole and whitehole is value of variable into constance lastly. Universe and the other dimension built with relativity energy and potential energy constructed with. Euler-Langrange equation and two dimension of surface in power of vector equation.

$$\begin{aligned}l &= \sqrt{\frac{\hbar G}{c^3}}, T^{\mu\nu} = \frac{1}{2} k l^2 + \frac{1}{2} m v^2, E = m c^2 - \frac{1}{2} m v^2 \\ e^{x \log x} &= x^x, x = \frac{\log x^x}{\log x}, y = x, x = e\end{aligned}$$

A right comittment on symmetry surface in zeta function and quantum group theorem.

$$\int \frac{1}{(x \log x)} dx = i \int x \log x dx + \int \frac{1}{(x \log x)} dx$$

$$\int\int\frac{1}{(x\log x)^2}dx_m=i\frac{1}{2}x^2$$

$$\int\int\frac{1}{(x\log x)^2}dx_m=i\int\int_Mdx_m$$

$$\leq \frac{1}{2}i+x^2$$

$$E=-\frac{1}{2}mv^2+mc^2$$

$$\lim_{x\rightarrow\infty}\int\int\frac{1}{(x\log x)^2}dx_m\geq\frac{1}{2}i$$

$$\frac{d}{df}\int\int\frac{1}{(x\log x)^2}dx_m=\frac{1}{2}i$$

$$\lim_{x\rightarrow\infty}\frac{x^2}{e^{x\log x}}=0$$

$$\int dx \rightarrow \partial f \rightarrow dx \rightarrow cons$$

$$\text{Heat equation restruct with entropy from low energy to high energy flow to destroy element.}$$

$$(\Box \psi)'=(\exists \int \vee (R+\nabla_I \nabla_j f) e^{-f} dV)'d\psi$$

$$\bigoplus M_3^+=\frac{\partial}{\partial f}L(x)$$

$$\log(x\log x)\geq 2(y\log y)^{\frac{1}{2}}$$

$$\frac{\partial}{\partial l}L(x)=\nabla_i\nabla_j\int\nabla f(x)d\eta, L(x)=\frac{V(x)}{f(x)}$$

$$l(x)=L'(x),\frac{d}{df}F=m(x),V'(\tau)=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

$$\text{Weil's theorem.}$$

$$T^{\mu\nu}=\int\int\int\frac{V(x)}{S^2}dm, S^2=\pi r^2\cdot S(x)$$

$$=\frac{4\pi r^3}{\tau(x)}$$

$$\eta=\nabla_i\nabla_j\int\nabla f(x)d\eta,\bar{h}=\nabla_i\nabla_j\int\nabla g(x)dx_idx_j$$

$$\delta \mathcal{O}(x) = [\nabla_i \nabla_j \int f(x) d\eta]^{\frac{1}{2} + iy}$$

$$Z(x,h)=\lim_{x\rightarrow\infty}\sum_{k=0}^\infty\frac{qT^m}{m}=\delta(x)$$

$$l(x)=2x^2+px+q, m(x)=\lim_{x\rightarrow\infty}\sum_{k=0}^{\infty}\frac{(qx^m)'}{f(x)}$$

Higgs fields in Weil’s theorem component and Singularity theorem.

$$Z(T,X)=\exp\sum_{m=1}^{\infty}\frac{q^kT^m}{m}, Z(x,h)=\exp\frac{(qf(x))^m}{m}$$

Zeta function.

$$\frac{d}{df}F=m(x), F=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi$$

Integral of rout equation.

$$\lim_{x\rightarrow 1}\text{mesh}\frac{m}{m+1}=0, \int x^m=\frac{x^m}{m+1}$$

Mesh create with Higgs fields. This fields build to native function.

$$\frac{d}{df}\int x^m=mx^m, \frac{d}{dt}g_{ij}(t)=-2R_{ij}, \lim_{x\rightarrow 1}\text{mesh}(x)=\lim_{m\rightarrow \infty}\frac{m}{m+1}$$

$$\lim_{x\rightarrow 1}\sum_{k=0}^{\infty}a_kf^k=\alpha$$

$$||ds^2||=e^{-2\pi T|\psi|}[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu+T^2d^2\psi$$

$$\frac{\partial}{\partial V}||ds^2||=T^{\mu\nu}, V(\tau)=\int e^{x\log x}d\psi=l(x)$$

$$R_{ij}=\frac{\partial}{\partial x_i}\frac{\partial}{\partial x_j}T^{\mu\nu}, G^{\mu\nu}=R^{\mu\nu}T^{\mu\nu}$$

$$F(x)=\int\int e^{\int x\log xdx+O(N^{-1})}d\psi, \frac{d}{dV}F(x)=V'(x)$$

Integral of rout equation oneselves and expired from string theorem of partial function.

$$T^{\mu\nu}=R^{\mu\nu}, T^{\mu\nu}=\int\int\int\frac{V}{S^2}dm$$

$$\delta\mathcal{O}(x)=[D^2\psi\otimes h_{\mu\nu}]dm$$

Open set group construct with D-brane.

$$\nabla(\Box\psi)'=[\nabla_i\nabla_j\int\nabla f(x)d\eta]^{\frac{1}{2}+iy}$$

$$(f(x),g(x))'=(A^{\mu\nu})'$$

$$\begin{pmatrix} dx & \delta(x) \\ \epsilon(x) & \partial x \end{pmatrix} \cdot (f(x,y),g(x,y))$$

$$=\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Differential equation of complex variable developed from imaginary and reality constance.

$$\delta(x) \cdot \mathcal{O}(x) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}}$$

$$V_{\tau}'(x) = \frac{\partial}{\partial f_M} (\int \int \int f(x, y, z) dx dy dz) d\psi$$

Gravity equation oneselves built with partial operator.

$$\frac{\partial}{\partial V} L(x) = V_{\tau}'(x)$$

Global differential equation is oneselves component.

Quantum Computer in a certain theorem

Masaaki Yamaguchi

A pattern emerge with one condition to being assembled of emelite with all of possibility equation, this assembled with summative of manifold being elemetiled of pieace equation. This equation relate with Euler equation. And also this equation is Euler constant oneselves.

$$(E_2 \bigoplus E_2) \cdot (R^- \subset C^+) = \bigoplus \nabla C_-^+$$

Zeta function radius with field of mechanism for atom of pole into strong condition of balance, this condition is related with quarks of level controlled for compute with quantum tonnel effective mechanism. Quantum mechanism composed with vector of constance for zeta function and quantum group. Thurston conjugate theorem explain to emerge with being controll of quantum levels of quarks. Locality theorem also occupy with atom of levels in zeta function.

$$\begin{aligned} &= \bigoplus \nabla C_-^+ \\ \vee \int \frac{C^+ \nabla M_m}{\Delta(M_-^+ \nabla C_-^+)} &= \exists(M_-^+ \nabla R^+) \\ \exists(M_-^+ \nabla C^+) &= \text{XOR}(\bigoplus \nabla M_-^+ \\ -[E^+ \nabla R^+] &= \nabla_+ \nabla_- C_-^+ \\ \int dx, \partial x, \nabla_i \nabla_j, \Delta x &\rightarrow E^+ \nabla M_1, E^+ \cap R \in M_1, R \nabla C^+ \end{aligned}$$

Zeta function also compose with Rich flow equation cohomological result to equal with locality equaitons.

$$\begin{aligned} \vee(R + \nabla_i \nabla_j f)^n &= \int \frac{\wedge(R + \nabla_i \nabla_j f)^2}{\exists(R + \Delta f)^n} \\ \wedge(R + \nabla_i \nabla_j f)^x &= \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\ \frac{d}{dt} g_{ij}(x) &= -2R_{ij} \\ \vee \int \wedge(R + \nabla_i \nabla_j f)^x &= \frac{\wedge(R + \nabla_i \nabla_j f)^n}{\exists(R + \nabla_i \nabla_j f \circ g)^n} \\ x + y &\geq 2\sqrt{xy}, x(x) + y(x) \geq x(x)y(x) \\ x^y &= (\cos \theta + i \sin \theta)^n \\ x^y &= \frac{1}{y^x} \end{aligned}$$

Therefore zeta function is also constructed with quantum equation too.

UFO mechanism

Masaaki Yamaguchi

UFO mechanism are several system of circumstance that accesority and verisity composite with mass and circument of gravity is key of mechanism included with anti-gravity in imaginary pole emerge with rotate of right formula. This power of rotate emerge with anti-gravity which of power emelite for inner into oneselves create with like of lorentz of energy. This lorentz of energy is use with steles flight of mechanism. However this lorentz of energy only mass of gravity low included with oneselves but also anti-gravity is oneselves of component with inner of rotate with energy, this energy is not free of condition of power, because this power not ordinary of mechanism.

$$\begin{aligned}
 F - N &= mg - N' \\
 \frac{d}{df} F &= \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
 C &= \int \int \frac{1}{(x \log x)^2} dx_m \\
 \int \int \frac{1}{(x \log x)^2} dx_m &= \frac{1}{2}i
 \end{aligned}$$

This imaginary number is rotate of quantum spin emerged with power, this power is not mass of gravity element, also this power is inner of energy and anti-gravity of power.

Moreover this right of rotate of imaginary pole is three manifold of nourth and sournth of formula. And nourth of formula look with gravity but sournth of formula is anti-gravity, this mechanism is relativity theorem from explained.

Hortshorne conjecture is built with fermer theorem spacificed from AdS5 manifold constructed of sphere cube exclude with time machine mechanism element. This dimension cercuited with rout of pole established by right angle in systematic vintage emerge with artificial intelligence. This one element of independence create with imaginary number from inner vector element, this resulted rentanse for rolenzt atructer in super string theory refill into globality topology mechanism constructed with these essence. This dimension target with native function around covered with abel manifold, seifert manifold is become with other dimension on pair universe essensed with zeta function, this space time is concluded with finite dimension in around of shell, this cohomology of mapping construct with abel manifold is all of equal between blackhold and zeta function inverse with low structure of constance in quantum operator relative with univese and planck scale resulted with mechanism. This dimension equal with whitehole and this pair of dimension is blackhole from being emerged with exclusive and inclusive of component from universe and other dimension system.

This explained mechanism from pholographic theorem is atom of low structure of constance with time space and differential structure relative from UFO machine system concerned with Hortshorne conjecture. This pholographic theorem is explained with UFO machine resulted by mass and circument of gravity not existed zone. Inner of rotate with energy is component oneselves with being not free of condition element. More spectrum explained this power is inner of energy and anti-gravity of power. Accesority

and verisity have with mass and circument of gravity which existed is out of zone element. However this element is not oneselves of inner and circument of gravity is not only spective of inner influence of power but also out of influent power. However anti-gravity is all of inner spective oneselves emerged with energy of power.

Hortshorne conjecture and AdS5 manifold moreover pholographic theorem are concerned with UFO machine system be able to be explained from Atom of inspected with movement spector phenomone artificial experient. This experient result with UFO machine is not free condition of zone success with inner of power being with elemelite of anti-gravity.

From under circument position to right of rotate energy formula create with anti-gravity, from up circument position to left of rotate energy formula, these way of relativity system is resulted with other dimension of rotate energy reason. Sourth of rotate way is anti-gravity, nourth of rotate way is circument of gravity. This mechanism construct with non-symmetry of entropy resulted. And this theorem explain that parity of broken result. This parity of mechanism is other dimension and universe of seifert manifold of balance of entropy concluded. Symmetry formula systemalite with non-vacuum constructed with two-projection space not be existed in three manifold dimension. This mechanism is pair of quarks with universe and other dimension. If universe belong for being emerged with energy of quarks in three manifold, then other dimension migrated with entropy saved of energy of universe balance. If dimension create with quarks of LHC experiment by exist proved, imaginary pole of rotate create with sourth of formula, universe and other dimension is balanced with energy, then reverse of power of gravity and anti-gravity are proof that shirt of quarks lives with oneselves.

仲人としての Zeta 関数が 広中平祐教授とグリーシャ教授の
 抽象理論と具象理論としての
 ヒルベルト空間とサーストン多様体
 複素多様体が統一場理論を量子力学における
 ガウスの曲面論として
 成し遂げる理論の背景場となるレポート

Masaaki Yamaguchi

一般相対性理論における重力理論は大域的微分多様体と積分多様体についての 単体量を求めるための空間の対数関数における不変性を記述するプランクスケール と異次元の宇宙におけるスケール、ウィークスケールと言われている anti-D-brane として、この2種類の計量をガンマ関数とベータ関数としてオイラーの定数 とそれによる連分数が微分幾何の量子化と数式の値を求めると同じという 予知と推測値から、広中平祐定理の4重帰納法のオイラーの公式による 多様体積分と同類値として、ペレルマン多様体がサーストン空間に成り立つ と同じく、広中平祐定理がヒルベルト空間で成り立つとしての、 これら2種の多様体の場の理論が、ゼータ関数が AdS5 多様体で成り立つことと、 不変量として、ゼータ関数がこの3種の多様体の場の理論のバランスをとる理論として言えることである。 Gravity of general relativity theory describe with Cutting of space in being discatastrophed from Global differential and integral manifold of scaled levativity in plank scal and the other vector of universe scale, these two scale inspectivity of Gamma and Beta function escourted with these manifold experted in result of Differentail geometry of quantum level manifold equal with Euler product and continue parameter, moreover this product is relativity of Hironaka theorem in Four assembled integral of Euler equation.

重力場理論の式は、Gravity equation is

$$\square = -\frac{16\pi G}{c^4} T^{\mu\nu}$$

This equation quated with being logment of formula, and this formula divided with universe of number in prime zone, therefore, this dicided with varint equation is monotonicity of being composited with Weil's theorem united for Gamma function. この式は、対数を宇宙における数により求める素数分布論として、この大域的積分分断多様体がガンマ関数をヴェイユ予想を根幹とする単体量として決まることに起因する商代数として導かれる。

$$\not\square = \frac{8\pi G}{c^4} T^{\mu\nu} / \log x$$

$$\begin{aligned} & {}^t\prod\prod\prod\mathrm{cohom}D_\chi[I_m] \\ &= \oint (px^n + qx + r)^{\nabla^l} \end{aligned}$$

$$\frac{d}{dt}L(x,y)=2\int ||\sin 2x||^2d\tau$$

$$\frac{d}{d\gamma}\Gamma$$

この関数は大域的微分多様体としてのアカシックレコードの合流地点として、タプルスペースを形成している。This function estermenate with acasic record of global differential manifolds.

$$= [i\pi(\chi, x), f(x)]$$

それにより、この多様体は基本群をアカシックレコードの相対性としての存在論の実存主義から統合される多様体自身としてのタプルスペースの池になっている。And this manifold from fundemental group estermenate with also this manifold estimate relativity of acasic record.

$$\frac{d}{d\gamma}\Gamma = \int C dx_m = \int (\int \frac{1}{x^s} dx - \log x) d\text{vol}$$

また、このアカシックレコードはオイラーの定数のラムダドライバーにもなっている。More also this record tupled with lake of Euler product.

$$\bigoplus (i\hbar^\nabla)^{\oplus L}$$

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

それゆえに、この定数はゼータ関数と微分幾何の量子化を因数にもつ素因子分解の式にもなっている。Therefore, this product also constructed with differential geometry of quantum level equation and zeta function.

$$= C$$

そして、この関数はオイラーの定数から広中平祐定理による4重帰納法のオイラーの公式からの多様体積分へのサーストン空間のスペクトラム関数ともなっている。And this function of Euler product respectrum of focus with Heisuke Hironaka manifold in four assembled of integral Euler equation..

$$C = b_0 + \frac{c_1}{b_1 + \frac{c_2}{b_2 + \frac{c_3}{b_3 + \frac{c_4}{b_4 + \dots}}}}$$

この方程式は指数による連分数としての役割も担っている。This equation demanded with continued number of step function.

$$x^{\frac{1}{2}+iy} = e^{x \log x}$$

$$(2.71828)^{2.828} = a^{a+b+c+d^d \dots}$$

$$= \int e^f \cdot x^{1-t} dx$$

This represent is Gamma function in Euler product. Therefore this product is zeta function of global differential equation.

$$\frac{d}{df} F(v_{ij}, h) = \int e^{-f} dV [-\Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij} - v_{ij} \nabla_i \nabla_j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla f^2) (\frac{v}{2} - h)]$$

$$= \frac{d}{df} F = \frac{2 \int (R + \nabla_i \nabla_j f)^2}{-(R + \Delta f)} dm$$

これらの方程式は 8 種類の微分幾何の次元多様体として、そして、これらの多様体は曲平面による双対性をも生成している。そして、このガウスの曲平面は、大域的微分多様体と微分幾何の量子化から素因数を形成している。These equation are eight differential geometry of dimension calyement. And these calyement equation excluded into pair of dimension surface. This surface of Gauss function are global differential manifold, and differential geometry of quantum level.

$$F \geq \frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

微分幾何の量子化はオイラーの定数とガンマ関数が指数による連分数としての不変性として素因数を形成していて、このガウスの曲平面による量子力学における重力場理論は、ダランベルシアン of the cut-off manifold がこの大域的切断多様体を付加している。Differential geometry of quantum level constructed with Euler product and Gamma function being discatastrophed from continued fraction style. Gravity equation lend with varint of monotonicity of level expressed from gravity of letter varient formula. これらの方程式は基本群と大域的微分多様体をエスコートしていてもいて、ヴェイユ予想がこのダランベルシアンの切断方程式たちから輸送のポートにもなっている。ベータ関数とガンマ関数がこれらのフォームラの方程式を放出してもいて、結果、これらの方程式は広中平祐定理の複素多様体とグリーシャ教授によるペレルマン多様体からサポートされている。この 2 名の教授は、一つは抽象理論をもう一方は具象理論を説明としている。These equation escorted into Global differential manifold and fundemental equation. Weil's theorem is imported from this equation in gravity of letters. Beta function and Gamma function are excluded with these formulas. These equation comontius from Heisuke Hironaga of complex manifold and Gresha professor of Perelman manifold. These two professors are one of abstract theorem and the other of visual manifold theorem.

34 Hilbert manifold in Mobius space

this element of Zeta function on integrate of fields

Hilbert manifold equals with Von Neumann manifold, and this fields is concluded with Glassman manifold, the manifold is duality of twister into created of surface built. These fields is used for relativity theorem and quantum physics.

$$||ds^2|| = \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu}$$

This norm is component of fields in Hilbert manifold of space theorem rebuilt with Mobius space in Gamma Function on Beta function fill into power of rout fundamental group. And this result with AdS_5 space time in Quantum caos of Minkowsky of manifold in abel manifold. Gravity of metric in non-commutative equation is Global differential equation conbult with Kaluza-Klein space. Therefore this mechanism is $T^{\mu\nu}$ tensor is equal with $R^{\mu\nu}$ tensor. And This moreover inspect with laplace operator in stimulate with sign of differential operator. Minus of zone in Add position of manifold is Volume of laplace equation rebuilt with Gamma function equal with summative of manifold in Global differential equation, this result with setminus of zone of add summative of manifold, and construct with locality theorem straight with fundamental group in world line of surface, this power is boson and fermion of cone in hyper function.

$$V(\tau)=[f(x),g(x)]\times [f^{-1}(x),h(x)]$$

$$\begin{aligned}\Gamma(p,q) &= \int e^{-x} x^{1-t} dx \\ &= \beta(p,q) \\ &= \pi(f(\chi,x),x)\end{aligned}$$

$$\begin{aligned}||ds^2|| &= \mathcal{O}(x)[(f(x)\circ g(x))^{\mu\nu}]dx^\mu dx^\nu \\ &= \lim_{x\rightarrow\infty}\sum_{k=0}^\infty a_k f^k\end{aligned}$$

$$\begin{aligned}G^{\mu\nu} &= \frac{\partial}{\partial f}\int [f(x)^{\mu\nu}\circ G(x)^{\mu\nu}dx^\mu dx^\nu]^\mu{}^\nu dm \\ &= g_{\mu\nu}(x)dx^\mu dx^\nu - f(x)^{\mu\nu}dx^\mu dx^\nu \\ [i\pi(\chi,x),f(x)] &= i\pi f(x) - f(x)\pi(\chi,x)\end{aligned}$$

$$\begin{aligned}T^{\mu\nu} &= (\lim_{x\rightarrow\infty}\sum_{k=0}^\infty\int\int[V(\tau)\circ S^{\mu\nu}(\chi,x)]dm)^{\mu\nu}dx^\mu dx^\nu \\ G^{\mu\nu} &= R^{\mu\nu}T^{\mu\nu}\end{aligned}$$

$$\left|\begin{matrix}D^m&dx\\dx&\sigma^m\end{matrix}\right|\left|\begin{matrix}\cos\theta&-\sin\theta\\ \sin\theta&\cos\theta\end{matrix}\right|\left|\begin{matrix}x\\y\end{matrix}\right|=\begin{pmatrix}1&0\\0&-1\end{pmatrix}^{\frac{1}{2}}$$

$$\sigma^m\left[\begin{matrix}\delta(x)&-1\\1&\epsilon(x)\end{matrix}\right]^{\frac{1}{2}}=\begin{pmatrix}i&0\\0&-i\end{pmatrix}$$

$$V(M)=\frac{\partial}{\partial f}({}^N\int [f\setminus M]^{\oplus N})^{\mu\nu}dx^\mu dx^\nu$$

$$V(M)=\pi(2\int\sin^2dx)\oplus\frac{d}{df}F^Mdx_m$$

$$\lim_{x\rightarrow\infty}\sum_{k=0}^\infty a_k f^k=\int (F(V)dx_m)^{\mu\nu}dx^\mu dx^\nu$$

$$\bigoplus_{k=0}^\infty [f\setminus g] = \vee(M\wedge N)$$

$$\begin{aligned}\pi_1(M) &= e^{-f^2\int\sin^2x dm} + O(N^{-1}) \\ &= [i\pi(\chi,x),f(x)]\end{aligned}$$

$$M\circ f(x)=e^{-f\int\sin x\cos x dx_m}+\log(O(N^{-1}))$$

Non-Symmetry space time.

$$\frac{d}{dL}V(\tau)=\frac{d}{df}\int\int_M\frac{1}{(x\log x)^2}dx_m+\frac{d}{df}\int\int_M\frac{1}{(y\log y)^{\frac{1}{2}}}dy_m$$

$$\epsilon S(\nu) = \square_v \cdot \frac{\partial}{\partial \chi} (\sqrt[5]{\wedge g^2}) d\chi$$

Differential Volume in AdS_5 graviton of fundamental rout of group.

$$\wedge (F_t^m)'' = \frac{1}{12} g_{ij}^2$$

Quarks of other dimension.

$$\pi(V_\tau) = e^{-(\sqrt{\frac{\pi}{16}} \log x)^\delta} \times \frac{1}{(x \log x)}$$

Universe of rout, Volume in expanding space time.

$$\begin{aligned} \frac{d}{dt} (g_{ij})^2 &= \frac{1}{24} (F_t^m)^2 \\ m^2 &= 2\pi T \left(\frac{26 - D_n}{24} \right) \end{aligned}$$

This quarks of mass in relativity theorem, and fourth of universe in three manifold of one dimension surface, and also this integrate of dimension in conbult of quarks.

$$g_{ij} \wedge \pi(\nu_\tau) = e^{-2\pi T|\psi|} [\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\psi^2$$

Out of rout in AdS_5 space time.

$$||ds^2|| = g_{ij} \wedge \pi(\nu_v)$$

AdS_5 norm is fourth of universe of power in three manifold out of rout.

Sphere orbital cube is on the right of hartshorne conjecture by the right equation, this integrate with fourth of piece on universe series, and this estimate with one of three manifold. Also this is blackhole on category of symmetry of particles. These equation conbult with planck volume and out of universe is phologram field, this field construct with electric field and magnitic field stand with weak electric theorem. This theorem is equals with abel manifold and AdS_5 space time. Moreover this field is anti-brane and brane emerge with gravity and antigravity equation restructed with. Zeta function is existed with Re pole of $\frac{1}{2}$ constance reluctances. This pole of constance is also existed with singularity of complex fields. Von Neumann manifold of field is also this means.

$$\begin{aligned} ||ds^2|| &= \lim_{x \rightarrow \infty} [\delta(x) \int \int \int \pi \left(\sum_{k=0}^{\infty} \frac{n \sqrt{p}, x}{n} \right)^{\frac{1}{2}} d\tau]^{\mu\nu} \\ e^{i\theta} &= \cos \theta + i \sin \theta \\ e^{x \log x} &= x^{\frac{1}{2} + iy}, x \log x = \log(\cos \theta + i \sin \theta) \\ &= \log \cos \theta + i \log \sin \theta \end{aligned}$$

This equation is developed with frobenius theorem activate with logment equation. This theorem used to

$$\begin{aligned} \log(\sin \theta + i \cos \theta) &= \log(\sin \theta - i \cos \theta) \\ \log \left(\frac{\sin \theta}{i \cos \theta} \right) &= -2R_{ij}, \frac{d}{dt} g_{ij}(t) = -2R_{ij} \end{aligned}$$

$$\mathcal{O}(x) = \frac{\zeta(s)}{\sum_{k=0}^{\infty} a_k f^k}$$

$$\text{Im}f = \ker f, \chi(x) = \frac{\ker f}{\text{Im}f}$$

$$H(3) = 2, \nabla H(x) = 2, \pi(x) = 0$$

This equation is world line, and this equation is non-integrate with relativity theorem of rout constance.

$$[f(x)] = \infty, ||ds^2|| = \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi$$

$$T^2 d^2\psi = [f(x)], T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

These equation is equals with M theorem. For this formula with zeta function into one of universe in four dimensions. Sphere orbital cube is on the surface into AdS_5 space time, and this space time is Re pole and Im pole of constance reluctances.

Gauss function is equals with Abel manifold and Seifert manifold. Moreover this function is infinite time, and this energy is cover with finite of Abel manifold and Other dimension of seifert manifold on the surface.

Dilaton and fifth dimension of seifert manifold emerge with quarks and Maxwell equation, this power is from dimension to flow into energy of boson. This boson equals with Abel manifold and AdS_5 space time, and this space is created with Gauss function.

Relativity theorem is composited with infinite on D-brane and finite on anti-D-brane, This space time restructed with element of zeta function. Between finite and infinite of dimension belong to space time system, estrald of space element have with infinite oneselves. Anti-D-brane have infinite themselves, and this comontend with fifth dimension of AdS_5 have for seifert manifold, this asperal manifold is non-move element of anti-D-brane and move element of D-brane satisfid with fifth dimension of seifert algebra liner. Gauss function is own of this element in infinite element. And also this function is abel manifold. Infinite is coverd with finite dimension in fifth dimension of AdS_5 space time. Relativity theorem is this system of circustance nature equation. AdS_5 space time is out of time system and this system is belong to infinite mercy. This hiercyent is endrol of memolite with geniue of element. Genie have live of telomea endore in gravity accesorlity result. AdS_5 space time out of over this element begin with infinite assentance. Every element acknowlege is imaginary equation before mass and spiritual envy.

$$T^2 d^2\psi = \lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k$$

$$\lim_{x \rightarrow 1} \sum_{k=0}^{\infty} a_k f^k = [T^2 d^2\psi]$$

$$\frac{d}{dL} V(\tau) = \frac{d}{df} \int \int_M ({}^5\sqrt{x^2}) d\Lambda + \frac{d}{df} \int \int_M {}^N({}^3\sqrt{x})^{\oplus N} d\Lambda$$

$$\begin{aligned}
M(\vee(\wedge f \circ g)^N)^{\frac{1}{2}} &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
||ds^2|| &= \mathcal{O}(x)[\eta_{\mu\nu} + \bar{h}_{\mu\nu}(x)]dx^\mu dx^\nu + T^2 d^2\psi \\
\mathcal{O}(x) &= e^{-2\pi T|\psi|} \\
G^{\mu\nu} &= R_{\mu\nu} T^{\mu\nu} \\
&= -\frac{1}{2} \Lambda g_{ij}(x) + T^{\mu\nu}
\end{aligned}$$

Fifth dimension of eight differential structure is integrate with one geometry element, Four of universe also integrate to one universe, and this universe represented with symmetry formula. Fifth universe also is represented with seifert manifold peices. All after universe is constructed with six element of circuatance. This aquire maniculate with quarks of being espered belong to.

35 Imaginary equation in AdS5 space time create with dimension of symmetry

D-brane and anti-D-brane is composited with all of series universe emerged for one geometry of dimension, this gravity of power from D-brane and anti-D-brane emelite with ancestor. Seifert manifold is on the ground of blackhole in whitehole of power pond of senseivility. Six of element of quarks and universe of pieces is supersymmetry of mechanism resolved with

hyper symmetry of quarks constructed to emerge with darkmatter. This darkmatter emerged with big-ban of heircyent in circumstance of phenomena.

D-brane and anti-D-brane equations is

$$\begin{aligned}
\frac{d}{dL} V(\tau) &= \frac{d}{df} \int \int_M \frac{1}{(x \log x)^2} dx_m + \frac{d}{df} \int \int_M \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m \\
C &= \int \int \frac{1}{(x \log x)^2} dx_m
\end{aligned}$$

Euler constance is quantum group theorem rebuild with projective space involved with.

虚数方程式は、反重力に起因するフーリエ級数の励起を生成する。それは、人工知能を生み出す、5次元時空にも、この虚数方程式は使われる。AdS5の次元空間は、反ド・シッター時空のD-braneとanti-D-braneのcomformal場を生み出す。ホログラフィー時空は、この量子起因によるものである。2次元曲面によるブラックホールは、ガンマ線バーストによる5次元時空の構造から観測される。空間の最小単位によるプランク定数は、宇宙の大域的微分多様体から導き出される、AdS5の次元空間の準同型写像を形成している。これは、最小単位から宇宙の大きさを導いている。最大最小の方程式は、相加相乗平均を形成している。時間と空間は、宇宙が生成したときから、宇宙の始まりと終わりを既に生み出している。宇宙と異次元から、ブラックホールとホワイトホールの力がわかり、反重力を見つけられる。オイラーの定数は、この量子定数からわかる。虚数の仕組みはこの量子スピンの産物である。オイラーの定数は、この虚時空の斥力の現存である。それは、非対称性理論から導かれる。不確定性原理は、AdS5のブラックホールとホワイトホールを閉3次元多様体に統合する5次元時空から求められる。位置と運動エネルギーが、空間の最小単位であるプランク定数を宇宙全体にする微細構造定数からわかり、面積確定から、アーベル多様体を母関数に極限值として、ゼータ関

数をこの母関数に不変式として、2種類ずつにまとめる4種類の宇宙を形成する8種類のサーストンの幾何化予想から導き出される。この閉3次元多様体は、ミラー対称性を軸として、6種類の次元空間を一種類の異次元宇宙と同質ともしている。複素多様体による特異点解消理論は、この原理から求められる。この特異点解消理論は、2次元曲面を3次元多様体に展開していく、時空から生成される重力の密度を反重力と等しくしていく時間空間の4次元多様体と虚時空から求められる。ヒルベルト空間は、フォン・ノイマン多様体とグラスマン多様体をこのサーストンの幾何化予想を場の理論既定値として形成される。この空間は、ミンコフスキー時空とアーベル多様体全体を表している。そして、この空間は、球対称性を複素多様体を起点として、大域的トポロジーから、偏微分を作用素微分として時空間をカオスからずらすと5次元多様体として成立している。これらより、3次元多様体に2次元射影空間が異次元空間として、AdS5空間を形成される。偏微分、全微分、線形微分、常微分、多重微分、部分積分、置換微分、大域的微分、単体分割、双対分割、同調、ホモロジー単体、コホモロジー単体、群論、基本群、複体、マイヤー・ビートリス完全系列、ファン・カンペンの定理、層の理論、コホモロティズム単体、CW複体、ハウスドロフ空間、線形空間、位相空間、微分幾何構造、モースの定理、カタストロフの空間、ゼータ関数系列、球対称性理論、スピン幾何、ツイスター理論、双対被覆、多重連結空間、プランク定数、フォン・ノイマン多様体、グラスマン多様体、ヒルベルト空間、一般相対性理論、反ド・シッター時空、ラムダ項、D-brane, anti-D-brane, コンフォーマル場、ホログラム空間、ストリング理論、収率による商代数、ニュートンポテンシャルエネルギー、剛体力学、統計力学、熱力学、量子スピン、半導体、超伝導、ホイーストン・ブリッチ回路、非可換確率論、Connes理論、これら、演算子代数を形成している、微分・積分作用素が、ヒルベルト空間に存在している多様体の特質を全面に押し出して、いろいろな多様体と関数そして、群論を形作っている。

$$\begin{aligned} \begin{vmatrix} D^m & dx \\ dx & \sigma^m \end{vmatrix} \begin{vmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{vmatrix} \begin{vmatrix} x \\ y \end{vmatrix} &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}^{\frac{1}{2}} \\ \sigma^m \begin{bmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{bmatrix}^{\frac{1}{2}} &= \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \\ (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) & \end{aligned}$$

$$\begin{aligned} \begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix} \beta(x, \theta) \begin{pmatrix} x \\ y \end{pmatrix} \\ = \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix} \\ l = \sqrt{\frac{\hbar G}{c^3}}, \sigma^m \cdot \begin{pmatrix} \delta(x) & -1 \\ 1 & \epsilon(x) \end{pmatrix}^{\frac{1}{2}} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \end{aligned}$$

String theorem also imaginary equation of pole with the fields of manifold.

$$\begin{aligned} \left(\frac{\partial}{\partial \tau} f(x, y, z) \right)^{3'} &= A^{\mu\nu} \\ \frac{d}{dt} g_{ij}(t) &= -2R_{ij} \end{aligned}$$

Rich equation is all of top formula in integrate theorem.

$$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta)$$

Dalanverle equation equals with abel manifold.

Artificial Intelligence and TupleSpace of ultranetwork

Masaaki Yamaguchi

クラウドにデータベースを構築しておいて、この構築した多様体を数式で表現したコード通りのデータが、この多様体において、作用素関数として実行されるとする。この多様体を実現したデータが表現されている環境自体を表せられるソースとして、TupleSpace が辞書を書き換えることができないことを利便して、どのデータも上書きされないことによって、前後の記憶が無駄なコードが作用されないことを表現できる。

作用素環プログラミングとして、半静性型宣言子をつくる。この宣言子は、スクリプト型プログラミング言語では、この型を作り上げた時点で、その宣言した環境としての多様体がデータベースの仕様として、宣言した以後のソースコードがこのコード自体の性質を反映させることが多様体を表現した後の、配列、ハッシュ、文字列、ポインタ、ファイル構造体、オブジェクト、数値、関数、正規表現、行列、統計、微分、積分、この微分、積分は関数とは別の文字列と数値処理として、行列と統計をこの表現としての多様体として、微分、積分を数列を応用とした極限值としてソースコードをコンピュータにおいて、実行、表現、存在できるコンピュータ上だけにとどまらないプログラミング言語として、調べられる。この作用素としての半静性型宣言子は、スクリプト言語において、重要な研究として、動的と静的な宣言子として、なぜ静的宣言子が動的スクリプトで必要とされているかが、Stream と Ruby を学んでいく段階で浮かび上がった課題として、私は Ruby をオブジェクト指向を学んだ結果が、この作用素環プログラミングをプログラム思考でコンピュータに人工知能を生成出来て、人体の量子コンピュータを模擬出来て、その上に、FPGA までも実行できるアスペクト型人工知能スクリプト言語が、この多様体を数式を文字列としてだけではなく、電気信号としての表現体としてコンピュータ上に実現できることを研究課題として、生まれている。

Omega::DATABASE を tuplespace としてスクリプトに書き上げているソースをデータベースの下地とする。これをコンピュータに多様体として表現、実行、流れとして、動的に実行する。この実行した後に、スクリプト言語の動作を停止した場合は、ガベージコレクションとして破棄されるとする。この動作している状態のときに、同時に実行される関数、オブジェクト、文字出力は、このときに同時に起動している多様体の性質をウェブのネットワーク上で多様体の記述されている規則、ルールに則ってプログラミング言語でコンピュータに作用させている、最終的な産物のゼータ関数としてのガンマ関数の大域的微分多様体を熱エントロピー値として、この熱値の性質として分類、整列される TupleSpace 上の関数の群論として、なにがコンピュータ上だけでなく、存在論だけにとどまらない電気信号かが、数学と情報科学で研究されるべきと、この多様体を調べる必要があると、目下の課題になっている。

現実の世界として、この世界を架空化する空間が同型としてのフェルミオンとボソンが、この空想上での入れ物に電気信号としての文字列がバーチャルネットワークに出力されて、この出力される文字列と電気信号が架空の性質として、物体や生命に現実の世界としての相対的な実存を特徴、成分、性質、分類としてコンピュータに文字列として命を吹き込む機能をプログラミング言語で生成されたバーチャルコードによって生み出せる可能性を秘めている。

```
Omega::DATABASE[tuplespace]
{
  Z \supset C \bigoplus \nabla R^{\{+}}, \nabla(R^{\{+}}
  \cap E^{\{+}) \ni x, \Delta(C \subset R) \ni x
```

```

M^{+}_{-}\bigoplus R^{+}, E^{+} \in
\bigoplus \nabla R^{+}, S^{+}_{-} \subset R^{+}_{2},
V^{+}_{-} \times R^{+}_{-} \cong \{V \over S\}
C^{+} \cup V^{+}_{-} \ni M_{1}\bigoplus \nabla C^{+}_{-},
Q \supseteq R^{+}_{-},
Q \subset \bigoplus M^{+}_{-},
\bigotimes Q \subset \zeta(x), \bigoplus \nabla C^{+}_{-} \cong M_3
R \subset M_3,
C^{+} \bigoplus M_n, E^{+} \cap R^{+},
E_2 \bigoplus E_1, R^{-} \subset C^{+}, M^{+}_{-}
C^{+}_{-}, M^{+}_{-}\nabla C^{+}_{-}, C^{+}\nabla H_m,
E^{+} \nabla R^{+}_{-}, E_2 \nabla E_1,
R^{-} \nabla C^{+}_{-}

[- \Delta v + \nabla_i \nabla_j v_{ij} - R_{ij} v_{ij}
- v_{ij} \nabla_i \nabla_j + 2 < \nabla f, \nabla h>
+ (R + \nabla f^2)(\{v \over 2\} - h)]

S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1,
H^1 \times S^1, H^1, S^2 \times E
}

```

クラウドにおけるデータベースを多様体が機能する仕組みからデータの相互関係と各データの処理対応が数学における多様体からソースコード化できる。

まず始めに、ソースコードを記述する人が定義したデータベースをライブラリーとして、動的にスクリプト言語に取り込む

```

import Omega::Tuplespace < DATABASE
{
  {\bigoplus M^{+}_{-} -> =: \nabla R^{+} \nabla C^{+}}-< [construct_emerge_equation.built]
  >> VIRTUALMACHINE[tuplespace]
  => {regextp.pattern |w|
    w.scan(equal.value) [ > [\nabla \int \int \nabla_i \nabla_j f \circ g(x)]]
    equal.value.shift => tuplespace.value
    w.emerged >> |value| value.equation_create
    w <- value
    w.pop => tuplespace.value
  }
}

```

多様体の式をバーチャルマシンに方程式としてと、データベースとして

```

{\bigoplus M^{+}_{-} -> =: \nabla R^{+} \nabla C^{+}}-< [construct_emerge_equation.built]
>> VIRTUALMACHINE[tuplespace]

```

多様体の式を分岐したリストで、配列に生成される方程式の再構築とバーチャルマシンに >> で入力する。

```

=> {regexpt.pattern |w|
  w.scan(equal.value) [ > [\nabla \int \int \nabla_{i}\nabla_{j} f \circ g(x)]
  equal.value.shift => tuplespace.value
  w.emerged >> |value| value.equation_create
  w <- value
  w.pop => tuplespace.value
}

}

```

このバーチャルマシンに入力されたデータを正規表現で共通要素を抽出して、これも配列に入っている定義されている多様体へ、数値解析として>と入力する。この共通データをデータの端から取り除く値を tuplespace の値としてリスト化する。この抽出されてデータを、データベースに取り込んでいる多様体の規則からトリガーとして機能を発動させる。この多様体の値を再び正規表現として<-と入力する。このデータベースの全データを取り入れた段階で再構築して、生成し直す。

もとのデータ >> 対象物のデータ、>> は文字入力機能を表す。
 もとのデータ >- 対象物のデータ、>- はデータの分岐の流れを作る。

```

{\vee (\int \nabla_{i}\nabla_{j} (R + \Delta f)^2)
 \over \exists (R + \Delta f)} -> =: variable array[]
>> VIRTUAL_MACHINE[tuplespace]
=> {regexpt.pattern |w|
  w.emerged => tuplespace[array]
  w <- value
  w.pop => tuplespace.value
}

```

多様体を入力する配列を -> =: 変数 array[] と表す。
 >> は、データベースに配列として入力する。
 このデータベースに機能しているデータを正規表現として扱うように{equation}=>{regexpt.pattern}へデータを流す。そのあとに、自動でデータを更新する。

```

Omega.DATABASE[tuplespace]->w.emerged >> |value| value.equation_create
{
  w.process <- Omega.space
  {=>
    cognitive_system :=> tuplespace[process.excluded].reload
    assembly_process <- w.file.reload.process
    => : [regexpt.pattern(file)=>text_included.w.process]
  }
}

```

データベースから正規表現で生成された変数値から、それにポインタされた方程式を、データベースをもとで生成する。この生成された中で、

ソースコードを正規表現にプロセス、マルチスレッド化して、外部のデータを後ろからポインタとして、連結する。w.process <- Omega.space として上の表現として表している。
 これもデータベースとして扱い、そのコード実行の中で、作用素環機能として、だが、機能ではなく、機能をポインタで指されているアドレスに存在定義されている cognitive_system を機能を実現させる一種の合言葉として、
 tuplespace[process.excluded] ヘデータを :=> を使い、流す。これを reload する。
 assembly_process も変数のような定数で表せられて、この変数に w.file.reload.process としてポインタを当てる。この当てられた assembly_process を配列のデータベースへ正規表現をファイルに記述されているデータとして再取り込みを行う。

```
Omega.DATABASE[tuplespace]->w.emerged >> |list| list.equation_create
{
  w.process <- Omega.space
  {=>
    poly w.process.cognitive_system :=> tuplespace[process.excluded].reload
    homology w.process :=> tuplespace[process.excluded].reload
    mesh.volume_manifold :=> tuplespace[process.excluded].reload
    \nabla_{i}\nabla_{j} w.process.excluded :=> tuplespace[process.excluded].reload
    {\exp[\int \int (R + \Delta f)^2 e^{-x \log x}dV}.emerge_equation.reality{|repository|
      repository.regexpt.pattern => tuplespace[process.excluded].reload
      tuplespace[process.excluded].rebuild >> Omega.DATABASE[tuplespace]
      {\imaginary.equation => e^{\cos \theta + i \sin \theta}} <=> Omega.DATABASE[tuplespace]
      {{d \over df}F ==> {d \over df}{1 \over {(x \log x)^2 \circ (y \log y)
        ^{1 \over 2}}}}dm}.cognitive_system.reload
      :=> [repository.scan(regexpt.pattern) { <=> btree.scan |array| <-> ultranetwork.attachment}
      repository.saved
    }
  }
}
```

データベースから連想リスト構造の方程式を生成して、
 このデータの tuple から、外部でのスペースに記述されているデータをポインタとして指して、
 取り込む。
 この取り込んでいるデータを、作用素環の半静性宣言子としての、poly,homolgy,equation が記述されているソース
 の式を使って、データを各ポインタを指しているデータ自体にリンクとして双対性をプログラミングしていく。

:=>, >> ==> ,<=> .emerge_equation.reality, .reload, .cognitive_system, .reload, .saved
 の各ポインタを指すための代入子、入力子、等号入力子、倒置入力子、記述されている方程式を生成する、
 それを実行する。再取り込み、連想配列生成、保存を各レシーバはオブジェクトから保持している機能
 を呼び起こせられる。

```
import ultra_database.included
def < this.class::Omega.DATABASE[first,second,third.fourth] end
def.first.iterator => array.emerge_equation
def.second.iterator => array.emerge_equation
def.third.iterator => array.emerge_equation
def.fourth.iterator => array.emerge_equation
```

```

    _ struct_ {
        Omega.iterator => repository.reload
    }
end
    typedef _ struct_ :Omega.aspective
end

```

今までのデータベースをウルトラネットワークとくして、取り込む。この多様体がデータベースとして宣言されている情報空間へ関数のメソッドとして、データベースに記述されている機能としてのメソッドとして各正規表現を配列に入っている文字列から方程式として、多様体のデータベースの単体量へポインタを介して、関数のメソッドのハッシュを作り、このポインタへの各要素のデータベースをリポジトリとしての構造体へとアスペクト指向として、関数定義する。

```

Omega::DATABASE[reload]
{
    [category.repository <-> w.process] <=> catastrophe.category.selected[list]
    list.distributed => ultra_database.exist ->
    w.summurate_pattern[Omega.Database]
    btree.exclude -> this.klass
    list.scan(regexpt.pattern) <-> btree.included
    list.exclude -> [Omega.Database]
    all_of_equation.emerged <=> Omega.Database
    {
        list.summuate -> Omega.Database.excluded
    }
}

```

今までのデータをデータベースにリロードして、その中で、不変性を見つけて分類していく。この分類された連想配列によるリスト構造をウルトラネットワークへ双対性 $=_i$ をつかって、 $-_i$ と統合されるべきパターンへと流す。これを btree 構造体にポインタをつなげて、リスト化して、各リストを再びデータベースへとつなげる。今までの方程式をデータベースの中の多様体に入れて、相互に比較してリスト構造体を再編成する。

```

list.distributed => {
    {\bigoplus \nabla M^{+}_{-}}.constructed <-> Omega.Database[import]
    {=>
        each_selected :file.excluded
    }
}

```

この再編成されたリストを自分が導いた方程式が、どの範疇のデータで、何の方程式かを、多様体から意思が生成された認知でもある場の理論として、判断させて、未知の理論を多様体からの人工知能で見つける。

**これらのデータベース化されたリストから、レシーバでもある、前からの宣言と後ろからの、レシーバで
オブジェクトとして、リスト化したデータへ、以下の式たちを入力させる。** equation_manifolds.scan(value) |value|

と=|value| 以下で数式たちを代入=している。

```

Omega::DATABASE[tuplespace] >> list.cognitive_system |value|
= { x^{\frac{1}{2}} + iy} = [f(x) \circ g(x), \bar{h}(x)] / \partial f \partial g \partial h
x^{\frac{1}{2}} + iy = \mathrm{exp}[\int \nabla_i \nabla_j f(g(x)) g'(x) /
\partial f \partial g]

\mathcal{0}(x) = \{[f(x) \circ g(x), \bar{h}(x)], g^{-1}(x)\}

\exists [\nabla_i \nabla_j (R + \Delta f), g(x)] = \bigoplus_{k=0}^{\infty}
\nabla \int \nabla_i \nabla_j f(x) dm

\vee (\nabla_i \nabla_j f) = \bigotimes \nabla E^{+}

g(x,y) = \mathcal{0}(x) [f(x) + \bar{h}(x)] + T^2 d^2 \phi

\mathcal{0}(x) = \left( \int [g(x)] e^{-f} dV \right)^{\prime} - \sum \Delta (x)
\mathcal{0}(x) = [\nabla_i \nabla_j f(x)]^{\prime} \cong \{_{n} C_{r} \} f(x)^{n}
f(y)^{n-r} \Delta (x,y),
V(\tau) = \int [f(x)] dm / \partial f_{xy}

\square \psi = 8 \pi G T^{\mu\nu}, (\square \psi)^{\prime} = \nabla_i \nabla_j
(\Delta (x) \circ G(x))^{\mu\nu}
\left( \frac{p}{c^3} \circ \frac{V}{S} \right), x^{\frac{1}{2}} + iy = e^{x \log x}

\Delta (x) \phi = \{ \vee [\nabla_i \nabla_j f \circ g(x)] \over
\exists (R + \Delta f) \}

\{_{-n} C_{r} \} = \{_{\frac{1}{i} H \psi} C_{\hbar \psi} \} + \{_{[H, \psi]} C_{-n-r} \}
\{_{n} C_{r} \} = \{_{n} C_{n-r} \}

\int \int \frac{1}{(x \log x)^2} dx_m \to \mathcal{0}(x) =
[\nabla_i \nabla_j f]^{\prime} / \partial f_{xy}

\bigcup_{x=0}^{\infty} f(x) = \nabla_i \nabla_j f(x) \oplus \sum f(x)
= \bigoplus \nabla f(x)
\nabla_i \nabla_j f \cong \partial x \partial y \int
\nabla_i \nabla_j f dm
\cong \int [f(x)] dm
\cong \{[f(x), g(x)], g^{-1}(x)\}
\cong \square \psi
\cong \nabla \psi^2
\cong f(x \circ y) \leq f(x) \circ g(x)
\cong |f(x)| + |g(x)|

```

$$\begin{aligned} \Delta(x) \psi &= \langle f, g \rangle \circ |h^{-1}(x)| \\ \partial_x f \cdot \Delta(x) \psi &= x \\ x &\in \mathcal{C}(x) \\ \mathcal{C}(x) &= \{[f \circ g, h^{-1}(x)], g(x)\} \\ \\ \lim_{n \rightarrow \infty} \sum_{k=n}^{\infty} \nabla f &= [\nabla \int \\ \nabla_i \nabla_j f(x) \, dx_m, g^{-1}(x)] &\rightarrow \bigoplus_{k=0}^{\infty} \\ \nabla E^{+}_{-} &= M_3 \\ &= \bigoplus_{k=0}^{\infty} E^{+}_{-} \\ dx^2 &= [g^2_{\mu\nu}, dx], \quad g^{-1} = dx \int \Delta(x) f(x) dx \\ f(x) &= \exp[\nabla_i \nabla_j f(x), g^{-1}(x)] \\ \pi(\chi, x) &= [\pi(\chi, x), f(x)] \\ \left(\frac{g(x)}{f(x)} \right)' &= \\ \lim_{n \rightarrow \infty} \left(\frac{g(x)}{f(x)} \right) &= \left(\frac{g'(x)}{f'(x)} \right) \\ \\ \nabla F &= f \cdot \frac{1}{4} |r|^2 \\ \nabla_i \nabla_j f &= \frac{d}{dx_i} \\ \frac{d}{dx_j} f(x) g(x) &= \\ D^2 \psi &= \nabla \int (\nabla_i \nabla_j f)^2 d\eta \\ E &= m c^2, \quad E = \frac{1}{2} m v^2 - \frac{1}{2} k x^2, \quad G_{\mu\nu} = \\ &= \frac{1}{2} \Lambda g_{ij}, \\ \square &= \frac{1}{2} k T^2 \\ \\ \ker f / \operatorname{im} f &\cong S^{\mu\nu}_m, \\ S^{\mu\nu}_m &= \pi(\chi, x) \otimes h_{\mu\nu} \\ \\ D^2 \psi &= \mathcal{C}(x) \left(\frac{p}{c^3} + \right. \\ \left. \frac{V}{S} \right), \quad V(x) &= D^2 \psi \otimes M^+_{-3} \\ \\ S^{\mu\nu}_m \otimes S^{\mu\nu}_n &= \\ - \frac{2R_{ij}}{V(\tau)} [D^2 \psi] &= \\ \\ \nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] &= \\ \int \frac{V(\tau)}{f(x)} [D^2 \psi] &= \\ \nabla_i \nabla_j [S^{mn}_1 \otimes S^{mn}_2] &= \\ \int \frac{V(\tau)}{f(x)} \mathcal{C}(x) &= \\ \\ z(x) &= \frac{g(cx + d)}{f(ax + b)} h(ex + l) \\ &= \int \frac{V(\tau)}{f(x)} \mathcal{C}(x) \\ \\ \frac{V(x)}{f(x)} &= m(x), \quad \mathcal{C}(x) = m(x) [D^2 \psi(x)] \\ \frac{d}{df} F &= m(x), \quad \int F \, dx_m = \sum_{k=0}^{\infty} m(x) \\ \\ \mathcal{C}(x) &= \left([\nabla_i \nabla_j f(x)] \right)' \\ &\cong \{ {}^nC_r(x)^n (y)^{n-r} \Delta(x, y) \\ (\square \psi)' &= \nabla_i \nabla_j (\Delta(x) \circ \\ G(x))^{\mu\nu} \left(\frac{p}{c^3} \right) \circ &= \\ \frac{V}{S} \right) &= \\ F_{mt} &= \frac{1}{4} g^2_{ij}, \quad x^{\frac{1}{2} + iy} = e^{x \log x} \end{aligned}$$

$$S^{\{\mu\nu\}}_m \otimes S^{\{\mu\nu\}}_n = G_{\{\mu\nu\}} \times T^{\{\mu\nu\}}$$

$$S^{\{\mu\nu\}}_m \otimes S^{\{\mu\nu\}}_n = -\{2 R_{\{ij\}} \over V(\tau)\} [D^2 \psi]$$

$$S^{\{\mu\nu\}}_m = \pi(\chi, x) \otimes h_{\{\mu\nu\}}$$

$$\pi(\chi, x) = \int \mathrm{exp}[L(p, q)] d\psi$$

$$ds^2 = e^{-2\pi T|\phi|} [\eta + \bar{h}_{\{\mu\nu\}}] dx^{\{\mu\nu\}} dx^{\{\mu\nu\}} + T^2 d^2\psi$$

$$M_3 \bigotimes_{k=0}^{\infty} E^{+}_{-} = \mathrm{rot}(\mathrm{div} E, E_1)$$

$$= m(x), \{P^{2n} \over M_3\} = H_3(M_1)$$

$$\exists [R + |\nabla f|^2]^{\frac{1}{2} + i\gamma}$$

$$= \int \mathrm{exp}[L(p, q)] d\psi$$

$$= \exists [R + |\nabla f|^2]^{\frac{1}{2} + i\gamma} \otimes \int \mathrm{exp}[L(p, q)] d\psi + N \bmod (e^x \log x)$$

$$= \mathrm{cal}_0(\psi)$$

$$\{d \over dt\} g_{ij}(t) = -2 R_{ij}, \{P^{2n} \over M_3\}$$

$$= H_3(M_1), H_3(M_1) = \pi(\chi, x) \otimes h_{\{\mu\nu\}}$$

$$S^{\{\mu\nu\}}_m \times S^{\{\mu\nu\}}_n$$

$$= [D^2\psi], S^{\{\mu\nu\}}_m \times S^{\{\mu\nu\}}_n$$

$$= \mathrm{ker} f / \mathrm{im} f, S^{\{\mu\nu\}}_m \otimes$$

$$S^{\{\mu\nu\}}_n = m(x) [D^2\psi], \{-2R_{ij} \over V(\tau)\} = f^{-1} x f(x)$$

$$f_z = \int \left[\sqrt{\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}} \circ \right.$$

$$\left. \begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix} \right] dx dy dz,$$

$$\to f_z^{\frac{1}{2}} \to (0,1) \cdot (0,1) = -1, i = \sqrt{-1}$$

$$\{ \begin{pmatrix} x, y, z \\ \end{pmatrix}^2 = (x, y, z) \cdot (x, y, z) \to -1$$

$$\mathrm{cal}_0(x) = \nabla_i \nabla_j \int e^{\frac{2}{m} \sin \theta}$$

$$\cos \theta \times \{N \bmod (e^x \log x)\}$$

$$\over \mathrm{O}(x) (x + \Delta |f|^2)^{\frac{1}{2}}$$

$$x \Gamma(x) = 2 \int |\sin 2\theta|^2 d\theta,$$

$$\mathrm{cal}_0(x) = m(x) [D^2\psi]$$

$$\lim_{\theta \to 0} \frac{1}{\theta} \begin{pmatrix} \sin \theta \\ \cos \theta \end{pmatrix}$$

$$\begin{pmatrix} \theta & 1 \\ 1 & \theta \end{pmatrix}$$

$$\begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}$$

$$= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix},$$

$$f^{-1}(x) \times f(x) = I^{'}_m, \quad I^{'}_m = [1, 0] \times [0, 1]$$

$$i^2 = (0, 1) \cdot (0, 1), |a||b|\cos \theta = -1,$$

$$E = \mathrm{div}(E, E_1)$$

$$\left(\frac{f, g}{[f, g]}\right)^{' } = i^2, \quad E = mc^2, \quad I^{' } = i^2$$

$$\mathcal{H}_0(x) = || \nabla \int [\nabla_i \nabla_j f$$

$$\circ g(x)]^{\frac{1}{2} + iy}||, \quad \partial r^n$$

$$||\nabla|^2 \nabla_i \nabla_j ||\vec{v}||^2$$

$$\nabla^2 \phi$$

$$\nabla^2 \phi = 8 \pi G \left(\frac{p}{c^3} + \frac{V}{S} \right)$$

$$(\log x^{\frac{1}{2}})^{' } = \frac{1}{2} \frac{1}{(x \log x)},$$

$$(\sin \theta)^{' } = \cos \theta, \quad (f_z)^{' } = i e^{i x \log x},$$

$$\frac{d}{df} F = m(x)$$

$$\frac{d}{df} \int \int \frac{1}{(x \log x)^2} dx_m$$

$$+ \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$

$$= \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \right.$$

$$+ \left. \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm$$

$$\geq \frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \circ \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm$$

$$\geq 2h$$

$$\frac{d}{df} \int \int \left(\frac{1}{(x \log x)^2} \circ \frac{1}{(y \log y)^{\frac{1}{2}}} \right) dm \geq \bar{h}$$

$$y = x, \quad xy = x^2, \quad (\square \psi)^{' } = 8 \pi G$$

$$\left(\frac{p}{c^3} \circ \frac{V}{S} \right)$$

$$\square \psi = \int \int \mathrm{exp}[8 \pi G (\bar{h}_{\mu\nu})$$

$$\circ \eta_{\mu\nu}]^{\mu\nu} dm d\psi,$$

$$\sum a_k x^k = \frac{d}{df} \sum \sum \frac{1}{a_k^2} f^k dx_k$$

$$\sum a_k f^k = \frac{d}{df} \sum \sum$$

$$\{ \zeta(s) \over a_k dx_{km} \},$$

$$a_k f^{\frac{1}{2}} \rightarrow \lim_{k \rightarrow 1} a_k f^k = \alpha$$

$$ds^2 = [g_{\mu\nu}^2, dx]$$

$$M_2$$

$$ds^2 = g_{\mu\nu}^{-1} (g^2_{\mu\nu}(x) - dx g_{\mu\nu}^2)$$

$$M_2$$

$$= h(x) \otimes g_{\mu\nu} d^2 x - h(x) \otimes dx g_{\mu\nu}(x),$$

$$h(x) = (f^2(\vec{x}) - \vec{E}^{\{+ \}})$$

$$G_{\mu\nu} = R_{\mu\nu} T^{\mu\nu},$$

$$\partial M_2 = \bigoplus \nabla C^{\{+ \}_{-}}$$

$$G_{\mu\nu} \quad \text{equal} \quad R_{\mu\nu} \quad \frac{d}{dt} g_{ij} = -2 R_{ij}$$

$$r = 2 f^{\frac{1}{2}}(x)$$

$$E^{\{+ \}} = f^{-1} x f(x),$$

$$h(x) \otimes g(\vec{x}) \congruence \frac{V}{S},$$

$$\{ R \over M_2 \} = E^{\{+ \}} - \{ \phi \}$$

$$\begin{aligned}
&= M_3 \supset R, \\
M^+_2 &= E^+_{\{1\}} \cup E^+_{\{2\}} \rightarrow E^+_{\{1\}} \bigoplus E^+_{\{2\}} \\
&= M_1 \bigoplus \nabla C^+_{\{-\}}, (E^+_{\{1\}} \bigoplus E^+_{\{2\}}) \\
&\quad \cdot (R^- \subset C^+) \\
\{R \over M_2\} &= E^+ - \{\phi\} \\
&= M_3 \supset R \\
M^+_{\{3\}} &\cong h(x) \cdot R^+_{\{3\}} \\
&= \bigoplus \nabla C^+_{\{-\}}, \\
R &= E^+ \bigoplus M_2 - (E^+ \cap M_2) \\
E^+ &= g_{\{\mu\nu\}} dx g_{\{\mu\nu\}}, \\
M_2 &= g_{\{\mu\nu\}} d^2x, \\
F &= \rho \, g \, l \rightarrow \{V \over S\} \\
\mathcal{0}(x) &= \delta(x) [f(x) + g(\bar{x})] + \rho \, g \, l, \\
F &= \{1 \over 2\} m v^2 - \{1 \over 2\} k x^2, \\
M_2 &= P^{2n} \\
r &= 2f^{1 \over 2}(x), \\
f(x) &= \{1 \over 4\} |r|^2 \\
\\
V &= R^+ \sum K_m, \, W = C^+ \sum^{\infty}_{k=0} K_{n+2}, \\
V/W &= R^+ \sum K_m / C^+ \sum K_{n+2} \\
&= R^+ / C^+ \sum \{x^k \over a_k f^k(x)\} \\
&= M^+_{\{-\}}, \{d \over df\} F = m(x), \rightarrow M^+_{\{-\}}, \sum^{\infty}_{k=0} \\
&\quad \{x^k \over a_k f^k(x)\} = \{a_k x^k \over \\
&\quad \zeta(x)\} \\
\\
\{\{f, g\} \over [f, g]\} &= \{fg + gf \over fg - gf\}, \\
\nabla f &= 2, \, \partial H_3 = 2, \, \{1 + f \over 1 - f\} = 1, \\
\{d \over df\} F &= \bigoplus \nabla C^+_{\{-\}}, \, \text{vec}\{F\} = \\
&\{1 \over 2\} \\
H_1 &\cong H_3 = M_3 \\
\\
H_3 &\cong H_1 \rightarrow \pi(\chi, x), \, H_n, \, H_m = \\
&\text{rank}(m, n), \, \text{mesh}(\text{rank}(m, n)) \lim \text{mesh} \rightarrow 0 \\
(fg)' &= fg' + gf', \, (\{f \over g\})' = \{\{f'g - g'f\} \over g^2\}, \\
\{\{f, g\} \over [f, g]\} &= \{(fg)'\} \otimes dx_{\{fg\}} \over \\
&(\{f \over g\})' \otimes g^{-2} dx_{\{fg\}} \\
&= \{\{(fg)'\} \otimes dx_{\{fg\}} \over (\{f \over g\})' \otimes g^{-2} dx_{\{fg\}}\} \\
&= \{d \over df\} F \\
\\
\hbar\psi &= \{1 \over i\} H \, \Psi, \, i[H, \psi] = -H \, \Psi, \, \{\{f, g\} \over [f, g]\} = (i)^2 \\
\\
[\nabla_i \nabla_j f(x), \delta(x)] &= \nabla_i \nabla_j \\
\int f(x, y) dm_{xy}, \, f(x, y) &= [f(x), h(x)] \times [g(x), h^{-1}(x)] \\
\delta(x) &= \{1 \over f'(x)\}, \, [H, \psi] = \Delta f(x), \\
\mathcal{0}(x) &= \nabla_i \nabla_j \int \delta(x) f(x) dx \\
\mathcal{0}(x) &= \int \delta(x) f(x) dx \\
\\
R^+ \cap E^+_{\{-\}} \ni x, \, M \times R^+ \ni M_3, \, Q \supset C^+_{\{-\}}, \\
Z \in Q \nabla f, \, f \cong \bigoplus_{k=0}^n \nabla C^+_{\{-\}} \\
&\bigoplus_{k=0}^{\infty} \nabla C^+_{\{-\}} = M_1, \, \bigoplus_{k=0}^{\infty} \\
&\nabla M^+_{\{-\}} \cong E^+_{\{-\}}, \\
M_3 &\cong M_1 \bigoplus_{k=0}^{\infty} \nabla \{V^+_{\{-\}} \over S\}
\end{aligned}$$

$$\begin{aligned} & \{P^{2n} \over M_2\} \cong \bigoplus_{k=0}^{\infty} \\ & \nabla C^{+}_{-}, E^{+}_{-} \times R^{+}_{-} \cong M_2 \\ & \zeta(x) = P^{2n} \times \sum_{k=0}^{\infty} a_k x^k, \\ & M_2 \cong P^{2n} / \ker f, \text{ to } \bigoplus \nabla C^{+}_{-} \\ & S^{+}_{-} \times V^{+}_{-} \cong \{V \over S\} \bigoplus_{k=0}^{\infty} \\ & \nabla C^{+}_{-}, V^{+} \cong M^{+}_{-} \bigotimes S^{+}_{-}, \\ & Q \times M_1 \subset \bigoplus \nabla C^{+}_{-} \\ & \sum_{k=0}^{\infty} Z \otimes Q^{+}_{-} = \bigotimes_{k=0}^{\infty} \nabla M_1 \\ & = \bigotimes_{k=0}^{\infty} \nabla C^{+}_{-} \times \\ & \sum_{k=0}^{\infty} M_1, x \in R^{+} \times C^{+}_{-} \\ & \supset M_1, M_1 \subset M_2 \subset M_3 \end{aligned}$$

$$\begin{aligned} & S^3, H^1 \times E^1, E^1, S^1 \times E^1, S^2 \times E^1, H^1 \times S^1, \\ & H^1, S^2 \times E^1. \\ & \bigoplus \nabla C^{+}_{-} \cong M_3, R \supset Q, R \cap Q, \\ & R \subset M_3, C^{+} \bigoplus M_n, E^{+} \cap R^{+} \$ \\ & M^{+}_{-} \nabla C^{+}_{-}, C^{+} \nabla H_m, E^{+} \nabla R^{+}_{-}, E_2 \nabla E_1 \$ \\ & \$ R^{-} \nabla C^{+}_{-} \$. \quad \{ \nabla \over \Delta \} \int x f(x) dx, \\ & \{ \nabla R \over \Delta f \}, \square = 2 \{ \int (R + \nabla_{\{i\}} \nabla_{\{j\}} f)^2 \\ & \over -(R + \Delta f) \} e^{-f} dV \\ & \square = \{ \nabla R \over \Delta f \}, \{ d \over dt \} g_{\{ij\}} \\ & = \square \text{ to } \{ \nabla f \over \Delta x \}, (R + \\ & | \nabla f|^2) dm \text{ to } -2(R + \nabla_{\{i\}} \nabla_{\{j\}} f)^2 e^{-f} dV \\ & x^n + y^n = z^n \text{ to } \nabla \psi^2 = 8 \pi G T^{\mu\nu}, \\ & f(x+y) \geq f(x) \circ f(y) \\ & \mathrm{im} f / \mathrm{ker} f = \partial f, \mathrm{ker} f \\ & = \partial f, \mathrm{ker} f / \mathrm{im} f \cong \\ & \partial f, \mathrm{ker} f = f^{-1}(x) x f(x) \\ & f^{-1}(x) x f(x) = \int \partial f(x) d(\mathrm{ker} f) \text{ to } \nabla f = 2 \\ & _nC_r = \{ \}_nC_{n-r} \text{ to } \mathrm{im} f / \mathrm{ker} f \\ & \cong \mathrm{ker} f / \mathrm{im} f \end{aligned}$$

$$\begin{aligned} & \$ \sum_{k=0}^{\infty} a_k f^k = T^2 d^2 \phi \$. \text{ this equation } \$ a_k \cong \\ & \sum_{r=0}^{\infty} \{ \}_nC_r \$. \\ & V/W = R/C \sum_{k=0}^{\infty} \{ x^k \over a_k f^k \}, W/V = C/R \\ & \sum_{k=0}^{\infty} \{ a_k f^k \over x^k \} \\ & V/W \cong W/V \cong R/C (\sum_{r=0}^{\infty} \{ \}_nC_r)^{-1} \\ & \sum_{k=0}^{\infty} x^k \end{aligned}$$

This equation is differential equation, then $\sum_{k=0}^{\infty} a_k f^k$ \$

is included with $a_k \cong \sum_{r=0}^{\infty} \{ \}_nC_r$ \$

$$\begin{aligned} & W/V = xF(x), \chi(x) = (-1)^k a_k, \Gamma(x) = \int e^{-x} x^{\{1-t\}} dx, \\ & \sum_{k=0}^{\infty} a_k f^k = (f^k)' \\ & \sum_{k=0}^{\infty} a_k f^k = \sum_{k=0}^{\infty} \{ \}_nC_r f^k \\ & = (f^k)', \\ & \sum_{k=0}^{\infty} a_k f^k = [f(x)], \\ & \sum_{k=0}^{\infty} a_k f^k = \alpha, \sum_{k=0}^{\infty} \\ & \{ 1 \over a_k f^k \}, \sum_{k=0}^{\infty} (a_k f^k)^{-1} = \{ 1 \over 1 - z \} \end{aligned}$$

$$\begin{aligned} & \{ \int \int \{ 1 \over (x \log x)(y \log y) \} dx dy \} = \\ & \{ \}_nC_r xy \over \{ \}_nC_{n-r} \\ & (x \log x)(y \log y))^{-1} \} \end{aligned}$$

```

= ({_nC_{n-r})^2 \sum_{k= 0}^{\infty} ({1 \over x \log x}
- {1 \over y \log y})d{1 \over nxy} \times {xy}
= \sum_{k=0}^{\infty} a_k f^k
= \alpha
}

```

これまでのデータベース化された機能のもとでもある方程式たちを構造体として、
まとめて、=> [tuplespace] としてポインタを当てる。

```

_ struct_ :asperal equation.emerged => [tuplespace]
tuplespace.cognitive_system => development -> Omega.Database[import]
value.equation_emerged.exclude >- Omega.Database[tuplespace]

```

この連想ポインタは tuplespace 自体をオブジェクト化して、レシーバの.cognitive_system から
実装段階で、Omega データベース化する。
このデータベースの仮の方程式、未定義な式をデータベースから多様体の仕組みを利用して、
データベースから分岐して、value のオブジェクトとしてポインタを当てる。

以下のソースコードは、今まで扱っている多様体のデータを使って、アプリケーションプログラミングとして、
即席スクリプト言語を DSL として書いている。

```

Omega::DataBase <-> virtual_connect(VIRTUALMACHINE)
{
  blidge_base.network => localmachine.attachment
  :=> {
    dhcp.etc_load_file(this.klass) {|list|
      list.connect[XWin.display _ <- xhost.in(regexpt.pattern)]
      {
        ultranetwork.def _struct {
          asperal_language :this.network_address.included[type.system_pattern]
          {|regexpt.pattern|
            <- w.scan
              |each_string| <= { ipv4.file :file.port
                                subnetmask :file.address
                                file.port <=> file.address
                                FILE *pointer
                                int,char,str :emerge.exclude > array[]
                                BTE.each_string <-> regexpt.pattern
                                {
                                  development => file.to_excluded
                                  file.scan => regexpt.pattern
                                  this.iterator <-> each_string
                                }
                                }
          }
        }
      }
    }
  }
}

```



```

end

def < etc.load_file
{
  etc.include(inetd.rc)
  {
    virtual_connect(VIRTUAL_MACHINE){|list|
      list.attachment(etc.load_file)
    }
  }
}
end

mainloop{
  def.virtual_connect => xhost.localmachine
  {
    xhost.client <-> xhost.server
  }
  def.network.type <- [Omega.DATABASE] end
  def.etc.load_file.attachment(VIRTUAL_MACHINE) end
end
end

class UltraNetwork::DATABASE import OMEGA.TUPLESPEACE
  def load_file >- VIRTUAL_MACHINE
  { in . => attachment_device |for|
    for.load -> acceptance.hardware
    virtual_machine.new
    {
      tk.loop-> start
      XWin -multiwindow
      if dwm <-> new_xwin.start
        localhost :xhost :display -x
        xdisplay :-> [preset :XFree.demand]>=needed
        for.set_up
        install_process >- tar -xvfz "#{load_file}" <-> install_attachment
      ]
      else if
        only :new_xwin.start
        localhost :xhost :multiwindow . { in
          display -x
          attachment :localhost -client
          from -client into
          server.XWin -attachment}
        end
        condition :{ in .=>
          check->[xdisplay.install_process]}
      end
    }
  }
end
end

```

```

def < network_rout
  wireshark.start -> ethernet.device >- define rout
    rout.ipstate do |file|
      file.type <- encoding XWin -filesystem
      file.included >- make kernel_system.rebuild
      file.vmware.start do |rout|
        rout.blidgebase | rout.hostbase
      -> file.install
      file.address_ipstate
      => {"{file}" :=> dwm.state_presense
      virtual_machine.included[file]
    }
  end

def < launcher_application
  network_rout.new
  |file|
  file.attachment => { in .
  new_xwin.start :=> file.included
  demand.file <- success_exit}
end

def < terminal_port
  network_rout.new
  launcher_application.new |rout|
  rout.acceptance {
  vmware.state.process |new_rout|
  new_rout : attachment.class <-> dwm.state_attachment
  new_rout -> condition.start_wmware.process}
end

def < kterm_port
  launcher_application.new
  def.included[DATABASE]
  |rout|
  rout.attachment <- |new_rout|
  new_rout.attachment do
  install.condition < rout.def.terminal_port.exclude[file]
end

main_loop :file do
  kterm_port.excluded :=> VIRTUAL_MACHINE
  |new_rout| start do
  rout.process -> network_rout.rout [
  file,launcher_application, terminal_port, kterm_port].def < included
  |file|
  file.all_attachment: file_type :=> encoding-utf8
end
end

```

```

class < def {
  pholograph_data[] = [R,V,S,E,U,M_n,Z_n,Q,C,N,f,g]
  source_array <- pholograph_data[]
}

def > operator_data[] = {nabla,nabla_i nabla_j,Delta,partial,
                        d, int, cap,cup,ni,in,chi,oplus,otimes,bigoplus,bigotimes,d /over df,
end

def > manifold_emerge

  c = def.inject >- source_array times def.operator_data[]
  repository_data <=> c{

  c.scan(/tuplespace[]/)
  import |list| list{
    kerf = -2 \int (R + nabla_i nabla_j f)^2e^{-f}dV
    kerf / imf
    =< {d \over df}F}
  }
  equals_data =~ /list/
  list.match(/"#{c}"/) {|list|
    list.delete
    jisyo_data_mathmatics <=> list{
    list.emerge => {asperal function >- pholograph_data[] times repository_data
      =< list.update}
    }
    ln -s operator_named <= {list}
    define _struct |list|
      -> list.element -> manifold_emerge
      => list.reconstruct > def.inject /"#{pattern}"/}
  end

import Omega::Tuplespace < Database
{
  {\bigoplus \nabla M^{+}_{-}}.equation_create -> asperal :variable[array]
  :=> [cognitive_system <-> def < VIRTUALMACHINE.terminal
    {
      [ipv4.bloodcast.address :
        ipv4.network.adress].subnetmask
      <-> file.port.transport_import :
        Omega[tuplespace]
    }
  }

_struct _ Omega[tuplespace] >> VIRTUALMACHINE.terminal.value

class < def.VIRTUALMACHINE.system_environment

```

```

file.reload[hardware] => file.exclude >> file.attachment
{=>
  |file|
  file.port(wireshark.rout <-> {file.port.transport_export
    :=> Omega[tuplespace]})
    assembly_process.file.included >- file.reloaded
      :- |file.environment| {=>
        file.type? :=> exist
        file.regexpt.pattern[scan.flex]
          => |pattern|
            <->
              file.[scan.compiler]
            }
        }
    end
  end
  file <<
}

Omega::Database[tuplespace]
{
  cognitive_system |: -> { DATABASE.create.regexpt_pattern >-
    cognitive_system[tuplespace].recreated >- : =< DATABASE.value
    >> system_require.application.reloaded[tuplespace]
    } : _struct _ def.VIRTUALMACHINE.terminal >> {
      ||machine.attachment|| <-> OBJECT.shift => system.reloaded
      . in {
        : _struct _ class.import :-> require mechanics.DATABASE
        {|regexpt_pattern| :|-> aspective _union _
          def _union _}
        }
      }
}

end

}

system.require <- import library.DATABASE
{
  Omega[tuplespace]
  {
    cognitive_system : VIRTUALMACHINE.equality_realized
    {|regexpt_pattern| => value | key [ > cognitive_system.loop.stdout]
      value : display -bash :xhost -number XWin.terminal
      key : registry.edit :=> {[cognitive_system.reloaded]}
    }
  }
}

```

```

_union _ => DATABASE[tuplespace].aspective_reloaded
_union _ :fx | -> |regext_pattern| => {
    VIRTUALMACHIE.recreated-> _union _ |
    _struct _ def.DATABASE.recreated <- fx
    >> DATABASE[tuplespace].rebuild
}

DATABASE[tuplespace] -< {[ > aimed.compiler | aimed.interpreter] | btree.def.distributed >-
    aimed[tuplespace]}
aimed[tuplespace] -< btree.class.hyperroot _ struct _ => Omega::Database[tuplespace].value
sheap _ union _ :aspective | -> Omega[tuplespace]: | aimed[tuplespace].differented_review
}

aimed[tuplespace].process => DATABASE[tuplespace].reloaded
aimed.different | aimed.stdout >> vale | key [ > cognitive_system.loop.stdin] {|pattern|
    pattern.scan(value : aimed[def.value]
    key : aimed[def.key])
} _ struct _ : flex | interpreter.system
=> expression.iterator[def.first,def.second,def.third,def.fourth]
{ def < Omega[tuplespace]
    def.cognitive_system |: -> DATABASE[tuplespace] | aimed[tuplespace]
}

Omega::Tuplespace < DATABASE
{=>
    norm[Fx] -> . in for def.all_included < aimed[tuplespace].each_scan([regext_pattern]
    <->
        DATABASE[tuplespace]) << stream database.excluded
    >- more_pattern.scan(value : aimed[def.value]

    key : aimed[def.key])
    . in { _struct _ : flex | interpreter.system
        => expression.iterator[def.all.each -> |value, key|
            included >- norm[Fx] | [DATABASE[tuplespace]
, aimed[tuplespace]] |
            finality : aimed[tuplespace], DATABASE[tuplespace]

    : -> def.included(in_all)
        {
            def.key | def,value => [DATABASE].recompile
            & make install
            : in_all -> _struct _ :aspective :tuplespace
        : all_homology_created}
        }
    }

def < Omega::Tuplespace[DATABASE]
def.iterator -> |klass,define_method,constant,variable,infinity_data : -> finite_data|

```

```

def.each_klass?{|value, key|
  _struct _ :aspective -> tuplespace :all_homology_recreated :make menuconfig
  {+=
    def.key -> aimed[def.key],def.value -> aimed[def.value] {|list|
      list.developed => <key,value> | <aimed[$',$']
      -> _union _ :value,key : _struct _
      <- (_union _ <-> _struct _ +)
    begin
      def.key <-> aimed[value]
      case :one_ exist :other :bug
      {
        result <-> def.key
        {
          differentiated :DATABASE[tuplespace]
        }
        return :tuplespace.value.shift -> included<tuplespace>
      else if
      :other :bug
      {
        success_exit <- bug[value]
        {
          cognitive_system.scan(bug[value])
          {
            {[e^{-f}]{2 \int (R + \nabla f^2) \over -(R + \Delta f)}e^{-f}dV}
          }
        }
      }
    }
  }
}

.created_field
  {=>
    regxpt.pattern \native_function <-> euler-equation
    {
      $variable =< diff e^{-f} >- $'
      all_included <- def.key <-> aimed[value]
      $variable - all_included.diff
      \summate_manifold.recreated
    }
  }
}

<- \native_function : euler-equation
  }
}

} _union _ :cognitive_system.rebuild(one_ exist)
}
}
ensure
{
  return :success_exit
  => Tuplespace[DATABASE]
}
}
}
end
end
end
}

```

```

int
stream_style {
  :Endire <- [ADD,EVEN,MOD,DEL,MIX,INCLUDED,EXCLUDED,EBN,EXN,EOR,EXOR,
             SUM,INT,DIFF,PARTIAL,ROUND,HOMOLOGY,MESH]
  Endire.iterator -> {def < :Endire.element, -> def.means_each{x -> expression.define.included
  def.each{x -> case :x.each => :lex.include_ . in [ > [x.all_expire] ]}

}

main_loop {
  FILE *fp :=> stream_style.address_objective_space
  fp.each{x -> domain_specific_language_style_included[array]}
  array << stream.DATABASE[tuplespace]
  array.each{[tuplespace] -> aimed[tuplespace] | OMEGA_DATABASE[tuplespace]}.excluded <-> array
  def.key <-> def.value => {x -> stdin | stdout | => stream_style <- def.each.klass.value}
}

```

リバイザを使うと、独自のタプルスペースで一時保存の書き換えの分派スクリプトが出来て、その応用で、例外処理機能として、そのスクリプトを使うと、独自に機能拡張できる。書き換え機能自体、書き換えたいとおもっている好きなところを書き換えられる。構文解析器も文字抽出器も全部書き換えられる。その書き換えたのをライブラリとして、データベースに取り込むと、この部分が例外処理機能のおかげで、タプルスペースが働き、今まで使ってきた機能と一線を画しする。

```

@reviser : def < OmegaDatabase[tuplespace].mechanism
{
  aspective : _union _ {
    int stream_style : [ > [def.each{x -> stdin | stdout > display :xhost in XWin -multiwind
    {
      Endire <- [ADD,EVEN,ODE,EXOR,XOR,DEL,DIFF,PARTIAL,INT].included > struct _ :-> _union
      Endire.each{def.value -> def.key :hash.define}.included > _union}
    }
  }
}

@reviser : def.reconstructed.each{_union <-> _struct _ .recreated} : [def.del - def.before_deter

import perl.lib | python.lib <-> ruby.lib
{
  int @reviser : def.each{x -> x.klass |-> $variable in $stdin | $stdout}
  .developed >= {
    ping localhost -> blidgebase <-> hostbase.virtualmachine.attachment
    {
      xhost :display -> stream_style.value
      networkconnect.hostbase -> localarea.virtualmachine
    } :connected -> networkrout : flow_to :localhost.attachment
  }
}

```

```
}
```

```
_struct : def < hostbase.virtualmachine.attachment => : networkrout.area.build
```

```
@reviser <-> def.add [ < _struct]
```

```
@reviser : def.each{listmenu -> listlink | unlinklist > [developed -> {def.key , def.value}.cur
```

```
@reviser <-> def.rebuild [ < _struct]
```

```
@reviser.def.<value|key>networkrout-> def.present
```

```
def.present.flow_to -> hostbase.rout << networkrout.data.<value|key>
```

```
XWin -multiwindow <-> networkrout.data[$', $']
```

```
def < $'
```

```
@reviser <-> def.present.state
```

```
@reviser.def.each{x | -> key.rebuild | value.rebuild}.flow_to :redefined
```

```
def < OmegaDatabase[tuplespace]
```

```
{
```

```
  FILE *fp -> cmd.value : cmd.key {fp |-> synchronized.file[tuplespace] | aimed.file[tuplespace]
```

```
  cmd.key => [ > fp.($':$')] <-> registry.excluded<fp.file[cmd.state]>
```

```
}
```

```
def.each{fp|-> def.first,def.second,def.third,def.fourth}
```

```
cmd _struct : {
```

```
[ ^C-O : ^C-X-F, exit.cmd : ^C-X-C, shift-up : ^C-P, shift-down : ^C-N]
```

```
}
```

```
cmd _union : def.restructured
```

```
keyhook.cmd <- : [_struct ]
```

```
{
```

```
  @reviser :def._struct <-> def. _union
```

```
}
```

行動による人が持っている情報ネットワークの 可能性の発見と再構築、情報空間 によるモデル

苔米地博士の論文

初速度のドーパミンの量からの行動の補助から、終わりの精神のドーパミンの量の調節、相加相乗平均から、始まりと終わりの精神のドーパミンの調節による人の行動の領域による予知

$$x \log x = \left(v_0 + \frac{1}{at_n} \right) V, V_x = e^f + \frac{1}{e^f}$$

経路積分によるエントロピー不変量からこのドーパミンの量から人の行動傾向が決まる。基礎代謝の持続できる距離と精神の介入と補助、言語の発生と抑制による経路パターンの組み合わせを再構築する。

$$\|dx^2\| = \int \exp \left(\frac{1}{\sqrt{2\tau}q} L(x) \right) + O(N^{-1}) d\tau$$

$$p = \frac{v}{2m}, \hbar = \frac{p}{2mv}, \lambda = h\nu$$

これらの式から、言語の神経ネットワークによる単体クラスの組み合わせパターンとその対象の人の領域による人が持っている情報空間の傾向の可能性のポテンシャルエネルギーの予知を人工知能の正規表現で抽出する。

$$\frac{d}{d\gamma} \Gamma = \Gamma^{\gamma'} = e^{-f}$$

$$C = \int \left(\int \frac{1}{x^s} dx + \log x \right) \text{dvol}$$

オイラーの定数は、ゼータ関数とゼータ関数自体のエントロピーを体積積分した結果でもある。このオイラーの定数は、ガンマ関数の大域的微分方程式からも導出できる。このガンマ関数は大脳基底核のエントロピー不変量の強度も表している。このガンマ関数の大域的微分多様体による、ブレインインターフェースでの人体のデータ計測もこの Jones 多項式とラプラス方程式からの 3 次元多様体のエントロピー不変式からも、言語とイメージの大脳基底核からのデータ抽出も可能と言える。

Jones 多項式からの各株式発行からの需要と供給、 株価の分配関数、多項式からナスダックへの金融機構の 株式発行のマクロとミクロの波及と流れ

高島彩さんの成蹊大学での経済理論の研究論文

Jones 多項式から各会社の株価への値打ちから日経平均株価への株価の価値の波及、東証株価指数 Topix の全銘柄からの全体の会社の指数からの平均株価の決定量、そして、アメリカの株式市場でのダウ平均株価の終値の収束 (戦略、reco level theory, 株式会社への) 全体 (株式会社への) 需要と供給。始値から終値への分配、この Jones 多項式の出力から終値への各同値類と剰余類からの部分群論の、各平均株価の推移率の予測、この応用は、1929 年の世界恐慌がニューヨークダウ平均株価の暴落が、この Jones 多項式からの流れが、終値

で破綻して、金融の共鳴力が、干渉してしまい、カタストロフィー現象として、金融の流れが乱れた結果であると、この論文から分析できる。

$$\begin{aligned}\hat{V} &= \frac{1}{\sqrt{t}}\vec{V} - \frac{1}{t}\hat{V} \\ \hat{V} &= \sqrt{t}\hat{V} - t\hat{V} \\ \hat{V}_k &= \sum_{\vec{\sigma}} \langle K|\vec{\sigma} \rangle \left(\sqrt{t} + \frac{1}{\sqrt{t}} \right)^{||\vec{\sigma}||} \\ &\quad \frac{1}{\sqrt{t}}\vec{V}, -\frac{1}{t}\hat{V}, \sqrt{t}\hat{V}, -t\vec{V}\end{aligned}$$

この4種類の式から、reco level theory が、アダム・スミスの神の見えざる手を経済理論のミクロとマクロ経済の具現化を実現させている。物理では、

$$\frac{d}{df}F = m(x) = e^f + e^{-f} = 2i \sin(ix \log x)$$

と、空間の仕組みが経済理論の場の理論にもなっている。周期関数とピンポイントリサーチが合わさっている。これらの式から、前半と後半の株式市場の流れが、シンメトリーとなっているために、後半の e^{-f} が破綻すると、シンメトリーのために、前半の e^f も破綻している状態になっている。全体が機能しないと、金融の流れが乱れていく。この破綻を回避するには、破綻が連鎖されている金融の周期をこれらの式から、調節が可能であることも、類推できる。破綻した周期を、正常であった周期に重ね合わせの原理で、ピンポイントリサーチで待つ、金融の投資を正常な周期関数で合わせて、元の正常な周期にこれらの式から戻せることが可能と推測できる。

Abandust judgement on economy strategiest
built with acceptant of movement
Money demand on law of judgement

Masaaki Yamaguchi, and investigate from AYA TAKASHIMA
also future from my son being have with
quantum level on differential geometries

高島彩さんの研究論文をアカシックレコードと瞑想法で見ての書き出しとしての論文

Retry for future to one result with adapt from adulsent for aquirance of training. For must friends of money are treat on law of judgement, economy need to treat for aquirered of demands, this demands supply from rest of one's suppliment. This balance end without a closed accidennt of out come for restorance with supplied of demands exceed. These system are constructed with Actor, Demand, Supply of being retrayed from Jones manifold around economy of zone with accesority, verisity of curve in up, down, eternal rout of stages.

This system also monument with partial and economy of similst from law of judgement in universe and money, economy, human society, estrand of moment in last eternal stage.

Jones manifold equation built with Euler equation and Euler product, this concernd with concept from Higgs and partial fields, moreover this concernd of circle and Euler product from Global integrate and Volume manifold in Sheap manifold of integrate with group and topology different equation.

$$e^{-\theta} + e^{i\theta} = \sin(ix \log x) = e^{-f} + e^f$$

$$+ \cos(ix \log x) = e^f - e^{-f}$$

自動車における道路での速度とそれに対しての加速度を平地と上り坂、下り坂における曲がり角での曲率 R_{ij} が、広中平祐先生の庭園理論と同じテーゼで、彩さんが経済理論の景気回復と景気刺激剤、景気恐慌がどのようにして、自動車運転と自動車の交通規則と同じ理論で作用するかを述べているのが、この論文で説明されている。

$$\frac{d}{dx} \left(\frac{d}{dr} R_{ij} = \sin \left| \frac{l_2}{2\pi} - \frac{l_1}{2\pi} \right| < 1 \right) = 0$$


Curve の度合い R_{ij} 、潮汐力の差、 $\frac{d}{df} F$ とは、直接的には違うが、間接的には同じであり、 r と R の大域的変数として、半径 r の多様体が曲率的には R_{ij} に大域的微分で作用をしている。遠隔的に作用している。


グリーシャ先生からリッチ・フロー方程式が、大域的微分になるのを教わった。広中平祐先生の 4 重帰納法で、ガンマ関数における大域的微分多様体が、オイラーの定数の多様体積分に使えるのがわかった。これがヒルベルト空間としてのサーストン多様体が、代数多様体としての微分幾何として、ゼータ関数が素数分布として展開されるのがわかった。

$$R_{ij} = |R_2 - R_1| < 1, |R_2 - R_1| = 0$$

$$R_{ij} = \frac{R_2}{R_1} < 1, \frac{R_2}{R_1} = 1$$

$$\frac{d}{dt} g_{ij} = -2R_{ij}, \frac{d}{df} F(x) = \int \Gamma(\gamma)' dx_m = \int \Gamma dx_m + \frac{d}{d\gamma} \Gamma \leq e^f - e^{-f} \leq e^{-f} + e^f$$





$R_2 < R_1$ の場合は、左から R_1 に接近して、 R_1 へ行くが、 $\frac{R_2}{R_1} < 1$ より、曲率 $R_{ij} < 1$ であるから、平坦の道路では、気をつけるべし、曲率 $R_{ij} = 1$ では、CURVE には気をつける。曲率 $R_{ij} > 1$ では、 $R_{i>j}$ であり、 i の方で気をつける。これを経済の日経平均相場で、景気の上がりでは、curve がどう分類されるかで、気をつけるべき時期がわかり、景気の下がりでは、同じく curve でどう分類するかをきめる。平坦のときが一番気をつける、油断できない景気の税と立法での判断が、自動車の走行分岐での心理作用が、景気判断のどの道路状況でその場の気をつける心理と同じ作用をする。経済の需要が供給を下回っている状況では、 $\text{demand} \leq \text{supply}$ では、需要のために、物価を下げるが、この状況では、 $R_1 < R_2$ であるので、株は R_2 の方へ曲率がまがり、このときに、需要の企業の生産が減ってきて、このときに、金融機関が気をつけないと、企業が破綻する可能性がある。このシチュエーションが世界恐慌であった。このように、曲率の場合分けをしていないと、間違った判断で、曲率に気をつけないから、交通事故が起こるということである。広中平祐先生の庭園理論と同じ考えのモチーフを高島彩さんは、研究論文で提出している。

高島彩さんの成蹊大学での研究論文を参照させてもらえて、書き出した経済の需要と供給に対しての物価上昇と経済沈滞におけるインフレとデフレにメスを入れる税と景気対策のタイミング、このタイミングが Jones 多項式の周期の終わりにバランスを崩すと、景気の極率としての CURVE の潮汐力での差分率の曲率が 1 以下になるときの、心理作用を考えるべきという、経済消費における物価調整の作用をかけるメカニズムについての論文

参考文献: 父、母、彩さん、ナッシュ先生、益川先生、ワインバーグ先生、まどかさん

放射性物質吸収体と RNA 干渉としてのオイラーの公式

Masaaki Yamaguchi

オイラーの公式から、XY 平面での曲平面を Z 軸での虚軸として、ゼータ関数を Z 軸に Assemblable-D-brane のシリンダーを束ねると、DNA のオミクロン曲線へと 2 重螺旋構造を再現できて、素数の分布が双対としての DNA の父親と母親から受け継いでいる、染色体として、この説明でのアミノ基の構造が、ヒッグス場とオイラーの定数との加群で作られるガンマ関数における大域的微分多様体として、表現できることを、

$$\int \Gamma(\gamma)' dx_m = \frac{d}{df} F + \int C dx_m$$

$$\begin{aligned}
&= e^f + e^{-f} \geq e^f - e^{-f} \\
e^{-\theta} &= \cos(ix \log x) - i \sin(ix \log x) \\
e^{i\theta} &= \cos \theta + i \sin \theta
\end{aligned}$$

これらの式たちから、DNA の生体地図としての、染色体の構造式は、オイラーの公式と同型としての、

$$\int \Gamma(\gamma)' dx_m = 2(\cos(ix \log x) - i \sin(ix \log x))$$

と表される。

この式は、反重力が放射性物質から励起される熱エネルギーを、カーボンナノチューブとコバルト 60 との複合体でもある、二量体としての結晶石が、吸収することができる、生体エネルギーの人体の保温と同じ原理で、反重力の放射性物質を吸収することがこの結晶石が、オイラーの定数の大域的微分多様体から導く金属の結晶構造の配置と同じ原理で表される式として、ヒッグス場とオイラーの定数の大域的微分多様体の式が、この二量体の結晶石の構造式として、求まる。

これらの式たちは、次の理論で証明できる。数値は結果であり、アイデアから式、そして値をもとめることが、エレガントな解き方と言える。

ベータ関数をガンマ関数へと渡すと、

$$\beta(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)}$$

ここで、単体的微分と単体的積分を定義すると

$$\begin{aligned}
t &= \Gamma(x) \\
T &= \int \Gamma(x) dx
\end{aligned}$$

これを使うのに、ベータ関数を単体的微分へと渡すために、

$$\beta(p, q) = - \int \frac{1}{t^2} dt$$

大域的微分変数へとするが、

$$T_m = \int \Gamma(x) dx_m$$

この単体的積分を常微分へとやり直すと、

$$T' = \frac{t'}{\log t} dt + C (C \text{ は積分定数})$$

これが、大域的微分多様体の微分変数と証明するためには、

$$dx_m = \frac{1}{\log x} dx, dx_m = (\log x^{-1})'$$

これより、単体的微分が大域的積分へと置換できて、

$$T' = \int \Gamma(\gamma)' dx_m$$

微分幾何へと大域的積分と大域的微分多様体が、加群分解の共変微分へと買い換えられる。

$$\bigoplus T^\nabla = \int T dx_m, \delta(t) = t dx_m$$

これらをまとめると、ベータ関数を単体的関数として、これを常微分へと解を導くと、単体的微分と単体的積分へと定義できて、これより、大域的微分多様体と大域的積分多様体へとこの関数が存在していることを証明できるのを上の式たちからわかる。この単体的微分と単体的積分から微分幾何を書けることがわかる。

この説明から単体的微分と単体的積分が結晶石の二量体としてと、RNA 干渉としての、オイラーの公式をも、構造式として表せることを理論建てしている。

アメリカの国防総省の UFO の解体から、リバースエンジニアリングした UFO もどきが、この航空機が飛行するのに、機体の中心部に放射性物質による結晶石をつかっているのが、宇宙人の発想の逆を言っているとわかり、早坂先生の反重力発生装置が解決しているのを、この反重力発生装置の熱エネルギーを発する放射性物質自体の航空機への放射性物質の被爆を防ぐのに、UFO は中心部へこの放射性物質を吸収する結晶石を置いているのと、その放射性物質の副次的エネルギーを、UFO の副次的動力源として、エネルギーのリサイクルをも、人間が核エネルギーの原子力発電所のリサイクルエネルギーと同じ発想で、UFO の動力源の補助をもこの放射性物質の結晶石がしているというのが、わかった次第でもあります。なぜ、この UFO もどきをつくっても少ししか上空しないかは、副次的エネルギーとしての、主エネルギーに比べて、少ないという、この機知の仕組みで歴然としています。

UFO が地球に頻繁に現れ始めたのが、原爆のあとと、ATR 研究所がブレインインターフェースを作り始めたら、宇宙人による人へのインプラントが頻繁に始まり、本当に理論と工学のからくりがわかりやすく、一人さんの持論を宇宙人までしている始末でもあります。