


CONTACT TRANSPORTER

異次元への輸送機 3 次元設計図書

量子プログラミング言語 Bada (contact_transporter/contact_blueprint.bada) による生成



原理: ジュネーブ CERN の衝突で粒子が消えた方向 = 異次元への扉

- × 特殊相対論 (地球の 1 秒 = 搭乗者の 18 時間, $\Gamma = 64800$)
- × 複素回転体(コマ)の幾何学 ($\Theta = \theta + i\phi$, $\phi = 11.7722$)
- × ガンマ関数におけるゼータ関数の大域的部分積分多様体 (Riemann-Siegel Z)
- × Jones 多項式 = 設計図生成機能

論文 15 本の全方程式: 2111 本 (数値評価 632 本 / 等式の数値検証 成立 231・不成立 181)

※ 本書は論文の方程式に基づく思索的・フィクションの設計図 (幾何的な可視化) です。

工学的に検証された装置ではなく、異次元への輸送を可能にするものではありません。

1. 設計パラメータ（Bada 実行結果）

- ① 衝突段

$$\sqrt{s} = 13.6 \text{ TeV (Linac4} \rightarrow \text{PSB} \rightarrow \text{PS} \rightarrow \text{SPS} \rightarrow \text{LHC)}$$
$$E_{T^{\text{miss}}} = 1348.29 \text{ GeV}$$
$$\text{扉の軸 } \hat{n} = (0.86964, -0.25101, 0.42511)$$
- ② 特殊相対論

$$\Gamma = 18 \text{ h} / 1 \text{ s} = 64800$$
$$1 - \beta = 1.19075\text{e-}10, \text{ラピディティ } \phi = \text{arcosh } \Gamma = 11.772200$$
$$\text{ベガ } 25.04 \text{ ly} \rightarrow \text{収縮距離 } 3.656\text{e+}09 \text{ km}, \text{片道 } 3.3874 \text{ h}$$
- ②' 輸送計量

$$ds^2 = e^{\{-2\pi T|\phi|\}}[\eta + \hbar]dx^\mu dx^\nu + T^2 d\phi^2$$
$$e^{\{-2\pi T|\phi|\}} = 1/\Gamma \rightarrow T|\phi| = 1.763290 \text{ (} x \log x = 1 \text{ の根 } 1.763220)$$
- ③ 複素回転体

$$\Theta = \theta + i\phi, \theta_0 = 0.523599 \text{ rad}$$
$$\omega(\text{外, 中, 内}) = 0.10472, 0.39241, 1.47043 \text{ rad/s}$$
$$\text{ポッド歳差 } \Omega = Mgl/(I_3\omega_3) = 3.70640 \text{ rad/s}$$
- ④ $\Gamma \cdot \zeta$ 多様体

$$\text{Riemann-Siegel } \theta(\phi) = -2.581360$$
$$Z(\phi) = e^{i\theta(\phi)} \zeta(\tfrac{1}{2} + i\phi) = -1.334150$$
- ⑤ Jones 多項式


$$3_1: -t^{-4} + t^{-3} + t^{-1} \mid -0.116342 + 2.30908i$$
$$4_1: t^{-2} - t^{-1} + 1 - t^1 + t^2 \mid 3.56478 + 0i$$
$$5_1: t^2 + t^4 - t^5 + t^6 - t^7 \mid -2.81527 + 1.09654i$$
- ⑦ 部品寸法 [m]

$$\text{外環 } R = 60.088, \text{中環 } R = 47.431, \text{内環 } R = 37.440, \text{管径} = 2.066$$
$$\text{ポッド } r = 5.021, \text{塔高} = 132.194, \text{井戸深さ} = 38.800$$
$$\text{Jones コイル: } 3_1 R = 70.269, 5_1 R = 56.215, 4_1 \text{ 床下}$$


$V(t^*)$ は $t^* = e^{i\theta(\phi)}$ での値。 $|V|$ がコイル寸法、 $|V(e^{i\alpha})|$ の極小が扉の共鳴窓の角度を決める。
寸法は各方程式群の数値エネルギー $\sum \log(1+|v|)$ で変調される (lib_registry.bada / main_blueprint.bada)。

2. 三面図 (単位 m)


上面図 (x-y)






正面図 (x-z)



側面図 (y-z)



3. ゼータ多様体・Jones 共鳴・方程式の割り当て



4. 全方程式レジストリ (2111 本) p.1/33

ACAFE.1	symb	$\nabla \Delta = \oint_{\mathbb{D}} \otimes \mathbb{M}(\square) \, \mathrm{d}\square$	[MANIFOLD]
ACAFE.2	symb	$\oint_{\mathbb{M}} \pi(\chi, x) = \oint_{\mathbb{M}} [i \pi(\chi, x), f(x)]$	[MANIFOLD]
ACAFE.3	differs	$\mathbb{M}(\square) = x^{\sim}y + y^{\sim}x + z^{\sim}a + u^{\sim}b + v^{\sim}c = 0$ (pypdf: 'xy+yx+za+ub+vc'; exponents lost) = 14 vs 0	[MANIFOLD]
ACAFE.4	holds	$1/y + 1/x + 1/a + 1/b + 1/c = 1$ = 1 vs 1	[MANIFOLD]
ACAFE.5	calc	$\nabla \Delta = \oint_{\mathbb{D}} \otimes \mathbb{M}(\square) \, \mathrm{d}m$, $T = \Gamma'(\gamma) \, \mathrm{d}x^{\sim}m$ = 0.422784	[GAMMA, MANIFOLD]
ACAFE.6	calc	$\square = 2(\sin(i \, x \, \log x) + i \, \cos(i \, x \, \log x))$ = 0 + 8i	[ROT, OTHER]
ACAFE.7	calc	$= \mathrm{d}/\mathrm{d}\gamma \, \Gamma$ = 0.422784	[GAMMA]
ACAFE.8	symb	$\mathbb{M} = [i \pi(\chi, x), f(x)]$	[MANIFOLD]
ACAFE.9	calc	$\square = -(16 \pi G/c^4) \, T_{\mu \nu}$ = -4.15329e-43	[OTHER]
ACAFE.10	calc	$= \kappa \, T_{\mu \nu}$ = 2.07665e-43	[OTHER]
ACAFE.11	symb	$\int E'(\sigma) \, \mathrm{d}\sigma = \nabla_{\perp} i \nabla_{\perp} j \otimes \mathbb{M}(\mathbb{H}(\sigma) \otimes \mathbb{K}(\sigma)) \, \nabla \eta \, \mathrm{d}\eta$	[MANIFOLD, QUANTUM]
ACAFE.12	symb	$\sigma = \int (h \nu) \, \nabla^{\sim}\{\otimes L\} \, \mathrm{d}\Psi$	[QUANTUM]
ACAFE.13	symb	$\nabla + (\square(\nabla \phi))^{\sim}\{\nabla \otimes L\} = \int \square'(\nabla \phi) \, \mathrm{d}x^{\sim}m = \boxplus \Psi$	[QUANTUM]
ACAFE.14	symb	$(1/x) \boxtimes y = \otimes \nabla w = (\boxtimes x)^{\sim}\{x+y\}$	[OTHER]
ACAFE.15	symb	$(\otimes \nabla w) \boxtimes (\otimes \nabla w) = \otimes (i \hbar \nabla)^{\sim}\{\otimes L\}$	[QUANTUM]
ACAFE.16	symb	$(\otimes (i \hbar \nabla)^{\sim}\{\otimes L\} + m)(\otimes (i \hbar \nabla)^{\sim}\{\otimes L\} + n) = L^{\sim}\{m+1\}/(m+1)$	[BETA, QUANTUM]
ACAFE.17	holds	$L^{\sim}\{m+1\}/(m+1) = \int (x-1)^{\sim}\{t-1\} \cdot t^{\sim}\{x-1\} \, \mathrm{d}t = \beta(p, q)$ = 0.0833333 vs 0.0833333	[BETA]
ACAFE.18	calc	$\square^{\sim}\{(x+y)/2\} - \sqrt{(x \cdot y)} = \square^{\sim}\{\ll o\}$ = 1	[OTHER]
ACAFE.19	symb	$\square \Psi \boxtimes \square \Psi$	[ZETA]
ACAFE.20	symb	$\boxtimes = F_{\perp}(\mathbb{T} \perp)$, $F_{\perp}(\mathbb{T} \perp)^{\sim}\{1/2\} = \nabla +$	[ZETA, JONES, TRANSPORT]
ACAFE.21	symb	$\square \int Z(\xi)/(\hbar \nu) \, \mathrm{d}x^{\sim}\xi$	[ZETA]
ACAFE.22	symb	$= \square \int Z(\xi)/(\log x) \, \mathrm{d}x^{\sim}\xi$	[ZETA]
ACAFE.23	calc	$S = \pi \oint r^2 \, \mathrm{d}r$ = 1.0472	[OTHER]
ACAFE.24	holds	$= \int \int e^{\sim}\{-x^2-y^2\} \, \mathrm{d}x \, \mathrm{d}y \cdot \oint r^2 \, \mathrm{d}r$ = 1.0472 vs 1.0472	[ZETA]
ACAFE.25	differs	$= 2 \pi S$ = 1.0472 vs 6.57974	[OTHER]
ACAFE.26	holds	$e^{\sim}\{2 \pi r\} = 1$ = 1 - 2.44929e-16i vs 1 + 0i	[ROT, ZETA]
ACAFE.27	symb	$\Gamma : r \rightarrow \chi = \nabla Y \rightarrow \square Y \rightarrow [\Delta, \nabla, \mathrm{d}, \partial, \delta, \mathrm{d}x^{\sim}m]^2 \rightarrow Y : m \rightarrow n$	[GAMMA]
ACAFE.28	calc	$\mathrm{d}/\mathrm{d}f \, F(x, y) = \otimes [\pi(\chi, x)/(x \, \log x) \, \mathrm{d}x^{\sim}m + i \pi(\chi, x)/(x \, \log x) \, \mathrm{d}y^{\sim}m] [\mathrm{Im}]$ = 0.0542868 + 0.0542868i	[GAMMA, OTHER]
ACAFE.29	symb	$\nabla + [\int \pi(\chi, x) \, \mathrm{d}x^{\sim}m + \int N \pi(\chi, x) \, \mathrm{d}y^{\sim}m] [\mathrm{dIm}]$	[MANIFOLD]
ACAFE.30	holds	$\int \mathrm{d}\boxtimes = \int f(x) \, \mathrm{d}(x \, \log x) = F$ = 2.71828 vs 2.71828	[ENTROPY]
ACAFE.31	calc	$\boxtimes = f/(x \, \log x)$ = 0.721348	[ENTROPY]
ACAFE.32	differs	$(e^{\sim}f + e^{\sim}\{-f\})/(e^{\sim}f - e^{\sim}\{-f\}) = \int \Gamma'(\gamma) \, \mathrm{d}x^{\sim}m / m(x)$ = 1.31304 vs 1	[GAMMA, JONES]
ACAFE.33	symb	$\mathrm{d}/\mathrm{d}f \, F(x, y) = m(x, y)$	[OTHER]
ACAFE.34	holds	$\square_{\mathbb{K}=\mathrm{prime}}^{\sim}\{\infty\} Z^{\sim}\{\ll D\} \boxtimes (\xi) [\mathrm{Im}] = \pi(\boxtimes \cdot \boxtimes, m)$ = 1.64484 vs 1.64493	[ZETA, MANIFOLD]
ACAFE.35	symb	$t \int \int \mathbb{D} \chi[\pi(\chi, x)] : x \rightarrow 2, : y \rightarrow \infty [\mathrm{dIm}] = \int \int \int [(\boxtimes x \cdot \boxtimes m - \boxtimes y \cdot \boxtimes n)/(t - t1)] \, \mathrm{d}\boxtimes m$	[BETA, MANIFOLD]
ACAFE.36	symb	$\beta(p, q) = \nabla + [\boxtimes m x, y \cdot \boxtimes x, y] \times [\sigma(\boxtimes)]$	[BETA]
ACAFE.37	symb	$E = \mathbb{K}(\sigma) \otimes \mathbb{H}(\sigma) = \beta^{\sim}\{-1\}(x) \times \beta(x)$	[BETA, MANIFOLD]
ACAFE.38	symb	$\nabla \nabla : \chi \rightarrow \nabla_{\perp} i \nabla_{\perp} j \, \Gamma'(\gamma) \, \mathrm{d}\gamma \, r^{\sim}m \rightarrow \Delta \nabla \square \Psi$	[GAMMA, MANIFOLD]
ACAFE.39	calc	$\boxtimes (\mathbb{H} \Psi \, \mathbb{D} \chi)^{\sim}\{\ll \mathbb{D} \phi\} = \oint Z'(\xi)/(2 \pi i) \, [\mathrm{dN}]$ = 1	[ZETA]
ACAFE.40	symb	$= [f(x)g'(x) - g(x)f'(x)] - [F(x)G(x)]$	[OTHER]
ACAFE.41	symb	$x \, \log x = i \pi(\chi, x)$	[ENTROPY]
ACAFE.42	symb	$\log x = \Delta f - \nabla g$	[OTHER]
ACAFE.43	symb	$= (\mathrm{d}/\mathrm{d}f \, F_g) \cdot (g(f))$	[OTHER]
ACAFE.44	symb	$\partial/\partial f \, F(x, y) = F_{\cdot} f(x, y) \, f'(x, y) = \partial/\partial f \, F(g(x, y))$	[OTHER]
ACAFE.45	symb	$\bigcirc \bigcirc \int^{\sim} N (\boxtimes \cdot \boxtimes)^{\sim} \mathbb{D} = \Delta \Delta \, \mathbb{D} \chi (* \nabla)^{\sim}\{\ll \mathbb{D}\}$	[JONES, QUANTUM, TRANSPORT]
ACAFE.46	symb	$t \int \int \mathbb{D} \pi(\chi, x) \, m = \ll \emptyset \square \gg$	[QUANTUM]
ACAFE.47	calc	$y = e^{\sim}n$ (n prime $\Rightarrow$ 'productivity number') = 20.0855	[ENTROPY]
ACAFE.48	calc	$e \cdot n^{\sim}$ Shannon entropy (text: 'neipia number times n') = 8.15485	[ENTROPY]
ACAFE.49	calc	$u = e^{\sim}n / \log 2$, $n = x \, \log x$ = 5.77078	[ZETA, ENTROPY]
ACAFE.50	differs	$x^{\sim}\{1/2 + iy\} = e^{\sim}\{x \, \log x\}$ = 1.08787 + 0.903628i vs 4	[ZETA]
ACAFE.51	holds	$\lim_{\{n \rightarrow \infty\}} (1 + 1/n)^{\sim}n = e$ = 2.71828 vs 2.71828	[OTHER]
ACAFE.52	differs	$C = 2e$, $C = e^{\sim}5$ (or e/5; garbled) = 5.43656 vs 148.413	[OTHER]
ACAFE.53	calc	$C = \lim_{\{x \rightarrow 2\}} e^{\sim}\{x \, \log x\}/\log 2$ = 5.77078	[ENTROPY]
ACAFE.54	symb	$\int C \, \mathrm{d}x^{\sim}m = n$	[OTHER]
ACAFE.55	holds	$\int \int e^{\sim}\{-x^2-y^2\} \, \mathrm{d}x \, \mathrm{d}y = \pi$ = 3.14159 vs 3.14159	[OTHER]
ACAFE.56	calc	$S = \pi r^2$ = 12.5664	[ROT]
ACAFE.57	calc	$V = \pi \int \sin 2x \, \mathrm{d}x ^2 \, \mathrm{d}x$ = 1.2337	[ROT]
ACAFE.58	differs	$V - \int \sin x \, \mathrm{d}x = \beta(p, q)$ = -0.766299 vs 0.166667	[BETA]
ACAFE.59	symb	$\beta(p, q) = \otimes (i \hbar \nabla)^{\sim}\{\otimes L\} = \mathrm{d}/\mathrm{d}f \, F(x, y)$	[BETA, QUANTUM]
ACAFE.60	symb	$= \int \Gamma'(\gamma) \, \mathrm{d}x^{\sim}m = (x, y) \cdot (x, y) = e^{\sim}\{ix\}$	[ROT, GAMMA]
ACAFE.61	holds	$e^{\sim}\{ix\} = \cos x + i \sin x$ = 0.866025 + 0.5i vs 0.866025 + 0.5i	[ROT]
ACAFE.62	differs	$\mathrm{d}/\mathrm{d}f \, e^{\sim}\{ix\} = \sin(i \, \log x) + i \cos(i \, \log x) = n$ = -0.909297 - 0.416147i vs 0 + 2i	[ROT, ZETA]
ACAFE.63	symb	$\pi \int [\Phi/\pi r^2] \, \mathrm{d}r$	[OTHER]
ACAFE.64	symb	$t \int \int \Delta(\pi(\chi, x)) [\mathrm{Im}]$	[MANIFOLD]
ACAFE.65	symb	$\Sigma(\sigma(\mathbb{H} \times \mathbb{K}))$	[OTHER]
ACAFE.66	symb	$[\square/\nabla]_{\mu \nu}$, $\nabla + \otimes \Delta$	[OTHER]
ACAFE.67	symb	$\nabla_{\perp} i \nabla_{\perp} j (\square \times \square) \, \mathrm{d}\tau$	[OTHER]
ACAFE.68	calc	$\sqrt{(x^{\sim}m \cdot y^{\sim}m)} = 16$	[OTHER]

4. 全方程式レジストリ (2111 本) p.2/33

ACAFE. 69	symb	$\int\!\!\int \square \operatorname{cohom} D\chi \text{ [Im]}$	[MANIFOLD]
ACAFE. 70	symb	$\otimes \left[\operatorname{S} D\chi \otimes h\nu \right]$	[QUANTUM]
ACAFE. 71	calc	$\ll i\hbar\phi*H\P \gg = 0 + 3.5i$	[ZETA, QUANTUM]
ACAFE. 72	symb	$\int \Delta(\xi) d\xi$	[ZETA]
ACAFE. 73	symb	$\bigcirc\bigcirc \int (\operatorname{Im})^{\wedge}\{\nabla L\}$	[MANIFOLD, QUANTUM]
ACAFE. 74	symb	$-2 \int \nabla_{-i}\nabla_{-j}(R + \nabla\nabla f) / (\Delta(R + \Delta)) \operatorname{dm}$	[MANIFOLD]
ACAFE. 75	symb	$\Delta(F(\Delta) \times \Delta(G(\Delta))) = -(F(\Delta) \cup F(\Delta)) + (F(\Delta) \cap F(\Delta))$	[MANIFOLD]
ACAFE. 76	symb	$\Sigma \square(\nabla)[\operatorname{Im}], \nabla+, [\nabla/\square], \pi \mid \mid \int \nabla_{-i}\nabla_{-j} f \nabla d\eta \mid \mid^2 = S^{\wedge}m \times S^{\wedge}\{m-1\}$	[JONES, MANIFOLD]
ACAFE. 77	holds	$(\pi r^2) \operatorname{dr}^{\wedge}m, \partial(e^{\wedge}f + e^{\wedge}\{-f\}) / d(e^{\wedge}f - e^{\wedge}\{-f\}) = 0.761594 \operatorname{vs} 0.761594$	[ROT, JONES]
ACAFE. 78	calc	$\xi = (a^{\wedge}t - t^{\wedge}n + a) = e^{\wedge}f = 1$	[ZETA, JONES]
ACAOS. 1	holds	$\int \int e^{\wedge}\{-x^2-y^2\} \operatorname{dx} \operatorname{dy} = \pi = 3.14159 \operatorname{vs} 3.14159$	[OTHER]
ACAOS. 2	differs	$\pi^{\wedge}e = (\int e^{\wedge}\{\cos \theta + i \sin \theta\} d\theta)^{\wedge}e = 22.4592 \operatorname{vs} 147.801 - 2.60195e-16i$	[ROT]
ACAOS. 3	differs	$= (\int e^{\wedge}\{i \theta\} d\theta)^{\wedge}e = 22.4592 \operatorname{vs} 2.40346e-42 - 4.05263e-42i$	[ROT]
ACAOS. 4	calc	$e^{\wedge}f \rightarrow f = 1, x \log x = 1 = 1.76322$	[JONES, ENTROPY]
ACAOS. 5	differs	$\int e^{\wedge}\{-i \theta\} d\theta = \pi^{\wedge}e = 5.81311e-16 + 2.32815e-16i \operatorname{vs} 22.4592$	[ROT]
ACAOS. 6	calc	$\square' = 2(\sin(i x \log x) + \cos(i x \log x)) = 4.25 + 3.75i$	[ROT, OTHER]
ACAOS. 7	holds	$\square = \cos(i x \log x) - i \sin(i x \log x) = 4 + 0i \operatorname{vs} 4$	[ROT]
ACAOS. 8	differs	$\int e^{\wedge}\{-\square\} d\square = \pi^{\wedge}e = 1 \operatorname{vs} 22.4592$	[OTHER]
ACAOS. 9	differs	$x = C/\log x, C = \int (1/x^{\wedge}s) \operatorname{dx} - \log x = 2 \operatorname{vs} -0.278652$	[ZETA]
ACAOS. 10	differs	$x^{\wedge}y = 1/y^{\wedge}x = 8 \operatorname{vs} 0.111111$	[OTHER]
ACAOS. 11	differs	$\pi^{\wedge}e = \int e^{\wedge}\{-\square\} d\square = e^{\wedge}\pi \int e^{\wedge}\{\square'\} d\square' = 22.4592 \operatorname{vs} 23.1407$	[BETA]
ACAOS. 12	symb	$\square = \square' \boxtimes \Psi \rightarrow \square = \Psi \boxtimes \square'$	[OTHER]
ACAOS. 13	symb	$(1/(d/dl)) \square(H\P)\nabla, \square d/dl (H\P)\nabla$	[MANIFOLD, QUANTUM]
ACAOS. 14	symb	$\beta^{\wedge}\{\square-\square'\} / \log x = \beta^{\wedge}\{\square-\square'\}$	[BETA]
ACAOS. 15	symb	$t\!\!\!\!\!\int (\nabla/\nabla l) \square(H\P)\nabla d\nabla$	[MANIFOLD]
ACAOS. 16	symb	$= \int \pi(\chi, \square) \nabla d\nabla$	[MANIFOLD]
ACAOS. 17	symb	$= \bigvee\!\!\!\int \pi(\square) d\nabla m$	[MANIFOLD]
ACAOS. 18	holds	$e^{\wedge}\pi = e^{\wedge}\{\int \int e^{\wedge}\{-x^2-y^2\} \operatorname{dx} \operatorname{dy}\} = 23.1407 \operatorname{vs} 23.1407$	[OTHER]
ACAOS. 19	differs	$= x^{\wedge}y = 1/y^{\wedge}x = \pi^{\wedge}e \cdot (1/e^{\wedge}\pi) = 23.1407 \operatorname{vs} 0.970548$	[OTHER]
ACAOS. 20	differs	$1/e^{\wedge}\pi = 1/\int e^{\wedge}\{-\square\} d\square = 0.0432139 \operatorname{vs} 0.999999$	[OTHER]
ACAOS. 21	differs	$e^{\wedge}\pi = \int e^{\wedge}\{-\square\} d\square \int e^{\wedge}\{\square'\} d\square' = 23.1407 \operatorname{vs} 1$	[OTHER]
ACAOS. 22	differs	$\pi^{\wedge}e = e^{\wedge}\pi \int e^{\wedge}\{\square'\} d\square' = 22.4592 \operatorname{vs} 23.1407$	[OTHER]
ACAOS. 23	differs	$e^{\wedge}\pi = \pi^{\wedge}e \int e^{\wedge}\{\square'\} d\square' = 23.1407 \operatorname{vs} 22.4592$	[OTHER]
ACAOS. 24	differs	$\pi^{\wedge}e = (\int e^{\wedge}\{-(\cos \theta + i \sin \theta)\} d\theta)^{\wedge}e = 22.4592 \operatorname{vs} 147.801 + 1.49055e-14i$	[ROT]
ACAOS. 25	differs	$\pi^{\wedge}e = 1/e^{\wedge}\pi = 1/\int e^{\wedge}\{-\square\} d\square = 22.4592 \operatorname{vs} 0.0432139$	[OTHER]
ACAOS. 26	calc	$\square' = 2(\sin(i x \log x) + \cos(i x \log x)) \text{ (repeat)} = 4.25 + 3.75i$	[ROT]
ACAOS. 27	holds	$\square = \cos(i x \log x) - i \sin(i x \log x) \text{ (repeat)} = 4 + 0i \operatorname{vs} 4$	[ROT]
ACAOS. 28	differs	$\int e^{\wedge}\{-\square\} d\square = \pi^{\wedge}e \text{ (repeat)} = 1 \operatorname{vs} 22.4592$	[OTHER]
ACAOS. 29	differs	$\log e^{\wedge}\pi = \log (\pi^{\wedge}e \int e^{\wedge}\{\square'\} d\square')^{\wedge}2 = 3.14159 \operatorname{vs} 6.2234$	[ZETA]
ACAOS. 30	differs	$\pi = \square/\square' = 3.14159 \operatorname{vs} 0.529183 - 0.466926i$	[OTHER]
ACAOS. 31	symb	$\pi = e^{\wedge}\beta \int \beta \operatorname{dx}^{\wedge}m$	[BETA, MANIFOLD]
ACAOS. 32	calc	$\pi = 1/\log x = \sqrt{g} = 1.3748$	[ZETA]
ACAOS. 33	holds	$= 1 = \pi r^2, r = 1/\sqrt{\pi} = 1 \operatorname{vs} 1$	[ROT]
ACAOS. 34	differs	$1/\sqrt{\pi} \pi = \int (1/\log x) \operatorname{dx}, \pi = \int \log x \operatorname{dx}^{\wedge}m = 0.56419 \operatorname{vs} 1.11842$	[ZETA, ENTROPY]
ACAOS. 35	symb	$\pi(\chi, x) = [i \pi(\chi, x), f(x)]$	[MANIFOLD]
ACAOS. 36	calc	$i \pi(\chi, x) \circ f(x) = i \int x \log x \operatorname{dx}, f(x) \circ \pi(\chi, x) = \int 1/(x \log x) \operatorname{dx} = 0 + 2.09726i$	[ENTROPY]
ACAOS. 37	symb	$\pi(\chi, x) = [i \pi(\chi, x), f(x)] \text{ (repeat)}$	[MANIFOLD]
ACAOS. 38	calc	$\pi(\chi, x) = \int x \log x \operatorname{dx} = 0.636294$	[ENTROPY]
ACAOS. 39	calc	$\pi(\chi, x) = \int 1/(x \log x) \operatorname{dx} = 0.693147$	[ENTROPY]
ACAOS. 40	symb	$\pi(\chi, x) = i \pi(\chi, x), i \int x \log x \operatorname{dx} = \int 1/(x \log x) \operatorname{dx}$	[ENTROPY]
ACAOS. 41	symb	$\int 1/(x \log x) \operatorname{dx} = i \int x \log x \operatorname{dx} - \int 1/(x \log x) \operatorname{dx}$	[ENTROPY]
ACAOS. 42	differs	$\int \int 1/(x \log x)^2 \operatorname{dx}^{\wedge}m = (1/2) i^{\wedge}3 = 0.281002 \operatorname{vs} 0 - 0.5i$	[OTHER]
ACAOS. 43	calc	$d/df F(x, y) = d/df \int \int 1/(x \log x)^2 \operatorname{dx}^{\wedge}m + d/df \int \int 1/(y \log y)^{\wedge}\{1/2\} \operatorname{dy}^{\wedge}m = 1.4353$	[MANIFOLD]
ACAOS. 44	holds	$i = x^{\wedge}\{90^{\circ}\}, x \sin 90^{\circ} = i = 6.12323e-17 + 1i \operatorname{vs} 0 + 1i$	[ROT]
ACAOS. 45	holds	$i = x^{\wedge}\{1/2\}, x = -1 = 6.12323e-17 + 1i \operatorname{vs} 0 + 1i$	[ROT]
ACAOS. 46	symb	$\pi(\chi, x) f(x) = i \int 1/(x \log x) \circ f(x) \operatorname{dx} = i \int x \log x \operatorname{dx}$	[ENTROPY]
ACAOS. 47	symb	$f(x) \pi(\chi, x) = f(x) \int 1/(x \log x) \operatorname{dx} = \int 1/(x \log x) \operatorname{dx}$	[ENTROPY]
ACAOS. 48	symb	$i y = x \sin 90^{\circ}$	[ROT]
ACAOS. 49	holds	$y = i, x = 1, i x = y = 0 + 1i \operatorname{vs} 0 + 1i$	[ROT]
ACAOS. 50	differs	$i \sin 90^{\circ} = -1 = 0 + 1i \operatorname{vs} -1 + 0i$	[ROT]
ACAOS. 51	differs	$1 \cdot \sin 90^{\circ} = i = 1 + 0i \operatorname{vs} 0 + 1i$	[ROT]
ACAOS. 52	differs	$\pi^{\wedge}e = e^{\wedge}\pi = 22.4592 \operatorname{vs} 23.1407$	[ZETA]
ACAOS. 53	symb	$[^{\wedge}\wedge V] : \chi \rightarrow ^{\wedge}\wedge\square]^{\wedge}\{\ll(p, q)\}$	[QUANTUM]
ACAOS. 54	symb	$= [^{\wedge}\wedge V] : y \rightarrow ^{\wedge}\wedge\square]^{\wedge}\{\ll\beta(\square, \square')\}$	[BETA, QUANTUM]
ACAOS. 55	symb	$\vee : \chi \rightarrow H\P D \rightarrow \otimes i\hbar \otimes x1 \otimes x2 \cdots$	[QUANTUM]
ACAOS. 56	symb	$\Phi(\delta(H\P))\nabla = \!\!\!\int\!\!\!\int \operatorname{cohom} D\chi(M)$	[MANIFOLD, QUANTUM]
ACAOS. 57	calc	$= \lim_{\{n \rightarrow \infty\}} (\operatorname{nPr} f(x)^{\wedge}n g(x)^{\wedge}\{n-r\}) \nabla = 1.18519$	[OTHER]
ACAOS. 58	symb	$\otimes(i\hbar\nabla)^{\wedge}\{\otimes L\} = ^{\wedge}\wedge H\P^{\wedge}\{i\hbar \operatorname{ih} \cdots\}$	[QUANTUM]

4. 全方程式レジストリ (2111 本) p.3/33

ACA0S.59	symb	$=H\Psi^{\wedge}\{i\hbar'\}$	[QUANTUM]
ACA0S.60	calc	$=i\hbar^{\wedge}\{i\hbar'\}\quad=0.20788+0i$	[QUANTUM]
ACA0S.61	calc	$=e^{\wedge}\{H\Psi\log H\Psi\}\quad=14$	[QUANTUM, ENTROPY]
ACAT.1	calc	$E(\sigma)=K(\sigma)\otimes H(\sigma)\quad=15$	[MANIFOLD, QUANTUM]
ACAT.2	differs	$d/d\gamma\int\Gamma=d/d\gamma\int\int1/(x\log x)^2dx^{\wedge}m+d/d\gamma\int\int1/(y\log y)^{\wedge}\{1/2\}dy^{\wedge}m\quad=0.422784\text{ vs }1.4353$	[GAMMA]
ACAT.3	symb	$H\Psi=\otimes(i\hbar)\nabla^{\wedge}\{\otimes L\}$	[QUANTUM]
ACAT.4	holds	$d/dV\int z^{\wedge}n\,dV=x^{\wedge}n+y^{\wedge}n\quad=25\text{ vs }25$	[MANIFOLD]
ACAT.5	holds	$d/dt\,g_{_ij}(t)=-2\,R_{_ij}\quad=0.5\text{ vs }0.5$	[MANIFOLD, TRANSPORT]
ACAT.6	symb	$\Box*=-2\cdot1/(R+\Delta)\, g_{_ij}+(R+\nabla_{_ij}) \,T_{_ij}+R_{_ij}$	[MANIFOLD]
ACAT.7	calc	$L(s)=\sqrt{x}\,e^{\wedge}\{i\,x\log x\}\quad=0.259447+1.39021i$	[ROT, ZETA]
ACAT.8	calc	$L(s)=(\int\Gamma(s)\int\int e^{\wedge}\{i\,x\log x\}\,)\,dx^{\wedge}m\quad=0.741545+0.545671i$	[GAMMA, ZETA]
ACAT.9	calc	$1/L(s)=\zeta(1-s)\,\zeta(s)\,\zeta(1+s)\cdots\quad=-0.164775$	[ZETA, TRANSPORT]
ACAT.10	symb	$=\int\mathbb{C}\,dx^{\wedge}m$	[OTHER]
ACAT.11	calc	$ ds^2 =8\pi G\,(\,p/c^3+V/S\,)\quad=1.67743e-09$	[SR, MANIFOLD, TRANSPORT]
ACAT.12	symb	$ ds^2 =H\Psi=(1/i)\,[H\Psi,\Psi H]$	[QUANTUM, TRANSPORT]
ACAT.13	symb	$ h_{_}\lambda ^2=H\Psi$	[MANIFOLD, ENTROPY]
ACAT.14	differs	$\int2\sin x\cos x\,dx\int e^{\wedge}\{2\sin x\cos x\}\,dx=\beta(p,q)\quad=3.10438\text{ vs }1$	[ROT, BETA]
ACAT.15	symb	$e^{\wedge}\beta\cdot\Box\,e^{\wedge}\{x\log x\}=\otimes(i\hbar\nabla)^{\wedge}\{\otimes L\}$	[BETA, QUANTUM]
ACAT.16	holds	$\int\int e^{\wedge}\{-x^2-y^2\}\,dx\,dy=\pi\quad=3.14159\text{ vs }3.14159$	[OTHER]
ACAT.17	differs	$\pi^{\wedge}e=(\int e^{\wedge}\{\cos\theta+i\sin\theta\}\,d\theta)^{\wedge}e\quad=22.4592\text{ vs }147.801-2.60195e-16i$	[ROT]
ACAT.18	differs	$=(\int e^{\wedge}\{i\theta\}\,d\theta)^{\wedge}e\quad=22.4592\text{ vs }2.40346e-42-4.05263e-42i$	[ROT]
ACAT.19	calc	$e^{\wedge}f\rightarrow f=1\,,\,x\log x=1\quad=1.76322$	[JONES, ENTROPY]
ACAT.20	differs	$\int e^{\wedge}\{-i\theta\}\,d\theta=\pi^{\wedge}e\quad=5.81311e-16+2.32815e-16i\text{ vs }22.4592$	[ROT]
ACAT.21	calc	$\Box'=2(\sin(i\,x\log x)+\cos(i\,x\log x)\,)\quad=4.25+3.75i$	[ROT, OTHER]
ACAT.22	holds	$\Box=\cos(i\,x\log x)-i\sin(i\,x\log x)\quad=4+0i\text{ vs }4$	[ROT]
ACAT.23	differs	$\int e^{\wedge}(-\Box)\,d\Box=\pi^{\wedge}e\quad=1\text{ vs }22.4592$	[OTHER]
ACAT.24	differs	$x=C/\log x\,,\,C=\int(1/x^{\wedge}s)\,dx-\log x\quad=2\text{ vs }-0.278652$	[ZETA]
ACAT.25	differs	$x^{\wedge}y=1/y^{\wedge}x\quad=8\text{ vs }0.111111$	[OTHER]
ACAT.26	differs	$\pi^{\wedge}e=\int e^{\wedge}(-\Box)\,d\Box=e^{\wedge}\pi\int e^{\wedge}\{\Box'\}\,d\Box'\quad=22.4592\text{ vs }23.1407$	[BETA]
ACAT.27	symb	$\Box=\Box'\,\boxtimes\Psi\rightarrow\Box=\Psi\,\boxtimes\Box'$	[OTHER]
ACAT.28	symb	$(1/(d/dl))\,\Box(H\Psi)\nabla\,,\,\Box\,d/dl\,(H\Psi)\nabla$	[MANIFOLD, QUANTUM]
ACAT.29	symb	$\beta^{\wedge}\{\Box-\Box'\}\,/\log x=\beta^{\wedge}\{\Box-\Box'\}$	[BETA]
ACAT.30	symb	$t\!\!\!\!\!\int\!\!\!\!\!\! (\nabla/\nabla l)\,\Box(H\Psi)\nabla\,d\nabla$	[MANIFOLD]
ACAT.31	symb	$=\int\pi(\chi,\Box)\,\nabla\,d\nabla$	[MANIFOLD]
ACAT.32	symb	$=\bigvee\!\!\!\!\!\int\!\!\!\!\!\pi(\Box)\,d\nabla m$	[MANIFOLD]
ACAT.33	holds	$e^{\wedge}\pi=e^{\wedge}\{\int\int e^{\wedge}\{-x^2-y^2\}\,dx\,dy\}\quad=23.1407\text{ vs }23.1407$	[OTHER]
ACAT.34	differs	$=x^{\wedge}y=1/y^{\wedge}x=\pi^{\wedge}e\cdot(1/e^{\wedge}\pi)\quad=23.1407\text{ vs }0.970548$	[OTHER]
ACAT.35	differs	$1/e^{\wedge}\pi=1/\int e^{\wedge}(-\Box)\,d\Box\quad=0.0432139\text{ vs }0.999999$	[OTHER]
ACAT.36	differs	$e^{\wedge}\pi=\int e^{\wedge}(-\Box)\,d\Box\int e^{\wedge}\{\Box'\}\,d\Box'\quad=23.1407\text{ vs }1$	[OTHER]
ACAT.37	differs	$\pi^{\wedge}e=e^{\wedge}\pi\int e^{\wedge}\{\Box'\}\,d\Box'\quad=22.4592\text{ vs }23.1407$	[OTHER]
ACAT.38	differs	$e^{\wedge}\pi=\pi^{\wedge}e\int e^{\wedge}\{\Box'\}\,d\Box'\quad=23.1407\text{ vs }22.4592$	[OTHER]
ACAT.39	differs	$\pi^{\wedge}e=(\int e^{\wedge}\{-(\cos\theta+i\sin\theta)\}\,d\theta)^{\wedge}e\quad=22.4592\text{ vs }147.801+1.49055e-14i$	[ROT]
ACAT.40	differs	$\pi^{\wedge}e=1/e^{\wedge}\pi=1/\int e^{\wedge}(-\Box)\,d\Box\quad=22.4592\text{ vs }0.0432139$	[OTHER]
ACAT.41	calc	$\Box'=2(\sin(i\,x\log x)+\cos(i\,x\log x)\,)\,(\text{repeat})\quad=4.25+3.75i$	[ROT]
ACAT.42	holds	$\Box=\cos(i\,x\log x)-i\sin(i\,x\log x)\,(\text{repeat})\quad=4+0i\text{ vs }4$	[ROT]
ACAT.43	differs	$\int e^{\wedge}(-\Box)\,d\Box=\pi^{\wedge}e\,(\text{repeat})\quad=1\text{ vs }22.4592$	[OTHER]
ACAT.44	calc	$\Box'=-2i\cosh dV\,(\text{pypdf}'\cos\text{hdV}')\quad=0-3.08616i$	[ROT, OTHER]
ACAT.45	differs	$\Box'=1/\Box\quad=4.25+3.75i\text{ vs }0.25+0i$	[OTHER]
ACAT.46	symb	$[^{\wedge}\vee :\chi\rightarrow^{\wedge}\Box]^{-}\{\ll(p,q)\}$	[QUANTUM]
ACAT.47	symb	$=[^{\wedge}\vee :y\rightarrow^{\wedge}\Box]^{-}\{\ll\beta(\Box,\Box')\}$	[BETA, QUANTUM]
ACAT.48	symb	$\vee :\chi\rightarrow H\Psi\mathcal{D}\rightarrow\otimes i\hbar\otimes x1\otimes x2\cdots$	[QUANTUM]
ACAT.49	symb	$\Phi(\delta(H\Psi))\nabla=\!\!\!\!\!\int\!\!\!\!\!\cohom\,D\chi(M)$	[MANIFOLD, QUANTUM]
ACAT.50	calc	$=\lim_{\{n\rightarrow\infty\}}(\,nPr\,f(x)^{\wedge}n\,g(x)^{\wedge}\{n-r\}\,)\,\nabla\quad=1.18519$	[OTHER]
ACAT.51	symb	$\otimes(i\hbar\nabla)^{\wedge}\{\otimes L\}=\text{\textasciitilde}\text{\textasciitilde}H\Psi^{\wedge}\{i\hbar\,i\hbar\cdots\}$	[QUANTUM]
ACAT.52	symb	$=H\Psi^{\wedge}\{i\hbar'\}$	[QUANTUM]
ACAT.53	calc	$=i\hbar^{\wedge}\{i\hbar'\}\quad=0.20788+0i$	[QUANTUM]
ACAT.54	calc	$=e^{\wedge}\{H\Psi\log H\Psi\}\quad=14$	[QUANTUM, ENTROPY]
ACATS.1	symb	$g_{_ij}[_{D\chi}]\,(\Box:x\vdash y)\rightarrow g_{_ij}[_{D\chi}]\,x\cdot y/\sim$	[MANIFOLD]
ACATS.2	symb	$x\cdot y/\Gamma=-2\int(R+\nabla_{_i}\nabla_{_j})/(R+\Delta)\,e^{\wedge}\{-f\}\,dV$	[GAMMA, MANIFOLD]
ACATS.3	symb	$\pi(\chi,x)=[\,i\pi(\chi,x),\,f(x)\,]$	[MANIFOLD]
ACATS.4	symb	$L(\Delta x) :x\vdash y\rightarrow g_{_ij}[_{D\chi}]\,l l'\cdot g g'/\sim$	[ZETA, MANIFOLD]
ACATS.5	symb	$L l'\cdot G g'/\Gamma=\beta(p,q)$	[GAMMA, ZETA, BETA]
ACATS.6	calc	$L[f(x):g(x)]= (f(x)+g(x))/2\geq\sqrt{(f(x)\cdot g(x))}\quad=0.21042$	[OTHER]
ACATS.7	symb	$\nabla\cdot\Delta=g_{_}\mu\,\nu(x)\,x\,g^{\wedge}\{-1\}_{_}\mu\,\nu(x)=e$	[OTHER]
ACATS.8	holds	$U^{\wedge}\{-1\}\,E\,U=[[1,\,0],\,[0,\,-1]]\quad=3.14018e-16\text{ vs }0$	[QUANTUM, TRANSPORT]
ACATS.9	calc	$H\Psi=\otimes(i\hbar\nabla)^{\wedge}\{\otimes L\}=e^{\wedge}\{iH\Psi-iH\Psi\}\quad=1+0i$	[QUANTUM]
ACATS.10	calc	$e^{\wedge}\{-iH\Psi\}\,f(x)\,e^{\wedge}\{iH\Psi\}=E_{_}n\quad=-0.416147-0.909297i$	[QUANTUM]
ACATS.11	symb	$E_{_}n=K_{_}n\times H_{_}n$	[MANIFOLD]

4. 全方程式レジストリ (2111 本) p.4/33

ACATS. 12	differs	$d/d\gamma \quad \Gamma = e^{-\{f\}} + e^{\{f\}} = i \sin(i \ x \ \log \ x) \quad = 4.25 + 0i \ \text{vs} \ -1.875 + 0i$	[ROT, GAMMA]
AGBZ. 1	symb	$\partial/\partial f \ F(x) = \iint \text{cohom} \ D_k(x) \ [Im]$	[MANIFOLD]
AGBZ. 2	symb	$\partial/\partial f \ F = t \iint \text{cohom} \ D_k(x)^{\wedge\{\llbracket p\rrbracket\}}$	[MANIFOLD]
AGBZ. 3	symb	$\oint r \ dx^m = 0(x,y)$	[OTHER]
AGBZ. 4	symb	$t \iiint D(\chi,x) \ \text{Hom}[D^2 \ \phi]^{\wedge\{\llbracket p\rrbracket\}} \equiv \text{vol}(V/S)$	[MANIFOLD]
AGBZ. 5	symb	$\partial/\partial f \ F(x) = t \iint \text{cohom} \ D_k(x)^{\wedge\{\llbracket p\rrbracket\}}$	[MANIFOLD]
AGBZ. 6	symb	$d/df \ F = (F)^{\wedge\{f'\}} \ , \ \int F \ dx^m = (F)^{\wedge f}$	[MANIFOLD]
AGBZ. 7	symb	$d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m = (\ \int \int 1/(x \ \log \ x)^2 \ dx^m)^{\wedge\{f'\}}$	[MANIFOLD]
AGBZ. 8	calc	$f' = 2x(\ \log \ x + \log(x+1) \) \quad = 7.16704$	[OTHER]
AGBZ. 9	differs	$\int \int F \ dx^m = (\ 1/(x \ \log \ x)^2)^{\wedge\{ \int 2x(\log \ x + \log(x+1)) \ dx\}} = e^{-\{f\}} \quad = 0.0701041 \ \text{vs} \ 0.0171043$	[MANIFOLD]
AGBZ. 10	differs	$d/df \ F = (\ 1/(x \ \log \ x)^2)^{\wedge\{2x(\log \ x + \log(x+1))\}} = e^{-\{f\}} \quad = 0.00926047 \ \text{vs} \ 58.4648$	[MANIFOLD]
AGBZ. 11	differs	$\log(x \ \log \ x) \geq 2 \ (y \ \log \ y)^{\wedge\{1/2\}} \quad = 0.00226757 \ \text{vs} \ 1$	[ENTROPY]
AGBZ. 12	differs	$\log(x \ \log \ x) \geq 2 \ \sqrt{(y \ \log \ y)} \quad = 0.00226757 \ \text{vs} \ 1$	[ENTROPY]
AGBZ. 13	symb	$\pi(\chi,x) = [\ i \ \pi(\chi,x), \ f(x) \]$	[MANIFOLD]
AGBZ. 14	symb	$\int 1/(x \ \log \ x) \ dx = i \ \int 1/(x \ \log \ x) \ dx^N + N \ i \ \int 1/(x \ \log \ x) \ dx$	[ENTROPY]
AGBZ. 15	symb	$\int 1/(x \ \log \ x) \ dx = i \ \int x \ \log \ x \ dx - \int 1/(x \ \log \ x) \ dx$	[ENTROPY]
AGBZ. 16	symb	$\int \int 1/(x \ \log \ x)^2 \ dx^m = i \ (1/2) \ x^2$	[OTHER]
AGBZ. 17	symb	$\int \int 1/(x \ \log \ x)^2 \ dx^m = i \ \iint_M dx^m \leq (1/2) \ i + x^2$	[OTHER]
AGBZ. 18	differs	$E = -(1/2) \ m \ v^2 + m \ c^2 \quad = 0.995 \ \text{vs} \ 1.00504$	[SR]
AGBZ. 19	symb	$\lim_{x \rightarrow \infty} \int \int 1/(x \ \log \ x)^2 \ dx^m \geq (1/2) \ i$	[OTHER]
AGBZ. 20	symb	$d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m = (1/2) \ i$	[OTHER]
AGBZ. 21	symb	$d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m = (\ \int \int 1/(x \ \log \ x)^2 \ dx^m)^{\wedge\{(f)'\}}$	[OTHER]
AGBZ. 22	differs	$(f)(f)' = (f \ f)'/2 = e^{\{x \ \log \ x\}} \quad = 2.3472 \ \text{vs} \ 1.57273$	[ENTROPY]
AGBZ. 23	holds	$\Gamma = \int e^{-\{x\}} \ x^{\wedge\{1-t\}} \ dx \ , \ d/df \ F = e^{-f} \ , \ \int F \ dx^m = e^{-\{f\}} \quad = 0.886227 \ \text{vs} \ 0.886227$	[GAMMA]
AGBZ. 24	symb	$\int x^{\wedge\{1-t\}} \ dx = d/df \ F \ , \ \int e^{-\{x\}} \ dx = \int F \ dx^m$	[GAMMA]
AGBZ. 25	holds	$e^{\{i \ \theta\}} = \cos \ \theta + i \ \sin \ \theta \quad = 0.866025 + 0.5i \ \text{vs} \ 0.866025 + 0.5i$	[ROT]
AGBZ. 26	symb	$x \perp y \rightarrow x \cdot y = \vec{0} \ , \ d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m = (1/2) \ i$	[OTHER]
AGBZ. 27	symb	$d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m = (\ \int \int \dots)^{\wedge\{(f)'\}}$	[OTHER]
AGBZ. 28	differs	$1/2 + (1/2) \ i = \vec{0} \ , \ (1/2) \cdot (1/2) \ i \quad = 0.5 + 0.5i \ \text{vs} \ 0 + 0i$	[ZETA]
AGBZ. 29	differs	$A + B = \vec{0} \ , \ A \cdot B = (1/4) \ i \quad = 0.707107 + 0.25i \ \text{vs} \ 0 + 0.25i$	[ZETA]
AGBZ. 30	symb	$\tan 90 \neq 0 \ , \ ds^2 = A + B$	[OTHER]
AGBZ. 31	holds	$\sin 0 = 0 \quad = 0 \ \text{vs} \ 0$	[ZETA]
AGBZ. 32	holds	$e^{\{i \ \theta\}} = \cos \ \theta + i \ \sin \ \theta \ (\text{repeat}) \quad = 0.362358 + 0.932039i \ \text{vs} \ 0.362358 + 0.932039i$	[ROT, QUANTUM]
AGBZ. 33	differs	$d/df \ F = d/df \ \int \int 1/(x \ \log \ x)^2 \ dx^m + d/df \ \int \int 1/(y \ \log \ y)^{\wedge\{1/2\}} \ dy^m = \Gamma(x,y) \quad = 0.319472 \ \text{vs} \ 0.406006$	[GAMMA]
AGBZ. 34	symb	$d/d\gamma \quad \Gamma = (e^{-\{x\}} \ x^{\wedge\{1-t\}})^{\wedge\{\gamma'\}} \rightarrow d/d\gamma \quad \Gamma = \Gamma(\gamma)'$	[GAMMA]
AGBZ. 35	holds	$\Gamma(s) = \int e^{-\{x\}} \ x^{\wedge\{s-1\}} \ dx \ , \ \Gamma'(s) = \int e^{-\{x\}} \ x^{\wedge\{s-1\}} \ \log \ x \ dx \quad = -0.577216 \ \text{vs} \ -0.577216$	[GAMMA]
AGBZ. 36	holds	$x^{\wedge\{s-1\}} = e^{\{(s-1) \ \log \ x\}} \quad = 2.82843 \ \text{vs} \ 2.82843$	[GAMMA]
AGBZ. 37	holds	$\partial/\partial s \ x^{\wedge\{s-1\}} = \partial/\partial s \ e^{-\{x\}} \cdot e^{\{(s-1) \ \log \ x\}} = e^{-\{x\}} \cdot \log \ x \cdot e^{\{(s-1) \ \log \ x\}} = e^{-\{x\}} \cdot \log \ x \cdot x^{\wedge\{s-1\}} \quad = 0.265327 \ \text{vs} \ 0.265327$	[GAMMA]
AGBZ. 38	holds	$d/d\gamma \quad \Gamma = \Gamma(\gamma)' \quad = 0.422784 \ \text{vs} \ 0.422784$	[GAMMA]
AGBZ. 39	symb	$d/d\gamma \quad \Gamma(s) = \int 0^{\infty} e^{-\{x\}} \ x^{\wedge\{s-1\}} \ dx^{\wedge\{(\int 0^{\infty} e^{-\{x\}} x^{\wedge\{s-1\}} \ \log \ x \ dx)'\}} = \Gamma(\Gamma \ \int \log \ x \ dx)' = e^{\{-x \ \log \ x\}}$	[GAMMA]
AGBZ. 40	holds	$\Gamma(s) = \int 0^{\infty} e^{-\{x\}} \ x^{\wedge\{s-1\}} \ dx \quad = 1.32934 \ \text{vs} \ 1.32934$	[GAMMA]
AGBZ. 41	holds	$\Gamma'(s) = \int 0^{\infty} e^{-\{x\}} \ x^{\wedge\{s-1\}} \ \log \ x \ dx \quad = 1.84557 \ \text{vs} \ 1.84557$	[GAMMA]
AGBZ. 42	symb	$d/df \ F = \int x^{\wedge\{s-1\}} \ dx \ , \ \int F \ dx^m = \int e^{-\{x\}} \ dx$	[GAMMA]
AGBZ. 43	symb	$d/df \ F = F^{\wedge\{(f)'\}} \ , \ \int F \ dx^m = F^{\wedge\{(f)\}}$	[MANIFOLD]
AGBZ. 44	calc	$H\Psi = \oplus(i\hbar\nabla)^{\wedge\{\oplus L\}} = \oplus H\Psi^{\wedge\{\nabla L\}} = e^{\{x \ \log \ x\}} = x^{\wedge\{(x)'\}} \quad = 4$	[QUANTUM]
AGBZ. 45	symb	$d/df \ F = m(x)$	[OTHER]
AGBZ. 46	symb	$d/dt \ \phi(t) = \hbar = (1/(2i)) \ e^{\{i\hbar\}}$	[QUANTUM]
AGBZ. 47	symb	$(i\hbar)' = (-e^{\{i\hbar\}})' = -i \ e^{\{i\hbar\}}$	[QUANTUM]
AGBZ. 48	calc	$\phi(x) = e^{\{-i\hbar t\}} \ , \ \oplus(i\hbar\nabla)^{\wedge\{\oplus L\}} = (1/2) \ e^{\{i\hbar(-i \ e^{\{i\hbar\}})\}} \quad = 0.571918 + 0.639942i$	[QUANTUM]
AGBZ. 49	differs	$= (1/(2f))^{\wedge\{-if\}} = (1/2)^{\wedge\{-if\}} \cdot e^{\{-x \ \log \ x\}} \cdot (f)^{\wedge i} \quad = -0.932687 + 0.360687i \ \text{vs} \ -0.121749 + 0.218351i$	[GAMMA, QUANTUM]
AGBZ. 50	holds	$= \int e^{-\{x\}} \ x^{\wedge\{t-1\}} \ dx \ , \ d/d\gamma \quad \Gamma = e^{-\{x \ \log \ x\}} \quad = 3.32335 \ \text{vs} \ 3.32335$	[GAMMA]
AGBZ. 51	differs	$f = x \ , \ i = t \ , \ 1/2 = a \ , \ \oplus a^{\{-t\}} \ x \ x^{\wedge t} \ [Im] \equiv \int e^{-\{x\}} \ x^{\wedge\{t-1\}} \ dx \quad = 32 \ \text{vs} \ 1$	[GAMMA, ZETA]
AGBZ. 52	holds	$ \phi(t)\rangle_s = e^{-\{i\hbar t\}} \ \Psi\rangle_H \ , \ \hat{A}_s = \hat{A}_H(0) \quad = 0 - 0.644218i \ \text{vs} \ 0 - 0.644218i$	[QUANTUM]
AGBZ. 53	holds	$ \Psi(t)\rangle_s \rightarrow d/dt \ ; \ i \ d/dt \ \phi(t)\rangle_s = \hat{H} \ \phi(t)\rangle_s \quad = 0 - 0.841471i \ \text{vs} \ 0 - 0.841471i$	[QUANTUM]
AGBZ. 54	holds	$\langle \hat{A}(t) \rangle = \langle \Psi(t) \hat{A}(0) \ \Psi(t) \rangle \quad = 0.362358 \ \text{vs} \ 0.362358$	[QUANTUM]
AGBZ. 55	holds	$d/dt \ \hat{A} = (1/i) \ [\hat{A}, \ H] \quad = 0 - 1.39341i \ \text{vs} \ 0 - 1.39341i$	[QUANTUM]
AGBZ. 56	holds	$\hat{A}(t) = e^{\{i\hbar t\}} \ \hat{A}(0) \ e^{-\{i\hbar t\}} \quad = 0 - 0.717356i \ \text{vs} \ 0 - 0.717356i$	[QUANTUM]
AGBZ. 57	calc	$\lim_{\theta \rightarrow 0} (\sin \ \theta, \cos \ \theta) \ [[\theta, \ 1], [1, \ \theta]] \ (\cos \ \theta, \sin \ \theta)^{\wedge T} = [[1, \ 0], [0, \ -1]] \quad = 1$	[ROT]
AGBZ. 58	symb	$f^{\{-1\}}(x) \ x \ f(x) = I'_{-m} \ , \ I'_{-m} = [1, \ 0] \times [0, \ 1]$	[OTHER]
AGBZ. 59	holds	$x + y \geq \sqrt{(x \ y)} \quad = 1 \ \text{vs} \ 1$	[OTHER]
AGBZ. 60	differs	$x^{\wedge\{1/2 + iy\}} / e^{\{x \ \log \ x\}} = 1 \quad = 0.271967 + 0.225907i \ \text{vs} \ 1 + 0i$	[ZETA]
AGBZ. 61	symb	$0(x) = \nabla_i \nabla_j \ \int e^{\{(2/m) \sin \ \theta \ \cos \ \theta\}} \times N \ \text{mod}(e^{\{x \ \log \ x\}})$	[ROT]
AGBZ. 62	symb	$0(x) \ (x + \Delta f ^2)^{\wedge\{1/2\}}$	[OTHER]
AGBZ. 63	differs	$x \ \Gamma(x) = 2 \ \int \sin 2\theta ^2 \ d\theta \ , \ 0(x) = m(x) [D^2 \ \phi] \quad = 1.32934 \ \text{vs} \ 1.5708$	[ROT, GAMMA]
AGBZ. 64	differs	$i^2 = (0,1) \cdot (0,1) \ , \ a b \ \cos \ \theta = -1 \quad = -1 + 0i \ \text{vs} \ 1$	[ROT]
AGBZ. 65	symb	$E = \text{div}(E, \ E_1)$	[OTHER]
AGBZ. 66	calc	$\{f,g\}/[f,g] = i^2 \ , \ E = m \ c^2 \ , \ I' = i^2 \quad = 8.98755e+16$	[SR, QUANTUM]

4. 全方程式レジストリ (2111 本) p.5/33

AGBZ, 67	differs	$d/d\gamma \Gamma = m(x) = e^{-x \log x} = 0.422784$ vs 0.25	[GAMMA, ZETA]
AGBZ, 68	differs	$\sin(ix) = (e^{-x} + e^x)/(2i) = 0 + 1.1752i$ vs $0 - 1.54308i$	[ROT]
AGBZ, 69	differs	$d/df F = m(x) = e^f + e^{-f} = 2i \sin(i x \log x) = 4.25 + 0i$ vs $-3.75 + 0i$	[ROT, BETA]
AGMK, 1	holds	$F(\text{vertex}) - N(\text{edge}) + E(\text{face}) = 2 = 2$ vs 2	[MANIFOLD]
AGMK, 2	symb	$\Gamma^{\{ (x,y,z) \}} = \int (F - N + E) + (F - E + N) + (N - E + F) dx^m$ (operators lost in pypdf)	[GAMMA, MANIFOLD]
AGMK, 3	symb	$\partial/\partial f S \Gamma^{\{ (x,y,z) \}}$	[GAMMA]
AGMK, 4	holds	$= \int (\int (1/x^s) dx - \log x) dvol = 0.577466$ vs 0.577216	[GAMMA, ZETA]
AGMK, 5	holds	$= \int e^{-x} x^{\{1-t\}} dx^m = 0.908639$ vs 0.908639	[GAMMA]
AGMK, 6	calc	$ ds^2 = e^{-2\pi T} \phi [\eta + \bar{h}(x)] dx^\mu dx^\nu + T^2 d^2\phi = 0.0432139$	[SR, MANIFOLD, TRANSPORT]
AGMK, 7	symb	$= 0(x)$	[OTHER]
AGMK, 8	calc	$\square = (8\pi G/c^4) T_\mu \nu = 2.07655e-43$	[OTHER]
AGMK, 9	calc	$\nabla\phi = 4\pi G\rho = 4.62448e-06$	[OTHER]
AGMK, 10	holds	$= \int (\int (1/x^s) dx - \log x) dvol$ (repeat) = 0.577341 vs 0.577216	[GAMMA, ZETA]
AGMK, 11	holds	$= \int e^{-x} x^{\{1-t\}} dx^m$ (repeat) = 0.897471 vs 0.897471	[GAMMA]
AGMK, 12	holds	$= d/df F = \int \Gamma(\gamma)' dx^m = 0.422784$ vs 0.422784	[GAMMA]
AGMK, 13	calc	$= \int C dx^m = (\int \Gamma dx^m - d/d\gamma \Gamma) / (\int \Gamma dx^m + d/d\gamma \Gamma) = 0.405694$	[GAMMA]
AGMK, 14	calc	$= e^{-\theta} + e^{\{i\theta\}} = 1.45841 + 0.5i$	[ROT]
AGMK, 15	holds	$= \square = 2(\cos(i x \log x) - i \sin(i x \log x)) = 8 + 0i$ vs $8 + 0i$	[ROT]
AGMK, 16	holds	$= (e^{-f} - e^{\{f\}}) / (e^{-f} + e^{\{f\}}) = -0.462117$ vs -0.462117	[JONES]
AGMK, 17	symb	$X \rightarrow Y, Y \rightarrow X$	[OTHER]
AGMK, 18	calc	$\otimes(i\hbar\nabla)^{\{\otimes L\}} = e^{-x \log x} = 0.25$	[QUANTUM, ENTROPY]
AGMK, 19	symb	$\square \rightarrow \nabla_j : x \rightarrow y, f(x) \rightarrow g(y), f^{-\{1\}}(x) \times f(x) \rightarrow g(y)$	[OTHER]
AGMK, 20	symb	$\square \rightarrow \nabla_j : f(x) = g(y)$	[OTHER]
AGMK, 21	holds	$d/dt g_{ij}(t) = -2 R_{ij}, F_t^m = -2 \int (R + \nabla_i \nabla_j f)/(\Delta + f) dm = 0.774613$ vs 0.774597	[MANIFOLD]
AGMK, 22	symb	$d/df F = m(x)$	[OTHER]
AGMK, 23	symb	$F_j = d/df F, F_t^m = e^{-\{f\}} dV$	[MANIFOLD]
AGMK, 24	calc	$\otimes(i\hbar\nabla)^{\{\otimes L\}} = e^{-x \log x}$ (repeat) = 0.037037	[QUANTUM, ENTROPY]
AGMK, 25	holds	$\beta(p,q) = \Gamma(p)\Gamma(q)/\Gamma(p+q) = 0.196349$ vs 0.19635	[GAMMA, BETA]
AGMK, 26	symb	$\Gamma(p)\Gamma(q)/\Gamma(p+q)$ [op] 1	[BETA]
AGMK, 27	symb	$\int e^{-x} x^{\{1-t\}} dx^m = \int e^{-x} x^{\{1-t\}} dx$ [op] $e^{-\theta}$	[GAMMA]
AGMK, 28	differs	$= \Gamma\gamma' = e^{-x \log x} = -2.51058$ vs 1.37328	[GAMMA, ENTROPY]
AGMK, 29	symb	$\{f,g\} \mapsto \{d,d\}$	[OTHER]
AGZE, 1	symb	$\partial/\partial f F(x) = \iint \text{cohom } D_k(x) [Im]$	[MANIFOLD]
AGZE, 2	symb	$\partial/\partial f F = t \iint \text{cohom } D_k(x)^{\{\ll p\}}$	[MANIFOLD]
AGZE, 3	symb	$\oint r dx^m = 0(x,y)$	[OTHER]
AGZE, 4	symb	$t \iiint D(x,x) \text{Hom}[D^2\phi]^{\{\ll p\}} \equiv \text{vol}(V/S)$	[MANIFOLD]
AGZE, 5	symb	$\partial/\partial f F(x) = t \iint \text{cohom } D_k(x)^{\{\ll p\}}$	[MANIFOLD]
AGZE, 6	symb	$d/df F = (F)^{\{f'\}}, \int F dx^m = (F)^{\sim f}$	[MANIFOLD]
AGZE, 7	symb	$d/df \int \int 1/(x \log x)^2 dx^m = (\int \int 1/(x \log x)^2 dx^m)^{\{f'\}}$	[MANIFOLD]
AGZE, 8	calc	$f' = 2x(\log x + \log(x+1)) = 7.16704$	[OTHER]
AGZE, 9	differs	$\int \int F dx^m = (1/(x \log x)^2)^{\{ \int 2x(\log x + \log(x+1)) dx = e^{-\{f\}} = 0.0701041$ vs 0.0171043	[MANIFOLD]
AGZE, 10	differs	$d/df F = (1/(x \log x)^2)^{\{ 2x(\log x + \log(x+1)) \}} = e^{-\{f\}} = 0.00926047$ vs 58.4648	[MANIFOLD]
AGZE, 11	differs	$\log(x \log x) \geq 2 (y \log y)^{\{1/2\}} = 0.00226757$ vs 1	[ENTROPY]
AGZE, 12	differs	$\log(x \log x) \geq 2 \sqrt{(y \log y)} = 0.00226757$ vs 1	[ENTROPY]
AGZE, 13	symb	$\pi(x,x) = [i \pi(x,x), f(x)]$	[MANIFOLD]
AGZE, 14	symb	$\int 1/(x \log x) dx = i \int 1/(x \log x) dx^N + N i \int 1/(x \log x) dx$	[ENTROPY]
AGZE, 15	symb	$\int 1/(x \log x) dx = i \int x \log x dx - \int 1/(x \log x) dx$	[ENTROPY]
AGZE, 16	symb	$\int \int 1/(x \log x)^2 dx^m = i (1/2) x^2$	[OTHER]
AGZE, 17	symb	$\int \int 1/(x \log x)^2 dx^m = i \iint_M dx^m \leq (1/2) i + x^2$	[OTHER]
AGZE, 18	differs	$E = -(1/2) m v^2 + m c^2 = 0.995$ vs 1.00504	[SR]
AGZE, 19	symb	$\lim_{x \rightarrow \infty} \int \int 1/(x \log x)^2 dx^m \geq (1/2) i$	[OTHER]
AGZE, 20	symb	$d/df \int \int 1/(x \log x)^2 dx^m = (1/2) i$	[OTHER]
AGZE, 21	symb	$d/df \int \int 1/(x \log x)^2 dx^m = (\int \int 1/(x \log x)^2 dx^m)^{\{(f)'\}}$	[OTHER]
AGZE, 22	differs	$(f)(f)' = (f f)'/2 = e^{\{x \log x\}} = 2.3472$ vs 1.57273	[ENTROPY]
AGZE, 23	holds	$\Gamma = \int e^{-x} x^{\{1-t\}} dx, d/df F = e^f, \int F dx^m = e^{-\{f\}} = 0.886227$ vs 0.886227	[GAMMA]
AGZE, 24	symb	$\int x^{\{1-t\}} dx = d/df F, \int e^{-x} dx = \int F dx^m$	[GAMMA]
AGZE, 25	holds	$e^{\{i\theta\}} = \cos \theta + i \sin \theta = 0.866025 + 0.5i$ vs $0.866025 + 0.5i$	[ROT]
AGZE, 26	symb	$x \perp y \rightarrow x \cdot y = \vec{0}, d/df \int \int 1/(x \log x)^2 dx^m = (1/2) i$	[OTHER]
AGZE, 27	symb	$d/df \int \int 1/(x \log x)^2 dx^m = (\int \int \dots)^{\{(f)'\}}$	[OTHER]
AGZE, 28	differs	$1/2 + (1/2) i = \vec{0}, (1/2) \cdot (1/2) i = 0.5 + 0.5i$ vs $0 + 0i$	[ZETA]
AGZE, 29	differs	$A + B = \vec{0}, A \cdot B = (1/4) i = 0.707107 + 0.25i$ vs $0 + 0.25i$	[ZETA]
AGZE, 30	symb	$\tan 90 \neq 0, ds^2 = A + B$	[OTHER]
AGZE, 31	holds	$\sin 0 = 0 = 0$ vs 0	[ZETA]
AGZE, 32	holds	$e^{\{i\theta\}} = \cos \theta + i \sin \theta$ (repeat) = $0.362358 + 0.932039i$ vs $0.362358 + 0.932039i$	[ROT, QUANTUM]
AGZE, 33	differs	$d/df F = d/df \int \int 1/(x \log x)^2 dx^m + d/df \int \int 1/(y \log y)^{\{1/2\}} dy^m = \Gamma(x,y) = 0.319472$ vs 0.406006	[GAMMA]
AGZE, 34	symb	$d/d\gamma \Gamma = (e^{-x} x^{\{1-t\}})^{\{\gamma'\}} \rightarrow d/d\gamma \Gamma = \Gamma(\gamma)'$	[GAMMA]
AGZE, 35	holds	$\Gamma(s) = \int e^{-x} x^{\{s-1\}} dx, \Gamma'(s) = \int e^{-x} x^{\{s-1\}} \log x dx = -0.577216$ vs -0.577216	[GAMMA]
AGZE, 36	holds	$x^{\{s-1\}} = e^{\{(s-1) \log x\}} = 2.82843$ vs 2.82843	[GAMMA]

4. 全方程式レジストリ (2111 本) p.6/33

AGZE, 37	holds	$\partial/\partial s\ x^{s-1} = \partial/\partial s\ e^{-x}\cdot e^{-\{(s-1)\log x\}} = e^{-x}\cdot \log x\cdot e^{-\{(s-1)\log x\}} = e^{-x}\cdot \log x\cdot x^{s-1}$	= 0.265327 vs 0.265327	[GAMMA]
AGZE, 38	holds	$d/d\gamma\ \Gamma = \Gamma(\gamma)'$	= 0.422784 vs 0.422784	[GAMMA]
AGZE, 39	symb	$d/d\gamma\ \Gamma(s) = \int_0^\infty e^{-x}\ x^{s-1}\ dx\cdot\{(\int_0^\infty e^{-x}\ x^{s-1}\ \log x\ dx)'\} = \Gamma(\int\ \log x\ dx)' = e^{-x\ \log x}$		[GAMMA]
AGZE, 40	holds	$\Gamma(s) = \int_0^\infty e^{-x}\ x^{s-1}\ dx$	= 1.32934 vs 1.32934	[GAMMA]
AGZE, 41	holds	$\Gamma'(s) = \int_0^\infty e^{-x}\ x^{s-1}\ \log x\ dx$	= 1.84557 vs 1.84557	[GAMMA]
AGZE, 42	symb	$d/df\ F = \int\ x^{s-1}\ dx,\ \int\ F\ dx^m = \int\ e^{-x}\ dx$		[GAMMA]
AGZE, 43	symb	$d/df\ F = F^{\{(f)'\}},\ \int\ F\ dx^m = F^{\{(f)\}}$		[MANIFOLD]
AGZE, 44	calc	$H\Psi = \otimes(i\hbar\nabla)^{\otimes L} = \otimes H\Psi^{\{\nabla L\}} = e^{-x\ \log x} = x^{\{(x)'\}} = 4$		[QUANTUM]
AGZE, 45	symb	$d/df\ F = m(x)$		[OTHER]
AGZE, 46	symb	$d/dt\ \phi(t) = \hbar = (1/(2i))\ e^{-i\hbar}$		[QUANTUM]
AGZE, 47	symb	$(i\hbar)' = (-e^{-i\hbar})' = -i\ e^{-i\hbar}$		[QUANTUM]
AGZE, 48	calc	$\phi(x) = e^{-i\hbar t},\ \otimes(i\hbar\nabla)^{\otimes L} = (1/2)\ e^{-i\hbar(-i\ e^{-i\hbar})}\ = 0.571918 + 0.639942i$		[QUANTUM]
AGZE, 49	differs	$= (1/(2f))^{\{-if\}} = (1/2)^{\{-if\}}\cdot e^{-\{-x\ \log x\}}\cdot (f)^{-i} = -0.932687 + 0.360687i\ \text{vs}\ -0.121749 + 0.218351i$		[GAMMA, QUANTUM]
AGZE, 50	holds	$= \int\ e^{-x}\ x^{t-1}\ dx,\ d/d\gamma\ \Gamma = e^{-x\ \log x}$	= 3.32335 vs 3.32335	[GAMMA]
AGZE, 51	differs	$f = x,\ i = t,\ 1/2 = a,\ \otimes a^{\{-t\}}\ x\ x^t\ [Im] \cong \int\ e^{-x}\ x^{t-1}\ dx = 32\ \text{vs}\ 1$		[GAMMA, ZETA]
AGZE, 52	holds	$ \phi(t)\rangle_s = e^{-i\hbar t}\ \Psi\rangle_H,\ \tilde{A}_s = \tilde{A}_H(0) = 0 - 0.644218i\ \text{vs}\ 0 - 0.644218i$		[QUANTUM]
AGZE, 53	holds	$ \Psi(t)\rangle_s \rightarrow d/dt;\ i\ d/dt\ \phi(t)\rangle_s = \hbar\ \phi(t)\rangle_s = 0 - 0.841471i\ \text{vs}\ 0 - 0.841471i$		[QUANTUM]
AGZE, 54	holds	$\langle\tilde{A}(t)\rangle = \langle\Psi(t) \ \tilde{A}(0)\ \Psi(t)\rangle = 0.362358\ \text{vs}\ 0.362358$		[QUANTUM]
AGZE, 55	holds	$d/dt\ \tilde{A} = (1/i)\ [\tilde{A}, H] = 0 - 1.39341i\ \text{vs}\ 0 - 1.39341i$		[QUANTUM]
AGZE, 56	holds	$\tilde{A}(t) = e^{-i\hbar t}\ \tilde{A}(0)\ e^{-i\hbar t} = 0 - 0.717356i\ \text{vs}\ 0 - 0.717356i$		[QUANTUM]
AGZE, 57	calc	$\lim_{\theta\rightarrow 0}\ (\sin\ \theta,\ \cos\ \theta)\ [[\theta,\ 1],\ [1,\ \theta]]\ (\cos\ \theta,\ \sin\ \theta)^T = [[1,\ 0],\ [0,\ -1]] = 1$		[ROT]
AGZE, 58	symb	$f^{-1}(x)\ x\ f(x) = I^{-1}_m,\ I^{-1}_m = [1,\ 0] \times [0,\ 1]$		[OTHER]
AGZE, 59	holds	$x + y \geq \sqrt{x\ y} = 1\ \text{vs}\ 1$		[OTHER]
AGZE, 60	differs	$x^{1/2 + iy}\ /\ e^{-x\ \log x} = 1 = 0.271967 + 0.225907i\ \text{vs}\ 1 + 0i$		[ZETA]
AGZE, 61	symb	$0(x) = \nabla_i\nabla_j\ \int\ e^{-\{(2/m)\sin\ \theta\ \cos\ \theta\}} \times N\ \text{mod}(e^{-x\ \log x})$		[ROT]
AGZE, 62	symb	$0(x)\ (x + \Delta f ^2)^{1/2}$		[OTHER]
AGZE, 63	differs	$x\ \Gamma(x) = 2\ \int\ \sin\ 2\theta ^2\ d\theta,\ 0(x) = m(x)[D^2\phi] = 1.32934\ \text{vs}\ 1.5708$		[ROT, GAMMA]
AGZE, 64	differs	$i^2 = (0,1)\cdot(0,1),\ a b \ \cos\ \theta = -1 = -1 + 0i\ \text{vs}\ 1$		[ROT]
AGZE, 65	symb	$E = \text{div}(E,\ E_1)$		[OTHER]
AGZE, 66	calc	$\{f,g\}/[f,g] = i^2,\ E = m\ c^2,\ I' = i^2 = 8.98755e+16$		[SR, QUANTUM]
AGZE, 67	differs	$d/d\gamma\ \Gamma = m(x) = e^{-x\ \log x} = 0.422784\ \text{vs}\ 0.25$		[GAMMA, ZETA]
AGZE, 68	differs	$\sin(ix) = (e^{-x} + e^x)/(2i) = 0 + 1.1752i\ \text{vs}\ 0 - 1.54308i$		[ROT]
AGZE, 69	differs	$d/df\ F = m(x) = e^f + e^{-f} = 2i\ \sin(i\ x\ \log x) = 4.25 + 0i\ \text{vs}\ -3.75 + 0i$		[ROT, BETA]
AQC4, 1	symb	$(E^2\ \otimes\ E^2)\cdot (R - C\ \mathbb{C}\pm) = \otimes\ \nabla\mathbb{C}\pm$		[ZETA]
AQC4, 2	symb	$= \otimes\ \nabla\mathbb{C}\pm$		[ZETA, QUANTUM]
AQC4, 3	symb	$\partial/\partial f\ F(x) = \int\!\!\!\int\ \text{cohom}\ D_k(x)\ [Im]$		[MANIFOLD]
AQC4, 4	symb	$\partial/\partial f\ F = t\int\!\!\!\int\ \text{cohom}\ D_k(x)^{\{\llbracket p\rrbracket\}}$		[MANIFOLD]
AQC4, 5	symb	$\oint\ r\ dx^m = 0(x,y)$		[OTHER]
AQC4, 6	symb	$t\int\!\!\!\int\ D(\chi,x)\ \text{Hom}[D^2\phi]^{\{\llbracket p\rrbracket\}} \cong \text{vol}(V/S)$		[MANIFOLD]
AQC4, 7	symb	$\partial/\partial f\ F(x) = t\int\!\!\!\int\ \text{cohom}\ D_k(x)^{\{\llbracket p\rrbracket\}}$		[MANIFOLD]
AQC4, 8	symb	$d/df\ F = (F)^{\{f'\}},\ \int\ F\ dx^m = (F)^{-f}$		[MANIFOLD]
AQC4, 9	symb	$d/df\ \int\ \int\ 1/(x\ \log x)^2\ dx^m = (\int\ \int\ 1/(x\ \log x)^2\ dx^m)^{\{f'\}}$		[MANIFOLD]
AQC4, 10	calc	$f' = 2x(\log x + \log(x+1)) = 7.16704$		[OTHER]
AQC4, 11	differs	$\int\ \int\ F\ dx^m = (\int\ 1/(x\ \log x)^2)^{\{2x(\log x + \log(x+1))\}}\ dx = e^{-\{f\}} = 0.0701041\ \text{vs}\ 0.0171043$		[MANIFOLD]
AQC4, 12	differs	$d/df\ F = (\int\ 1/(x\ \log x)^2)^{\{2x(\log x + \log(x+1))\}} = e^{\{f\}} = 0.00926047\ \text{vs}\ 58.4648$		[MANIFOLD]
AQC4, 13	differs	$\log(x\ \log x) \geq 2\ (y\ \log y)^{1/2} = 0.00226757\ \text{vs}\ 1$		[ENTROPY]
AQC4, 14	differs	$\log(x\ \log x) \geq 2\ \sqrt{y\ \log y} = 0.00226757\ \text{vs}\ 1$		[ENTROPY]
AQC4, 15	symb	$\pi(\chi,x) = [i\ \pi(\chi,x),\ f(x)]$		[MANIFOLD]
AQC4, 16	symb	$\int\ 1/(x\ \log x)\ dx = i\ \int\ 1/(x\ \log x)\ dx^N + N\ i\ \int\ 1/(x\ \log x)\ dx$		[ENTROPY]
AQC4, 17	symb	$\int\ 1/(x\ \log x)\ dx = i\ \int\ x\ \log x\ dx - \int\ 1/(x\ \log x)\ dx$		[ENTROPY]
AQC4, 18	symb	$\int\ \int\ 1/(x\ \log x)^2\ dx^m = i\ (1/2)\ x^2$		[OTHER]
AQC4, 19	symb	$\int\ \int\ 1/(x\ \log x)^2\ dx^m = i\ \int\!\!\!\int\ M\ dx^m \leq (1/2)\ i + x^2$		[OTHER]
AQC4, 20	differs	$E = -(1/2)\ m\ v^2 + m\ c^2 = 0.995\ \text{vs}\ 1.00504$		[SR]
AQC4, 21	symb	$\lim_{x\rightarrow\infty}\ \int\ \int\ 1/(x\ \log x)^2\ dx^m \geq (1/2)\ i$		[OTHER]
AQC4, 22	symb	$d/df\ \int\ \int\ 1/(x\ \log x)^2\ dx^m = (1/2)\ i$		[OTHER]
AQC4, 23	symb	$d/df\ \int\ \int\ 1/(x\ \log x)^2\ dx^m = (\int\ \int\ 1/(x\ \log x)^2\ dx^m)^{\{(f)'\}}$		[OTHER]
AQC4, 24	differs	$(f)(f)' = (f\ f)'/2 = e^{-x\ \log x} = 2.3472\ \text{vs}\ 1.57273$		[ENTROPY]
AQC4, 25	holds	$\Gamma = \int\ e^{-x}\ x^{\{1-t\}}\ dx,\ d/df\ F = e^{-f},\ \int\ F\ dx^m = e^{-\{f\}} = 0.886227\ \text{vs}\ 0.886227$		[GAMMA]
AQC4, 26	symb	$\int\ x^{\{1-t\}}\ dx = d/df\ F,\ \int\ e^{-x}\ dx = \int\ F\ dx^m$		[GAMMA]
AQC4, 27	holds	$e^{-i\theta} = \cos\ \theta + i\ \sin\ \theta = 0.866025 + 0.5i\ \text{vs}\ 0.866025 + 0.5i$		[ROT]
AQC4, 28	symb	$x\perp y\rightarrow x\cdot y = \vec{0},\ d/df\ \int\ \int\ 1/(x\ \log x)^2\ dx^m = (1/2)\ i$		[OTHER]
AQC4, 29	symb	$d/df\ \int\ \int\ 1/(x\ \log x)^2\ dx^m = (\int\ \int\ \dots)^{\{(f)'\}}$		[OTHER]
AQC4, 30	differs	$1/2 + (1/2)\ i = \vec{0},\ (1/2)\cdot(1/2)\ i = 0.5 + 0.5i\ \text{vs}\ 0 + 0i$		[ZETA]
AQC4, 31	differs	$A + B = \vec{0},\ A\cdot B = (1/4)\ i = 0.707107 + 0.25i\ \text{vs}\ 0 + 0.25i$		[ZETA]
AQC4, 32	symb	$\tan\ 90 \neq 0,\ ds^2 = A + B$		[OTHER]
AQC4, 33	holds	$\sin\ 0 = 0 = 0\ \text{vs}\ 0$		[ZETA]
AQC4, 34	holds	$e^{-i\theta} = \cos\ \theta + i\ \sin\ \theta\ (\text{repeat}) = 0.362358 + 0.932039i\ \text{vs}\ 0.362358 + 0.932039i$		[ROT, QUANTUM]

4. 全方程式レジストリ (2111 本) p.7/33

AQC4.35	differs	$d/d\mathfrak{f}\;F=d/d\mathfrak{f}\;\int\int1/(x\log x)^2\;dx^m+d/d\mathfrak{f}\;\int1/(y\log y)^{\{1/2\}}\;dy^m=\Gamma(x,y)\quad=0.319472\;\text{vs}\;0.406006$	[GAMMA]
AQC4.36	symb	$d/d\gamma\;\Gamma=(\mathfrak{e}^{\{-x\}}\;x^{\{1-t\}})^{\{\gamma'\}}\rightarrow d/d\gamma\;\Gamma=\Gamma(\gamma)'$	[GAMMA]
AQC4.37	holds	$\Gamma(s)=\int\mathfrak{e}^{\{-x\}}\;x^{\{s-1\}}\;dx,\;\Gamma'(s)=\int\mathfrak{e}^{\{-x\}}\;x^{\{s-1\}}\;\log x\;dx\quad=-0.577216\;\text{vs}\;-0.577216$	[GAMMA]
AQC4.38	holds	$x^{\{s-1\}}=\mathfrak{e}^{\{(s-1)\log x\}}\quad=2.82843\;\text{vs}\;2.82843$	[GAMMA]
AQC4.39	holds	$\partial/\partial s\;x^{\{s-1\}}=\partial/\partial s\;\mathfrak{e}^{\{-x\}}\cdot\mathfrak{e}^{\{(s-1)\log x\}}=\mathfrak{e}^{\{-x\}}\cdot\log x\cdot\mathfrak{e}^{\{(s-1)\log x\}}=\mathfrak{e}^{\{-x\}}\cdot\log x\cdot x^{\{s-1\}}\quad=0.265327\;\text{vs}\;0.265327$	[GAMMA]
AQC4.40	holds	$d/d\gamma\;\Gamma=\Gamma(\gamma)'\quad=0.422784\;\text{vs}\;0.422784$	[GAMMA]
AQC4.41	symb	$d/d\gamma\;\Gamma(s)=\int0^\infty\mathfrak{e}^{\{-x\}}\;x^{\{s-1\}}\;dx^{\{(\int0^\infty\mathfrak{e}^{\{-x\}}x^{\{s-1\}}\;\log x\;dx)'\}}=\Gamma(\int\log x\;dx)'=\mathfrak{e}^{\{-x\log x\}}$	[GAMMA]
AQC4.42	holds	$\Gamma(s)=\int0^\infty\mathfrak{e}^{\{-x\}}\;x^{\{s-1\}}\;dx\quad=1.32934\;\text{vs}\;1.32934$	[GAMMA]
AQC4.43	holds	$\Gamma'(s)=\int0^\infty\mathfrak{e}^{\{-x\}}\;x^{\{s-1\}}\;\log x\;dx\quad=1.84557\;\text{vs}\;1.84557$	[GAMMA]
AQC4.44	symb	$d/d\mathfrak{f}\;F=\int x^{\{s-1\}}\;dx,\;\int F\;dx^m=\int\mathfrak{e}^{\{-x\}}\;dx$	[GAMMA]
AQC4.45	symb	$d/d\mathfrak{f}\;F=F^{\{(f)'\}},\;\int F\;dx^m=F^{\{(f)\}}$	[MANIFOLD]
AQC4.46	calc	$H\Psi=\mathfrak{e}(i\hbar\nabla)^{\{\mathfrak{e}L\}}=\mathfrak{e}\;H\Psi^{\{\nabla L\}}=\mathfrak{e}^{\{x\log x\}}=x^{\{(x)'\}}\quad=4$	[QUANTUM]
AQC4.47	symb	$d/d\mathfrak{f}\;F=m(x)$	[OTHER]
AQC4.48	symb	$d/dt\;\phi(t)=\hbar=(1/(2i))\;\mathfrak{e}^{\{i\hbar\}}$	[QUANTUM]
AQC4.49	symb	$(i\hbar)'=(-\mathfrak{e}^{\{i\hbar\}})'=-i\;\mathfrak{e}^{\{i\hbar\}}$	[QUANTUM]
AQC4.50	calc	$\phi(x)=\mathfrak{e}^{\{-i\hbar t\}},\;\mathfrak{e}(i\hbar\nabla)^{\{\mathfrak{e}L\}}=(1/2)\;\mathfrak{e}^{\{i\hbar(-i\;\mathfrak{e}^{\{i\hbar\}})\}}\quad=0.571918+0.639942i$	[QUANTUM]
AQC4.51	differs	$= (1/(2\mathfrak{f}))^{\{-i\mathfrak{f}\}}=(1/2)^{\{-i\mathfrak{f}\}}\cdot\mathfrak{e}^{\{-x\log x\}}\cdot(f)^{-i}\quad=-0.932687+0.360687i\;\text{vs}\;-0.121749+0.218351i$	[GAMMA, QUANTUM]
AQC4.52	holds	$=\int\mathfrak{e}^{\{-x\}}\;x^{\{t-1\}}\;dx,\;d/d\gamma\;\Gamma=\mathfrak{e}^{\{-x\log x\}}\quad=3.32335\;\text{vs}\;3.32335$	[GAMMA]
AQC4.53	differs	$f=x,\;i=t,\;1/2=a,\;\mathfrak{e}\;a^{\{-t\}}\;x\;x^t\;[Im]\equiv\int\mathfrak{e}^{\{-x\}}\;x^{\{t-1\}}\;dx\quad=32\;\text{vs}\;1$	[GAMMA, ZETA]
AQC4.54	holds	$ \phi(t)\rangle_s=\mathfrak{e}^{\{-i\hbar t\}}\; \Psi\rangle_H,\;\hat{A}_s=\hat{A}_H(0)\quad=0-0.644218i\;\text{vs}\;0-0.644218i$	[QUANTUM]
AQC4.55	holds	$ \Psi(t)\rangle_s\rightarrow d/dt\;i\;d/dt\; \phi(t)\rangle_s=\hat{H}\; \phi(t)\rangle_s\quad=0-0.841471i\;\text{vs}\;0-0.841471i$	[QUANTUM]
AQC4.56	holds	$\langle\hat{A}(t)\rangle=\langle\Psi(t) \;\hat{A}(0)\; \Psi(t)\rangle\quad=0.362358\;\text{vs}\;0.362358$	[QUANTUM]
AQC4.57	holds	$d/dt\;\hat{A}=(1/i)\;[\hat{A},\;H]\quad=0-1.39341i\;\text{vs}\;0-1.39341i$	[QUANTUM]
AQC4.58	holds	$\hat{A}(t)=\mathfrak{e}^{\{i\hbar t\}}\;\hat{A}(0)\;\mathfrak{e}^{\{-i\hbar t\}}\quad=0-0.717356i\;\text{vs}\;0-0.717356i$	[QUANTUM]
AQC4.59	calc	$\lim_{\{\theta\rightarrow0\}}(\sin\theta,\;\cos\theta)\;[[\theta,\;1],[1,\;\theta]](\cos\theta,\;\sin\theta)^T= [[1,\;0],[0,\;-1]]\quad=1$	[ROT]
AQC4.60	symb	$f^{\{-1\}}(x)\;x\;f(x)=I'_m,\;I'_m=[1,\;0]\times[0,\;1]$	[OTHER]
AQC4.61	holds	$x+y\geq\sqrt{x\;y}\quad=1\;\text{vs}\;1$	[OTHER]
AQC4.62	differs	$x^{\{1/2+iy\}}/\mathfrak{e}^{\{x\log x\}}=1\quad=0.271967+0.225907i\;\text{vs}\;1+0i$	[ZETA]
AQC4.63	symb	$0(x)=\nabla_{-i}\nabla_{-j}\;\int\mathfrak{e}^{\{(2/m)\sin\theta\cos\theta\}}\times N\;\text{mod}(\mathfrak{e}^{\{x\log x\}})$	[ROT]
AQC4.64	symb	$0(x)\;(x+\Delta f ^2)^{\{1/2\}}$	[OTHER]
AQC4.65	differs	$x\;\Gamma(x)=2\;\int \sin2\theta ^2\;d\theta,\;0(x)=m(x)[D^2\phi]\quad=1.32934\;\text{vs}\;1.5708$	[ROT, GAMMA]
AQC4.66	differs	$i^2=(0,1)\cdot(0,1),\; a b \cos\theta=-1\quad=-1+0i\;\text{vs}\;1$	[ROT]
AQC4.67	symb	$E=\text{div}(E,\;E_1)$	[OTHER]
AQC4.68	calc	$\{f,g\}/[f,g]=i^2,\;E=m\;c^2,\;I'=i^2\quad=8.98755e+16$	[SR, QUANTUM]
AQC4.69	differs	$d/d\gamma\;\Gamma=m(x)=\mathfrak{e}^{\{-x\log x\}}\quad=0.422784\;\text{vs}\;0.25$	[GAMMA, ZETA]
AQC4.70	differs	$\sin(ix)=(\mathfrak{e}^{\{-x\}}+\mathfrak{e}^{\{x\}})/(2i)\quad=0+1.1752i\;\text{vs}\;0-1.54308i$	[ROT]
AQC4.71	differs	$d/d\mathfrak{f}\;F=m(x)=\mathfrak{e}^f+\mathfrak{e}^{\{-f\}}=2i\;\sin(i\;x\log x)\quad=4.25+0i\;\text{vs}\;-3.75+0i$	[ROT, BETA]
AQC4.72	symb	$(E^\pm\oplus E^\pm)\cdot(R-\subset\mathbb{C}\pm)=\mathfrak{e}\;\nabla\mathbb{C}\pm\;(\text{repeat})$	[ZETA]
AQC4.73	symb	$=\mathfrak{e}\;\nabla\mathbb{C}\pm\;(\text{repeat})$	[ZETA]
AQC4.74	symb	$\partial/\partial\mathfrak{f}\;F=t\!\!\!\int\!\!\!\int\;\text{cohom}\;D_k(x)^{\{\ll p\}}=\nabla\nabla\;\int\;\nabla f\;dx^m=\mathfrak{e}(i\hbar\nabla)^{\{\mathfrak{e}L\}}$	[MANIFOLD, QUANTUM]
AQC4.75	symb	$\bigvee\;\int\;\mathbb{C}\pm\;\nabla\;M^m$	[MANIFOLD]
AQC4.76	symb	$\Delta(M\pm\;\nabla\mathbb{C}\pm)=\exists(M\pm\;\nabla R+)$	[MANIFOLD]
AQC4.77	symb	$\exists(M\pm\;\nabla\mathbb{C}+)=XOR(\;\mathfrak{e}\;M\pm\;)$	[QUANTUM]
AQC4.78	symb	$-[E+\;\nabla R+]=\nabla+\;\nabla-\;\mathbb{C}\pm$	[OTHER]
AQC4.79	symb	$\int\;dx,\;\partial x,\;\nabla_{-i}\nabla_{-j},\;\Delta x\rightarrow E+\nabla M1,\;E+\cap R\in M1,\;R\;\nabla\mathbb{C}+$	[MANIFOLD]
AQC4.80	symb	$\bigvee(R+\nabla_{-i}\nabla_{-j}\;f)^n=\int\;\bigwedge(R+\nabla_{-i}\nabla_{-j}\;f)^2/\;\exists(R+\Delta f)^n$	[ZETA, MANIFOLD]
AQC4.81	symb	$\bigwedge(R+\nabla_{-i}\nabla_{-j}\;f)^n x=d/d\mathfrak{f}\;\int\int1/(y\log y)^{\{1/2\}}\;dy^m$	[MANIFOLD]
AQC4.82	holds	$d/dt\;g_{ij}(x)=-2\;R_{ij}\quad=0.774613\;\text{vs}\;0.774597$	[MANIFOLD]
AQC4.83	symb	$\bigvee\;\int\;\bigwedge(R+\nabla_{-i}\nabla_{-j}\;f)^n x=\bigwedge(R+\nabla_{-i}\nabla_{-j}\;f)^n/\;\exists(R+\nabla_{-i}\nabla_{-j}\;f\circ g)^n$	[MANIFOLD]
AQC4.84	differs	$x+y\geq2\sqrt{xy}),\;x(x)+y(x)\geq x(x)\;y(x)\quad=0.414966\;\text{vs}\;1$	[OTHER]
AQC4.85	holds	$x^ny=(\cos\theta+i\sin\theta)^n\quad=0.362358+0.932039i\;\text{vs}\;0.362358+0.932039i$	[ROT]
AQC4.86	differs	$x^ny=1/y^nx\quad=8\;\text{vs}\;0.111111$	[ZETA]
AHODGE.1	holds	$\Delta_p=d\delta+\delta d\quad=16\;\text{vs}\;16$	[MANIFOLD]
AHODGE.2	holds	$b_p=\dim\ker\;\Delta_p=n0(\Delta_p)\quad=1\;\text{vs}\;1$	[MANIFOLD]
AHODGE.3	symb	$\ker\;\Delta_p\equiv H^p(M,R)$	[MANIFOLD]
AHODGE.4	holds	$\zeta_p(s)=\sum_{\{\lambda>0\}}\lambda^{-\{s\}}\quad=3.28947\;\text{vs}\;3.28987$	[ZETA]
AHODGE.5	holds	$\zeta_p(0)=a_1^{-\{(p)\}}-b_p\;(d=2)\quad=-0.666333\;\text{vs}\;-0.666667$	[ZETA, MANIFOLD]
AHODGE.6	holds	$\zeta_p(0)=-b_p=-C(d,p)\;(\text{flat torus})\quad=-2\;\text{vs}\;-2$	[ZETA]
AHODGE.7	holds	$\sum_{\{p=0\}}^{\{d\}}(-1)^p\;\text{Tr}\;\mathfrak{e}^{\{-t\Delta_p\}}=\chi(M)\;(a\text{ll}\;t>0)\quad=2\;\text{vs}\;2$	[MANIFOLD]
AHODGE.8	holds	$K_0(t)=\sum_{\{\ell\geq0\}}(2\ell+1)\;\mathfrak{e}^{\{-t\ell(\ell+1)\}},\;K_1(t)=\sum_{\{\ell\geq1\}}2(2\ell+1)\;\mathfrak{e}^{\{-t\ell(\ell+1)\}},\;K_2=K_0\;(S^2)\quad=20.3367+38.6734i\;\text{vs}\;20.3367+38.6734i$	[MANIFOLD]
AHODGE.9	holds	$T^2\colon\Delta_1\equiv\Delta_0\oplus\Delta_0\;;\;K_0(0.05)=62.831853\quad=62.8319\;\text{vs}\;62.8319$	[MANIFOLD]
AHODGE.10	holds	$\chi=6(\zeta_0(0)+1)\quad=2.002\;\text{vs}\;2$	[ZETA, MANIFOLD]
AHODGE.11	symb	$H^k(X,\mathbb{C})=\mathfrak{e}\;\{p+q=k\}\;H^{\{p,q\}}(X)$	[MANIFOLD]
AHODGE.12	symb	$\text{Hodge class}\in H^{\{2p\}}(X,Q)\cap H^{\{p,p\}}(X)\;;\;[Z]\in H^{\{2p\}}(X,Q)\cap H^{\{p,p\}}$	[MANIFOLD]
AHODGE.13	symb	$\Omega^p=\ker\;\Delta_p\oplus\text{im}\;d\oplus\text{im}\;\delta$	[MANIFOLD]
AHODGE.14	holds	$X=\mathbb{C}^n/\Lambda,\;\Lambda\cong Z^{\{2n\}},\;\ker\;\Delta_k\cong\Lambda^k(R^{\{2n\}})^*,\;b_k=\mathbb{C}(2n,k)\quad=6\;\text{vs}\;6$	[MANIFOLD]

4. 全方程式レジストリ (2111 本) p.8/33

AHODGE, 15	holds	$J^2 = -1$, $h^{\sim}\{p,q\} = C(n,p) \ C(n,q)$ = 6 vs 6	[MANIFOLD]
AHODGE, 16	holds	$\Omega(Jx, Jy) = \Omega(x, y) \iff J^{\sim}T \ \Omega \ J = \Omega$ = 0 vs 0	[ROT, MANIFOLD]
AHODGE, 17	symb	$\Lambda = Z \cdot 1 \oplus Z \cdot \tau 1 \oplus Z \cdot 1 \oplus Z \cdot \tau 2$; e1=(1,0), e2=(τ 1,0), e3=(0,1), e4=(0, τ 2)	[MANIFOLD]
AHODGE, 18	holds	$\tau = x + iy : J \ e1 = (-x/y) \ e1 + (1/y) \ e2$, $J \ e2 = (-y - x^2/y) \ e1 + (x/y) \ e2$ = -1 + 0i vs -1 + 0i	[ROT, MANIFOLD]
AHODGE, 19	holds	$9^{\sim}6 = 531441$ = 531441 vs 531441	[OTHER]
AHODGE, 20	holds	$\rho(E \tau 1 \times E \tau 2) = 2 + \text{rank Hom}(E \tau 1, E \tau 2)$ = 4 + 2i vs 4 + 2i	[MANIFOLD]
AHODGE, 21	holds	#solutions $\approx 9^{\sim}\rho$ (81 = 9 ² , 6561 = 9 ⁴) = 81 vs 81	[OTHER]
AHODGE, 22	holds	$H^2(X, Z) \cong Z^{\sim}6$, $b_2 = 6$ = 6 vs 6	[MANIFOLD]
AHODGE, 23	symb	$0 \rightarrow Z \rightarrow 0 \rightarrow 0* \rightarrow 0$; $H^1(0*) \rightarrow H^2(Z)$	[MANIFOLD]
AHODGE, 24	holds	$J_{\cdot} \theta = \cos \theta \cdot I + \sin \theta \cdot K$, $J_{\cdot} \theta^2 = -1$, $J_{\cdot} \theta^{\sim}T \ J_{\cdot} \theta = 1$ = 0 + 0i vs 0 + 0i	[ROT, MANIFOLD]
AHODGE, 25	holds	$b_p = (1, 4, 6, 4, 1)$ = 0 vs 0	[MANIFOLD]
AHODGE, 26	holds	$\rho(\theta) = 4$ if $\cos \theta : \sin \theta \in Q$; $\rho(\theta) = 3$ otherwise = 4 + 3i vs 4 + 3i	[ROT, MANIFOLD]
AHODGE, 27	symb	$\Lambda^2 = \Lambda^+ \oplus \Lambda^-$	[MANIFOLD]
AHODGE, 28	holds	(1,1) real forms = $\Lambda^- \oplus R \cdot \omega_{\cdot} \theta$, $\omega_{\cdot} \theta = \cos \theta \cdot \omega_{\cdot} I + \sin \theta \cdot \omega_{\cdot} K$ = 0 vs 0	[ROT, MANIFOLD]
AHODGE, 29	holds	$I = [[0, -1, 0, 0], [1, 0, 0, 0], [0, 0, 0, -1], [0, 0, 1, 0]]$, $K = [[0, 0, 0, -1], [0, 0, -1, 0], [0, 1, 0, 0], [1, 0, 0, 0]]$ = 0 + 0i vs 0 + 0i	[ROT]
AHODGE, 30	holds	$ J^{\sim}T \ \Omega \ J - \Omega _{\text{max}} < 1e-8$ = 0 vs 0	[OTHER]
AHODGE, 31	differs	$\zeta 3 = 1/2 + i \sqrt{3}/2$ = -1 + 1.22465e-16i vs 1 + 0i	[OTHER]
AHODGE, 32	holds	$M(\square) = x^{\sim}y + y^{\sim}x + z^{\sim}a + u^{\sim}b + v^{\sim}c = 0$, $1/y + 1/x + 1/a + 1/b + 1/c = 1$ (quoted from cafe) = -4.54747e-12 + 1i vs 0 + 1i	[MANIFOLD]
AHODGE, 33	holds	$\Sigma \ x_i^{\sim}d = 0$ (d = number of variables) = 0 + 1.22465e-16i vs 0 + 0i	[MANIFOLD]
AHODGE, 34	symb	$c_1 = 0$	[MANIFOLD]
AHODGE, 35	symb	$\nabla_{\cdot} i \nabla_{\cdot} j (R + \nabla \nabla f)$ (quoted)	[MANIFOLD]
AHODGE, 36	symb	$\zeta(s) = \beta / \log x$, $H\Psi = \oplus (i \hbar \nabla)^{\sim}\{=L\}$ (quoted)	[ZETA, QUANTUM]
AHODGE, 37	holds	$s = \Sigma_p (-1)^{\sim}p \ \text{heat_trace}(\text{spec_p}, t)$ = 0 vs 0	[MANIFOLD]
ST, 1	symb	$d/df \ F = d/df \ \int \int 1/(x \log x)^2 \ dx_m + d/df \ \int \int 1/(y \log y)^{\sim}\{1/2\} \ dy_m (\times 3)$	[ZETA, MANIFOLD]
ST, 2	holds	$\log(x \log x) \geq 2(y \log y)^{\sim}\{1/2\} (\times 8)$ = 1 vs 1	[MANIFOLD]
ST, 3	symb	$d/df \ F = 2 \int (R + \nabla_{\cdot} i \nabla_{\cdot} j \ f)^2 / (- (R + \Delta f)) \ dm (\times 4)$	[MANIFOLD, ENTROPY]
ST, 4	symb	$d/df \ F \cdot t^{\sim}m = (1/4) \ g_{ij}^2$, $4V_{\cdot} \tau = g_{ij}^2$	[MANIFOLD]
ST, 5	symb	$ ds^2 = e^{\sim}\{-2\pi T \phi \} [\eta_{\cdot} \mu \nu + \bar{h}_{\cdot} \mu \nu] dx^{\sim} \mu \ dx^{\sim} \nu + T^2 \ d^2 \phi (\times 4)$	[SR, MANIFOLD, TRANSPORT]
ST, 6	symb	$\delta(x) \cdot 0(x) = ds^2 $, $\eta_{\cdot} \mu \nu = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla f(x) d\eta]^{\sim}\{1/2\}$	[MANIFOLD]
ST, 7	symb	$\bar{h}_{\cdot} \mu \nu = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla g(x) \ dx_i \ dx_j]^{\sim}\{iy\}$	[ZETA, MANIFOLD]
ST, 8	symb	$\eta_{\cdot} \mu \nu + \bar{h}_{\cdot} \mu \nu = \int [D^2 \phi \oplus h_{\cdot} \mu \nu] \ dm$	[SR, MANIFOLD]
ST, 9	symb	$\iiint V/S^2 \ dm = 0(x) = \int \int e^{\sim}\{f x \log x \ dx + 0(N^{-1})\} \ d\phi$	[MANIFOLD]
ST, 10	symb	$(\nabla \phi / \square \phi)^{\sim} = (1/2)^{\sim} = 0$	[OTHER]
ST, 11	symb	$(\eta_{\cdot} \mu \nu / \bar{h}_{\cdot} \mu \nu) = 1/i$	[MANIFOLD, QUANTUM]
ST, 12	symb	$\hbar \phi = (1/i) \ H\Psi$	[QUANTUM]
ST, 13	symb	$8\pi G(p/c^3 + V/S) = ds^2 $	[SR, TRANSPORT]
ST, 14	symb	$\lim_{\{x \rightarrow 1\}} \Sigma_{\{k=0\}^{\sim} \infty} a_k \ f^{\sim}k = T^2 \ d^2 \phi$	[MANIFOLD, ENTROPY]
ST, 15	symb	$H_3(x) = 0$, $\chi(3) = 2$, $\pi(\chi, x) = \int \int 1/(x \log x)^2 \ dx_m = (1/2)i$	[ZETA, MANIFOLD]
ST, 16	symb	$\eta_{\cdot} \mu \nu = \nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla f(x) d\eta$, $\bar{h}_{\cdot} \mu \nu = \nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla g(x) \ dx_i \ dx_j$	[MANIFOLD]
ST, 17	symb	$\eta_{\cdot} \mu \nu = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla f(x) d\eta]^{\sim}\{1/2\}$, $\bar{h}_{\cdot} \mu \nu = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla g(x) \ dx_i \ dx_j]^{\sim}\{iy\}$	[ZETA, MANIFOLD]
ST, 18	symb	$ ds^2 = e^{\sim}\{-2\pi T \phi \} [\eta_{\cdot} \mu \nu + \bar{h}_{\cdot} \mu \nu(x)] dx^{\sim} \mu \ dx^{\sim} \nu + T^2 \ d^2 \phi = \delta 0(x) [f(x) \circ g(x)] dx^{\sim} \mu \ dx^{\sim} \nu + \lim_{\{x \rightarrow \infty\}} \Sigma_{\{k=0\}^{\sim} \infty} a_k \ f^{\sim}k$	[ZETA, MANIFOLD, TRANSPORT]
ST, 19	symb	$(\square \phi)^{\sim} = \nabla_{\cdot} i \nabla_{\cdot} j (\delta(x) \circ G(x))^{\sim}\{\mu \nu\} \ (p/c^3 \circ V/S) (\times 3)$	[MANIFOLD]
ST, 20	symb	$\square \phi = 8\pi G \ T^{\sim}\{\mu \nu\}$, $0(x) = [\nabla_{\cdot} i \nabla_{\cdot} j \ f(x)]^{\sim}$	[MANIFOLD, ENTROPY]
ST, 21	symb	$\equiv nC_r \ f(x)^{\sim}n \ f(y)^{\sim}\{n-r\} \ \delta(x, y)$, $V(\tau) = \int [f(x)] \ dm / \ \partial f_{xy}$	[MANIFOLD]
ST, 22	symb	$p/c^3 \circ V/S = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla f(x) d\eta \circ \nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla g(x) \ dx_i \ dx_j]$	[MANIFOLD]
ST, 23	symb	$ ds^2 = (\delta(x) \circ G(x))^{\sim}\{\mu \nu\} \rightarrow \nabla_{\cdot} i \nabla_{\cdot} j (\delta(x) \circ G(x))^{\sim}\{\mu \nu\}$	[ZETA, MANIFOLD]
ST, 24	symb	$d/df \ F(v_{ij}, h) = [-\Delta v + \nabla_{\cdot} i \nabla_{\cdot} j \ v_{ij} - R_{ij} \ v_{ij} - v_{ij} \nabla_{\cdot} i \nabla_{\cdot} j + 2<\nabla f, \nabla h> + (R + \nabla f^2)(v/2 - h)]$	[MANIFOLD, ENTROPY]
ST, 25	symb	$e^{\sim}\{-f\} \circ e^{\sim}\{-f\} \rightarrow -2R_{ij}$, $e^{\sim}\{-f\} \rightarrow e^{\sim}\{-2\pi T \phi \} = [\nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla f(x) d\eta \circ \nabla_{\cdot} i \nabla_{\cdot} j \ \int \nabla g(x) \ dx_i \ dx_j]$	[MANIFOLD]
ST, 26	holds	$(x + y)/2 \geq \sqrt[{\sim}]{xy}$ = 1 vs 1	[OTHER]
ST, 27	differs	$x^{\sim}\{1/2 + iy\} / e^{\sim}\{x \log x\} = 1$ = 0.271967 + 0.225907i vs 1 + 0i	[ZETA]
ST, 28	symb	$\pi(\chi, x) = i \pi(\chi, x) \circ f(x) - f(x) \circ \pi(\chi, x)$	[MANIFOLD, QUANTUM]
ST, 29	symb	$\int 1/(x \log x) \ dx = i \int x \log x \ dx - \int 1/(x \log x) \ dx$	[MANIFOLD]
ST, 30	symb	$\int \int 1/(x \log x)^2 \ dx_m \geq (1/2)i$	[ZETA, MANIFOLD]
ST, 31	symb	$y = x (\times 3)$	[OTHER]
ST, 32	symb	$\int \int 1/(y \log y)^{\sim}\{1/2\} \ dy_m \geq 1/2$	[ZETA, MANIFOLD]
ST, 33	symb	$F \cdot t^{\sim}m \geq \int_M (R + \nabla_{\cdot} i \nabla_{\cdot} j \ f) \ e^{\sim}\{-f\} \ dV$	[MANIFOLD, ENTROPY]
ST, 34	symb	$F_t \geq (2/n) \ f^2$	[MANIFOLD]
ST, 35	symb	$d/df \ F_t = (1/4) \ g_{ij}^2$	[MANIFOLD]
ST, 36	symb	$f(r) = (1/2) \ m \ \sqrt[{\sim}]{(1 + f^{\sim}(r))} / f(r) - m \ g \ f(r)$	[OTHER]
ST, 37	symb	$m = 1$, $g = 1$	[OTHER]
ST, 38	calc	$f(r) = (1/4) r ^2 (\times 2)$ = 1	[MANIFOLD]
ST, 39	symb	$\int \int 1/(x \log x)^2 \ dx_m + \int \int 1/(y \log y)^{\sim}\{1/2\} \ dy_m = 0$	[ZETA, MANIFOLD]
ST, 40	symb	$x^{\sim}\{1/2 + iy\} = e^{\sim}\{x \log x\}$	[ZETA]
ST, 41	symb	$d/df \ F(v_{ij}, h) = \int e^{\sim}\{-f\} [-\Delta v + \nabla_{\cdot} i \nabla_{\cdot} j \ v_{ij} - R_{ij} \ v_{ij} - v_{ij} \nabla_{\cdot} i \nabla_{\cdot} j + 2<\nabla f, \nabla h> + (R + \nabla f^2)(v/2 - h)] (\times 2)$	[MANIFOLD, ENTROPY]
ST, 42	symb	$F \geq d/df \ \int \int 1/(x \log x)^2 \ dx_m + d/df \ \int \int 1/(y \log y)^{\sim}\{1/2\} \ dy_m$	[ZETA, MANIFOLD]
ST, 43	symb	$\pi(\chi, x) = [i \pi(\chi, x), f(x)] (\times 3)$	[MANIFOLD, QUANTUM]

4. 全方程式レジストリ (2111 本) p.9/33

ST.44	symb	$\int \int 1/(x \log x)^2 \, dx_m + \int \int 1/(y \log y)^{\{1/2\}} \, dy_m \geq 0$	[ZETA, MANIFOLD]
ST.45	symb	$\Delta x \Delta p \geq (1/4) i$	[QUANTUM]
ST.46	symb	$\delta g/L^2 \sim (G/c^4) (\delta E/L^3) (\times 2)$	[SR, QUANTUM]
ST.47	symb	$\delta E \geq \hbar/T \equiv \hbar c/L (\times 2)$	[QUANTUM]
ST.48	calc	$\delta g \geq L_p^2/L^2 (\times 2) = 2.61216e-70$	[QUANTUM]
ST.49	holds	$\sqrt{(hG/c^4)} \equiv 1.616 \times 10^{-33} = 1.00014 \text{ vs } 1$	[QUANTUM]
ST.50	holds	$C = 0.5772156\dots = 0.577216 \text{ vs } 0.577216$	[GAMMA, ZETA]
ST.51	symb	$F = d/df \int 1/(x \log x)^2 \, dx_m + d/df \int 1/(y \log y)^{\{1/2\}} \, dy_m$	[ZETA, MANIFOLD]
ST.52	symb	$f_z = \int [\sqrt{((x_1 x_2 x_3; y_1 y_2 y_3) \circ (x_1 x_2 x_3; y_1 y_2 y_3))}] \, dx \, dy \, dz (\times 3)$	[QUANTUM]
ST.53	symb	$F_t^{\sim m} = (1/4) (g_{ij})^2$	[MANIFOLD]
ST.54	symb	$\Delta E = -2(T-t) R_{ij} + \nabla_i \nabla_j f - g_{ij}/(2(T-t)) ^2 (\times 2)$	[MANIFOLD, ENTROPY]
ST.55	holds	$A = BQ + R = 17 \text{ vs } 17$	[OTHER]
ST.56	symb	$[x] = A$	[OTHER]
ST.57	symb	$dx^{\sim n} = \sum_{\{k=0\}}^{\sim \infty} x^{\sim n} \, dx$	[OTHER]
ST.58	calc	$R_n = n!/(n-r)! (x^{\sim n})' = 1600$	[OTHER]
ST.59	holds	$\beta(p,q) = \int_0^1 x^{\{p-1\}} (1-x)^{\{q-1\}} \, dx = 0.0833333 \text{ vs } 0.0833333$	[BETA]
ST.60	holds	$Z(T,X) = \exp(\sum_{m=1}^{\sim \infty} (q^{\sim k} T)^{\sim m} / m) = 1.66667 \text{ vs } 1.66667$	[ZETA]
ST.61	symb	$Z(T,X) = P_1(T)P_3(T)\cdots P_{\{2n-1\}} / (P_0(T)P_2(T)P_4(T)\cdots P_{\{2n\}})$	[ZETA]
ST.62	symb	$ v = \int (\pi r^2 + \P) dx ^2$	[OTHER]
ST.63	symb	$\Delta E = \int (\operatorname{div}(\operatorname{rot} E) \cdot e^{\{-ix \log x\}}) dx$	[ROT, ENTROPY]
ST.64	symb	$(\nabla \phi)^2 = \int t \, f(t) \, df(x)/(e^{\{-x\}} t^{\{x-1\}}) \, dx$	[GAMMA]
ST.65	symb	$(\nabla \phi)^2 = \int t \, f(t) (\Gamma(t) \, df(x)) \, dx$	[GAMMA]
ST.66	calc	$(\nabla \phi)^2 = 1/\Gamma(x+y) = 0.300901$	[GAMMA]
ST.67	symb	$C = \int 1/x^{\sim s} \, dx - \log x$	[GAMMA, ZETA]
ST.68	symb	$\int \int 1/(x \log x)^2 \, dx_m = (1/2) i (\times 2)$	[ZETA, MANIFOLD]
ST.69	holds	$\beta(p,q) = \Gamma(p) \Gamma(q)/\Gamma(p+q) (\times 2) = 0.196349 \text{ vs } 0.19635$	[ROT, GAMMA, BETA, TRANSPORT]
ST.70	symb	$\beta(p,q) \geq -\int 1/t^2 \, dt$	[BETA]
ST.71	symb	$ds^2 = g_{\mu \nu}(x) dx^{\sim \mu} \, dx^{\sim \nu} + \phi^2(x) (\kappa^2 A_{\mu \nu}(x) dx^{\sim \mu})^2 (\times 2)$	[SR, MANIFOLD, TRANSPORT]
ST.72	symb	$G_{\mu \nu} + \Lambda R_{\mu \nu} = \kappa^2 T_{\mu \nu}$	[SR]
ST.73	symb	$\nabla \phi^2 = 8 \pi G(p/c^3 + V/S) (\times 3)$	[TRANSPORT, ENTROPY]
ST.74	symb	$d/dt \, g_{ij}(t) = -2R_{ij} (\times 4)$	[MANIFOLD]
ST.75	symb	$ds^2 = -N(r)^2 dt^2 + \phi^2(r) (dr^2 + r^2 d\theta^2)$	[SR, MANIFOLD]
ST.76	calc	$ds^2 = -dt^2 + r^{\{-8 \pi Gm\}} (dr^2 + r^2 d\theta^2) = 0.312654$	[SR, TRANSPORT]
ST.77	symb	$dx^2 = g_{\mu \nu}(x) (g_{\mu \nu}(x) dx^2 - dx \, g_{\mu \nu}(x))$	[MANIFOLD, TRANSPORT]
ST.78	symb	$dx = (g_{\mu \nu}(x)^2 dx^2 - g_{\mu \nu}(x) dx \, g_{\mu \nu}(x))^{\{1/2\}} (\times 3)$	[MANIFOLD, TRANSPORT]
ST.79	symb	$\pi(X,x) = i \pi(X,x) f(x) - f(x) \pi(X,x) (\times 2)$	[ZETA, QUANTUM]
ST.80	symb	$G_{\mu \nu} + \Lambda R_{\mu \nu} = \kappa^2 T_{\mu \nu}$	[SR]
ST.81	calc	$X(3) = (-1)^3 \langle a_0 a_1 a_2 a_3 \rangle + (-1)^2 (\langle a_0 a_1 a_2 , a_1 a_2 a_3 , a_0 a_1 a_3 \rangle) + (-1) \langle a_1 a_2, a_0 a_3, a_2 a_3 \rangle + (-1)^0 \langle a_0, a_1 a_2, a_3 \rangle = 3$	[MANIFOLD]
ST.82	symb	$H_n(x) = \ker f / \operatorname{im} f$	[MANIFOLD]
ST.83	symb	$X(x) = r(H_n(x))$	[MANIFOLD]
ST.84	holds	$X(3) = H_3(x) = 0 = 0 \text{ vs } 0$	[MANIFOLD]
ST.85	symb	$H_3(\Pi) = Z (\times 2)$	[MANIFOLD]
ST.86	holds	$\log x(\log x) \geq 2(y \log y)^{\{1/2\}} = 1 \text{ vs } 1$	[ZETA, MANIFOLD, TRANSPORT]
ST.87	symb	$\pi(X,X) = [i \pi(X,x), f(x)]$	[QUANTUM, ENTROPY]
ST.88	calc	$\pi(X,x) = \int 1/(x \log x)^2 \, dx = 0.281002$	[MANIFOLD, ENTROPY]
ST.89	symb	$R_{\mu \nu} - (1/2) R \, g_{\mu \nu} = \kappa^2 T_{\mu \nu}$	[SR]
ST.90	calc	$L_p = \sqrt{(hG/c^3)} = 1.61622e-35$	[QUANTUM]
ST.91	symb	$V(\tau) = \int \tau(p)^{\{-n/2\}} \exp(-(1/\sqrt{(2 \tau(q))}) L(x) dx) + O(N^{-1})$	[MANIFOLD, ENTROPY]
ST.92	symb	$(1/\tau)(N/2 + \tau(2\Delta F - \nabla f ^2 + R) + f) \bmod N^{-1}$	[MANIFOLD, ENTROPY]
ST.93	symb	$d/df \, F = m(x) (\times 2)$	[MANIFOLD, QUANTUM]
ST.94	symb	$\pi(K'') \equiv \pi(K'' , 0)$	[JONES, MANIFOLD]
ST.95	symb	$l(\langle P,Q \rangle), \, l(\langle Q,R \rangle), \, l(\langle R,S \rangle), \, l(\langle P,R \rangle), \, l(\langle P,S \rangle), \, l(\langle Q,S \rangle) = \alpha, \, \beta, \, \gamma, \, \delta, \, \xi, \, \eta$	[JONES]
ST.96	symb	$e = \alpha \beta \gamma, \, \gamma = \beta^{-1} \alpha^{-1}, \, \xi = \alpha \gamma, \, \delta = \eta \gamma^{-1}, \, \eta = \beta \xi$	[JONES, MANIFOLD]
ST.97	symb	$\beta^5 = (\beta \theta)^2 = \theta^3$	[JONES, MANIFOLD]
ST.98	symb	$[\beta] = [\theta] = 0$	[MANIFOLD]
ST.99	symb	$ds^2 = e^{\{-2kT(x) \phi \}} [\eta_{\mu \nu} + \hbar_{\mu \nu}(x)] dx^{\sim \mu} \, dx^{\sim \nu} + T^2(x) d\phi^2 (\times 2)$	[SR, MANIFOLD, TRANSPORT]
ST.100	calc	$\log x(\log x) = 2(y \log y)^{\{1/2\}} = 4.60304$	[ZETA, MANIFOLD]
ST.101	symb	$ f(x) \rightarrow [f, f^{-1}] \times [g, h]$	[QUANTUM, TRANSPORT]
ST.102	holds	$(x-1)(2y-b) \geq z = 1 \text{ vs } 1$	[TRANSPORT]
ST.103	holds	$(1/(x-1))(1/(2y-b)) \leq 1/z = 1 \text{ vs } 1$	[TRANSPORT]
ST.104	calc	$f(x) = 1/(x-1) + 1/(2y-b) = 0.833333$	[ZETA, TRANSPORT]
ST.105	symb	$d/df \geq (d/df) \, f$	[OTHER]
ST.106	symb	$dx \leq dF$	[OTHER]
ST.107	calc	$ f(x) = (1/4) r ^2 = 2.25$	[MANIFOLD]
ST.108	differs	$\Gamma(x) = \int x^{\{1-t\}} e^{\{-x\}} \, dx = 1.77245 \text{ vs } 0.885996$	[GAMMA, MANIFOLD]
ST.109	symb	$f(x) = [f, f^{-1}] \times [g, h]$	[OTHER]
ST.110	symb	$V(x) = \int (1/\sqrt{(2 \tau(q))} (\exp \int L(x) dx) + O(N^{-1})$	[MANIFOLD, ENTROPY]
ST.111	symb	$Zeta(x,h) = \exp((q \, f(x))^{\sim m} / m) (\times 2)$	[ZETA]

4. 全方程式レジストリ (2111 本) p.10/33

ST.112	symb	$\mathrm{d}/\mathrm{d}f\;F(x)=m(x)$	[MANIFOLD, QUANTUM]
ST.113	symb	$\mathrm{d}s^2=g_{\mu\nu}\,\mathrm{d}x^\mu\,\mathrm{d}x^\nu+\phi^2(x)(\kappa^2A_{\mu\nu}(x)\mathrm{d}x^\mu\mathrm{d}x^\nu)^2$	[SR, MANIFOLD, TRANSPORT]
ST.114	differs	$\Sigma_{\{k=0\}^\infty} x_k+y_k ^2=\Sigma x ^2+2\Sigma xy +\Sigma y ^2\leq\Sigma x ^2+\Sigma y ^2=0$ vs 1	[OTHER]
ST.115	symb	$f(x+y)=f(x)+f(y)\geq f(x \circ y)$	[OTHER]
ST.116	holds	$\chi(x)=H_3(x)=0=0$ vs 0	[MANIFOLD]
ST.117	symb	$V(x)/f(x)=m(x)\;(\times 2)$	[MANIFOLD]
ST.118	symb	$F(x)=0$	[MANIFOLD]
ST.119	symb	$\mathrm{d}/\mathrm{d}f\;F(x)\geq 0$	[MANIFOLD, ENTROPY]
ST.120	symb	$\lim_{\{x\rightarrow\infty\}\;\mathrm{mesh}}F(x)/f(x)\rightarrow 0$	[MANIFOLD]
ST.121	symb	$\nabla f(x)=2$	[OTHER]
ST.122	symb	$f(x)=\int 1/x^s\,\mathrm{d}x-\log x$	[GAMMA, ZETA]
ST.123	symb	$\mathrm{d}/\mathrm{d}f\;(x-y)^n=\Pi(x-y)^n/\partial f_{xy}$	[OTHER]
ST.124	symb	$\Delta E=\int\mathrm{div}(\mathrm{rot}\;E)\;e^{-ix\log x}\,\mathrm{d}x$	[ROT, ENTROPY]
ST.125	symb	$f(x)=\Sigma_{\{k=0\}^\infty}a_kx^k$	[OTHER]
ST.126	symb	$f(x)=\chi-\{x\}\;(\times 2)$	[MANIFOLD]
ST.127	symb	$\int_{-\beta}^{-\alpha}(x-\alpha)(x-\beta)\,\mathrm{d}x=2\int_Mf(x)\mathrm{d}x$	[OTHER]
ST.128	differs	$\int_{-\beta}^{-\alpha}a(x-\alpha)(x-\beta)\,\mathrm{d}x=-a(\beta-\alpha)^3/6=1.33333$ vs -1.33333	[OTHER]
ST.129	symb	$\mathrm{d}/\mathrm{d}f\;F(x)=0$	[MANIFOLD]
ST.130	calc	$\log(x\log x)-2(y\log y)^{\{1/2\}}=[d/f\;F(x)]=0.781797$	[ZETA, MANIFOLD]
ST.131	symb	$\nabla\phi^2=8\pi Gh+8\pi V/S$	[ENTROPY]
ST.132	symb	$\square_v=2\sqrt{(2\pi Gh)}+2\sqrt{(2\pi V/S)}\;(\times 2)$	[ENTROPY]
ST.133	symb	$\sqrt{(2\pi T)}=2\sqrt{(2\pi V/S)}$	[ENTROPY]
ST.134	differs	$\log(x\log y)\geq 2(xe^x)^{\{1/2\}}=0$ vs 1	[MANIFOLD]
ST.135	symb	$\log x+\log\log y=2(ye^x)^{\{1/2\}}$	[OTHER]
ST.136	symb	$e^x+y=2(ye^x)^{\{1/2\}}\rightarrow 1/2$	[OTHER]
ST.137	differs	$e^{\{x+1\}}=2(xe^x)^{\{1/2\}}=7.38906$ vs 3.29744	[OTHER]
ST.138	holds	$e^{\{2(x+1)\}}/(4xe^x)\geq 1=1$ vs 1	[OTHER]
ST.139	symb	$F_{\mathbf{t}^m}=(1/4)\;g_{ij}^2$	[MANIFOLD]
ST.140	calc	$(x+1)/(4x)=y^2=0.612372$	[OTHER]
ST.141	differs	$1/4+1/x=y^2=0.75$ vs 0.375	[OTHER]
ST.142	differs	$\log(xy)\geq 2(yx)^{\{1/2\}}=0$ vs 1	[MANIFOLD]
ST.143	symb	$(\log(xy))'\geq 4(yx)^{-1/2}$	[OTHER]
ST.144	differs	$1/(xy)\geq 4(yx)^{-1/2}=0$ vs 1	[OTHER]
ST.145	holds	$\log x+\log y\geq 4(yx)^{-1/2}=1$ vs 1	[OTHER]
ST.146	differs	$1/x+1/y\geq 4(yx)^{-1/2}=0$ vs 1	[OTHER]
ST.147	differs	$((x+y)/(xy))\;y^x^{\{1/2\}}\geq 4=0$ vs 1	[OTHER]
ST.148	differs	$xy^{\{1/2\}}/(x+y)\leq 1/4=0$ vs 1	[OTHER]
ST.149	differs	$\Gamma(xy)/\Gamma(x+y)\leq 1/4=0$ vs 1	[GAMMA]
ST.150	holds	$x+y\geq 2\sqrt{(xy)}=1$ vs 1	[OTHER]
ST.151	symb	$\Delta x\Delta y\geq (1/4)\;i$	[QUANTUM]
ST.152	symb	$e^{\{2(yx)^{\{1/2\}}\}}=g_{ij}^2$	[MANIFOLD]
ST.153	symb	$xy=e^{\{2(yx)^{\{1/2\}}\}}\rightarrow g_{ij}^2$	[MANIFOLD]
ST.154	differs	$1+1/x=4y=1.5$ vs 2.44949	[OTHER]
ST.155	holds	$(e^x+x)^2\geq 4xe^x=1$ vs 1	[OTHER]
ST.156	calc	$r^2= x0x-2x0\;e^x+e^{\{2x0\;x\}} \;/\;\sqrt{(x0^2+y0^2)}=0.0559751$	[OTHER]
ST.157	symb	$r^2(x0^2+y0^2)=[(e^x+1)(e^x-1)+(x+i)(x-i)]+2xe^{\{x^2\}}$	[OTHER]
ST.158	calc	$r= 1+x-4xy \;/\;\sqrt{(x0^2+y0^2)}=14.8492$	[OTHER]
ST.159	symb	$r^2(x0^2+y0^2)=[(x-1)(x+1)+(x-i)(x+i)]+2x0\;e^{\{2x\}}$	[OTHER]
ST.160	symb	$F_{\mathbf{t}^m}=[f(x)]$	[MANIFOLD]
ST.161	symb	$[f(x)]=0(x)$	[MANIFOLD]
ST.162	symb	$X\in 0(x)$	[MANIFOLD]
ST.163	symb	$0\rightarrow[d/f\;F]\rightarrow\infty$	[OTHER]
ST.164	differs	$\log xy\geq 2(xe^x)^{\{1/2\}}=0$ vs 1	[OTHER]
ST.165	symb	$(y+x)^2-4i(y+x)(xe)^{\{1/2\}}-4xe$	[OTHER]
ST.166	symb	$r^2(y^2+x^2)=[(y+x)^2-4i(y+x)xe^{\{1/2\}}]-4xe$	[OTHER]
ST.167	symb	$r^2(x^2+y^2)=[(\log y+\log x)^2-4i(\log y+\log x)xe^{\{1/2\}}]-4xe$	[OTHER]
ST.168	symb	$\delta(f)=\int(g(c+d)/f(a+b))\;z(f+g)\;dz$	[OTHER]
ST.169	symb	$y=f(y)$	[OTHER]
ST.170	symb	$\mathrm{d}s^2=e^{-2\pi T \phi }\big[\tau_{\mu\nu}+\bar{h}_{\mu\nu}\big]\mathrm{d}x^\mu\,\mathrm{d}x^\nu+T^2\;d^2\theta$	[MANIFOLD, TRANSPORT]
ST.171	symb	$[x\;y;\;a\;b]\rightarrow[z]$	[OTHER]
ST.172	symb	$f[z]\rightarrow\mathrm{d}x$	[OTHER]
ST.173	symb	$x\cdot y=0$	[OTHER]
ST.174	symb	$\infty\rightarrow[f(x)]$	[OTHER]
ST.175	symb	$\chi-\{x\}=\infty$	[MANIFOLD]
ST.176	symb	$f(x)+0(x)\rightarrow\mathrm{finite}$	[MANIFOLD, TRANSPORT]
ST.177	symb	$f(x)=[F(x)]$	[OTHER]
ST.178	symb	$\mathrm{d}s^2=e^{-2\pi T \phi }\big[\eta_{\mu\nu}+\bar{h}_{\mu\nu}\big]\mathrm{d}x^\mu\,\mathrm{d}x^\nu+T^2\;d^2\phi$	[SR, MANIFOLD, TRANSPORT]
ST.179	symb	$\nabla^2\phi=8\pi Gh+8\pi V/S$	[ENTROPY]

4. 全方程式レジストリ (2111 本) p.11/33

ST.180	symb	$2\sqrt{2\pi V/S} = \int \int 1/(x \log x)^2 \, dx_m$	[MANIFOLD, TRANSPORT, ENTROPY]
ST.181	calc	$m = \sqrt{2\pi T} = 2.50663$	[ENTROPY]
ST.182	symb	$f(x) = 0$	[OTHER]
ST.183	symb	$F = \int \int 1/(x \log x)^2 \, dx_m + \int \int 1/(y \log y)^{1/2} \, dy_m$	[ZETA, MANIFOLD]
ST.184	symb	$F = 0, \, x \cdot y = 0, \, F(x) = 0, \, G(x) = 0, \, 0 \in X, \, X \cong 0, \, x - y = 0, \, F(x,y) = 0, \, x^n + y^n = z^n, \, F(x) = x^n + y^n, \, F(x) = 0$	[OTHER]
ST.185	calc	$ds^2 = r/(d + e \cos \theta) = 0.8$	[ROT, TRANSPORT]
ST.186	symb	$0 \in X \rightarrow \infty, \, F(x) = z^n$	[OTHER]
ST.187	symb	$x^n + y^n = z^n$	[OTHER]
ST.188	symb	$F(x) = 0(\delta(x))[(x - f(x))(x + f(x)) + (y - \check{f}(x))(y + \check{f}(x))] + 0(N^{-1})$	[OTHER]
ST.189	symb	$f(x) = \sum_{k=0}^{\infty} a_k f^k(x)$	[OTHER]
ST.190	symb	$\int f(x), \, \partial f(x), \, dx \rightarrow [f(x)]$	[OTHER]
ST.191	symb	$0(x) = \mathcal{X}(x) - \{x\}$	[MANIFOLD]
ST.192	symb	$\sum_{k=0}^{\infty} a_k x^k = f(x) + \{x\}$	[OTHER]
ST.193	symb	$\int dx, \, \partial x, \, dx, \, [f(x)] \rightarrow F(x) \rightarrow \int dx \rightarrow f(x) \rightarrow \partial x \rightarrow x \rightarrow x \rightarrow \text{const} \rightarrow [f(x)] \rightarrow \infty$	[MANIFOLD]
ST.194	symb	$(E_2 \otimes E_2) \cdot (R^- \subset C^+) = \otimes \nabla C \pm$	[QUANTUM]
ST.195	symb	$\bigvee \int C^+ \nabla M_m / \Delta (M \pm \nabla C \pm) = \exists (M \pm \nabla R^+)$	[QUANTUM]
ST.196	symb	$\exists (M \pm \nabla C^+) = X0R(\otimes \nabla M \pm)$	[QUANTUM]
ST.197	symb	$-[E^+ \nabla R^+] = \nabla_+ \nabla_- C \pm$	[QUANTUM]
ST.198	symb	$\int dx, \, \partial x, \, \nabla_i \nabla_j, \, \Delta x \rightarrow E^+ \nabla M_1, \, E^+ \cap R \in M_1, \, R \nabla C^+ (\times 3)$	[QUANTUM]
ST.199	symb	$\bigvee (R + \nabla_i \nabla_j f)^n = \int \wedge (R + \nabla_i \nabla_j f)^2 / \exists (R + \Delta f)^n$	[MANIFOLD, QUANTUM]
ST.200	symb	$\wedge (R + \nabla_i \nabla_j f)^n x = d/df \int \int 1/(y \log y)^{1/2} \, dy_m$	[ZETA, QUANTUM]
ST.201	symb	$d/dt \, g_{ij}(x) = -2R_{ij}$	[MANIFOLD]
ST.202	symb	$\bigvee \int \wedge (R + \nabla_i \nabla_j f)^n x = \wedge (R + \nabla_i \nabla_j f)^n / \exists (R + \nabla_i \nabla_j f \circ g)^n$	[MANIFOLD, QUANTUM]
ST.203	differs	$x + y \geq 2\sqrt{xy}, \, x(x) + y(x) \geq x(x)y(x) = 0 \text{ vs } 1$	[OTHER]
ST.204	holds	$x^ny = (\cos \theta + i \sin \theta)^n = 6.12323e-17 + 1i \text{ vs } 6.12323e-17 + 1i$	[ROT, QUANTUM]
ST.205	differs	$x^ny = 1/y^nx = 8 \text{ vs } 0.111111$	[ZETA]
ST.206	calc	$F - N = mg - N' = 21.6$	[ROT, TRANSPORT]
ST.207	symb	$C = \int \int 1/(x \log x)^2 \, dx_m (\times 2)$	[MANIFOLD, TRANSPORT]
ST.208	symb	$y = (\nabla \phi)^{1/2}$	[OTHER]
ST.209	symb	$\Delta E / (1/(2\sqrt{2\pi G})) = p/c^3 + \rho$	[ENTROPY]
ST.210	symb	$ds = (g_{\mu\nu}(x)dx^\mu dx^\nu + \phi^2(x)(\kappa^2 A_{\mu\nu}(x)dx^\mu dx^\nu)^{1/2}$	[SR, TRANSPORT]
ST.211	symb	$\Delta E / (1/(2\sqrt{2\pi G})) - \rho = c^3, \, p/(2\pi) = c^3$	[ENTROPY]
ST.212	symb	$\partial/\partial f \int (\sin 2x)^2 \, dx = x - y ^2$	[ROT]
ST.213	symb	$x = \bar{x} / x $	[QUANTUM]
ST.214	symb	$ \bar{x} = \sqrt{x}, \, i = \sqrt{-1}, \, \bar{x} x = \bar{x}, \, \bar{x} = 1x, \, \bar{x} ^2 = -1$	[QUANTUM]
ST.215	symb	$(x / \bar{x})^2 = 1/(-1), \, \sqrt{x} = i$	[QUANTUM]
ST.216	holds	$x \bmod N = 0 = 0 \text{ vs } 0$	[OTHER]
ST.217	symb	$\sum_{M=0}^{\infty} \int_M dm \rightarrow \sum_{\{x=0\}}^{\infty} F_x = \int_m dm = F$	[MANIFOLD]
ST.218	symb	$(x - y)/a = (y - z)/b = (z - x)/c$	[OTHER]
ST.219	symb	$\Pi(x - y)/\partial y$	[OTHER]
ST.220	symb	$ \int f(x) ^2 \rightarrow \int \pi r^2 \, dx \cong V(\tau)$	[MANIFOLD, ENTROPY]
ST.221	symb	$\int f(x) ^2/f(x) \, dx$	[ENTROPY]
ST.222	symb	$V(\tau) \rightarrow \text{mesh}$	[MANIFOLD]
ST.223	calc	$\int x_1 x_2 x_3; y_1 y_2 y_3; z_1 z_2 z_3 \, dv(\tau) = 1$	[OTHER]
ST.224	calc	$ x_1 x_2 x_3; y_1 y_2 y_3; z_1 z_2 z_3 _{[dx=v]} = 18$	[OTHER]
ST.225	symb	$z_y = a_x + b_y + c_z$	[OTHER]
ST.226	symb	$dz_y = d(z_y)$	[OTHER]
ST.227	symb	$[f, f^{-1}] = ff^{-1} - f^{-1}f$	[QUANTUM]
ST.228	symb	$V(\tau) = \int \tau(q)^{-n/2} \exp(-(1/\sqrt{2\tau(q)}) L(x)dx) + 0(N^{-1})$	[MANIFOLD, ENTROPY]
ST.229	symb	$(1/\tau)(N/2 + \tau(2\Delta f - \nabla f ^2 + R) + f) \bmod N^{-1}$	[MANIFOLD, ENTROPY]
ST.230	symb	$ds^2 = -N(r)^2 dt^2 + \phi^2(r)(dr^2 + r^2 d\theta^2)$	[SR, MANIFOLD]
ST.231	symb	$\sum_{n=0}^{\infty} a_1 x^1 + a_2 x^2 \cdots a_{n-1} x^{n-1} \rightarrow \sum_{n=0}^{\infty} a_n x^n \rightarrow \alpha$	[OTHER]
ST.232	calc	$f = n \nu \lambda, \, \lambda = x/l, \, \int dn \nu \lambda = f(x), \, xf(x) = F(x), \, [f(x)] = \nu h (\times 2) = 3.31304e-19$	[QUANTUM]
ST.233	symb	$0(x) = ([\nabla_i \nabla_j f(x)])' (\times 2)$	[MANIFOLD]
ST.234	symb	$\cong {}_nC_r(x)^n(y)^{n-r} \delta(x,y) (\times 2)$	[MANIFOLD]
ST.235	symb	$F_t{}^m = (1/4) g_{ij}^2, \, x^{1/2 + iy} = e^{\{x \log x\}} (\times 2)$	[ZETA, MANIFOLD]
ST.236	symb	$S_m^{\sim}\{\mu\nu\} \otimes S_n^{\sim}\{\mu\nu\} = G_{\mu\nu} \times T^{\sim}\{\mu\nu\} (\times 2)$	[SR, MANIFOLD]
ST.237	symb	$S_m^{\sim}\{\mu\nu\} \otimes S_n^{\sim}\{\mu\nu\} = -(2R_{ij}/V(\tau))[D^2\phi] (\times 4)$	[MANIFOLD]
ST.238	symb	$S_m^{\sim}\{\mu\nu\} = \pi(\chi,x) \otimes h_{\mu\nu} (\times 2)$	[MANIFOLD]
ST.239	symb	$\pi(\chi,x) = \int \exp[L(p,q)] \, d\phi (\times 2)$	[QUANTUM]
ST.240	symb	$ds^2 = e^{-2\pi T \phi } [\eta + \bar{h}_{\mu\nu}] dx^{\sim}\{\mu\nu\} dx^{\sim}\{\mu\nu\} + T^2 d^2\phi (\times 2)$	[MANIFOLD, TRANSPORT]
ST.241	symb	$M_3 \otimes_{k=0}^{\infty} E \pm = \text{rot}(\text{div } E, E_1) = m(x), \, P^{\sim}\{2n\}/M_3 = H_3(M_1) (\times 2)$	[ROT, MANIFOLD]
ST.242	symb	$\exists [R + \nabla f ^2]^{1/2 + iy} = \int \exp[L(p,q)] \, d\phi = \exists [R + \nabla f ^2]^{1/2 + iy} \otimes \int \exp[L(p,q)] \, d\phi + N \bmod (e^{\{x \log x\}}) = 0(\phi) (\times 2)$	[ZETA, MANIFOLD, QUANTUM]
ST.243	symb	$d/dt \, g_{ij}(t) = -2R_{ij}, \, P^{\sim}\{2n\}/M_3 = H_3(M_1), \, H_3(M_1) = \pi(\chi,x) \otimes h_{\mu\nu} (\times 2)$	[MANIFOLD, QUANTUM]
ST.244	symb	$S_m^{\sim}\{\mu\nu\} \times S_n^{\sim}\{\mu\nu\} = [D^2\phi], \, S_m^{\sim}\{\mu\nu\} \times S_n^{\sim}\{\mu\nu\} = \ker f/\text{im } f, \, S_m^{\sim}\{\mu\nu\} \otimes S_n^{\sim}\{\mu\nu\} = m(x)[D^2\phi], \, -2R_{ij}/V(\tau) = f^{-1} x f(x) (\times 2)$	[MANIFOLD]
ST.245	symb	$f_z = \int \sqrt{((x y z; u v w) \circ (x y z; u v w))} \, dx \, dy \, dz \rightarrow f_z^{1/2} \rightarrow (0,1) \cdot (0,1) = -1, \, i = \sqrt{-1} (\times 2)$	[QUANTUM]

ST. 246	symb	$(x,y,z)^2=(x,y,z)\cdot(x,y,z)\rightarrow-1\ (\times 2)$	[QUANTUM]
ST. 247	symb	$0(x)=\nabla_i\nabla_j\int e^{\{-2/m\}\sin\theta\cos\theta}\times N\mathrm{mod}(e^{\{x\log x\}}/\{0(x)(x+\Delta f ^2)^{1/2}\})\ (\times 2)$	[ROT, MANIFOLD]
ST. 248	differs	$x\Gamma(x)=2\int \sin2\theta ^2d\theta, \, 0(x)=m(x)[D^2\phi]\ (\times 2) \qquad =1.32934\ \mathrm{vs}\ 1.5708$	[ROT, GAMMA]
ST. 249	calc	$\lim_{\{\theta\rightarrow 0\}}(1/\theta)(\sin\theta,\cos\theta)(\theta\,1;1\,\theta)(\cos\theta;\sin\theta)^T=(1\,0;0\,-1),\,f^{-1}(x)xf(x)=I'_{-m},\,I'_{-m}=[1,0]\times[0,1]\ (\times 2)$ $=100.02$	[ROT, QUANTUM, TRANSPORT]
ST. 250	symb	$i^2=(0,1)\cdot(0,1),\, a b \cos\theta=-1,\,E=\mathrm{div}(E,E_1)\ (\times 2)$	[ROT, QUANTUM]
ST. 251	calc	$(\{f,g\}/[f,g])' = i^2,\,E=mc^2,\,I' = i^2\ (\times 2) \qquad =8.98755e+16$	[SR, QUANTUM]
ST. 252	symb	$0(x)= \nabla\int[\nabla_i\nabla_j\,f\circ g(x)]^{\{-1/2+iy\}} ,\,\partial r^{\mathrm{n}} \nabla ^2\rightarrow\nabla_i\nabla_j \nabla ^2\ (\times 2)$	[ZETA, MANIFOLD]
ST. 253	symb	$\nabla^2\phi=8\pi G(p/c^3+V/S)\ (\times 2)$	[ENTROPY]
ST. 254	symb	$(\log x^{\{1/2\}})'=(1/2)(1/(x\log x)),\,(\sin\theta)'=\cos\theta,\,(f_z)'=ie^{\{ix\log x\}},\,d/df\,F=m(x)\ (\times 2)$	[ROT, ZETA]
ST. 255	symb	$d/df\int\int1/(x\log x)^2\,dx_m+d/df\int\int1/(y\log y)^{\{1/2\}}\,dy_m=d/df\int\int1/(x\log x)^2+1/(y\log y)^{\{1/2\}}\,dm\geq d/df\int\int$ $(1/((x\log x)^2\circ(y\log y)^{\{1/2\}}))\,dm\geq 2h\ (\times 2)$	[ZETA, MANIFOLD, QUANTUM]
ST. 256	symb	$d/df\int\int(1/((x\log x)^2\circ(y\log y)^{\{1/2\}}))\,dm\geq h\ (\times 2)$	[MANIFOLD, QUANTUM]
ST. 257	symb	$y=x,\,xy=x^2,\,(\Box\phi)'=8\pi G(p/c^3\circ V/S)\ (\times 2)$	[TRANSPORT]
ST. 258	symb	$\Box\phi=\int\exp[8\pi G(\hbar\mu\nu\circ\eta_\mu)^{\sim\nu}]\,dm\,d\phi,\,\Sigma_{a,k}x^{\sim k}=d/df\,\Sigma\Sigma(1/(a_k^2f^{\sim k}))\,dx_k\ (\times 2)$	[QUANTUM, TRANSPORT]
ST. 259	symb	$\Sigma_{a,k}f^{\sim k}=d/df\,\Sigma\Sigma(\zeta(s)/a_k)\,dx_{\{k_m\}},\,a_k^2f^{\sim\{1/2\}}\rightarrow\lim_{\{k\rightarrow 1\}}a_kf^{\sim k}=\alpha\ (\times 2)$	[ZETA]
ST. 260	symb	$0(x)=D^2\phi\otimes h_\mu\nu,\,ds^2=e^{\{-2\pi T \phi \}}[\eta_\mu\nu+\hbar_\mu\nu]dx^\mu dx^\nu+T^2\,d^2\phi$	[MANIFOLD, TRANSPORT]
ST. 261	holds	$f(x)+f(y)\geq 2\sqrt{f(x)f(y)},\,(1/4)(f(x)+f(y))^2\geq f(x)f(y) \qquad =1\ \mathrm{vs}\ 1$	[OTHER]
ST. 262	symb	$T^{\{\mu\nu\}}=(p/c^3+V/S)^{-1},\,E\pm=f^{-1}\times f(x),\,E=mc^2$	[SR, ENTROPY]
ST. 263	symb	$0(x)=\Box\iiint(\nabla_i\nabla_j\,f\circ g(x))^2/V(x)\,dm$	[MANIFOLD]
ST. 264	symb	$ds^2=g_\mu\nu^2\,d^2x+g_\mu\nu\,dx\,g_\mu\nu(x),\,E\pm=f^{-1}\times f(x),\,V/S=AA^{-1},\,AA^{-1}=E$	[MANIFOLD]
ST. 265	calc	$0(x)=\iiint(f(x^2,\,y^3,\,z^3)\,dx\,dy\,dz,\,S(r)=\pi r^2,\,V(r)=4\pi r^3 \qquad =100.531$	[OTHER]
ST. 266	holds	$E\pm=f(x)\cdot e^{\{-x\log x\}},\,\Sigma_{\{k=0\}}^{\sim\infty}a_kf^{\sim k}=f(x)\cdot e^{\{-x\log x\}} \qquad =0.25\ \mathrm{vs}\ 0.25$	[MANIFOLD, QUANTUM]
ST. 267	symb	$P^{\{2n\}}/M_3=E\pm-\phi,\,\zeta(x)/\Sigma_{\{k=0\}}^{\sim\infty}a_kx^{\sim k}=0(x)$	[ZETA, MANIFOLD]
ST. 268	symb	$0(N^{-1})=1/0(x),\,\Box=(8\pi G/c^3)T^{\{\mu\nu\}}$	[SR]
ST. 269	symb	$\partial^2f(\Box\phi)=-2\Box\iiint V/S^2\,dm,\,d/dt\,g_{ij}(t)=-2R_{ij}$	[MANIFOLD]
ST. 270	symb	$E\pm=e^{\{-2\pi T \phi \}}[\eta_\mu\nu+\hbar_\mu\nu]dx^\mu dx^\nu+T^2\,d^2\phi$	[MANIFOLD, TRANSPORT]
ST. 271	calc	$S_1^{\{mn\}}\otimes S_2^{\{mn\}}=D^2\phi\otimes h_\mu\nu,\,S_m^{\sim\{\mu\nu\}}\otimes S_n^{\sim\{\mu\nu\}}=\int[D^2\phi]dm=\nabla_i\nabla_j\int f(x)dm,\,h=p/(mv),\,h\lambda=f,\,f\lambda=h\nu \qquad =$ $7.2742e-10$	[QUANTUM]
ST. 272	symb	$ ds^2 =S_m^{\sim\{\mu\nu\}}\otimes S_n^{\sim\{\mu\nu\}},\,E\pm=f^{-1}(x)\times f(x)$	[MANIFOLD]
ST. 273	symb	$R^+\subset\mathbb{C}\pm,\,\nabla R^+\rightarrow\otimes Q\pm$	[QUANTUM]
ST. 274	symb	$\nabla_i\nabla_j\,R\pm=\otimes\nabla Q\pm$	[QUANTUM]
ST. 275	symb	$M_1=R_3^+/E\pm-\{\phi\},\,M_3\cong M_2,\,M_3\cong M_1$	[MANIFOLD]
ST. 276	symb	$H_3(\Pi)=Z_1\otimes Z_1,\,H_3(M_1)=0$	[MANIFOLD]
ST. 277	symb	$\nabla\phi^2=8\pi G(p/c^3+V/S)\ (\times 2)$	[TRANSPORT, ENTROPY]
ST. 278	symb	$(\partial\gamma^{\mathrm{n}}+m^2)\cdot\phi=\int[D^2\phi\otimes h_\mu\nu]\,dm=0$	[QUANTUM]
ST. 279	symb	$\Box=\pi(\chi,x)\otimes h_\mu\nu=D^2\phi\otimes h_\mu\nu$	[MANIFOLD, TRANSPORT]
ST. 280	symb	$\int[D^2\phi]\,dm=\pi(M_1),\,H_n(m_1)=D^2\phi-\pi(\chi,x)=\ker f/\mathrm{im}\,f$	[MANIFOLD, ENTROPY]
ST. 281	symb	$\int\,Dq\exp[L(x)]\,d\phi+0(N^1)=\pi(\chi,x)\otimes h_\mu\nu=D^2\phi\otimes h_\mu\nu$	[QUANTUM, ENTROPY]
ST. 282	symb	$\lim_{\{x\rightarrow 1\}}\Sigma_{\{k=0\}}^{\sim\infty}\zeta(x)/(a_kf^{\sim k})=\int [D^2\phi\otimes h_\mu\nu] \,dm$	[ZETA, MANIFOLD]
ST. 283	symb	$\nabla\phi^2=\Box\iiint V/S^2\,dm$	[MANIFOLD]
ST. 284	symb	$0(x)=D^2\phi\otimes h_\mu\nu$	[MANIFOLD]
ST. 285	symb	$dx<\partial x<\nabla\phi<\Box v$	[OTHER]
ST. 286	symb	$\otimes<\Sigma<\otimes<\int<\wedge$	[OTHER]
ST. 287	symb	$(\delta\phi(x))^2=\iiint V(x)/S^2\,dm,\,\delta\phi(x)=(\iiint V(x)/S^2\,dm)^{\sim\{1/2\}}$	[QUANTUM]
ST. 288	symb	$\nabla\phi^2=-4R\int\delta(V\cdot S^{-3})\,dm$	[MANIFOLD]
ST. 289	symb	$\nabla\phi=2R\zeta(s)_i$	[ZETA]
ST. 290	symb	$\Sigma_{\{k=0\}}^{\sim\infty}(a_kx^{\sim k}/(m\,dx))\,f^{\sim k}(x)=(m/n!)\,f^{\mathrm{n}}(x)=((\zeta(s))^{\sim k}/df)\,m(x),\,(\delta(x))^{\sim\{1/2\}}=(x\log x/x^{\mathrm{n}})^{\mathrm{n}}$	[ZETA]
ST. 291	symb	$0(x)=\int[D^2\phi\otimes h_\mu\nu]\,dm/e^{\{x\log x\}}$	[MANIFOLD]
ST. 292	symb	$0(x)=V(x)/\int[D^2\phi\otimes h_\mu\nu]\,dm$	[MANIFOLD]
ST. 293	calc	$M_3=e^{\{x\log x\}},\,x^{\{1/2+iy\}}=e^{\{x\log x\}},\,(x)=M_3/e^{\{x\log x\}}=nE_x \qquad =27$	[ZETA, QUANTUM]
ST. 294	calc	$V/S^2=p/c^3,\,h\nu\neq p/(mv),\,E_1=h\nu,\,E_2=mc^2,\,E_1\cong E_2 \qquad =1.23554e+20$	[SR, QUANTUM]
ST. 295	symb	$l=\sqrt{(hG/c^3)}\cdot\int\int(1/(y\log y)^{\{1/2\}})\,dy_m/\int\int(1/(x\log x)^2)\,dx_m=1/i$	[ZETA, MANIFOLD, QUANTUM]
ST. 296	symb	$i\hbar c=0,\,\hbar c=G/i,\,(1/2)/((1/2)i)=\nabla_1/\nabla_2\leq 1$	[QUANTUM]
ST. 297	symb	$A=BQ+R,\,ds^2=e^{\{-2\pi T \phi \}}[\eta_\mu\nu+\hbar_\mu\nu(x)]dx^\mu dx^\nu+T^2\,d^2\phi$	[TRANSPORT]
ST. 298	holds	$ds^2=g_\mu\nu\,dx^\mu dx^\nu+\kappa^2(A^{\sim\{\mu\nu\}})^2,\,\int\int e^{\{-x^2-y^2\}}\,dx\,dy=\pi \qquad =3.14159\ \mathrm{vs}\ 3.14159$	[SR, TRANSPORT]
ST. 299	symb	$\Gamma(x)=\int e^{\{-x\}}x^{\{1-t\}}\,dx=\delta(x)\pi(x)f^{\mathrm{n}}(x)$	[GAMMA]
ST. 300	symb	$ds^2=[T^2\,d^2\phi]$	[MANIFOLD]
ST. 301	symb	$0(x)=[x]$	[MANIFOLD]
ST. 302	symb	$\nabla\phi^2=8\pi G(p/c^3\circ V/S)$	[ENTROPY]
ST. 303	symb	$pV/S=h$	[QUANTUM]
ST. 304	differs	$\beta(p,q)=\Gamma(p)\Gamma(q)/\Gamma(p+q)\cong\Gamma(p+q)/(\Gamma(p)\Gamma(q)) \qquad =0.0833333\ \mathrm{vs}\ 12$	[GAMMA, BETA]
ST. 305	symb	$\ker f/\mathrm{im}\,f\cong\mathrm{im}\,f/\ker f\ (\times 2)$	[MANIFOLD]
ST. 306	symb	$\otimes\nabla g(x)=[\nabla_i\nabla_j\int\nabla f(x)d\eta],\,\bigcup_{\{k=0\}}^{\sim\infty}(\otimes\nabla f(x))=\Box\iiint\nabla g(x)d\eta$	[MANIFOLD]
ST. 307	calc	$a'=\sqrt{(v/(1-(v/c)^2))},\,F=ma' \qquad =16764.7$	[SR]
ST. 308	symb	$\nabla f(x)=\int_M\Box(\otimes\nabla f(x))^{\mathrm{n}}\,dm$	[MANIFOLD]
ST. 309	symb	$\Box=2(T-t) R_{ij}+\nabla\nabla f-g_{ij}^2/(2(T-t)) $	[MANIFOLD, ENTROPY]
ST. 310	symb	$(\Box+m)\cdot\phi=0$	[SR, QUANTUM]

4. 全方程式レジストリ (2111 本) p.13/33

ST.311	symb	$\square \times \square = (\square + m^2) \cdot \phi, \ (\partial \gamma^{\wedge} n + \delta \phi) \cdot \phi = 0$	[SR, QUANTUM]
ST.312	symb	$\nabla_{\neg} i \nabla_{\neg} j \int \int_{\neg} M \nabla f(t) dt = \square (\cup_{\neg} \{k=0\}^{\neg} \infty \otimes \nabla g(x) d \eta)$	[MANIFOLD]
ST.313	symb	$\int_{\neg} M \left(l \times l \right) dm = \sum l \otimes l d \eta$	[MANIFOLD]
ST.314	symb	$= [D^2 \phi \otimes h_{\neg} \mu \nu]^{\neg} \{1/2 + i y\} = H_3(M_{\neg} 1)$	[ZETA, MANIFOLD]
ST.315	symb	$y = \nabla_{\neg} i \nabla_{\neg} j \int \nabla g(x) dx, \ y' - y + (d/dx)z + [n] = 0$	[OTHER]
ST.316	holds	$z = \cos x + i \sin x = e^{\neg} i \theta \} \quad = 0.866025 + 0.5i \text{ vs } 0.866025 + 0.5i$	[ROT]
ST.317	symb	$T^{\neg} \{ \mu \nu \} = -2R_{\neg} ij, \ d/dt \ g_{\neg} ij(t) = -2R_{\neg} ij$	[SR, MANIFOLD]
ST.318	symb	$G_{\neg} \mu \nu = R_{\neg} \mu \nu \ T^{\neg} \{ \mu \nu \}, \ \sigma(x) \otimes \delta(x) = (E_{\neg} n^+ \times H_m)$	[SR, MANIFOLD]
ST.319	symb	$G_{\neg} \mu \nu = [\partial R_{\neg} ij / \partial f]^2, \ \delta(x) \cdot V(x) = \lim_{\neg} \{n \rightarrow 1\} \ \delta(x)$	[SR, MANIFOLD]
ST.320	calc	$\lim_{\neg} \{n \rightarrow \infty\} \ mesh \ V(x) = m/(m + 1) \quad = 0.9$	[MANIFOLD]
ST.321	symb	$V(x) = \sigma \cdot S^2(x), \ \iiint V(x)/S^2(x) \ dm = [D^2 \phi \otimes h_{\neg} \mu \nu]$	[MANIFOLD]
ST.322	symb	$g(x) \lfloor_{\neg} \{ \delta(x,y) \} = d/dt \ g_{\neg} ij(t), \ \sigma(x,y) \cdot g(x) \lfloor_{\neg} \{ \delta(x,y) \} = R_{\neg} ij \lfloor_{\neg} \{ \sigma(x,y) \} = \int \ R_{\neg} ij^{\neg} \{ a(x-y)^{\neg} n + r^{\neg} n \}$	[MANIFOLD]
ST.323	symb	$(ux + vy + wz)/\Gamma = \int \ R_{\neg} ij^{\neg} \{ (x-u)(y-v)(z-w) \} \ dV$	[GAMMA, MANIFOLD]
ST.324	symb	$(\square + m) \cdot \phi = 0, \ E = mc^2, \ \partial / \partial f \ \square \phi = 4 \pi G \rho$	[SR, QUANTUM]
ST.325	differs	$(\partial \gamma^{\wedge} n + m) \cdot \phi = 0, \ E = mc^2 - (1/2)mv^2 = (- (1/2)(v/c)^2 + m) \cdot c^2 = (- (1/2)a^2 + m) \cdot c^2, \ F = ma, \ \int \ a \ dx = (1/2)a^2 + C \quad = 1.69751e+17 \text{ vs } 1.74751e+17$	[SR, QUANTUM, TRANSPORT]
ST.326	symb	$T^{\neg} \{ \mu \nu \} = - (1/2)a^2, \ (e^{\neg} \{ i \theta \})' = ie^{\neg} \{ i \theta \}$	[ROT, SR]
ST.327	symb	$Z \supset C \otimes \nabla R^+, \ \nabla(R^+ \cap E^+) \ni x, \ \Delta(C \subset R) \ni x, \ M \pm \otimes R^+, \ E^+ \in \otimes \nabla R^+, \ S \pm \subset R_{\neg} 2, \ V \pm \times R \pm \equiv V/S, \ C^+ \cup V \pm \ni M_{\neg} 1 \otimes \nabla C \pm, \ Q \supseteq R \pm$ $Q \subset \otimes M \pm, \ \otimes Q \subset \xi(x), \ \otimes \nabla C \pm \equiv M_{\neg} 3, \ R \subset M_3, \ C^+ \otimes M_{\neg} n, \ E^+ \cap R^+, \ E_2 \otimes E_1, \ R^{-} \subset C^+ (\times 2)$	[ZETA, MANIFOLD, QUANTUM]
ST.328	symb	$[-\Delta v + \nabla_{\neg} i \nabla_{\neg} j \ v_{\neg} ij - R_{\neg} ij \ v_{\neg} ij - v_{\neg} ij \nabla_{\neg} i \nabla_{\neg} j + 2 \langle \nabla f, \nabla h \rangle + (R + \nabla^2 f)(v/2 - h)] \ (\times 2)$	[MANIFOLD, ENTROPY]
ST.329	symb	$S^3, \ H' \times E', \ E', \ S' \times E', \ S^2 \times E', \ H' \times S', \ H', \ S^2 \times E \ (\times 3)$	[MANIFOLD]
ST.330	symb	$V(\int \nabla_{\neg} i \nabla_{\neg} j (R + \Delta f)^2) \ / \ \exists (R + \Delta f)$	[MANIFOLD]
ST.331	symb	$\exp[\int \int (R + \Delta f)^2 \ e^{\neg} \{-x \log x\} \ dV]$	[MANIFOLD, QUANTUM]
ST.332	calc	$imaginary.equation \Rightarrow e^{\neg} \{ \cos \theta + i \sin \theta \} \quad = 2.0864 + 1.13981i$	[ROT]
ST.333	symb	$d/df \ F \Longrightarrow d/df \ \int \int \ 1/((x \log x)^2 \circ (y \log y)^{\neg} \{1/2\}) \ dm$	[ZETA, MANIFOLD]
ST.334	symb	$x^{\neg} \{1/2 + i y\} = [f(x) \circ g(x), \ h(x)] \ / \ \partial f \partial g \partial h \ (\times 2)$	[ZETA]
ST.335	symb	$x^{\neg} \{1/2 + i y\} = \exp[\int \nabla_{\neg} i \nabla_{\neg} j \ f(g(x)) \ g' \ (x) \ / \ \partial f \partial g] \ (\times 2)$	[ZETA]
ST.336	symb	$0(x) = \{[f(x) \circ g(x), \ h(x)], \ g^{-1}(x)\} \ (\times 2)$	[MANIFOLD]
ST.337	symb	$\exists [\nabla_{\neg} i \nabla_{\neg} j (R + \Delta f), \ g(x)] = \otimes_{\neg} \{k=0\}^{\neg} \infty \ \nabla \int \nabla_{\neg} i \nabla_{\neg} j \ f(x) \ dm \ (\times 2)$	[MANIFOLD]
ST.338	symb	$V(\nabla_{\neg} i \nabla_{\neg} j \ f) = \otimes \nabla E^+ \ (\times 2)$	[MANIFOLD]
ST.339	symb	$g(x,y) = 0(x)[f(x) + \bar{h}(x)] + T^2 \ d^2 \phi \ (\times 2)$	[MANIFOLD, TRANSPORT]
ST.340	symb	$0(x) = (\int [g(x)] \ e^{\neg} \{-f\} \ dV)' \ - \ \Sigma \ \delta(x) \ (\times 2)$	[MANIFOLD]
ST.341	symb	$0(x) = [\nabla_{\neg} i \nabla_{\neg} j \ f(x)]' \ \equiv n C_r \ f(x)^{\neg} n \ f(y)^{\neg} \{n-r\} \ \delta(x,y), \ V(\tau) = \int [f(x)] \ dm / \partial f_{.xy} \ (\times 2)$	[MANIFOLD]
ST.342	symb	$\square \phi = 8 \pi G \ T^{\neg} \{ \mu \nu \}, \ (\square \phi)' = \nabla_{\neg} i \nabla_{\neg} j (\delta(x) \circ G(x))^{\neg} \{ \mu \nu \} (p/c^2 \circ V/S), \ x^{\neg} \{1/2 + i y\} = e^{\neg} \{x \log x\} \ (\times 2)$	[SR, ZETA]
ST.343	symb	$\delta(x) \phi = V[\nabla_{\neg} i \nabla_{\neg} j \ f \circ g(x)] \ / \ \exists (R + \Delta f) \ (\times 2)$	[MANIFOLD]
ST.344	symb	$-n C_r = _{\neg} \{ (1/i) H \phi \} C_{\neg} \{ h \phi \} + _{\neg} \{ [H, \phi] \} C_{\neg} \{ -n-r \} \ (\times 2)$	[QUANTUM]
ST.345	holds	$n C_r = n C_{\neg} \{n-r\} \ (\times 2) \quad = 35 \text{ vs } 35$	[OTHER]
ST.346	symb	$\int \int \ 1/(x \log x)^2 \ dx_m \rightarrow 0(x) = [\nabla_{\neg} i \nabla_{\neg} j \ f]' \ / \partial f_{.xy} \ (\times 2)$	[MANIFOLD]
ST.347	symb	$\cup_{\neg} \{x=0\}^{\neg} \infty \ f(x) = \nabla_{\neg} i \nabla_{\neg} j \ f(x) \otimes \Sigma f(x) = \otimes \nabla f(x) \ (\times 2)$	[MANIFOLD]
ST.348	symb	$\nabla_{\neg} i \nabla_{\neg} j \ f \equiv \partial x \partial y \int \nabla_{\neg} i \nabla_{\neg} j \ f \ dm \equiv \int [f(x)] dm \equiv \{[f(x), g(x)], \ g^{-1}(x)\} \equiv \square \phi \equiv \nabla \phi^2 \equiv f(x \circ y) \leq f(x) \circ g(x) \equiv f(x) + g(x) \ (\times 2)$	[MANIFOLD]
ST.349	symb	$\delta(x) \phi = \langle f, g \rangle \circ h^{-1}(x) \ (\times 2)$	[QUANTUM]
ST.350	symb	$\partial f_{.x} \cdot \delta(x) \phi = x \ (\times 2)$	[OTHER]
ST.351	symb	$x \in 0(x) \ (\times 2)$	[MANIFOLD]
ST.352	symb	$0(x) = \{[f \circ g, \ h^{-1}(x)], \ g(x)\} \ (\times 2)$	[MANIFOLD]
ST.353	symb	$\lim_{\neg} \{n \rightarrow \infty\} \ \Sigma_{\neg} \{k=n\}^{\neg} \infty \ \nabla f = [\nabla \int \nabla_{\neg} i \nabla_{\neg} j \ f(x) \ dx_m, \ g^{-1}(x)] \rightarrow \otimes_{\neg} \{k=0\}^{\neg} \infty \ \nabla E \pm = M_3 = \otimes_{\neg} \{k=0\}^{\neg} \infty \ E \pm \ (\times 2)$	[MANIFOLD]
ST.354	symb	$dx^2 = [g^2_{\neg} \mu \nu, \ dx], \ g^{-1} = dx \int \ \delta(x) f(x) dx \ (\times 2)$	[MANIFOLD]
ST.355	symb	$f(x) = \exp[\nabla_{\neg} i \nabla_{\neg} j \ f(x), \ g^{-1}(x)] \ (\times 2)$	[MANIFOLD]
ST.356	symb	$(g(x)/f(x))' = \lim_{\neg} \{n \rightarrow \infty\} \ g(x)/f(x) = g' \ (x)/f' \ (x) \ (\times 2)$	[OTHER]
ST.357	symb	$\nabla F = f \cdot (1/4) r ^2 \ (\times 2)$	[MANIFOLD]
ST.358	symb	$\nabla_{\neg} i \nabla_{\neg} j \ f = (d/dx_i)(d/dx_j) \ f(x) g(x) \ (\times 2)$	[OTHER]
ST.359	symb	$D^2 \phi = \nabla \int (\nabla_{\neg} i \nabla_{\neg} j \ f)^2 \ d \eta \ (\times 2)$	[MANIFOLD]
ST.360	calc	$E = mc^2, \ E = (1/2)mv^2 - (1/2)kx^2, \ G^{\neg} \{ \mu \nu \} = (1/2) \Lambda g_{\neg} ij, \ \square = (1/2)kT^2 \ (\times 2) \quad = 7$	[SR, OTHER]
ST.361	symb	$ker \ f/im \ f \equiv S_m^{\neg} \{ \mu \nu \}, \ S_m^{\neg} \{ \mu \nu \} = \pi(\chi, x) \otimes h_{\neg} \mu \nu \ (\times 2)$	[MANIFOLD]
ST.362	symb	$D^2 \phi = 0(x)(p/c^3 + V/S), \ V(x) = D^2 \phi \otimes M^+_3 \ (\times 2)$	[MANIFOLD, ENTROPY]
ST.363	symb	$\nabla_{\neg} i \nabla_{\neg} j [S_1^{\neg} \{mn\} \otimes S_2^{\neg} \{mn\}] = \int \ (V(\tau)/f(x)) [D^2 \phi] \ (\times 2)$	[MANIFOLD]
ST.364	symb	$\nabla_{\neg} i \nabla_{\neg} j [S_1^{\neg} \{mn\} \otimes S_2^{\neg} \{mn\}] = \int \ (V(\tau)/f(x)) \ 0(x) \ (\times 2)$	[MANIFOLD]
ST.365	symb	$z(x) = (g(cx + d)/f(ax + b)) \ h(ex + l) = \int \ (V(\tau)/f(x)) \ 0(x) \ (\times 2)$	[MANIFOLD]
ST.366	symb	$V(x)/f(x) = m(x), \ 0(x) = m(x) [D^2 \phi(x)] \ (\times 2)$	[MANIFOLD]
ST.367	symb	$d/df \ F = m(x), \ \int F \ dx_m = \Sigma_{\neg} \{k=0\}^{\neg} \infty \ m(x) \ (\times 2)$	[MANIFOLD]
ST.368	symb	$ds^2 = [g^2_{\neg} \mu \nu, \ dx] \ (M_2) \ (\times 2)$	[MANIFOLD]
ST.369	symb	$ds^2 = g^{-1}_{\neg} \mu \nu (g^2_{\neg} \mu \nu(x) - dx \ g^2_{\neg} \mu \nu) = h(x) \otimes g_{\neg} \mu \nu \ d^2 x - h(x) \otimes dx \ g_{\neg} \mu \nu(x), \ h(x) = (f^2(\mathfrak{X}) - \vec{E}^+) \ (M_2) \ (\times 2)$	[MANIFOLD, TRANSPORT]
ST.370	symb	$G_{\neg} \mu \nu = R_{\neg} \mu \nu \ T^{\neg} \{ \mu \nu \}, \ \partial M_2 = \otimes \nabla C \pm \ (\times 2)$	[SR, MANIFOLD]
ST.371	symb	$G_{\neg} \mu \nu = R_{\neg} \mu \nu, \ d/dt \ g_{\neg} ij = -2R_{\neg} ij \ (\times 2)$	[SR, MANIFOLD]
ST.372	holds	$r = 2f^{\neg} \{1/2\}(x) \ (\times 2) \quad = 3 \text{ vs } 3$	[MANIFOLD]
ST.373	symb	$E^+ = f^{-1} x f(x), \ h(x) \otimes g(\mathfrak{X}) \equiv V/S, \ R/M_2 = E^+ - \{ \phi \} = M_3 \supset R \ (\times 2)$	[MANIFOLD]
ST.374	symb	$M^+_2 = E^+_1 \cup E^+_2 \rightarrow E^+_1 \otimes E^+_2 = M_1 \otimes \nabla C \pm, \ (E^+_1 \otimes E^+_2) \cdot (R^{-} \subset C^+) \ (\times 2)$	[MANIFOLD]
ST.375	symb	$R/M_2 = E^+ - \{ \phi \} = M_3 \supset R \ (\times 2)$	[MANIFOLD]
ST.376	symb	$M^+_3 \equiv h(x) \cdot R^+_3 = \otimes \nabla C \pm, \ R = E^+ \otimes M_2 - (E^+ \cap M_2) \ (\times 2)$	[MANIFOLD]

ST. 377	symb	$E^+=g_{\mu\nu}\,dx_{\mu}\,dx_{\nu},\,M_2=g_{\mu\nu}\,d^2x\,(\times 2)$	[MANIFOLD]
ST. 378	calc	$F=\rho\,g\,l\rightarrow V/S\,(\times 2)\quad=19600$	[OTHER]
ST. 379	symb	$0(x)=\delta(x)[f(x)+g(\bar{x})]+\rho\,g\,l\,(\times 2)$	[OTHER]
ST. 380	calc	$F=(1/2)mv^2-(1/2)kx^2,\,M_2=P^{\{2n\}}\,(\times 2)\quad=7$	[MANIFOLD, OTHER]
ST. 381	calc	$f(x)=(1/4) r ^2\,(\times 2)\quad=4$	[MANIFOLD]
ST. 382	symb	$V=R^+\Sigma K_m,\,W=C^+\Sigma_{\{k=0\}^\infty}K_{\{n+2\}},\,V/W=R^+\Sigma K_m/\,C^+\Sigma K_{\{n+2\}}=(R^+/C^+)\Sigma\,x^{\textstyle k}/(a_k\,f^{\textstyle k}(x))=M\pm\,(\times 2)$	[MANIFOLD]
ST. 383	symb	$d/df\,F=m(x)\rightarrow M\pm,\,\Sigma_{\{k=0\}^\infty}x^{\textstyle k}/(a_k\,f^{\textstyle k}(x))=a_k\,x^{\textstyle k}/\zeta(x)\,(\times 2)$	[ZETA]
ST. 384	symb	$\{f,g\}/[f,g]=(fg+gf)/(fg-gf)\,(\times 2)$	[QUANTUM]
ST. 385	symb	$\nabla f=2,\,\partial H_3=2,\,(1+f)/(1-f)=1\,(\times 2)$	[MANIFOLD]
ST. 386	symb	$d/df\,F=\otimes\nabla C\pm,\,\bar{F}=1/2\,(\times 2)$	[MANIFOLD]
ST. 387	symb	$H_1\cong H_3=M_3\,(\times 2)$	[MANIFOLD]
ST. 388	symb	$H_3\cong H_1\rightarrow\pi(\chi,x),\,H_n,H_m=\text{rank}(m,n),\,\text{mesh}(\text{rank}(m,n))\,\lim\text{mesh}\rightarrow 0\,(\times 2)$	
ST. 389	symb	$(fg)'=fg'+gf',\,(f/g)'=(f'\,g-g'\,f)/g^2\,(\times 2)$	[OTHER]
ST. 390	symb	$\{f,g\}/[f,g]=(fg)'\,\otimes dx_{fg}/\,((f/g)'\,\otimes g^{-2}dx_{fg})=d/df\,F\,(\times 2)$	[QUANTUM]
ST. 391	symb	$\hbar\phi=(1/i)H\Psi,\,i[H,\phi]=-H\Psi,\,\{f,g\}/[f,g]=(i)^2\,(\times 2)$	[QUANTUM]
ST. 392	symb	$[\nabla_i\nabla_j\,f(x),\,\delta(x)]=\nabla_i\nabla_j\,\int\,f(x,y)\,d\mathbf{m}_{xy},\,f(x,y)=[f(x),h(x)]\times[g(x),h^{-1}(x)]\,(\times 2)$	[MANIFOLD, QUANTUM]
ST. 393	symb	$\delta(x)=1/f'(x),\,[H,\phi]=\Delta f(x)\,(\times 2)$	[QUANTUM]
ST. 394	symb	$0(x)=\nabla_i\nabla_j\,\int\,\delta(x)f(x)\,dx\,(\times 2)$	[MANIFOLD]
ST. 395	symb	$0(x)=\int\,\delta(x)f(x)\,dx\,(\times 2)$	[MANIFOLD]
ST. 396	symb	$R^+\cap E\pm\supset x,\,M\times R^+\supset M_3,\,Q\supset C\pm,\,Z\in Q\nabla f,\,f\equiv\otimes_{\{k=0\}^n}\nabla C\pm\,(\times 2)$	[MANIFOLD]
ST. 397	symb	$\otimes_{\{k=0\}^\infty}\nabla C\pm=M_1,\,\otimes_{\{k=0\}^\infty}\nabla M\pm\equiv E\pm\,(\times 2)$	[MANIFOLD]
ST. 398	symb	$M_3\equiv M_1\otimes_{\{k=0\}^\infty}\nabla(V\pm/S)\,(\times 2)$	[MANIFOLD]
ST. 399	symb	$P^{\{2n\}}/M_2\equiv\otimes_{\{k=0\}^\infty}\nabla C\pm,\,E\pm\times R\pm\equiv M_2\,(\times 2)$	[MANIFOLD]
ST. 400	symb	$\zeta(x)=P^{\{2n\}}\times\Sigma_{\{k=0\}^\infty}a_k\,x^{\textstyle k}\,(\times 2)$	[ZETA, MANIFOLD]
ST. 401	symb	$M_2\equiv P^{\{2n\}}/\ker f\rightarrow\otimes\nabla C\pm\,(\times 2)$	[MANIFOLD]
ST. 402	symb	$S\pm\times V\pm\equiv(V/S)\,\otimes_{\{k=0\}^\infty}\nabla C\pm,\,V^+\equiv M\pm\otimes S\pm\,(\times 2)$	[MANIFOLD]
ST. 403	symb	$Q\times M_1\subset\otimes\nabla C\pm\,(\times 2)$	[MANIFOLD]
ST. 404	symb	$\Sigma_{\{k=0\}^\infty}Z\otimes Q\pm\equiv\otimes_{\{k=0\}^\infty}\nabla M_1\equiv\otimes_{\{k=0\}^\infty}\nabla C\pm\times\Sigma_{\{k=0\}^\infty}M_1,\,x\in R^+\times C\pm\supset M_1,\,M_1\subset M_2\subset M_3\,(\times 2)$	[MANIFOLD]
ST. 405	symb	$\otimes\nabla C\pm\equiv M_3,\,R\supset 0,\,R\cap 0,\,R\subset M_3,\,C^+\otimes M_n,\,E^+\cap R^+,\,M\pm\nabla C\pm,\,C^+\nabla H_m,\,E^+\nabla R\pm,\,E_2\nabla E_1,\,R^-\nabla C\pm\,(\times 2)$	[MANIFOLD]
ST. 406	symb	$(\nabla/\Delta)\int xf(x)dx,\,\nabla R/\Delta f,\,\square=2\int\,((R+\nabla_i\nabla_j\,f)^2/(-(R+\Delta f)))\,e^{-\{f\}}\,dV\,(\times 2)$	[MANIFOLD, ENTROPY]
ST. 407	symb	$\square=\nabla R/\Delta f,\,d/dt\,g_{ij}=\square\rightarrow\nabla f/\Delta x,\,(R+ \nabla f ^2)d\mathbf{m}\rightarrow-2(R+\nabla_i\nabla_j\,f)^2\,e^{-\{f\}}\,dV\,(\times 2)$	[MANIFOLD]
ST. 408	symb	$x^{\textstyle n}+y^{\textstyle n}=z^{\textstyle n}\rightarrow\nabla\phi^2=8\pi G\,T^{\{\mu\nu\}},\,f(x+y)\geq f(x)\circ f(y)\,(\times 2)$	[SR, OTHER]
ST. 409	symb	$\text{im}\,f/\ker f=\partial f,\,\ker f=\partial f,\,\ker f/\text{im}\,f\equiv\partial f,\,\ker f=f^{-1}(x)xf(x)\,(\times 2)$	[MANIFOLD]
ST. 410	symb	$f^{-1}(x)xf(x)=\int\partial f(x)\,d(\ker f)\rightarrow\nabla f=2\,(\times 2)$	[MANIFOLD]
ST. 411	symb	$n\mathbb{C}_r=n\mathbb{C}_{\{n-r\}}\rightarrow\text{im}\,f/\ker f\equiv\ker f/\text{im}\,f\,(\times 2)$	[MANIFOLD]
ST. 412	symb	$\Sigma_{\{k=0\}^\infty}a_k\,f^{\textstyle k}=T^2\,d^2\phi,\,a_k\equiv\Sigma_{\{r=0\}^\infty}n\mathbb{C}_r\,(\times 2)$	[MANIFOLD, TRANSPORT]
ST. 413	symb	$V/W=(R/C)\Sigma_{\{k=0\}^\infty}x^{\textstyle k}/(a_k\,f^{\textstyle k}),\,W/V=(C/R)\Sigma_{\{k=0\}^\infty}a_k\,f^{\textstyle k}/x^{\textstyle k}\,(\times 2)$	[MANIFOLD]
ST. 414	symb	$V/W\equiv W/V\equiv(R/C)(\Sigma_{\{r=0\}^\infty}n\mathbb{C}_r)^{-1}\,\Sigma_{\{k=0\}^\infty}x^{\textstyle k}\,(\times 2)$	[MANIFOLD]
ST. 415	symb	$W/V=xF(x),\,\chi(x)=(-1)^k\,a_k,\,\Gamma(x)=\int e^{-\{x\}}\,x^{\textstyle\{1-t\}}\,dx\,(\times 2)$	[GAMMA, MANIFOLD]
ST. 416	symb	$\Sigma_{\{k=0\}^n}a_k\,f^{\textstyle k}=(f^{\textstyle k})'\,=\Sigma_{\{k=0\}^\infty}n\mathbb{C}_r\,f^{\textstyle k}\,(\times 2)$	[OTHER]
ST. 417	holds	$\Sigma_{\{k=0\}^\infty}a_k\,f^{\textstyle k}=[f(x)],\,\Sigma\,a_k\,f^{\textstyle k}=\alpha,\,\Sigma\,1/(a_k\,f^{\textstyle k}),\,\Sigma(a_k\,f^{\textstyle k})^{-1}=1/(1-z)\,(\times 2)\quad=2\,\text{vs}\,2$	[OTHER]
ST. 418	symb	$\int\int\,1/(x\log x)(y\log y)\,dxy=n\mathbb{C}_r\,xy/\,(n\mathbb{C}_{\{n-r\}}(x\log x)(y\log y))^{-1}=(n\mathbb{C}_{\{n-r\}})^2\,\Sigma_{\{k=0\}^\infty}(1/(x\log x)-1/(y\log y))\,d(1/(nxy))\times xy=\Sigma\,a_k\,f^{\textstyle k}=\alpha\,(\times 2)$	[MANIFOLD]
ST. 419	symb	$\ker f=-2\int(R+\nabla_i\nabla_j\,f)^2\,e^{-\{f\}}\,dV,\,\ker f/\text{im}\,f\leq d/df\,F$	[MANIFOLD, ENTROPY]
ST. 420	symb	$e^{-\{f\}}[2\int(R+\nabla f^2)/(-(R+\Delta f))\,e^{-\{f\}}\,dV]$	[MANIFOLD, ENTROPY]
ST. 421	symb	$p_x=mv,\,\lambda(x)=\text{esperial}\,f(x),\,[F(x)]=1/(1-z),\,0<\tan\theta<i,\,-1<\tan\theta<i,\,-1\leq\sin\theta\leq 1,\,-1\leq\cos\theta\leq 1$	[ROT, QUANTUM]
ST. 422	symb	$R\nabla E^+=f(x)\nabla e^{\{x\log x\}}\,(\times 2)$	[MANIFOLD]
ST. 423	symb	$Q\nabla C^+=(d/df)F(x)\,\nabla\int\delta(x)f(x)dx$	[MANIFOLD]
ST. 424	symb	$E^+\nabla f=e^{\{x\log x\}}\,\nabla n!\,f(x)/E(X)$	[GAMMA]
ST. 425	symb	$(u+v+w)(x+y+z)/\Gamma$	[GAMMA, MANIFOLD]
ST. 426	symb	$\exp(\nabla(R^+\cap E^+),\,\Delta(C\supset R))=\pi(R,\,C\nabla E^+)=\text{rot}(E_1,\,\text{div}\,E_2)\,(\times 3)$	[ROT, MANIFOLD]
ST. 427	symb	$xf(x)=F(x)\,(\times 3)$	[OTHER]
ST. 428	symb	$\square x=\int\,f(x)/\nabla(R^+\cap E^+)\,d\square x=\int\,(\Delta f(x)\circ E^+)/\nabla(R^+\cap E^+)\,\square x$	[QUANTUM, TRANSPORT]
ST. 429	symb	$\exp(\nabla(R^+\cap E^+),\,\Delta(C\supset R))$	[MANIFOLD]
ST. 430	symb	$d(R\nabla E^+)=\Delta f(x)\circ E^+(x)$	[MANIFOLD]
ST. 431	symb	$\square x=\int\,d(R\nabla E^+)/\nabla(R^+\cap E^+)\,d\square x$	[MANIFOLD]
ST. 432	symb	$\square x=\int\,\nabla_i\nabla_j(R+E^+)/\nabla(R^+\cap E^+)\,d\square x$	[MANIFOLD]
ST. 433	symb	$x^{\textstyle n}+y^{\textstyle n}=z^{\textstyle n}\rightarrow\square x=\int\,\nabla_i\nabla_j(R+E^+)/\nabla(R^+\cap E^+)\,d\square x$	[MANIFOLD, TRANSPORT]
ST. 434	symb	$\square=-2(T-t) R_{ij}+\nabla\nabla f+g_{ij}^2/(-2(T-t)) $	[MANIFOLD, ENTROPY]
ST. 435	symb	$\square=-2\int\,((R+\nabla_i\nabla_j\,f)/(-(R+\Delta f)))\,e^{-\{f\}}\,dV$	[MANIFOLD, ENTROPY]
ST. 436	symb	$d/dt\,g_{ij}=-2R_{ij},\,d/df\,F=m(x)$	[MANIFOLD]
ST. 437	symb	$R\nabla E^+=\Delta f(x)\circ E^+(x),\,d(R\nabla E^+)=\nabla_i\nabla_j(R+E^+)$	[MANIFOLD]
ST. 438	symb	$R\nabla E^+=f(x)\nabla e^{\{x\log x\}},\,R\nabla E^+=f(x),\,f(x)=M_1$	[MANIFOLD]
ST. 439	symb	$-2(T-t) R_{ij}+\nabla\nabla f=\int\int\,1/(y\log y)^{\{1/2\}}\,dy_m$	[ZETA, ENTROPY]
ST. 440	symb	$(1/(-2(T-t))) g_{ij}^2 =\int\int\,1/(x\log x)^2\,dx_m$	[ZETA, ENTROPY]
ST. 441	symb	$(\square+m^2)\phi=0,\,(\gamma^{\textstyle n}\partial_n+m^2)\phi=0$	[SR, QUANTUM]
ST. 442	calc	$\square=(\delta\phi+m^2)\phi,\,(\delta\phi+m^2c)\phi=0,\,E=mc^2,\,E=-(1/2)mv^2+mc^2,\,\square\phi^2=(\partial\phi+m^2)\phi\quad=8.98755\text{e}+16$	[SR, QUANTUM, TRANSPORT]
ST. 443	symb	$\square\phi^2=(8\pi G/c^4)T^{\{\mu\nu\}},\,\nabla\phi^2=8\pi G(p/c^3+V/S),\,d/dt\,g_{ij}=-2R_{ij},\,f(x)+g(x)\geq f(x)\circ g(x)$	[SR, ENTROPY]

4. 全方程式レジストリ (2111 本) p.15/33

ST.444	symb	$\int_{-1}^1 \beta^a x^a (x-1)(y-1) \geq 2 \int f(x)g(x)dx$	[OTHER]
ST.445	calc	$m^2 = 2\pi T(26 - D_n)/24, r_n = 1/(1-z) = 5.75959$	[QUANTUM]
ST.446	symb	$(\partial \phi + m^2 c) \phi = 0, T^{\{\mu \nu\}} = (1/2)mv^2 - (1/2)kx^2$	[QUANTUM]
ST.447	symb	$\int [f(x)]dx = \int f(x)dx , \int \nabla x \, dx = \Delta \Sigma f(x)dx, (e^{\{-i \theta\}})' = ie^{\{i \theta\}}$	[ROT]
ST.448	calc	$T^{\{\mu \nu\}} = nh \nu = 3.31304e-18$	[QUANTUM]
ST.449	differences	$T^{\{\mu \nu\}} = (1/2)mv^2 - (1/2)kx^2 \geq mc^2 - (1/2)mv^2 = 0 \text{ vs } 1$	[SR, TRANSPORT]
ST.450	symb	$Z \in R \cap Q, R \subset M_+, C^+ \otimes \{k=0\}^n H_m, E^+ \cap R^+ (\times 2)$	[MANIFOLD]
ST.451	symb	$M_+ = \Sigma_{\{k=0\}^n} C^+ \otimes H_M, M_+ = \Sigma_{\{k=0\}^n} C^+ \cup H_+ (\times 2)$	[MANIFOLD]
ST.452	symb	$E_2 \otimes E_1, R^- \subset C^+, M_{\pm}, G_{\pm}; M_{\pm} \nabla C_{\pm}, C^+ \nabla H_m, E^+ \nabla R^+; E_1 \nabla E_2, R^- \nabla C^+, \otimes \nabla M_{\pm}, \otimes \nabla C_{\pm}, R \supset Q (\times 2)$	[MANIFOLD]
ST.453	symb	$d/df \, F = \otimes \nabla M_{\pm}, \otimes \nabla C_{\pm}$	[MANIFOLD]
ST.454	symb	$\nabla \circ \Delta = \nabla \Sigma f(x)$	[MANIFOLD]
ST.455	symb	$\Delta \rightarrow mesh \, f(x)dx, \partial x$	[MANIFOLD]
ST.456	symb	$\nabla \rightarrow \partial_{xy}, (d/dx)f(x)(d/dx)g(x)$	[OTHER]
ST.457	symb	$\Box x = -[f, g]$	[QUANTUM]
ST.458	symb	$d/dt \, g_{ij} = -2R_{ij}$	[MANIFOLD]
ST.459	symb	$\lim_{x \rightarrow \infty} \Sigma f(x)dx = \int dx, \nabla \int dx$	[OTHER]
ST.460	symb	$(E_2 \otimes_{\{k=0\}^n} E_1) \cdot (R^- \subset C^+) = \otimes_{\{k=0\}^n} \nabla C_{\pm} (\times 2)$	[MANIFOLD, QUANTUM]
ST.461	symb	$\bigvee \int C_{\pm} \nabla H_m / \Delta (M_{\pm} \nabla C_{\pm}) = \wedge M_{\pm} \otimes_{\{k=0\}^n} C_{\pm} (\times 2)$	[MANIFOLD]
ST.462	symb	$\exists (M_{\pm} \nabla C_{\pm}) = XOR(\otimes_{\{k=0\}^n} \nabla M_{\pm}) (\times 2)$	[QUANTUM]
ST.463	symb	$-[E^+ \nabla R_{\pm}] = \nabla_+ \nabla_- C_{\pm} (\times 2)$	[QUANTUM]
ST.464	differences	$(\cos x \sin x; \sin x - \cos x)(x; y) = (1 \, 0; 0 \, -1) = 0.95345 - 0.604226i \text{ vs } 1 - 1i$	[ROT]
ST.465	differences	$(\cos x - 1; 1 - \sin x)(\cos x; \sin x) = (1 \, 0; 0 \, -1) = 0.25 + 0.616025i \text{ vs } 1 - 1i$	[ROT]
ST.466	holds	$\Sigma_{\{k=0\}^n} \cos k \theta = (\sin((n+1)\theta/2)/\sin(\theta/2)) \cos(n\theta/2) = -0.448716 \text{ vs } -0.448716$	[ROT]
ST.467	holds	$\Sigma_{\{k=0\}^n} \sin k \theta = (\sin((n+1)\theta/2)/\sin(\theta/2)) \sin(n\theta/2) = 2.47708 \text{ vs } 2.47708$	[ROT]
ST.468	holds	$\sin \theta = (e^{\{i \theta\}} - e^{\{-i \theta\}})/(2i), \cos \theta = (e^{\{i \theta\}} + e^{\{-i \theta\}})/2, \tan \theta = (e^{\{i \theta\}} - e^{\{-i \theta\}})/(i(e^{\{i \theta\}} + e^{\{-i \theta\}})) = 0 \text{ vs } 0$	[ROT]
ST.469	differences	$\lim_{\{\theta \rightarrow 0\}} \sin \theta / \theta \rightarrow 1, \lim_{\{\theta \rightarrow 0\}} \cos \theta / \theta \rightarrow 1 = 1 + 10000i \text{ vs } 1 + 1i$	[ROT]
ST.470	differences	$(e^{\{-i \theta\}})' = 2, \nabla f = 2, e^{\{i \theta\}} = \cos \theta + i \sin \theta = 1 + 0.866025i \text{ vs } 2 + 0.866025i$	[ROT]
ST.471	calc	$(\cos x - 1; 1 - \sin x)(\cos x; \sin x) = (1 \, 0; 0 \, -1) \rightarrow [\cos^2 \theta + \sin \theta + \cos \theta - 2 \sin^2 \theta] \, 1 \, 0; 0 \, -1 = -1.61603$	[ROT]
ST.472	differences	$2 \sin \theta \cos \theta = 2n \lambda \sin \theta = 0.866025 \text{ vs } 0.5$	[ROT, QUANTUM]
ST.473	calc	$\lim_{\{\theta \rightarrow 0\}} (1/\theta)(\sin \theta, \cos \theta)(\theta \, 1; 1 \, 0)(\cos \theta; \sin \theta)^T = (1 \, 0; 0 \, -1), f^{-1}(x)xf(x) = 1 = 100.02$	[ROT, QUANTUM, TRANSPORT]
ST.474	calc	$\sin \theta = \lambda/d, -py_1 \sin 90^\circ \leq \sin \theta \leq py_2 \sin 90^\circ, \lambda = h/(mv) = 0.72742$	[QUANTUM]
ST.475	symb	$\lambda_2/\lambda_1 = (\sin \theta / \sin(\theta/2)) h, \lambda' \geq 2h, \int \sin 2\theta = x - y $	[ROT, QUANTUM]
ST.476	symb	$\int \nabla \phi^2 \, d\nabla \phi = \Box \phi$	[TRANSPORT]
ST.477	symb	$\Box \phi = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm$	[MANIFOLD, TRANSPORT]
ST.478	symb	$0(x) = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\{1/2\}}$	[MANIFOLD]
ST.479	symb	$\delta \cdot 0(x) = [\nabla_i \nabla_j \int \nabla g(x) dx_{ij}]^{\{i y\}}$	[ZETA, MANIFOLD]
ST.480	symb	$ ds^2 = e^{\{-2\pi T \phi \}}[0(x) + \delta 0(x)]dx^{\sim \mu} dx^{\sim \nu} + \lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}}^{\infty} a_k f^{\sim k}$	[MANIFOLD, TRANSPORT]
ST.481	holds	$\log(x \log x) \geq 2(y \log y)^{\{1/2\}}, (y \log y)^{\{1/2\}}/\log(x \log x) \leq 1/2 = 1 \text{ vs } 1$	[MANIFOLD, TRANSPORT, ENTROPY]
ST.482	symb	$\delta(x) = \text{reality of value} / \text{exist of value} \leq 1$	[TRANSPORT]
ST.483	calc	$\text{expanding of universe} = \text{exist of value} \rightarrow \log(x \log x) = \Box \phi = 3.13662$	[TRANSPORT, ENTROPY]
ST.484	calc	$\text{freeze out of universe} = \text{reality of value} \rightarrow (y \log y)^{\{1/2\}} = \nabla \phi^2 = 1.17741$	[ROT, TRANSPORT]
ST.485	symb	$l(x) = 2x^2 + qx + r = (ax + b)(cx + d) + r, e^{\{l(x)\}} = d/df \, L(x), G_{\mu \nu} = g(x) \wedge f(x)$	[OTHER]
ST.486	symb	$V(\tau) = \int \int \exp[L(x)] \, dx_m + 0(N^{-1}), g(x) = L(x) \circ \int \int e^{\{2x^2 + 2x + r\}} \, dx_m$	[MANIFOLD]
ST.487	symb	$ ds^2 = d/df \, L(x) , \eta = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{\{1/2\}}$	[MANIFOLD]
ST.488	symb	$\tilde{h} = [\nabla_i \nabla_j \int \nabla g(x) d_{ij}]^{\{i y\}}, d/dt \, g_{ij}(t) = -2R_{ij}$	[MANIFOLD, TRANSPORT]
ST.489	symb	$(1/\tau)(N/2 + r(2\Delta F - \nabla f ^2 + R) + f) \bmod N^{-1}$	[MANIFOLD, ENTROPY]
ST.490	calc	$(x^2/a) \cos x + (y^2/b) \sin x = r^2 = 1.39211$	[ROT]
ST.491	symb	$S_m^2 = \int \pi r^2 \, dr ^2, V(\tau) = r^2 \times S_m^2, S_1^{\{mn\}} \otimes S_2^{\{mn\}} = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm$	[MANIFOLD]
ST.492	symb	$ ds^2 = e^{\{-2\pi T \phi \}}[\eta_{\mu \nu} + \tilde{h}_{\mu \nu}(x)]dx^{\sim \mu} dx^{\sim \nu} + T^2 \, d^2 \phi$	[SR, MANIFOLD, TRANSPORT]
ST.493	symb	$V(x) = \int (1/\sqrt{(2\tau q)})(\exp L(x)dx) + 0(N^{-1})$	[MANIFOLD, ENTROPY]
ST.494	symb	$V(x) = 2\int ((R + \nabla_i \nabla_j f)/(-(R + \Delta f))) e^{\{-f\}} \, dV, V(\tau) = \int \tau(p)^{\{-n/2\}} \exp(-(1/\sqrt{(2\tau q)}) L(x)dx) + 0(N^{-1})$	[MANIFOLD, ENTROPY]
ST.495	symb	$ (x \, y \, z; u \, v \, w) ^2_{[g_{\mu \nu}(x)]} = (f(x)dx^{\sim \mu} dx^{\sim \nu}, f'(y)dy^{\sim \mu} dy^{\sim \nu}, f''(z)dz^{\sim \mu} dz^{\sim \nu}) \cdot (u, v, w) = \text{diag}(1, 1, i) \equiv (g(x, y, z)/f(a, b, c)) \cdot h^{-1}(u, v, w)$	[MANIFOLD, QUANTUM]
ST.496	symb	$d/df \, \iiint \Box \phi \, d\phi \, d_{xy} = V(\Box \phi), \lim_{\{n \rightarrow \infty\}} \Sigma_{\{k=0\}}^{\infty} V_k(\Box \phi) = (\partial/\partial f) \, i h c$	[QUANTUM, TRANSPORT]
ST.497	symb	$\lim_{\{n \rightarrow \infty\}} \Sigma_{\{k=0\}}^{\infty} G_{\mu \nu} = f(x) \circ m(x)$	[MANIFOLD]
ST.498	symb	$(a_k f^{\sim k})' = nC_0 a_0 f^{\sim n} + nC_1 a_1 f^{\sim \{n-1\}} \cdots nC_{\{r-1\}} a_n f^{\sim \{n-1\}}$	[MANIFOLD]
ST.499	symb	$\int a_k f^{\sim k} \, dx_k = (a_{\{n+1\}}/(k+1)) f^{\sim \{n+1\}} + (a_{\{k+2\}}/(k+2)) f^{\sim \{k+2\}} + \cdots (a_0/k) f^{\sim k}$	[OTHER]
ST.500	symb	$\partial/\partial f \, \Box \phi = (1/4) g_{ij}^2 (\times 2)$	[MANIFOLD]
ST.501	symb	$(\nabla \phi^2/\Box \phi)' = 0 (\times 2)$	[MANIFOLD]
ST.502	symb	$(y \log y)^{\{1/2\}}/\log(x \log x) = (1/2)/((1/2)i)$	[ZETA, MANIFOLD]
ST.503	symb	$\{f, g\}/[f, g] = 1/i, (\{f, g\}/[f, g])' = i^2$	[QUANTUM]
ST.504	symb	$(i)^2 \rightarrow (1/4) g_{ij}, F_t^{\sim m} = (1/4) g_{ij}^2, f(r) = (1/4) r ^2, 4f(r) = g_{ij}^2$	[MANIFOLD, QUANTUM]
ST.505	symb	$(1/y) \cdot (1/y') \cdot (y''/y') \cdot (y'''/y'') \cdots = nC_r y^2 \cdot y^3 \cdots / (nC_r y^1 y^2 \cdots)$	[OTHER]
ST.506	symb	$(\partial y/\partial x) \cdot (\partial/\partial y) f(y) = y' \cdot f'(y)$	[OTHER]
ST.507	symb	$\int l \times l \, dm = (l \otimes l)_m$	[MANIFOLD]
ST.508	symb	$= (d/dx^{\sim \mu})(d/dx^{\sim \nu}) f^{\sim \{\mu \nu\}} \cdot \nabla \phi^2 = \Box \phi$	[MANIFOLD]
ST.509	calc	$\nabla \phi^2/\Box \phi = 1/2, l = 2\pi r, V = 4/(\pi r^3) = 12.5664$	[MANIFOLD]

4. 全方程式レジストリ (2111 本) p. 16/33

ST.510	symb	$S_{\cdot}(4\pi r^3)/(2\pi r^2)\cdot(\pi r^2) = \pi r^2, H_3 = 2, \pi(H_3) = 0$	[MANIFOLD, ENTROPY]
ST.511	symb	$f(r) = (1/2)\sqrt{(1+f'(r))/f(r)} + mg f(r)$	[ENTROPY]
ST.512	symb	$i\hbar c = G, \hbar c = G/i$	[QUANTUM]
ST.513	symb	$S_{\cdot}n^{\cdot}m = S_{2S_1} - S_{1S_2} $	[MANIFOLD]
ST.514	symb	$\square\phi = V\cdot S_{\cdot}n^{\cdot}m$	[MANIFOLD]
ST.515	symb	$G^{\cdot}\{\mu\nu\} \equiv R^{\cdot}\{\mu\nu\}$	[SR, MANIFOLD]
ST.516	symb	$\nabla_{\cdot i}\nabla_{\cdot j}(\square\phi)d\phi_{\cdot xy} = (\partial/\partial f)\square\phi\cdot T^{\cdot}\{\mu\nu\}$	[MANIFOLD]
ST.517	symb	$= (8\pi G/c^4)(T^{\cdot}\{\mu\nu\})^2, (\chi\otimes\pi) = \iiint\square\phi\ d^3\phi = \text{div}(\text{rot } E, E_1)\cdot e^{\cdot}\{-ix\log x\}$	[ROT, SR]
ST.518	symb	$\sigma(H_n\otimes K_m) = E_n\times H_m$	[MANIFOLD]
ST.519	symb	$H_n = [\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla g(x)dx_{\cdot i}dx_{\cdot j}]^{\cdot}\{iy\}, K_m = [\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f(x)d\eta]^{\cdot}\{1/2\}$	[ZETA, MANIFOLD]
ST.520	symb	$ ds^2 = \sigma(H_n\times K_m), \sigma(\chi,x)\times\pi(\chi,x) = (\partial/\partial f)L(x), V'_{\cdot}\tau(x) = (\partial/\partial V)L(x)$	[MANIFOLD]
ST.521	symb	$(\square\phi)' = (\partial/\partial f_M)(\iiint f(x,y,z)dx\ dy\ dz)'d\phi$	[MANIFOLD]
ST.522	symb	$(\partial/\partial V)L(x) = V(\tau)\int\int e^{\cdot}\{\int x\log x\ dx + 0(N^{-1})\}d\phi$	[MANIFOLD]
ST.523	symb	$(d/df)\sum_{\{k=1\}}^{\cdot}n\sum_{\{k=0\}}^{\cdot}\infty a_{\cdot k}f^{\cdot}k = (d/df)m(x), V(x)/f(x) = m(x)$	[MANIFOLD]
ST.524	symb	$4V'_{\cdot}\tau(x) = g_{\cdot ij}^2, (d/dl)L(x) = \sigma(\chi,x)\times V_{\cdot}\tau(x)$	[MANIFOLD]
ST.525	symb	$ ds^2 = e^{\cdot}\{-2\pi T \phi \}[\eta_{\cdot\mu\nu} + \hbar_{\cdot\mu\nu}(x)]dx^{\cdot\mu}dx^{\cdot\nu} + T^2d\phi^2$	[MANIFOLD, TRANSPORT]
ST.526	symb	$f^{\cdot}\{(2)\}(x) = [\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f^{\cdot}\{(5)\}d\eta]^{\cdot}\{1/2\} = [f^{\cdot}\{(2)\}(x)d\eta]^{\cdot}\{1/2\}$	[MANIFOLD, TRANSPORT]
ST.527	symb	$\nabla_{\cdot i}\nabla_{\cdot j}\int F(x)d\eta = (\partial/\partial f)F$	[MANIFOLD]
ST.528	symb	$\nabla f = (d/dx) f$	[OTHER]
ST.529	symb	$\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f\ d\eta = (\partial/\partial x_{\cdot i})(\partial/\partial x_{\cdot j})((d/dx) f)$	[OTHER]
ST.530	symb	$(z_{3z_2} - z_{2z_3})/(z_{2z_1} - z_{1z_2}) = \omega$	[OTHER]
ST.531	symb	$(\bar{z}_{3z_2} - \bar{z}_{2z_3})/(\bar{z}_{2z_1} - \bar{z}_{1z_2}) = \bar{\omega}$	[OTHER]
ST.532	symb	$\omega\cdot\bar{\omega} = 0, z_{\cdot n} = \omega - \{x\}, z_{\cdot n}\cdot\bar{z}_{\cdot n} = 0, \bar{Z}^{\cdot}n\cdot\bar{Z}_{\cdot}n = 0$	[QUANTUM]
ST.533	symb	$[f, g]\times [g, f] = fg + gf = \{f, g\}$	[QUANTUM]
ST.534	symb	$V(\tau) = \int\int e^{\cdot}\{\int x\log x\ dx + 0(N^{-1})\}d\phi, V'_{\cdot}\tau(x) = (\partial/\partial f_M)(\iiint f(x,y,z)dx\ dy\ dz)'d\phi$	[MANIFOLD]
ST.535	symb	$(\square\phi)' = 4\bar{V}(x), (\partial/\partial V)L(x) = m(x), V(\tau) = \int(1/\sqrt{(2\tau q)})\exp[L(x)]d\phi + 0(N^{-1})$	[MANIFOLD]
ST.536	symb	$V(\tau) = \iiint V/S^2\ dm, f(r) = (1/2)\sqrt{(1+f'(r))/f(r)} + mg f(r), \log(x\log x) \geq 2(y\log y)^{1/2}, F_{\cdot}t^{\cdot}m = (1/4)g_{\cdot ij}^2, d/dt$ $g_{\cdot ij}(t) = -2R_{\cdot ij}$	[MANIFOLD]
ST.537	calc	$\nabla_{\cdot i}\nabla_{\cdot j}v = (1/2)mv^2 + mc^2, \int\nabla_{\cdot i}\nabla_{\cdot j}v\ dv = (\partial/\partial f)L(x) = 8.98755e+16$	[SR]
ST.538	symb	$(\square\phi)^2 = -2\int\nabla_{\cdot i}\nabla_{\cdot j}v\ d^2v, (\square\phi)^2 = (\nabla\phi^2/\square\phi)'$	[MANIFOLD]
ST.539	symb	$= d/df\int\int 1/(x\log x)^2\ dm, \oplus\nabla^+_3 = \int\bigvee(R + \nabla_{\cdot i}\nabla_{\cdot j}f)^2/\exists(R + \Delta f)\ dV = \langle x,y,z\rangle\cdot\langle u,v,w\rangle/\Gamma$	[GAMMA, MANIFOLD]
ST.540	symb	$\oplus\pm = \int\exp[\int\nabla_{\cdot i}\nabla_{\cdot j}f\ d\eta]d\phi = L(x)\cdot(\partial/\partial l)F(x) = (\square\phi)^2$	[MANIFOLD, QUANTUM]
ST.541	calc	$l = \sqrt{(hG/c^3)}, T^{\cdot}\{\mu\nu\} = (1/2)kl^2 + (1/2)mv^2, E = mc^2 - (1/2)mv^2 = 4.05126e-35$	[SR, QUANTUM, TRANSPORT]
ST.542	holds	$e^{\cdot}\{x\log x\} = x^{\cdot}x, x = \log x^{\cdot}x / \log x, y = x, x = e = 9.88212\ \text{vs}\ 9.88212$	[ZETA]
ST.543	symb	$\int 1/(x\log x)\ dx = i\int x\log x\ dx + \int 1/(x\log x)\ dx$	[ZETA, QUANTUM]
ST.544	symb	$\int\int 1/(x\log x)^2\ dx_{\cdot m} = i(1/2)x^2$	[ZETA, MANIFOLD]
ST.545	symb	$\int\int 1/(x\log x)^2\ dx_{\cdot m} = i\int\int_{\cdot M}dx_{\cdot m} \leq (1/2)i + x^2$	[ZETA, MANIFOLD]
ST.546	calc	$E = -(1/2)mv^2 + mc^2 = 8.98755e+16$	[SR, TRANSPORT]
ST.547	symb	$\lim_{\{x\rightarrow\infty\}}\int\int 1/(x\log x)^2\ dx_{\cdot m} \geq (1/2)i$	[ZETA, MANIFOLD]
ST.548	symb	$d/df\int\int 1/(x\log x)^2\ dx_{\cdot m} = (1/2)i$	[ZETA, MANIFOLD]
ST.549	holds	$\lim_{\{x\rightarrow\infty\}}x^2/e^{\cdot}\{x\log x\} = 0 = 3.8147e-24\ \text{vs}\ 0$	[OTHER]
ST.550	symb	$\int dx \rightarrow \partial f \rightarrow dx \rightarrow \text{cons}$	[ENTROPY]
ST.551	symb	$(\square\phi)' = (\exists\int\bigvee(R + \nabla_{\cdot i}\nabla_{\cdot j}f)e^{\cdot}\{-f\}dV)'d\phi$	[MANIFOLD]
ST.552	symb	$\oplus M^+_3 = (\partial/\partial f)L(x)$	[MANIFOLD]
ST.553	symb	$(\partial/\partial l)L(x) = \nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f(x)d\eta, L(x) = V(x)/f(x)$	[MANIFOLD]
ST.554	symb	$l(x) = L'(x), d/df F = m(x), V'(\tau) = \int\int e^{\cdot}\{\int x\log x\ dx + 0(N^{-1})\}d\phi$	[MANIFOLD]
ST.555	symb	$T^{\cdot}\{\mu\nu\} = \iiint V(x)/S^2\ dm, S^2 = \pi r^2\cdot S(x) = 4\pi r^2/\tau(x)$	[SR, MANIFOLD]
ST.556	symb	$\eta = \nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f(x)d\eta, \hbar = \nabla_{\cdot i}\nabla_{\cdot j}\int\nabla g(x)dx_{\cdot i}dx_{\cdot j}$	[MANIFOLD]
ST.557	symb	$\delta 0(x) = [\nabla_{\cdot i}\nabla_{\cdot j}\int f(x)d\eta]^{\cdot}\{1/2 + iy\}$	[ZETA, MANIFOLD]
ST.558	symb	$Z(x,h) = \lim_{\{x\rightarrow\infty\}}\sum_{\{k=0\}}^{\cdot}\infty qT^{\cdot}m/m = \delta(x)$	[ZETA]
ST.559	symb	$l(x) = 2x^2 + px + q, m(x) = \lim_{\{x\rightarrow\infty\}}\sum_{\{k=0\}}^{\cdot}\infty (qx^{\cdot}m)' / f(x)$	[ZETA, QUANTUM]
ST.560	calc	$Z(T,x) = \exp\sum_{\{m=1\}}^{\cdot}\infty q^{\cdot}kT^{\cdot}m/m, Z(x,h) = \exp((q f(x))^{\cdot}m/m) = 1.5625$	[ZETA]
ST.561	symb	$d/df F = m(x), F = \int\int e^{\cdot}\{\int x\log x\ dx + 0(N^{-1})\}d\phi$	[ZETA, MANIFOLD]
ST.562	symb	$\lim_{\{x\rightarrow 1\}}\text{mesh } m/(m+1) = 0, \int x^{\cdot}m = x^{\cdot}m/(m+1)$	[OTHER]
ST.563	holds	$d/df\int x^{\cdot}m = mx^{\cdot}m, d/dt g_{\cdot ij}(t) = -2R_{\cdot ij}, \lim_{\{x\rightarrow 1\}}\text{mesh}(x) = \lim_{\{m\rightarrow\infty\}}m/(m+1) = 0.9999\ \text{vs}\ 1$	[MANIFOLD]
ST.564	symb	$\lim_{\{x\rightarrow 1\}}\sum_{\{k=0\}}^{\cdot}\infty a_{\cdot k}f^{\cdot}k = \alpha$	[OTHER]
ST.565	symb	$(\partial/\partial V) ds^2 = T^{\cdot}\{\mu\nu\}, V(\tau) = \int e^{\cdot}\{x\log x\}d\phi = l(x)$	[SR, MANIFOLD]
ST.566	symb	$R_{\cdot ij} = (\partial/\partial x_{\cdot i})(\partial/\partial x_{\cdot j})T^{\cdot}\{\mu\nu\}, G^{\cdot}\{\mu\nu\} = R^{\cdot}\{\mu\nu\}T^{\cdot}\{\mu\nu\}$	[SR, MANIFOLD]
ST.567	symb	$F(x) = \int\int e^{\cdot}\{\int x\log x\ dx + 0(N^{-1})\}d\phi, (d/dV)F(x) = V'(x)$	[MANIFOLD]
ST.568	symb	$T^{\cdot}\{\mu\nu\} = R^{\cdot}\{\mu\nu\}, T^{\cdot}\{\mu\nu\} = \iiint V/S^2\ dm$	[SR, MANIFOLD]
ST.569	symb	$\delta 0(x) = [D^2\phi\otimes h_{\cdot\mu\nu}]\ dm$	[MANIFOLD]
ST.570	symb	$\nabla(\square\phi)' = [\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f(x)d\eta]^{\cdot}\{1/2 + iy\}$	[ZETA, MANIFOLD]
ST.571	symb	$(f(x), g(x))' = (A^{\cdot}\{\mu\nu\})'$	[OTHER]
ST.572	symb	$(dx\ \delta(x); \varepsilon(x)\ \partial x)\cdot(f(x,y), g(x,y)) = (1\ 0; 0\ -1)$	[ROT, QUANTUM]
ST.573	calc	$\delta(x)\cdot 0(x) = (1\ 0; 0\ -1)^{\cdot}\{1/2\} = 6.12323e-17 + 1i$	[ROT, QUANTUM]
ST.574	symb	$V'_{\cdot}\tau(x) = (\partial/\partial f_M)(\iiint f(x,y,z)dx\ dy\ dz)d\phi$	[MANIFOLD]
ST.575	symb	$(\partial/\partial V)L(x) = V'_{\cdot}\tau(x)$	[MANIFOLD]
ST.576	symb	$\int\int 1/(x\log x)^2\ dx_{\cdot m} = [\nabla_{\cdot i}\nabla_{\cdot j}\int\nabla f(x)d\eta]\times U(r)$	[MANIFOLD, ENTROPY]

4. 全方程式レジストリ (2111 本) p. 17/33

ST. 577	symb	$S_1^{\wedge \{mn\}} \otimes S_2^{\wedge \{mn\}} = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm$	[ZETA, MANIFOLD]
ST. 578	symb	$U(r) = (1/2)\sqrt{(1 + f'(r))/f(r)} + mgr$	[OTHER]
ST. 579	calc	$F_{\mathrm{t}^m} = (1/4) r ^2 = 1$	[MANIFOLD]
ST. 580	symb	$\int \int 1/(y \log y)^{\{1/2\}} \, dy_m = [\nabla_i \nabla_j \int \nabla f(x) d\eta] \times E \pm$	[ZETA, MANIFOLD]
ST. 581	symb	$E \pm = \exp[L(x)] \, dm \, d\phi + 0(N^-)$	[QUANTUM]
ST. 582	calc	$d/df \, F = [\nabla_i \nabla_j \int \nabla f(x) d\eta](U(r) + E \pm) = (1/2)mv^2 + mc^2 = 1.79751e+17$	[SR, MANIFOLD]
ST. 583	symb	$x^{\{1/2 + iy\}} = \exp[\int \nabla_i \nabla_j f(g(x)) \, g'(x) \, \partial f \partial g]$	[ZETA]
ST. 584	symb	$\nabla_i \nabla_j \int \nabla f(x) d\eta$	[MANIFOLD]
ST. 585	symb	$\nabla_i \nabla_j \otimes M_3 = M_3/P^{\{2n\}} \cong P^{\{2n\}}/M_3 \cong M_1 = [M_1]$	[MANIFOLD]
ST. 586	symb	$[f(x)] = \Sigma_{\{k=0\}^{\infty}} a_k f^k, \lim_{\{n \rightarrow 1\}} [f(x)] = \lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = \alpha, e^{\{i \theta\}} = \cos \theta + i \sin \theta, H_3(M_1) = 0$	[ROT, ZETA, MANIFOLD]
ST. 587	differs	$(x^2 \cos \theta)/a + (y^2 \sin \theta)/b = r^2, (\cos \theta - \sin \theta; \sin \theta \cos \theta)(x; y) = (1 \, 0; 0 \, -1) = 0.366025 + 1.36603i \, \text{vs} \, 1 - 1i$	[ROT]
ST. 588	symb	$\chi(x) = \Sigma_{\{k=0\}^{\infty}} (-1)^n r^n, d/df \, F = d/df \, \Sigma \Sigma a_k f^k$	[MANIFOLD]
ST. 589	holds	$= a_{1a_2 \cdots a_n} - a_{1 \cdots a_{n-1}} - a_n \cdots a_1 , \lim_{\{n \rightarrow 0\}} \chi(x) = 2 = 2 \, \text{vs} \, 2$	[MANIFOLD]
ST. 590	symb	$\lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = nC_r f(x)^n f(y)^{\{n-r\}} \delta(x,y)$	[MANIFOLD]
ST. 591	symb	$\lim_{\{n \rightarrow \infty\}} nC_r f(x)^n f(y)^{\{n-r\}} \delta(x,y)$	[OTHER]
ST. 592	calc	$\lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} (1/(n+1))^s = \lim_{\{n \rightarrow 1\}} Z^r = 1/z = 1.64493$	[ZETA]
ST. 593	differs	$\beta(p,q) = \Gamma(p)\Gamma(q)/\Gamma(p+q) \cong \Gamma(p+q)/(\Gamma(p)\Gamma(q)), \lim_{\{n \rightarrow 1\}} a_k f^k \cong \lim_{\{n \rightarrow \infty\}} \zeta(s)/(a^k f^k) = 0.19635 \, \text{vs} \, 5.09296$	[GAMMA, ZETA, BETA]
ST. 594	symb	$\lim_{\{n \rightarrow 1\}} \zeta(s) = 0, 0(x) = \zeta(s)$	[ZETA]
ST. 595	symb	$\Sigma_{\{x=0\}^{\infty}} f(x) \rightarrow \otimes_{\{k=0\}^{\infty}} \nabla f(x) = \int_M \delta(x) f(x) dx$	[MANIFOLD]
ST. 596	symb	$\iiint_M (V/S^2) e^{\{-f\}} \, dV = \int \int_D -(f(x,y)^2, g(x,y)^2) - \int \int_D (g(x,y)^2, f(x,y)^2)$	[MANIFOLD]
ST. 597	symb	$\lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm = \int \exp[L(x)] \, d\phi \, dm \times E \pm = S_1^{\wedge \{mn\}} \otimes S_1^{\wedge \{mn\}}$	[MANIFOLD]
ST. 598	symb	$= Z_1 \otimes Z_1 = M_1$	[MANIFOLD]
ST. 599	symb	$H_n^m(\chi, h) = \int \int_M (V/(R + \Delta f)) e^{\{-f\}} \, dV, V/S^2 = \iiint_M [D^2 \phi \otimes h_{\mu \nu}] \, dm$	[MANIFOLD]
ST. 600	symb	$\iiint_M (V/S^2) \, dm = \int_D (l \times l) \, dm$	[MANIFOLD]
ST. 601	symb	$\int \int_D -g(x,y)^2 \, dm - \int \int_D -f(x,y)^2 \, dm = -2R_{ij}, [f(x), g(x)] \times [h(x), g^{-1}(x)]$	[MANIFOLD]
ST. 602	symb	$ D^m \, dx; dx \, \partial^m \cos \theta - \sin \theta; \sin \theta \cos \theta x; y = 1 \, 0; 0 \, -1 ^{\{1/2\}} (\times 3)$	[ROT, QUANTUM]
ST. 603	symb	$(D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) (\times 2)$	[ROT, MANIFOLD]
ST. 604	symb	$(1 \, 0; 0 \, i) \beta(x, \theta) (x; y) = (i \, 0; 0 \, -1) (\times 2)$	[ROT, BETA, QUANTUM]
ST. 605	calc	$l = \sqrt{(hG/c^3)}, \sigma^m \cdot (\delta(x) - 1; 1 \, \varepsilon(x))^{\{1/2\}} = (i \, 0; 0 \, -i) (\times 2) = 4.05126e-35$	[ROT, QUANTUM]
ST. 606	symb	$((\partial/\partial \varpi) f(x,y,z))^{\{3'\}} = A^{\{\mu \nu\}} (\times 2)$	[OTHER]
ST. 607	symb	$= (D^m, dx) \cdot (\cos \theta, \sin \theta) \times (dx, \partial^m) \cdot (\cos \theta, \sin \theta) (\times 2)$	[ROT, MANIFOLD]
ST. 608	symb	$\lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = \lim_{\{n \rightarrow 1\}} a_n/a_{n-1} \cong \alpha$	[MANIFOLD]
ST. 609	symb	$\lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k/a_{k+1} = \Sigma_{\{k=0\}^{\infty}} a_k f^k$	[OTHER]
ST. 610	symb	$\square = (8\pi G/c^4) T^{\{\mu \nu\}}$	[SR]
ST. 611	symb	$\int x \log x \, dx = \int \int_M (e^{\{x \log x\}} \sin \theta \, d\theta) \, dx \, d\theta = (\log \sin \theta \, d\theta)^{\{1/2\}} - (\delta(x) \cdot \varepsilon(x))^{\{1/2\}}$	[ROT, TRANSPORT]
ST. 612	symb	$T^{\{\mu \nu\}} = \int \int_M [\nabla_i \nabla_j e^{\{\int x \log x \, dx + 0(N^{-1})\}}] \, dm \, d\phi $	[SR, MANIFOLD]
ST. 613	symb	$(dx \, \delta(x); \varepsilon(x) \, \partial^m(x)) (f_{mn}(x), g_{\mu \nu}(x)) = (1 \, 0; 0 \, -1)^{\{1/2\}}$	[ROT, QUANTUM]
ST. 614	symb	$ ds^2 = H(x)_{mn} \otimes K(y)_{mn}$	[MANIFOLD]
ST. 615	symb	$ H(x) ^{\{-1/2 + iy\}} \subset R \pm \cup C \pm \cong M_3$	[ZETA, MANIFOLD]
ST. 616	symb	$ H(x) ^{\{1/2 + iy\}} \subseteq M_3$	[ZETA, MANIFOLD]
ST. 617	symb	$K(y) = (x \, y \, z; a \, b \, c) ^2_{\{g_{\mu \nu}(x)\}} \cong (f(x,y,z)/g(a,b,c)) h^{-1}(u,v,w)$	[MANIFOLD]
ST. 618	symb	$V = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm, S^2 = \int e^{\{-2 \sin \theta \cos \theta\}} \cdot \log \sin \theta \, dx \, d\theta + 0(N^{-1}), 0(x) = \zeta(s)/\lim_{\{x \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = \alpha$	[ROT, ZETA, MANIFOLD]
ST. 619	symb	$0(x) = T^{\{\mu \nu\}}, \lim_{\{x \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f^k = T^{\{\mu \nu\}}$	[SR, MANIFOLD]
ST. 620	symb	$G_{\mu \nu} = R_{\mu \nu} T^{\{\mu \nu\}}, M_3 = \iiint (V/S^2) \, dm, -(1/(2(T-t))) R_{ij} = \square \phi$	[SR, MANIFOLD]
ST. 621	symb	$m(x) = [f(x)]$	[MANIFOLD]
ST. 622	symb	$f(x) = \int \int e^{\{x \log x \, dx + 0(N^{-1})\}} + T^2 \, d^2 \phi$	[MANIFOLD, TRANSPORT]
ST. 623	symb	$F(x) = \nabla_i \nabla_j \int \nabla f(x,y) \, dm ^{\{1/2 + iy\}}$	[ZETA, MANIFOLD]
ST. 624	symb	$G_{\mu \nu} = [\nabla_i \nabla_j \int \nabla g(x,y,z) dx \, dy \, dz] ^{\{(m/2) \sin \theta \cos \theta\}} \log \sin \theta \, d\theta \, d\phi$	[ROT, SR]
ST. 625	symb	$G_{\mu \nu} = R_{\mu \nu} T^{\{\mu \nu\}}$	[SR, MANIFOLD]
ST. 626	symb	$T^{\{\mu \nu\}} = F(x), 2(T-t) g_{ij}^2 = \int \int 1/(x \log x)^2 \, dx_m$	[SR, ZETA, MANIFOLD]
ST. 627	symb	$\phi \delta(x) = [m(x)], \nabla(\square \phi) = \nabla_i \nabla_j \int \nabla g(x,y) d\eta$	[QUANTUM]
ST. 628	symb	$\nabla \cdot (\square \phi) = (1/4)g_{ij}^2, \square \phi = (8\pi G/c^4) T^{\{\mu \nu\}}$	[SR, MANIFOLD]
ST. 629	symb	$pV/c^3 = S \circ h_{\mu \nu} = h$	[QUANTUM]
ST. 630	calc	$T^{\{\mu \nu\}} = \hbar \nu/S, T^{\{\mu \nu\}} = \iiint (V/S^2) \, dm, \langle d/df \rangle m(x) = V(x)/F(x) = 5.27286e-16$	[SR, QUANTUM]
ST. 631	symb	$y = x, d/df \, F = m(x), R_{ij} _{\{g_{\mu \nu}(x)\}} = [\nabla_i \nabla_j g(x,y)]^{\{1/2 + iy\}}$	[ZETA, MANIFOLD, QUANTUM]
ST. 632	symb	$\nabla \cdot (\square \phi) = (\partial/\partial f) F = \int \int \nabla_i \nabla_j f(x) \, d\eta_{\mu \nu}$	[MANIFOLD]
ST. 633	symb	$\int [\nabla_i \nabla_j g(x,y)] \, dm = (\partial/\partial f) R_{ij} _{\{g_{\mu \nu}(x)\}}$	[ZETA, MANIFOLD]
ST. 634	symb	$G(x) = \nabla_i \nabla_j f + R_{ij} _{\{g_{\mu \nu}(x)\}} + \nabla(\square \phi) + (\square \phi)^2$	[SR, MANIFOLD]
ST. 635	symb	$G_{\mu \nu} + \Lambda g_{ij} = T^{\{\mu \nu\}}, T^{\{\mu \nu\}} = (d/dx_{\mu})(d/dx_{\nu}) f_{\mu \nu} + -2(T-t) R_{ij} + f'' + (f')^2 = \int \exp[L(x)] \, dm + 0(N^{-1}) e^{\{(2/m) \sin \theta \cos \theta\}} \cdot \log(\sin \theta) \, dx + 0(N^{-1})$	[ROT, SR, MANIFOLD]
ST. 636	symb	$(\partial/\partial f) F = (\nabla_i \nabla_j)^{-1} \circ F(x)$	[MANIFOLD]
ST. 637	symb	$0(x) = \int [\nabla_i \nabla_j \int \nabla f(x) dm] \, d\phi = \int [\nabla_i \nabla_j f(x) d\eta_{\mu \nu}] \, d\phi$	[MANIFOLD]
ST. 638	symb	$\nabla f = \int \nabla_i \nabla_j [\int (S^{-3}/\delta(x)) \, dV] \, dm$	[MANIFOLD]
ST. 639	symb	$ \int [\nabla_i \nabla_j f] \, dm ^{\{1/2 + iy\}} = \text{rot}(\text{div} \, E, E_1)$	[ROT, ZETA]
ST. 640	symb	$= 2\langle f, h \rangle, V(x)/f(x) = \rho(x)$	[OTHER]
ST. 641	symb	$\int_M \rho(x) dx = \square \phi, -2\langle g, h \rangle = \text{div}(\text{rot} \, E, E_1) = -2R_{ij}$	[ROT, MANIFOLD]
ST. 642	symb	$0(x) = (\nabla_i \nabla_j/S^2) \int \nabla_i \nabla_j f \cdot g(x) dx \, dy] \, d\phi = \int (\delta(x))^{\{2 \sin \theta \cos \theta\}} \log \sin \theta \, d\theta \, d\phi$	[ROT, QUANTUM]
ST. 643	symb	$\delta \cdot 0(x) = [\nabla_i \nabla_j f \, d\eta \int e^{\{2 \sin \theta \cos \theta\}} \cdot \log \sin \theta \, d\theta]$	[ROT, MANIFOLD]

ST.644	symb	$i\hbar\phi= \int(\partial/\partial z)[i(xy+\bar{y}x)/(z-\bar{z})]dm\,d\phi $	[QUANTUM]
ST.645	symb	$\delta(x)\int_{\mathbb{C}R^2}\sin\theta\cos\theta\}=\Gamma(p+q)$	[ROT, GAMMA]
ST.646	symb	$(\Box+m)\cdot\phi=(\nabla_{-i}\nabla_{-j}f _{\{g_{-}\mu\nu(x)\}}+v\nabla_{-i}\nabla_{-j})$	[QUANTUM]
ST.647	symb	$\int[m(x)(\operatorname{rot}\cdot\operatorname{div}(E,\,E_1))]dm\,d\phi$	[ROT]
ST.648	symb	$G_{-}\mu\nu=\Box\iiint(x,y,z)^3\,dx\,dy\,dz=(8\pi G/c^4)T^{\sim}\{\mu\nu\}$	[SR]
ST.649	symb	$(d/dV)F=\delta(x)\int\nabla_{-i}\nabla_{-j}f\,d\eta_{-}\mu\nu$	[MANIFOLD]
ST.650	symb	$x^{\sim}n+y^{\sim}n=z^{\sim}n,\,\delta(x)\int z^{\sim}n=(d/dV)z^3,\,(x,y)\cdot(\delta^{\sim}m,\,\partial^{\sim}m)=(x,y)\cdot(z^{\sim}n,\,f),\,n\bot x,\,n\bot y=0$	[OTHER]
ST.651	symb	$\vee(\nabla_{-i}\nabla_{-j}f)\cdot X0R(\Box\phi)=(d/df)\int_{-}M\,F\,dV$	[MANIFOLD, QUANTUM]
ST.652	calc	$\partial(x)\int z^3=(d/dV)z^2,\,\Sigma_{\{k=0\}^{\infty}}1/(n+1)^{\sim}s=0(x)\qquad=1.20206$	[ZETA]
ST.653	symb	$\int 0(x)dx=\delta(x)\pi(x)f(x)$	[ZETA]
ST.654	symb	$0(x)=\int\sigma(x)^{\sim}\{2\sin\theta\cos\theta\}\log\sin\theta\,d\theta\,d\phi$	[ROT]
ST.655	symb	$0(x)= (\nabla_{-i}\nabla_{-j}f/S^{\sim})\int[\nabla_{-i}\nabla_{-j}f\circ g(x)dx\,dy] $	[MANIFOLD]
ST.656	holds	$\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_k/a_{-}\{k+1\}=(\log\sin\theta\,dx)^{\sim}=\cos\theta/\sin\theta=x/y\qquad=1.37638\,\text{vs}\,1.37638$	[ROT]
ST.657	holds	$\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k=1/(1-z)\qquad=1.42857\,\text{vs}\,1.42857$	[OTHER]
ST.658	symb	$\Box\phi=(8\pi G/c^4)T^{\sim}\{\mu\nu\}$	[SR]
ST.659	calc	$(\partial/\partial f)\Box\phi=4\pi G\rho\qquad=4.61274e-06$	[SR]
ST.660	symb	$\int\rho(x)=\Box\phi,\,V(x)/f(x)=\rho(x)$	[OTHER]
ST.661	symb	$(\partial^{\sim}n/\partial f^{\sim}\{n-1\})F=\int[D^2\phi\otimes h_{-}\mu\nu]dm=P_{-}P_{-}3\cdots P_{-}\{2n-1\}/(P_{-}0P_{-}2\cdots P_{-}\{2n+2\})=\otimes\nabla M_{-}1$	[ZETA, MANIFOLD]
ST.662	symb	$\otimes\nabla M_{-}1=\sigma_{-}n(\chi,x)\otimes\sigma_{-}\{n-1\}(\chi,x)=\{f,h\}\circ[f,h]^{-1}$	[MANIFOLD, QUANTUM]
ST.663	symb	$=g^{-1}(x)_{-}\mu\nu\,dx\,g_{-}\mu\nu(x),\,\Sigma_{\{k=0\}^{\infty}}\nabla^{\sim}n_{\mathbb{C}_-}rf^{\sim}n(x)\equiv\Sigma_{\{k=0\}^{\infty}}\nabla^{\sim}n\nabla^{\sim}\{n-1\}_{\mathbb{C}_-}rf^{\sim}n(x)\,g^{\sim}\{n-r\}(x)$	[MANIFOLD]
ST.664	symb	$\Sigma_{\{k=0\}^{\infty}}(\partial^{\sim}n/\partial^{\sim}\{n-1\}f)\circ\zeta(x)/n!=\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k$	[ZETA]
ST.665	symb	$(f)^{\sim}n=n_{\mathbb{C}_-}rf^{\sim}n(x)\,g^{\sim}\{n-r\}(x)\cdot\delta(x,y)$	[OTHER]
ST.666	symb	$(e^{\sim}\{i\theta\})^{\sim}=ie^{\sim}\{i\theta\},\,\int e^{\sim}\{i\theta\}=(1/i)e^{\sim}\{i\theta\},\,ihc=G,\,hc=(1/i)G$	[ROT, QUANTUM]
ST.667	calc	$(\Box\phi,\,\nabla^2)^{\sim}\cdot(8\pi G/c^4)T^{\sim}\{\mu\nu\},\,4\pi G\rho=(-(1/2)mv^2+mc^2,\,(1/2)kT^2+(1/2)mv^2)\cdot(\cos\theta\,-\sin\theta;\sin\theta\,\cos\theta)=(1\,0;0\,i)$ $=6.35516e+16-6.35516e+16i$	[ROT, SR, TRANSPORT]
ST.668	symb	$(\{f,g\}/[f,g])^{\sim}=i^2,\,\nabla^2/\Box\phi=1/2$	[QUANTUM]
ST.669	symb	$\int\int\{f,g\}/[f,g]=(1/2)i,\,\int\int1/(y\log y)^{\sim}\{1/2\}\,dy_{-}m=1/2$	[ZETA, QUANTUM]
ST.670	symb	$\int\int1/(x\log x)^2\,dx_{-}m=(1/2)i,\,d/dt\,g_{-}ij=-2R_{-}ij$	[ZETA, MANIFOLD]
ST.671	symb	$(\partial/\partial x)(f(x)g(x))^{\sim}=\otimes\nabla_{-i}\nabla_{-j}f(x)g(x)$	[OTHER]
ST.672	symb	$\int f^{\sim}(x)g(x)dx=[f(x)g(x)]-\int f(x)g^{\sim}(x)dx$	[OTHER]
ST.673	symb	$A-G,\,T-C$	[OTHER]
ST.674	symb	$d/dL\,V(\tau)=d/df\int f_{-}M\,1/(x\log x)^2\,dx_{-}m+d/df\int f_{-}M\,1/(y\log y)^{\sim}\{1/2\}\,dy_{-}m(\times2)$	[ZETA, MANIFOLD, TRANSPORT]
ST.675	symb	$ ds^2 =\lim_{\{x\rightarrow\infty\}}[\delta(x)\iiint\pi(\Sigma_{\{k=0\}^{\infty}}(\eta\sqrt{p},\,x)/n)^{\sim}\{1/2\}\,d\tau]^{\sim}\{\mu\nu\}\,(\times2)$	[ZETA, MANIFOLD]
ST.676	symb	$V(\tau)=[f(x),\,g(x)]\times[f^{-1}(x),\,h(x)]$	[MANIFOLD]
ST.677	symb	$\Gamma(p,q)=\int e^{\sim}\{-x\}\,x^{\sim}\{1-t\}\,dx=\beta(p,q)=\pi(f(\chi,x),\,x)$	[GAMMA, BETA, MANIFOLD]
ST.678	symb	$ ds^2 =0(x)[(f(x)\circ g(x))^{\sim}\{\mu\nu\}]dx^{\sim}\mu\,dx^{\sim}\nu=\lim_{\{x\rightarrow\infty\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k$	[MANIFOLD]
ST.679	symb	$G^{\sim}\{\mu\nu\}=(\partial/\partial f)\int[f(x)^{\sim}\{\mu\nu\}\circ G(x)^{\sim}\{\mu\nu\}]dx^{\sim}\mu\,dx^{\sim}\nu]^{\sim}\{\mu\nu\}dm=g_{-}\mu\nu(x)dx^{\sim}\mu\,dx^{\sim}\nu-f(x)^{\sim}\{\mu\nu\}dx^{\sim}\mu\,dx^{\sim}\nu$	[SR, MANIFOLD]
ST.680	symb	$[i\pi(\chi,x),\,f(x)]=i\pi f(x)-f(x)\pi(\chi,x)$	[QUANTUM]
ST.681	symb	$T^{\sim}\{\mu\nu\}=(\lim_{\{x\rightarrow\infty\}}\Sigma_{\{k=0\}^{\infty}}\int\int[V(\tau)\circ S^{\sim}\{\mu\nu\}(\chi,x)]dm)^{\sim}\{\mu\nu\}\,dx^{\sim}\mu\,dx^{\sim}\nu$	[SR, MANIFOLD]
ST.682	symb	$G^{\sim}\{\mu\nu\}=R^{\sim}\{\mu\nu\}T^{\sim}\{\mu\nu\}$	[SR, MANIFOLD]
ST.683	symb	$\sigma^{\sim}m[\delta(x)^{-1};\,1\,\varepsilon(x)]^{\sim}\{1/2\}=(i\,0;0\,-i)$	[ROT, QUANTUM]
ST.684	symb	$V(M)=(\partial/\partial f)(^{\sim}N\int[f\setminus M]^{\sim}\{\otimes N\})^{\sim}\{\mu\nu\}\,dx^{\sim}\mu\,dx^{\sim}\nu$	[MANIFOLD]
ST.685	symb	$V(M)=\pi(2\int\sin^2dx)\otimes(d/df)F^{\sim}M\,dx_{-}m$	[ROT, MANIFOLD]
ST.686	symb	$\lim_{\{x\rightarrow\infty\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k=\int(F(V)dx_{-}m)^{\sim}\{\mu\nu\}\,dx^{\sim}\mu\,dx^{\sim}\nu$	[MANIFOLD]
ST.687	symb	$\otimes_{\{k=0\}^{\infty}}[f\setminus g]=\vee(M\wedge N)$	[MANIFOLD]
ST.688	symb	$\pi_{-}1(M)=e^{\sim}\{-f\,2\int\sin^2x\,dm\}+0(N^{-1})=[i\pi(\chi,x),\,f(x)]$	[ROT, MANIFOLD]
ST.689	symb	$M\circ f(x)=e^{\sim}\{-f\int\sin x\cos x\,dx_{-}m\}+\log(0(N^{-1}))$	[ROT, MANIFOLD]
ST.690	symb	$\varepsilon S(\nu)=\Box_{-}v\cdot(\partial/\partial\chi)^{(5\sqrt{\cdot}/(\wedge g^2))}\,d\chi$	[TRANSPORT, ENTROPY]
ST.691	symb	$\wedge(F_t^{\sim}m)^{\sim}=(1/12)\,g_{-}ij^2$	[MANIFOLD]
ST.692	calc	$\pi(V_{-}\tau)=e^{\sim}\{-(\sqrt{\cdot}/(\pi/16)\log x)^{\sim}\delta\}\times1/(x\log x)\qquad=0.239394$	[ZETA, TRANSPORT]
ST.693	symb	$d/dt\,(g_{-}ij)^2=(1/24)(F_{-}t^{\sim}m)^2$	[MANIFOLD]
ST.694	calc	$m^2=2\pi T((26-D_{-}n)/24)\qquad=4.18879$	[QUANTUM]
ST.695	symb	$g_{-}ij\wedge\pi(V_{-}\tau)=e^{\sim}\{-2\pi T \phi \}[\eta_{-}\mu\nu+\bar{h}_{-}\mu\nu(x)]dx^{\sim}\mu\,dx^{\sim}\nu+T^2\,d\phi^2$	[MANIFOLD, TRANSPORT]
ST.696	symb	$ ds^2 =g_{-}ij\wedge\pi(V_{-}\nu)$	[MANIFOLD, TRANSPORT]
ST.697	holds	$e^{\sim}\{i\theta\}=\cos\theta+i\sin\theta\qquad=0.453596+0.891207i\,\text{vs}\,0.453596+0.891207i$	[ROT]
ST.698	symb	$e^{\sim}\{x\log x\}=x^{\sim}\{1/2+iy\},\,x\log x=\log(\cos\theta+i\sin\theta)=\log\cos\theta+i\log\sin\theta$	[ROT, ZETA]
ST.699	symb	$\log(\sin\theta+i\cos\theta)=\log(\sin\theta-i\cos\theta)$	[ROT]
ST.700	symb	$\log(\sin\theta/(i\cos\theta))=-2R_{-}ij,\,d/dt\,g_{-}ij(t)=-2R_{-}ij$	[ROT, MANIFOLD]
ST.701	symb	$0(x)=\zeta(s)/\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k$	[ZETA]
ST.702	symb	$\operatorname{Im}f=\ker f,\,\chi(x)=\ker f/\operatorname{Im}f$	[MANIFOLD]
ST.703	symb	$H(3)=2,\,\nabla H(x)=2,\,\pi(x)=0$	[MANIFOLD]
ST.704	symb	$[f(x)]=\infty,\, ds^2 =0(x)[\eta_{-}\mu\nu+\bar{h}_{-}\mu\nu(x)]dx^{\sim}\mu\,dx^{\sim}\nu+T^2\,d^2\phi$	[SR, MANIFOLD, TRANSPORT]
ST.705	symb	$T^2\,d^2\phi=[f(x)],\,T^2\,d^2\phi=\lim_{\{x\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k$	[MANIFOLD, TRANSPORT]
ST.706	symb	$T^2\,d^2\phi=\lim_{\{x\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k$	[TRANSPORT]
ST.707	symb	$\lim_{\{x\rightarrow1\}}\Sigma_{\{k=0\}^{\infty}}a_kf^{\sim}k=[T^2\,d^2\phi]$	[TRANSPORT]
ST.708	symb	$d/dL\,V(\tau)=d/df\int f_{-}M\,(\sqrt[5]{\cdot}/(x^2))\,d\wedge+d/df\int f_{-}M\,^{\sim}N(\sqrt[5]{\cdot}/x)^{\sim}\{\otimes N\}\,d\wedge$	[MANIFOLD, TRANSPORT]
ST.709	symb	$^{\sim}M(\vee(\wedge f\circ g)^{\sim}N)^{\sim}\{1/2\}=d/df\int f_{-}M\,1/(x\log x)^2\,dx_{-}m+d/df\int f_{-}M\,1/(y\log y)^{\sim}\{1/2\}\,dy_{-}m$	[ZETA, MANIFOLD]
ST.710	symb	$ ds^2 =0(x)[\eta_{-}\mu\nu+\bar{h}_{-}\mu\nu(x)]dx^{\sim}\mu\,dx^{\sim}\nu+T^2\,d^2\phi,\,0(x)=e^{\sim}\{-2\pi T \phi \}$	[MANIFOLD, TRANSPORT]

4. 全方程式レジストリ (2111 本) p.19/33

ST.711	symb	$G^{\{\mu\nu\}} = R_{\mu\nu} T^{\{\mu\nu\}} = -(1/2) \Lambda g_{ij}(x) + T^{\{\mu\nu\}}$	[SR]
ST.712	symb	$[id_x - t]_{\{v=0\}} = \int e^{\{\partial x_m\}} dx_m + L(p,q), r ^2 = \bar{x} x , \Gamma(x) = \int e^{\{-x\}} x^{\{1-t\}} dx = (\bar{x} - i)(x + 1), \Pi(\chi, x) = ie^{\{x \log x\}}$	[SR, GAMMA, TRANSPORT]
UF0.1	calc	$U = GMm/r = 7.50723e+11$	[TRANSPORT]
UF0.2	differs	$\square_{ag}(x) = 2(\sin(i \cdot x \log x) + \cos(i \cdot x \log x)) = 2\cosh(x \log x) = 4.25 + 3.75i$ vs $4.25 + 0i$	[ROT, TRANSPORT]
UF0.3	calc	$E_{\perp} = mc^2 - (1/2)mv^2 = 1.07851e+21$	[SR, TRANSPORT]
UF0.4	holds	$\square_{dal}(x) = \cos(i \cdot x \log x) - i \sin(i \cdot x \log x) = e^{\{x \log x\}} = x^{\bar{x}} = 4 + 0i$ vs $4 + 0i$	[ZETA, TRANSPORT]
UF0.5	calc	$d\mu(x) = 1/(x \log x)^2 = 0.520342$	[MANIFOLD]
UF0.6	holds	$x \log x \rightarrow 0 (x \rightarrow 1) = 0.000100005$ vs 0	[MANIFOLD]
UF0.7	symb	$\int \int 1/(x \log x)^2 dx_m$	[MANIFOLD]
UF0.8	calc	$\zeta(s) = \beta(p,q)/\log x = 0.120225$	[ZETA, BETA]
UF0.9	holds	$\beta(p,q) = \Gamma(p)\Gamma(q)/\Gamma(p+q) = 0.0833333$ vs 0.0833333	[GAMMA, BETA]
UF0.10	calc	$\Xi = \beta(H+1, M+1)/\log(N+1) = 0.0056407$	[ROT, BETA, ENTROPY]
UF0.11	holds	$\alpha_{ag}(x) \equiv \text{Re}[\square_{ag}(x)]/2 = \cosh(x \log x) \geq 1 = 22.8259$ vs 22.8259	[TRANSPORT]
UF0.12	holds	$E_{vac} = \rho \cdot x^{\bar{x}} = 1$ vs 1	[TRANSPORT]
UF0.13	calc	$E_{ag} = U_{grav} \cdot \alpha_{ag}(x) = (GMm/r) \cdot \cosh(x \log x) = 1.59529e+12$	[TRANSPORT]
UF0.14	holds	$x = \text{manifold_coord}(r0/r), r0 = 6.371 \times 10^6 \text{ m} = 2$ vs 2	[TRANSPORT]
UF0.15	holds	$E_{ag} \geq U_{grav} = 1$ vs 1	[TRANSPORT]
UF0.16	holds	$E_{\perp} \approx 1.079 \times 10^{21} \text{ J} (m = 1.2 \times 10^4 \text{ kg}); (1/2)mv^2 = 6 \times 10^{11} \text{ J} (v = 10^4 \text{ m/s}) = 0.999542 + 1i$ vs $1 + 1i$	[SR, TRANSPORT]
UF0.17	calc	$E_{vac}(x) = \rho_{vac} \cdot x^{\bar{x}}, x^{\bar{x}} \rightarrow \infty (x \rightarrow \infty) = 1.5625e-06$	[TRANSPORT]
UF0.18	holds	$e^{\bar{\pi}} \pi \approx \pi^{\bar{e}} e (\pi^{\bar{e}} \pi = 23.1407, \pi^{\bar{e}} e = 22.4592, \Delta = 0.6815) = 0.681535$ vs 0.6815	[OTHER]
UF0.19	calc	$a = (L-1) \cdot g_{eff}, L = E_{ag}/U_{grav} = \cosh(x \log x), g_{eff} = GM/r^2 = 11.047$	[TRANSPORT]
UF0.20	symb	$\leftarrow : \pi(\chi, x)$	[QUANTUM]
UF0.21	symb	$\leftarrow : \int \int 1/(x \log x)^2$	[MANIFOLD]
UF0.22	calc	$\succ : e^{\{-x \log x\}} = 0.25$	[QUANTUM]
UF0.23	holds	$L = 2.125 \text{ (ground)}; L(1 \text{ km}) = 2.12450, L(5 \text{ km}) = 2.12251, L(20 \text{ km}) = 2.11510, L(100 \text{ km}) = 2.07677 = 6.03102e-06$ vs 0	[TRANSPORT]
UF0.24	holds	$\text{altitude after 10 steps} = 607.6 \text{ m}, v = 110.46 \text{ m/s} = 607.538 + 110.454i$ vs $607.6 + 110.46i$	[TRANSPORT]
UF0.25	holds	$\text{reservoir} \geq 99.9\% \text{ of initial after 10 draws of } 5 \times \text{ current level} = 1$ vs 1	[TRANSPORT]
UF0.26	holds	$\Delta < 1 = 1$ vs 1	[OTHER]
INV.1	holds	$\int \int e^{\{-x^2-y^2\}} dx dy = \pi = 3.14159$ vs 3.14159	[GAMMA, OTHER]
INV.2	differs	$\pi^{\bar{e}} = (\int e^{\{\cos \theta + i \sin \theta\}} d\theta)^{\bar{e}} = 22.4592$ vs $147.801 - 2.60195e-16i$	[ROT, OTHER]
INV.3	differs	$\pi^{\bar{e}} = (\int e^{\{i \theta\}} d\theta)^{\bar{e}} = 22.4592$ vs $2.40346e-42 - 4.05263e-42i$	[ROT, OTHER]
INV.4	calc	$e^{\{f \rightarrow f\}} = 1, x \log x = 1 = 1.76322$	[ENTROPY, OTHER]
INV.5	symb	$\int e^{\{-i \theta\}} d\theta = \pi^{\bar{e}}$	[ROT, OTHER]
INV.6	calc	$\square' = 2(\sin(i x \log x) + \cos(i x \log x)) = 4.25 + 3.75i$	[ROT, ENTROPY]
INV.7	holds	$\square = \cos(i x \log x) - i \sin(i x \log x) = 4 + 0i$ vs 4	[ROT, ENTROPY]
INV.8	symb	$\int e^{\{-\square\}} d\square = \pi^{\bar{e}}$	[OTHER]
INV.9	symb	$1/x = C / \log x, C = \int_1 x^{\bar{s}} dx - \log x$	[ZETA]
INV.10	differs	$x^{\bar{y}} = 1/y^{\bar{x}}, \pi^{\bar{e}} = \int e^{\{-\square\}} d\square = e^{\bar{\pi}} \pi / \int e^{\{-\square'\}} d\square' = 22.4592$ vs 0.0432139	[OTHER]
INV.11	symb	$\square = \square' \boxtimes \Psi \rightarrow \square' = \Psi \boxtimes \square$	[QUANTUM]
INV.12	symb	$(d/dl) \square(H\Psi) \nabla, \square (d/dl)(H\Psi) \nabla$	[QUANTUM]
INV.13	symb	$\beta \square - \square' / \log x = \beta \square - \square'$	[BETA]
INV.14	symb	$\int \int \int (\nabla / \nabla_{\perp}) \square(H\Psi) \nabla d\nabla = \int \pi(\chi, \square) \nabla d\nabla = \int \int \pi(\square) d\nabla^{\bar{m}}$	[MANIFOLD, QUANTUM]
INV.15	differs	$e^{\bar{\pi}} \pi = e^{\{\int e^{\{-x^2-y^2\}} dx dy\}} = x^{\bar{y}} = 1/y^{\bar{x}} = \pi^{\bar{e}} = 23.1407$ vs 22.4592	[GAMMA, OTHER]
INV.16	symb	$1/e^{\bar{\pi}} \pi = 1 / \int e^{\{-\square\}} d\square, e^{\bar{\pi}} \pi = \int e^{\{-\square\}} d\square / \int e^{\{-\square'\}} d\square'$	[OTHER]
INV.17	symb	$\pi^{\bar{e}} = e^{\bar{\pi}} \int e^{\{-\square'\}} d\square'; e^{\bar{\pi}} \pi = \pi^{\bar{e}} / \int e^{\{-\square'\}} d\square'$	[OTHER]
INV.18	differs	$\pi^{\bar{e}} = (\int e^{\{-(\cos \theta + i \sin \theta)\}} d\theta)^{\bar{e}} = 22.4592$ vs $147.801 + 1.49055e-14i$	[ROT, OTHER]
INV.19	symb	$\pi^{\bar{e}} = 1/e^{\bar{\pi}} \pi = 1/\int e^{\{-\square\}} d\square$	[OTHER]
INV.20	symb	$\log e^{\bar{\pi}} \pi = \log(\pi^{\bar{e}} / \int e^{\{-\square'\}} d\square')$	[OTHER]
INV.21	symb	$\pi = \square / \square'$	[OTHER]
INV.22	symb	$\pi = e^{\bar{\beta}} \beta \int dx^{\bar{m}}$	[BETA, MANIFOLD]
INV.23	holds	$\pi = 1/\log x = \sqrt{g} = 1 = \pi r^2, r = 1/\sqrt{\pi} = 1$ vs 1	[ZETA, MANIFOLD]
INV.24	differs	$1/\sqrt{\pi} \pi = \int_1 \log x dx, \pi = \int \log x dx^{\bar{m}} = 0.56419$ vs -0.999913	[ENTROPY]
INV.25	symb	$\pi(\chi, x) = [i\pi(\chi, x), f(x)]$	[MANIFOLD, QUANTUM]
INV.26	calc	$i\pi(\chi, x) \circ f(x) = i \int x \log x dx, f(x) \circ \pi(\chi, x) = \int_1 (x \log x) dx = -0.25$	[MANIFOLD, ENTROPY]
INV.27	holds	$\pi(\chi, x) = \int x \log x dx; \pi(\chi, x) = \int_1 x \log x dx = 0.636294$ vs 0.636294	[MANIFOLD, ENTROPY]
INV.28	symb	$\pi(\chi, x) = i\pi(\chi, x), i \int x \log x dx \equiv \int_1 (x \log x) dx$	[ENTROPY]
INV.29	symb	$\int_1 (x \log x) dx = i \int x \log x dx - \int_1 (x \log x) dx$	[ENTROPY]
INV.30	differs	$\int \int_1 (x \log x)^2 dx^{\bar{m}} = 1/(2i) = 0.0740741 + 0i$ vs $0 - 0.5i$	[ENTROPY]
INV.31	calc	$(d/df) F(x,y) = (d/df) \int \int_1 (x \log x)^2 dx^{\bar{m}} + (d/df) \int \int_1 (y \log y)^{\bar{(1/2)}} dy^{\bar{m}} = 0.0740741 + 0.482402i$	[MANIFOLD, ENTROPY]
INV.32	symb	$i = x^{\{90^\circ\}}, x \sin 90^\circ = i$	[ROT]
INV.33	holds	$i = x^{\{1/2\}}, x = -1 = 6.12323e-17 + 1i$ vs $0 + 1i$	[ROT]
INV.34	symb	$\pi(\chi, x) f(x) = i \int_1 (x \log x) \circ f(x) dx = i \int x \log x dx$	[MANIFOLD, ENTROPY]
INV.35	symb	$f(x) \pi(\chi, x) = f(x) \int_1 (x \log x) dx = \int_1 (x \log x) dx$	[MANIFOLD, ENTROPY]
INV.36	symb	$i y = x \sin 90^\circ; y = i, x = 1, i x = y$	[ROT]
INV.37	symb	$i \sin 90^\circ = -1, 1 \sin 90^\circ = i$	[ROT]
INV.38	differs	$\pi^{\bar{e}} e \equiv e^{\bar{\pi}} \pi = 22.4592$ vs 23.1407	[OTHER]

4. 全方程式レジストリ (2111 本) p.20/33

INV.39	holds	$2^2 = e^{\wedge}(x \log x) = 4 \ ; \ 3^2 = e^{\wedge}(x \log x) = 27 \ ; \ 4^4 = 256 \ ; \ 5^5 = 3125 \quad = 3412 \text{ vs } 3412$	[ENTROPY]
INV.40	symb	$y = x \log x \ , \ \log x \rightarrow (\log x)^{\wedge}(-1)$	[ENTROPY]
INV.41	symb	$x^{\wedge}(1/2 + iy) = x \log x \ / \ \log x \ ; \ x^{\wedge}(1/2 + iy) = e^{\wedge}(x \log x)$	[ZETA, ENTROPY]
INV.42	differs	$x^{\wedge}(1/2 + iy) = \log_x e^{\wedge}(x \log x) = \log_x 4, \log_x 27, \log_x 256, \log_x 3125 \quad = 1.08787 + 0.903628i \text{ vs } 2 + 0i$	[ZETA]
INV.43	symb	$y = e^{\wedge}(x \log x) = \sqrt{a} \ , \ e^{\wedge}(e^{\wedge}(x \log x)) = a \ , \ e^{\wedge}((x \log x)^2)$	[ENTROPY]
INV.44	symb	$x^2 = \pm a \ , \ \lim_{\{n \rightarrow \infty\}} (x - y) = e^{\wedge}(x \log x)$	[OTHER]
INV.45	calc	$ ds^2 = 8\pi G \ (\ p/c^3 + V/S \) \quad = 5.5912e-10$	[SR, MANIFOLD]
INV.46	symb	$y = x \log x \ / \ (\log x) = x \ ; \ x \log x = a \Rightarrow a/(\log x) \rightarrow x$	[ENTROPY]
INV.47	symb	$x^{\wedge}(1/2 + iy) = e^{\wedge}(x \log x) \ , \ x^{\wedge}(1/2+iy) = x \log x \ / \ (\log x) = x$	[ZETA]
INV.48	symb	$x = 1/2 \ , \ iy = 0$	[ZETA]
INV.49	symb	$\int \mathbb{C} \, dx^{\wedge}m = 0 \ , \ (d/df) \int \mathbb{C} \, dx^{\wedge}m = 0 \ , \ \theta' = e^{\wedge}(x \log x)$	[MANIFOLD]
INV.50	symb	$H\Psi = \otimes_L \ (i\hbar\nabla) \otimes L = \otimes a \, f \, x^{\wedge}(1-f) \ [Im] = \int e^{\wedge}x \, x^{\wedge}(1-t) \, dx^{\wedge}m = e^{\wedge}(x \log x)$	[GAMMA, QUANTUM]
INV.51	calc	$H = -K \sum p \log p \quad = 1.03972$	[ENTROPY]
INV.52	calc	$\int \Gamma(\gamma)' \, dx^{\wedge}m = (e^{\wedge}f + e^{\wedge}(-f)) \geq (e^{\wedge}f - e^{\wedge}(-f)) \quad = 0.735759$	[GAMMA]
INV.53	differs	$(e^{\wedge}f + e^{\wedge}(-f))(e^{\wedge}f - e^{\wedge}(-f)) = 0 \quad = 7.25372 \text{ vs } 0$	[OTHER]
INV.54	symb	$(d/df) \, F \cdot \int \mathbb{C} \, dx^{\wedge}m \geq 0$	[MANIFOLD]
INV.55	symb	$y = f(g(x)') \, dx = \int f(x)' \, g(x)' \, dx$	[OTHER]
INV.56	symb	$y = f(\log x)' \, dx = f'(x) \cdot (1/x)$	[OTHER]
INV.57	holds	$y = f'(x)/x = 2(\cos(i \, x \log x) - i \sin(i \, x \log x)) = \mathbb{C} \quad = 8 + 0i \text{ vs } 8$	[ROT, ENTROPY]
INV.58	symb	$c = f(x) \cdot \log x \ , \ dx^{\wedge}m = (\log x)^{\wedge}(-1)$	[ZETA]
INV.59	symb	$\mathbb{C} = (d/d\gamma) \, \Gamma = \otimes (i\hbar\nabla) \otimes L$	[GAMMA, QUANTUM]
INV.60	symb	$e^{\wedge}\theta \cdot (d/d\gamma) \, \Gamma = \Gamma \, \gamma' = \int \Gamma(\gamma)' \, dx^{\wedge}m$	[GAMMA]
INV.61	symb	$y = f(x) \log x \ , \ y' = f'(x) + f'(x)/x$	[OTHER]
INV.62	holds	$(\sin(\log x))' \, dx = \cos(\log x) \cdot (1/x) \quad = 0.384619 \text{ vs } 0.384619$	[ROT]
INV.63	symb	$\sin y \ / \ x = \sin \bar{u} = a + t \sin \bar{u}$	[ROT]
INV.64	symb	roots: $i, -i, 2i, -2i$	[ROT]
INV.65	symb	$\lim_{\{n \rightarrow \infty\}} (\ f(b) - f(a)) = f'(c)(b - a)$	[OTHER]
INV.66	symb	$\sin y \ / \ x = 2(\cos(i \, x \log x) - i \sin(i \, x \log x)) = e^{\wedge}\theta$	[ROT]
INV.67	symb	$\int\!\!\!\int \Delta(\pi(x,x)) \ [Im]$	[MANIFOLD]
INV.68	symb	$[\Box/\nabla]_{\mu\nu} \ , \ \nabla + \otimes \Delta \ , \ F^{\top}\bot \ , \ \Sigma \ (\ \sigma(\, H(\delta) \times K(\delta) \,) \)$	[MANIFOLD]
INV.69	symb	$\nabla_{\perp} i \nabla_{\perp} j \ (\Box \times \Box') \, d\tau \ , \ \sqrt{(x^{\wedge}m \, y^{\wedge}m)}$	[MANIFOLD]
INV.70	symb	$\int\!\!\!\int \Box \, \text{cohom } D_{\chi} \ [Im] \otimes [\, S \, D_{\chi} \otimes h\nu \,]$	[MANIFOLD, QUANTUM]
INV.71	symb	$\ll i\hbar\phi \ \ * \ \, H\Psi \gg \ , \ \int \Delta(\xi) \, d\xi$	[ZETA, QUANTUM]
INV.72	symb	$\oint\!\!\!\oint R \ (Im) \, \nabla_{\perp} L$	[MANIFOLD]
INV.73	symb	$-2 \int \nabla_{\perp} i \nabla_{\perp} j \ (R + \nabla_{\perp} i \nabla_{\perp} j \, f) \ / \ \Delta(R + \Delta) \, dm$	[MANIFOLD]
INV.74	symb	$\Delta(\, F(\Delta) \times \Delta(G(\Delta)) \,) = -(\, F(\Delta) \cup F(\Delta) \,) + (\, F(\Delta) \cap F(\Delta) \,)$	[MANIFOLD]
INV.75	symb	$\Sigma \ \Box(\nabla)[Im] \, \nabla + \ , \ [\nabla/\Box] \ , \ (\nabla + +) \ , \ \chi(x)$	[MANIFOLD]
INV.76	symb	$\pi \ \ \int \nabla_{\perp} i \nabla_{\perp} j \ \int \nabla f \, d\eta \ ^2 = S^{\wedge}m \times S^{\wedge}(m-1)$	[MANIFOLD]
INV.77	symb	$\pi \, r^2 \, dr^{\wedge}m \ , \ (a \, t - t^{\wedge}n + a) = e^{\wedge}f \rightarrow \partial/df$	[JONES, MANIFOLD]
INV.78	calc	$(d/d(e^{\wedge}f + e^{\wedge}(-f))) \ (e^{\wedge}(-f) - e^{\wedge}f) \quad = -1.31304$	[OTHER]
INV.79	symb	$e^{\wedge}f = a \, t^{\wedge}n - t^{\wedge}(n-1) + nCr \, x^{\wedge}n \, y^{\wedge}(n-1)$	[GAMMA, JONES]
INV.80	symb	$(1/2) \, m \, t^2 - \in x^{\wedge}n \, y^{\wedge}(n-1) \, dx^{\wedge}m \, dy^{\wedge}m$	[OTHER]
INV.81	symb	$\ll i\hbar\phi \ \ * \ \, H\Psi \gg = f \ \Box' (\Box\Psi) \, dm$	[QUANTUM]
INV.82	symb	$\times([\pi(x,x), \downarrow]) = \chi(y \ \ : \rightarrow x) = \chi^{\wedge}(-1)(x) \times \chi(y) \ , \ \Box\Psi = \nabla + \pi \ \ \int [\times \ \ : \ x \rightarrow y] \ ^2 \, d\tau$	[MANIFOLD, QUANTUM]
INV.83	symb	$\Box \ \ : \ x \rightarrow f(x) \ , \ \Psi([\pi(x,x), \downarrow]) = (\downarrow, \downarrow, \rightarrow, \leftarrow)$	[QUANTUM]
INV.84	symb	$\sigma(\, H(\delta) \times K(\delta) \,) = \Box' \int\!\!\!\int d \bowtie$	[MANIFOLD]
INV.85	symb	$m \otimes (i\hbar\nabla) \otimes L + n \otimes (i\hbar\nabla) \otimes L = (m+n) \otimes (i\hbar\nabla) \otimes L$	[QUANTUM]
INV.86	symb	$m \otimes (i\hbar\nabla) \otimes L - n \otimes (i\hbar\nabla) \otimes L = (m-n) \otimes (i\hbar\nabla) \otimes L$	[QUANTUM]
INV.87	symb	$\otimes (i\hbar\nabla) \otimes L^{\wedge}m \times \otimes (i\hbar\nabla) \otimes L^{\wedge}n = \otimes (i\hbar\nabla) \otimes L^{\wedge}(m+n)$	[QUANTUM]
INV.88	symb	$\otimes (i\hbar\nabla) \otimes L^{\wedge}m \ / \ \otimes (i\hbar\nabla) \otimes L^{\wedge}n = \otimes (i\hbar\nabla) \otimes L^{\wedge}(m/n)$ (sic, should be m-n)	[QUANTUM]
INV.89	symb	$d/df \ \otimes (i\hbar\nabla) \otimes L = \otimes (i\hbar\nabla) \otimes L \otimes (i\hbar\nabla) \otimes L' = \otimes (i\hbar\nabla) \otimes L'$	[QUANTUM]
INV.90	symb	$\int \ \otimes (i\hbar\nabla) \otimes L \, dx^{\wedge}m \ , \ \otimes (i\hbar\nabla) \otimes L = n$	[QUANTUM]
INV.91	symb	$n^{\wedge}(L+1) \ / \ (n+1) = \int e^{\wedge}x \, x^{\wedge}(1-t) \, dx$	[GAMMA]
INV.92	holds	$(\ \otimes (i\hbar\nabla) \otimes L + m \otimes (i\hbar\nabla) \otimes L + n \) = n^{\wedge}(L+1)/(n+1) = t \int (1-x)^{\wedge}n \, x^{\wedge}(m-1) \, dx = \int x^{\wedge}(m-1)(1-x)^{\wedge}(n-1) \, dx \quad = 0.0833333 \text{ vs } 0.0833333$	[GAMMA, BETA, MANIFOLD]
INV.93	symb	$\nabla(i\hbar\nabla) \otimes L \ , \ \otimes (i\hbar\nabla) \otimes L \ , \ \Box(i\hbar\nabla) \otimes L \ , \ \boxtimes(i\hbar\nabla) L \, dx^{\wedge}m \ , \ \boxplus(i\hbar\nabla L \, dx^{\wedge}m)$	[QUANTUM]
INV.94	symb	$\int f(x) \, dx = \int \Gamma(\gamma)' \, dx^{\wedge}m = 2(\cos(i \, x \log x) - i \sin(i \, x \log x))$	[ROT, GAMMA]
INV.95	symb	$(\ \int f(x)dx \ / \ \log x \) = \lim_{\{\theta \rightarrow \infty\}} (\ \int f(x)dx \ / \ \theta \) = 0, \ 1$	[OTHER]
INV.96	holds	$e^{\wedge}(i \, \theta) = \cos \, \theta + i \sin \, \theta \quad = 0.866025 + 0.5i \text{ vs } 0.866025 + 0.5i$	[ROT]
INV.97	symb	$(\ \int f(x) \, dx \)' = 2(\, i \sin(i \, x \log x) - \cos(i \, x \log x) \,) = 2(\, -\cos(i \, x \log x) + i \sin(i \, x \log x) \,)$	[ROT]
INV.98	symb	$(\cos(i \, x \log x) - i \sin(i \, x \log x) \)' = (d/de^{\wedge}(i\theta)) \ (\cos, -\sin) \cdot (\sin, \cos) \)$	[ROT]
INV.99	symb	$\int \Gamma(\gamma)'' \, dx^{\wedge}m = \int \Gamma(\gamma)' \, dx^{\wedge}m \ / \ \nabla L$	[GAMMA]
INV.100	symb	$\int \Gamma \, dx^{\wedge}m \cdot (d/d\gamma)\Gamma \ / \ \nabla L \leq (\ \int \Gamma \, dx^{\wedge}m + (d/d\gamma)\Gamma \) \ / \ \nabla L \leq \ e^{\wedge}f - e^{\wedge}(-f) \leq e^{\wedge}(-f) + e^{\wedge}f \ = 0, \ 1$	[GAMMA]
INV.101	symb	$f = \Pi_{\perp} f_{\perp}$ (conductor of abelian K/k)	[ZETA]
INV.102	symb	$(\ \alpha, \, K/k \ / \ p \) = (\, K/k \ / \ b \) \ (\in G)$	[ZETA]
INV.103	symb	$\alpha/\alpha_o \equiv 1 \ (\text{mod } f_{\perp}) \ , \ \alpha_o \equiv 1 \ (\text{mod } f \, f_{\perp}^{\wedge}(-1)) \rightarrow \alpha \in k \ , \ (\alpha_o) = p^{\wedge}\alpha \, b$	[ZETA]
INV.104	symb	$(\ \alpha, \, K/k \ / \ p_{\perp}^{\wedge}(\infty) \) = 1 \text{ or } 0$	[ZETA]
INV.105	symb	$\Pi_{\perp} \ (\ \alpha, \, b \ / \ p \) = 1$	[ZETA]

4. 全方程式レジストリ (2111 本) p.21/33

INV. 106	symb	$ ds^2 =\lim_{\{x\rightarrow\infty}\left[\delta(x)\int\int\pi\left(\sum_{\{k=0\}^{\infty}}(n\sqrt{p},x/n)\right)^{(1/2)}d\tau\right]_{\mu\nu}$	[ZETA, MANIFOLD]
INV. 107	symb	$p^{\left(\alpha\right)}_n=n\sqrt{p}/n,\text{ }n\sqrt{p}=\otimes(i\hbar\nabla)\otimes L=n\text{ }p^{\left(1/n\right)}=n-n\text{ }p$	[ZETA, QUANTUM]
INV. 108	calc	$\int\Gamma(\gamma)^{\prime}dx^{\prime}m=e^{\left(-x\log x\right)}=0.25$	[GAMMA, ENTROPY]
INV. 109	calc	$\lim_{\{s\rightarrow 1+0\}}\sum_{\{p\in M\}}(1/N(p)^{\prime}s)/\log(1/(s-1))=\text{density of }M=0.954504$	[ZETA]
INV. 110	symb	$\alpha(f\text{ }d/dt,\text{ }g\text{ }d/dt)=\int_S\left f^{\prime}\text{ }f^{\prime\prime};\text{ }g^{\prime}\text{ }g^{\prime\prime}\right dt$	[OTHER]
INV. 111	symb	$B(f\text{ }d/dt,\text{ }g\text{ }d/dt,\text{ }h\text{ }d/dt)=\int_S\left f\text{ }f^{\prime}\text{ }f^{\prime\prime};\text{ }g\text{ }g^{\prime}\text{ }g^{\prime\prime};\text{ }h\text{ }h^{\prime}\text{ }h^{\prime\prime}\right dt$	[OTHER]
INV. 112	symb	$E_0\times S^2$	[MANIFOLD]
INV. 113	calc	$e^{\left(-x\log x\right)}\leq y\leq e^{\left(x\log x\right)},\text{ }y\neq 0=2.25$	[ENTROPY]
INV. 114	symb	$(\partial/\partial f)F(x)=\int\!\!\!\int\text{cohom }D_k(x)\left[Im\right]$	[MANIFOLD]
INV. 115	symb	$(\partial/\partial f)F=\text{ }^t\!\!\!\int\!\!\!\int\text{cohom }D_k(x)\ll p=\square^{\prime}$	[MANIFOLD]
INV. 116	symb	$\nabla_i\nabla_j\int f(x)\text{ }d\eta=\partial^2/(\partial x\partial y)\int\square^{\prime}d\eta$	[MANIFOLD]
INV. 117	calc	$R^{\prime}_{\mu\nu}+(1/2)\wedge g^{\prime}_{ij}=\int\left(i\text{ }v/\sqrt{(1-(v/t)^2)}+v/\sqrt{(1-(v/t)^2)}\right)dvol=0.2+0.2i$	[SR, MANIFOLD]
INV. 118	symb	$=\int C\text{ }dx^{\prime}m=\int\kappa\text{ }T_{\mu\nu}\text{ }dx^{\prime}m=T_{\mu\nu}\text{ }T^{\mu\nu}$	[MANIFOLD]
INV. 119	calc	$\square^{\prime}=(8\pi G/c^4)T_{\mu\nu}/\log x=2.99583e-43$	[SR, MANIFOLD]
INV. 120	symb	$\text{ }^t\!\!\!\int\!\!\!\int\text{cohom }D_{\mathcal{X}}\left[Im\right]=\oint(p\text{ }x^{\prime}n+q\text{ }x+r)\nabla_{\perp}$	[MANIFOLD]
INV. 121	calc	$(d/dl)L(x,y)=2\int \sin 2x ^2d\tau,\text{ }(d/d\gamma)\Gamma=3.14159$	[ROT, GAMMA]
INV. 122	symb	$\oint\!\!= ds^2 =\lim_{\{x\rightarrow\infty}\left[\delta(x)\int\int\pi\left(\sum n\sqrt{p},x/n\right)^{(1/2)}d\tau\right]_{\mu\nu}$	[ZETA, MANIFOLD]
INV. 123	symb	$e^{\left(-2\pi T\right)} \phi \left[\eta+\hbar\right]dx^{\prime}\mu\text{ }dx^{\prime}\nu+T^2\text{ }d^2\phi$	[MANIFOLD]
INV. 124	symb	$-(16\pi G/c^4)T_{\mu\nu}/\log x=\square^{\prime};\text{ }-(16\pi G/c^4)T_{\mu\nu}/e^{\left(-2\pi T\right)} \phi =4\pi G\rho$	[SR, MANIFOLD]
INV. 125	symb	$\partial/(\partial x\partial y)=\nabla_i\nabla_j;\square\int\!\!\!\int=\text{ }^t\!\!\!\int\!\!\!\int;\text{ }(\partial/\partial x)\int\!\!\!\int=\nabla_i\nabla_j\int\nabla f(x)\text{ }d\eta;\text{ }(\partial/\partial f)\int\!\!\!\int=\square\int\!\!\!\int$	[MANIFOLD]
INV. 126	symb	$\Gamma \Gamma,\text{ }B B,\text{ }E E,\text{ }C C,\text{ }F F,\text{ }\beta \beta,\text{ }D D,\text{ }\text{ }^t\!\!\!\int\!\!\!\int\equiv\otimes D\otimes L$	[GAMMA, BETA]
INV. 127	symb	$\square+\square^{\prime}=\phi,\text{ }\square^{\prime} \phi=\square$	[OTHER]
INV. 128	calc	$[[1,0],[0,-i]]^{(1/2)}\cdot[[1,0],[0,-i]]=-0.707107-0.707107i$	[QUANTUM]
INV. 129	symb	$x^{(1/2)}\equiv x,\text{ }\phi^{(1/2)}=\phi$	[OTHER]
INV. 130	symb	$\oint r\text{ }dx^{\prime}m=0(x,y)$	[MANIFOLD]
INV. 131	symb	$\text{ }^t\!\!\!\int\!\!\!\int D(\mathcal{X},x)\text{ Hom}[D^2\phi]\ll p\equiv vol\text{ } (V/S)$	[MANIFOLD]
INV. 132	symb	$(d/df)F=(F)\text{ }f^{\prime},\text{ }\int F\text{ }dx^{\prime}m=(F)\text{ }f$	[OTHER]
INV. 133	symb	$(d/df)\int\int_1(x\log x)^2\text{ }dx^{\prime}m=\int\int_1(x\log x)^2\text{ }dx^{\prime}m$	[ENTROPY]
INV. 134	symb	$f^{\prime}/f^{\prime}=2x(\log x+\log(x+1))$	[OTHER]
INV. 135	symb	$\!\!\!\int\!\!\!\int F\text{ }dx^{\prime}m=(1/(x\log x)^2)\left(\int 2x(\log x+\log(x+1))\text{ }dx\right)=e^{\left(-f\right)}$	[ENTROPY]
INV. 136	calc	$(d/df)F=(1/(x\log x)^2)\left(2x(\log x+\log(x+1))\right)=e^{\left(-f\right)}=3.72931$	[ENTROPY]
INV. 137	calc	$\log(x\log x)\geq 2\left(y\log y\right)^{(1/2)};\log(x\log x)\geq 2\sqrt{(y\log y)}=-2.43823$	[ENTROPY]
INV. 138	symb	$\int ds^{\prime} \text{ }dx^{\prime}m=\int 8\pi G(p/c^3+V/S)\text{ }dvol=\int 8\pi G(p/c^3+V/S)\text{ }dy^{\prime}m=\int_1 8\pi G(p/c^3+V/S)\text{ }dx^{\prime}m$	[SR, MANIFOLD]
INV. 139	symb	$(d/df)F=m(x),\text{ }\otimes i\hbar\nabla\otimes L=\nabla_i\nabla_j\int\nabla f(x)\text{ }d\eta$	[MANIFOLD, QUANTUM]
INV. 140	symb	$dx^{\prime}m=y/\log x,\text{ }dy^{\prime}m=x/\log x$	[ZETA]
INV. 141	symb	$e^{\left(-f\right)}dV=dy^{\prime}m=dvol$	[MANIFOLD]
INV. 142	symb	$\beta(p,q)=(\text{number})+\text{Abel manifold}=\text{AdS}_5\text{ manifold}=(d/df)F+\int C\text{ }dx^{\prime}m=\int\Gamma(\gamma)^{\prime}\text{ }dx^{\prime}m$	[GAMMA, BETA, MANIFOLD]
INV. 143	symb	$ ds^2 =e^{\left(-2\pi G \phi \right)}\left[\eta+\hbar(x)\right]dx^{\prime}\mu\text{ }dx^{\prime}\nu+T^2\text{ }d^2\phi$	[MANIFOLD]
INV. 144	symb	$\int\text{ }dvol=\square\phi,\text{ }\int\nabla\phi^2\text{ }d\nabla\phi=\square\phi$	[MANIFOLD]
INV. 145	symb	$\text{expanding of universe}=\text{exist of value}=\log(x\log x)=\square\phi$	[ENTROPY]
INV. 146	symb	$\text{freeze out of universe}=\text{reality of value}=(y\log y)^{(1/2)}=\nabla\phi$	[ENTROPY]
INV. 147	symb	$x^{\prime}n+y^{\prime}n=z^{\prime}n,\text{ }\beta(p,q)=x^{\prime}n+y^{\prime}n-\delta(x)=z^{\prime}n-\delta(x)$	[BETA]
INV. 148	symb	$(d/dt)g_{ij}=-2R_{ij}$	[MANIFOLD]
INV. 149	differs	$\cos x/\left((\cos x)^{\prime}\cdot(\sin x)^{\prime}\right)=z^{\prime}n,\text{ }z^{\prime}n=-2e^{\left(x\log x\right)}=-1.1884\text{ vs }-2$	[ROT]
INV. 150	calc	$z=e^{\left(-f\right)}+e^{\left(-f\right)}-y,\text{ }\beta(p,q)=e^{\left(-f\right)}+e^{\left(-f\right)}=3.08616$	[BETA]
INV. 151	symb	$\square x=\int^{\prime}\{f(x)\}\nabla(R+\cap E+)\text{ }d\square x=\int^{\prime}\{\Delta f(x)\circ E+\}\nabla(R+\cap E+)\square x$	[MANIFOLD]
INV. 152	symb	$\square x=\int^{\prime}\{d(R\nabla E+)\}\nabla(R+\cap E+)\text{ }d\square x=\int^{\prime}\{\nabla i\nabla j(R+E+)\}\nabla(R+\cap E+)\text{ }d\square x$	[MANIFOLD]
INV. 153	symb	$\exp(\nabla(R+\cap E+),\Delta(C\supset R))=\pi(R1\subset\nabla E+)=\text{rot}(E1,\text{div }E2)$	[ROT, MANIFOLD]
INV. 154	holds	$x\text{ }f(x)=F(x),\text{ }s\text{ }\Gamma(s)=\Gamma(s+1)=3.32335\text{ vs }3.32335$	[GAMMA]
INV. 155	symb	$Q\nabla C+=\left(d/df\right)F(x)\nabla\int\delta(s)\text{ }f(x)\text{ }dx$	[OTHER]
INV. 156	symb	$E+\nabla f=e^{\left(x\log x\right)}\nabla n!\text{ }f(x)$	[GAMMA]
INV. 157	symb	$E(x)\left(d/df\right)F\text{ }f^{\prime}=e^{\left(x\log x\right)};\left(C\nabla\right)\otimes Q=e^{\left(x\log x\right)}$	[ENTROPY]
INV. 158	symb	$R\nabla E+=f(x)\nabla e^{\left(x\log x\right)},\text{ }\left(d/df\right)F=F\text{ }f^{\prime}=e^{\left(x\log x\right)}$	[ENTROPY]
INV. 159	symb	$\int dx=x+C\text{ } (C\text{ integration constant})$	[OTHER]
INV. 160	holds	$\beta(p,q)=\int x^{(1-t)}(1-x)^{\text{ }^t\!}dx=\int t^{\prime}x^{(1-t)}(x-1)\text{ }dt=0.392697\text{ vs }0.392699$	[BETA]
INV. 161	holds	$0\leq y\leq 1,\int_0^1x^{*10}(1-x)^{*20}dx=B(11,21)=\Gamma(11)\Gamma(21)/\Gamma(32)=10!\cdot20!/31!=1/931395465=1\text{ vs }1$	[GAMMA, BETA]
INV. 162	symb	$1/931395465=1/9=1/(1-x)\text{ (sic)}$	[OTHER]
INV. 163	holds	$1/(1-z)=\sum_{\{k=0\}^{\infty}}z^{\prime}k,\text{ }1/(1+z^2)=\sum_{\{k=0\}^{\infty}}(-1)^{\prime}k\text{ }z^{\prime}(2k)=2.8\text{ vs }2.8$	[OTHER]
INV. 164	symb	$f(x)=\sum_{\{k=0\}^{\infty}}a_k\text{ }z^{\prime}k$	[OTHER]
INV. 165	symb	$d^{\prime}ny/dx^{\prime}n=n!\text{ }y^{\prime}(n+1),\text{ }f^{\prime}(0)(0)=n!\text{ }f(0)^{\prime}(n+1)=n!$	[OTHER]
INV. 166	symb	$f(x)\equiv\sum_{\{k=0\}^{\infty}}x^{\prime}n=1/(1-x)$	[OTHER]
INV. 167	holds	$dy/dx=y^2,\text{ }(1/y^2)(dy/dx)=1,\int(1/y^2)(dy/dx)\text{ }dx=\int dy/y^2=-1/y,\text{ }-1/y=x-C,\text{ }y=1/(C-x)=4\text{ vs }4$	[OTHER]
INV. 168	symb	$\exists x=0,\text{ }y=1\Rightarrow y=1/(1-x)$	[OTHER]
INV. 169	calc	$(dx,\partial x)\cdot(\varepsilon x,\delta x)=[[1,0],[0,-1]]^{(1/2)}=6.12323e-17+1i$	[QUANTUM]
INV. 170	symb	$\pi\left(\sum_{\{k=0\}^{\infty}}(n\sqrt{p},x/n)\right)^{(1/2)}d\tau]_{\mu\nu}=e^{\left(-f\right)}dV$	[ZETA, MANIFOLD]
INV. 171	symb	$V=\int\!\!\!\int\pi\left(e^{\left(-f\right)}dV\right)\text{ }dx^{\prime}m,\text{ }\delta V=M,\text{ }(d/dM)V=m(x)$	[MANIFOLD]
INV. 172	symb	$\nabla\phi^2=8\pi G(p/c^3+V/S)$	[SR, ENTROPY]
INV. 173	symb	$y=x;\text{ }y=(\nabla\phi)^{(1/2)}$	[OTHER]

INV. 174	symb	$\Delta E^{\wedge}(1/2) \ / \ \sqrt{(2 \pi G)} = p/c^2 + \rho$	[SR]
INV. 175	symb	$ds^2 = g_{\mu \nu}(x) \, dx^{\wedge} \mu \, dx^{\wedge} \nu + \phi^2(x) \, (\kappa^2 \, A_{\mu \nu}(x) \, dx^{\wedge} \mu)^2$	[SR, MANIFOLD]
INV. 176	symb	$ds = (\, g_{\mu \nu}(x) \, dx^{\wedge} \mu \, dx^{\wedge} \nu + \phi^2(x) (\kappa^2 \, A_{\mu \nu}(x) \, dx^{\wedge} \mu)^2 \,)^{\wedge}(1/2)$	[SR, MANIFOLD]
INV. 177	symb	$\Delta E^{\wedge}(1/2)/\sqrt{(2 \pi G)} - \rho = c^3 \, , \, p/(2 \pi) = c^3$	[SR]
INV. 178	symb	$ds^2 = e^{\wedge}(-2kT(x) \phi) \, [\eta_{\mu \nu} + \bar{h}_{\mu \nu}(x)] \, dx^{\wedge} \mu \, dx^{\wedge} \nu + T^2(x) \, d\phi^2$	[MANIFOLD]
INV. 179	symb	$f_{,z} = \int \, \sqrt{(\, x1 \, x2 \, x3; \, y1 \, y2 \, y3 \, \circ \, x1 \, x2 \, x3; \, y1 \, y2 \, y3 \,) \, } \, dx \, dy \, dz$	[MANIFOLD]
INV. 180	symb	$(\partial/\partial f) \int (\sin 2x)^2 \, dx = x - y ^2$	[ROT]
INV. 181	symb	$x = \overline{x} \ / \ \overline{x} $	[OTHER]
INV. 182	symb	$ \overline{x} = \sqrt{x} \, , \, i = \sqrt{-1} \, , \, \overline{x} \, x = \overline{x} \, , \, \overline{x} = 1x \, , \, \overline{x} ^2 = -1$	[ROT]
INV. 183	symb	$(\, x \ / \ \overline{x} \,)^2 = 1/(-1) \, , \, \sqrt{x} = i$	[ROT]
INV. 184	symb	$x \bmod N = 0$	[OTHER]
INV. 185	symb	$\Sigma_{\{M=0\}}^{\wedge} \infty \, \int_{_M} dm \rightarrow \Sigma_{\{x=0\}}^{\wedge} \infty \, F_{_x} = \int_{_m} dm = F$	[OTHER]
INV. 186	symb	$(x-y)/a = (y-z)/b = (z-x)/c \, , \, \Pi(x-y) \ / \ \partial xy$	[OTHER]
INV. 187	symb	$ \int f(x) \, ^2 \rightarrow \int \pi \, r^2 \, dx \equiv V(\tau)$	[MANIFOLD]
INV. 188	symb	$\int f(x) ^2 \, / \, f(x) \, dx$	[ENTROPY]
INV. 189	calc	$V(\tau) \rightarrow mesh \int x1 \, x2 \, x3; \, y1 \, y2 \, y3; \, z1 \, z2 \, z3 \, dv(\tau) \, , \, x1..z3 \, dx = v \quad = 1$	[MANIFOLD]
INV. 190	symb	$z \, y = a \, x + b \, y + c \, z \, , \, d \, z \, y = d(zy)$	[OTHER]
INV. 191	symb	$[f, \, f^{\wedge}(-1)] = f \, f^{\wedge}(-1) - f^{\wedge}(-1) \, f$	[QUANTUM]
INV. 192	symb	$V(\tau) = \int \, \tau \, (q)^{\wedge}(-n/2) \, \exp(\, -(1/\sqrt{(2 \, \tau \, q))}) \, L(x) \, dx \,) + 0(N^{\wedge}(-1))$	[MANIFOLD, ENTROPY]
INV. 193	symb	$(1/\tau) \, (\, N/2 + \, \tau \, (2\Delta f - \nabla f ^2 + R) + f \,) \bmod \, N^{\wedge}(-1)$	[MANIFOLD, ENTROPY]
INV. 194	symb	$\Delta E = -2(T-t) \, \, R_{ij} + \nabla_i \nabla_j \, f - g_{ij}/(2(T-t)) \, ^2$	[MANIFOLD, ENTROPY]
INV. 195	symb	$(d/df) \, F = 2 \, \int \, (R + \nabla_i \nabla_j \, f)^2 \, / \, -(R + \Delta f) \, dm$	[MANIFOLD, ENTROPY]
INV. 196	symb	$(d/dt) \, g_{ij}(t) = -2 \, R_{ij}$	[MANIFOLD]
INV. 197	symb	$dx = (\, g_{\mu \nu}(x)^2 \, dx^2 - g_{\mu \nu}(x) \, dx \, g_{\mu \nu} \,)^{\wedge}(1/2)$	[MANIFOLD]
INV. 198	symb	$ds^2 = -N(r)^2 \, dt^2 + \phi^2(r)(dr^2 + r^2 \, d\theta^2)$	[SR, MANIFOLD]
INV. 199	symb	$\Sigma_{\{n=0\}}^{\wedge} \infty \, (a1 \, x1 + a2 \, x2 \, \ldots \, a_{\{n-1\}} \, x_{\{n-1\}}) \rightarrow \Sigma_{\{n=0\}}^{\wedge} \infty \, a_n \, x^{\wedge} n \rightarrow \alpha$	[OTHER]
INV. 200	symb	$f = n \, \nu \, \lambda \, , \, \lambda = x/l \, , \, \int \, d \, n \, \nu \, \lambda = f(x) \, , \, x \, f(x) = F(x) \, , \, [f(x)] = \nu \, h$	[QUANTUM]
INV. 201	symb	$0(x) = ([\nabla_i \nabla_j \, f(x)])^{\wedge} \equiv nCr \, (x)^{\wedge} n \, (y)^{\wedge} (n-r) \, \delta(x,y)$	[MANIFOLD]
INV. 202	symb	$(\Box \phi)^{\wedge} = \nabla_i \nabla_j \, (\delta(x) \circ G(x))_{\mu \nu} \, (\, p/c^3 \, \circ \, V/S \,)$	[MANIFOLD]
INV. 203	symb	$F_{m^{\wedge} t} = (1/(4g^2)) \, ij \, , \, x^{\wedge}(1/2+iy) = e^{\wedge}(x \log x)$	[ZETA, QUANTUM]
INV. 204	symb	$S^{\wedge} \mu \, \nu_{_m} \circ S^{\wedge} \mu \, \nu_{_n} = G_{\mu \nu} \, \times \, T_{\mu \nu}$	[MANIFOLD]
INV. 205	symb	$S^{\wedge} \mu \, \nu_{_m} \circ S^{\wedge} \mu \, \nu_{_n} = -2R_{ij} \, / \, V(\tau) \, [D^2 \phi]$	[MANIFOLD]
INV. 206	symb	$S^{\wedge} \mu \, \nu_{_m} = \pi(\chi,x) \circ h_{\mu \nu} \, , \, \pi(\chi,x) = \int \, \exp[L(p,q)] \, d\phi$	[MANIFOLD, QUANTUM]
INV. 207	symb	$ds^2 = e^{\wedge}(-2\pi T \phi) \, [\eta + \bar{h}_{\mu \nu}] \, dx^{\wedge} \mu \, \nu \, dx^{\wedge} \mu \, \nu + T^2 \, d^2 \phi$	[MANIFOLD]
INV. 208	symb	$M3 \circ_{\{k=0\}}^{\wedge} \infty \, Et+ = \text{rot}(\text{div} \, E, \, E1) = m(x) \, , \, P^{*2n} M3 = H3(M1)$	[ROT, MANIFOLD]
INV. 209	symb	$\exists [R + \nabla f ^2]^{\wedge}(1/2+iy) = f \, \exp[L(p,q)] \, d\phi = \exists [R + \nabla f ^2]^{\wedge}(1/2+iy) \circ f \, \exp[L(p,q)] \, d\phi + N \bmod(e^{\wedge}(x \log x)) = 0(\phi)$	[ZETA, QUANTUM]
INV. 210	symb	$P^{*2n} M3 = H3(M1) \, , \, H3(M1) = \pi(\chi,x) \circ h_{\mu \nu}$	[MANIFOLD]
INV. 211	symb	$S^{\wedge} \mu \, \nu_{_m} \times S^{\wedge} \mu \, \nu_{_n} = [D^2 \phi] \, , \, S \times S = \ker f \, / \, \text{im} \, f \, , \, S \circ S = m(x)[D^2 \phi] \, , \, -2R_{ij} \, / \, V(\tau) = f^{\wedge}(-1) \, x \, f(x)$	[MANIFOLD]
INV. 212	symb	$f_{,z} = \int \, [\, \sqrt{(\, x \, y \, z; \, u \, v \, w \, \circ \, x \, y \, z; \, u \, v \, w \,) \, } \,] \, dx \, dy \, dz \rightarrow f^{\wedge}(1/2)$	[MANIFOLD]
INV. 213	symb	$z \rightarrow (0,1) \cdot (0,1) = -1 \, , \, i = \sqrt{-1}$	[ROT]
INV. 214	symb	$(x,y,z)^2 = (x,y,z) \cdot (x,y,z) \rightarrow -1$	[ROT]
INV. 215	symb	$0(x) = \nabla_i \nabla_j \, \int \, e^{\wedge}((2/m) \sin \theta \cos \theta) \, \times \, N \bmod(e^{\wedge}(x \log x))$	[ROT]
INV. 216	symb	$0(x) \, (x + \Delta f ^2)^{\wedge}(1/2)$	[OTHER]
INV. 217	differs	$x \, \Gamma(x) = 2 \, \int \, \sin 2\theta \, ^2 \, d\theta \, , \, 0(x) = m(x)[D^2 \phi] \quad = 2 \, \text{vs} \, 3.14159$	[ROT, GAMMA]
INV. 218	symb	$\lim_{\{\theta \rightarrow 0\}} (1/\theta) \, [\sin \theta; \cos \theta] \, [[\theta,1],[1,\theta]] \, [\cos \theta; \sin \theta] = [[1,0],[0,-1]]$	[ROT, QUANTUM]
INV. 219	symb	$f^{\wedge}(-1)(x) \, x \, f(x) = I'_{_m} \, , \, I'_{_m} = [1,0] \, \times \, [0,1]$	[QUANTUM]
INV. 220	symb	$i^2 = (0,1) \cdot (0,1) \, , \, a b \cos \theta = -1 \, , \, E = \text{div}(E, \, E1)$	[ROT]
INV. 221	calc	$\{f,g\} \, / \, [f,g]^{\wedge} = i^2 \, , \, E = mc^2 \, , \, I' = i^2 \quad = 8.98755e+16$	[SR, QUANTUM]
INV. 222	symb	$0(x) = \, \nabla \, \int \, [\nabla_i \nabla_j \, f \circ g(x)]^{\wedge}(1/2+iy) \, \, , \, \partial r^{\wedge} n \, \nabla ^2 \rightarrow \nabla_i \nabla_j \, \nabla ^2$	[ZETA, MANIFOLD]
INV. 223	symb	$\nabla^2 \phi = 8 \pi G \, (\, p/c^3 + V/S \,)$	[SR, MANIFOLD]
INV. 224	symb	$(\log x^{\wedge}(1/2))^{\wedge} = (1/2)(1/(x \log x)) \, (\text{sic}) \, , \, (\sin \theta)^{\wedge} = \cos \theta \, , \, (f,z)^{\wedge} = i \, e^{\wedge}(i \, x \log x) \, , \, (d/df)F = m(x)$	[ROT]
INV. 225	symb	$(d/df) \int \int_{_1} (x \log x)^2 \, dx^{\wedge} m + (d/df) \int \int_{_1} (y \log y)^{\wedge}(1/2) \, dy^{\wedge} m = (d/df) \int \int \, (\, 1/(x \log x)^2 + 1/(y \log y)^{\wedge}(1/2) \,) \, dm \geq$ $(d/df) \int \int \, (\, 1/((x \log x)^2 \, \circ \, (y \log y)^{\wedge}(1/2)) \,) \, dm \geq 2h$	[QUANTUM, ENTROPY]
INV. 226	symb	$(d/df) \int \int \, (\, 1/((x \log x)^2 \, \circ \, (y \log y)^{\wedge}(1/2)) \,) \, dm \geq \hbar$	[QUANTUM, ENTROPY]
INV. 227	symb	$y = x \, , \, xy = x^2 \, , \, (\Box \phi)^{\wedge} = 8 \pi G \, (\, p/c^3 \, \circ \, V/S \,)$	[MANIFOLD]
INV. 228	symb	$\Box \phi = \iint \exp[\, 8 \pi G \, (\, \bar{h}_{\mu \nu} \, \circ \, \eta_{\mu \nu}) \, \nu \,] \, dm \, d\phi$	[MANIFOLD]
INV. 229	symb	$\Sigma \, a_k \, x^{\wedge} k = (d/df) \, \Sigma \Sigma \, (1/a_k^2) \, f_{_k} \, dx_{_k} \, ; \, \Sigma \, a_k \, f_{_k} = (d/df) \, \Sigma \Sigma \, \xi'(s)/a_k \, dx_{_k} \, m^{\wedge} ; \, a_k^2 \, f^{\wedge}(1/2) \rightarrow \lim_{\{k \rightarrow 1\}} a_k \, f_{_k} = \alpha$	[ZETA]
INV. 230	symb	$0(x) = D^2 \phi \circ h_{\mu \nu} \, (\times 3) \, , \, ds^2 = e^{\wedge}(-2\pi T \phi)[\eta_{\mu \nu} + \bar{h}_{\mu \nu}] \, dx^{\wedge} \mu \, dx^{\wedge} \nu + T^2 \, d^2 \phi$	[MANIFOLD]
INV. 231	calc	$f(x) + f(y) \geq 2\sqrt{(\, f(x) \, f(y) \,)} \, , \, (1/4)(\, f(x) + f(y) \,)^2 \geq f(x) \, f(y) \quad = 2$	[OTHER]
INV. 232	symb	$T_{\mu \nu} = (\, p/c^3 + V/S \,)^{\wedge}(-1) \, , \, Et = f^{\wedge}(-1) \, x \, f(x) \, , \, E = mc^2$	[SR]
INV. 233	symb	$0(x) = \Box \, \iiint \, (\nabla_i \nabla_j \, f \circ g(x))^2 \, / \, V(x) \, dm$	[MANIFOLD]
INV. 234	symb	$ds^2 = g^2_{\mu \nu} \, d^2 x + g_{\mu \nu} \, dx \, g_{\mu \nu}(x) \, , \, Et = f^{\wedge}(-1) \, x \, f(x) \, , \, V/S = A \, A^{\wedge}(-1) \, , \, A \, A^{\wedge}(-1) = E$	[MANIFOLD]
INV. 235	calc	$0(x) = \iiint \, f(x^2, \, y^3, \, z^3) \, dx \, dy \, dz \, , \, S(r) = \pi \, r^2 \, , \, V(r) = 4 \pi \, r^3 \, (\text{sic}; \, 4/3 \, \pi r^3) \quad = 12.5664$	[OTHER]
INV. 236	symb	$Et+ = f(x) \cdot e^{\wedge}(-x \log x) \, , \, \Sigma_{\{k=0\}}^{\wedge} \infty \, a_k \, f_{_k} = f(x) \cdot e^{\wedge}(-x \log x)$	[ENTROPY]
INV. 237	symb	$P^{*2n} M3 = Et - \phi \, , \, \xi'(x) \, / \, \Sigma_{\{k=0\}}^{\wedge} \infty \, a_k \, x^{\wedge} k = 0(x)$	[ZETA]
INV. 238	symb	$0(N^{\wedge}(-1)) = 1/0(x) \, , \, \Box = (8 \pi G/c^3) \, T_{\mu \nu}$	[SR, MANIFOLD]
INV. 239	symb	$\partial^2 f(\Box \phi) = -2 \Box \, \iiint \, V/S^2 \, dm$	[MANIFOLD]
INV. 240	symb	$Et+ = e^{\wedge}(-2\pi T \phi)[\eta_{\mu \nu} + \bar{h}_{\mu \nu}] \, dx^{\wedge} \mu \, dx^{\wedge} \nu + T^2 \, d^2 \phi$	[MANIFOLD]

4. 全方程式レジストリ (2111 本) p.23/33

INV.241

symb

$S^{\text{mn}}_1\otimes S^{\text{mn}}_2=D^2\phi\otimes h_{\mu\nu}\,,\,S^{\text{}}_{\mu\nu\text{m}}\otimes S^{\text{}}_{\mu\nu\text{n}}=\int\left[D^2\phi\right]dm=\nabla_{\text{i}}\nabla_{\text{j}}\int f(x)\,dm$

[MANIFOLD]

INV.242

symb

$h=p/(mv)\,\,(\text{sic};\,\lambda=h/p)\,,\,h\lambda=f\,,\,f\lambda=h\nu$

[QUANTUM]

INV.243

symb

$||ds^2||=S^{\text{}}_{\mu\nu\text{m}}\otimes S^{\text{}}_{\mu\nu\text{n}}\,,\,E_{+-}=f^{\text{}}(-1)(x)\times f(x)$

[MANIFOLD]

INV.244

symb

$Rt\subset C_{+-}\,,\,\nabla R_{+}\rightarrow\otimes Q_{+-}\,,\,\nabla_{\text{i}}\nabla_{\text{j}}R_{+-}=\otimes\nabla Q_{+-}$

[MANIFOLD]

INV.245

symb

$M1=R+3\,\,/\,\,E_{+-}-\{\phi\}\,,\,\,M3\cong M2\,,\,\,M3\cong M1$

[MANIFOLD]

INV.246

symb

$H3(\Pi)=Z1\otimes Z1\,,\,\,H3(M1)=0$

[MANIFOLD]

INV.247

symb

$ds^2=e^{\text{}}(-2\pi T|\phi|)[\eta_{\mu\nu}+\bar{h}_{\mu\nu}(x)]\,dx^{\text{}}_{\mu}\,dx^{\text{}}_{\nu}+T^2\,d^2\phi$

[MANIFOLD]

INV.248

symb

$\nabla\phi^2=8\pi G(\,\,p/c^3+V/S\,\,)$

[SR]

INV.249

symb

$(\partial\,\gamma_{\text{n}}+m^2)\cdot\phi=\int\left[D^2\phi\otimes h_{\mu\nu}\right]dm=0$

[SR, QUANTUM]

INV.250

symb

$\square=\pi(\chi,x)\otimes h_{\mu\nu}=D^2\phi\otimes h_{\mu\nu}$

[MANIFOLD]

INV.251

symb

$\int\left[D^2\phi\right]dm=\pi(M1)\,,\,\,H_{\text{n}}(m1)=D^2\phi-\pi(\chi,x)=\ker f\,\,/\,\,\text{im}\,f$

[MANIFOLD]

INV.252

symb

$\int D\,q\,\exp[L(x)]\,d\phi+O(N^{\text{}}1)=\pi(\chi,x)\otimes h_{\mu\nu}=D^2\phi\otimes h_{\mu\nu}$

[QUANTUM, ENTROPY]

INV.253

symb

$\lim_{\{x\rightarrow 1\}}\,\,\Sigma_{\{k=0\}}^{\text{}}\infty\,\,\zeta(x)/(a_k\,f_k)=\int\,||\left[D^2\phi\otimes h_{\mu\nu}\right]||\,dm$

[ZETA]

INV.254

symb

$\nabla\phi^2=\square\,\,\text{fff}\,\,V/S^2\,dm$

[MANIFOLD]

INV.255

symb

$dx<\partial x<\nabla\phi<\square v\,;\,\otimes<\Sigma<\otimes<\int<\wedge$

[OTHER]

INV.256

symb

$(\delta\phi(x))^2=\text{fff}\,V(x)/S^2\,dm\,,\,\,\delta\phi(x)=(\,\,\text{fff}\,V(x)/S^2\,dm\,\,)^{\text{}}(1/2)$

[MANIFOLD]

INV.257

symb

$\nabla\phi^2=-4R\int\,\,\delta(V\cdot S^{\text{}}(-3))\,dm$

[MANIFOLD]

INV.258

symb

$\nabla\phi=2R\,\,\zeta(s)\,\,i\,\,\Sigma_{\{k=0\}}^{\text{}}\infty\,\,a_k\,x^{\text{}}k$

[ZETA]

INV.259

symb

$m\,d\,x\,f_k(x)=m\,\,/\,\,(n!\,f_n(x))=(\zeta(s))^{\text{}}k\,\,/\,\,df\,m(x)\,,\,\,(\delta(x))^{\text{}}(1/2)=(x\log x\,\,/\,\,x^{\text{}}n)\,n$

[ZETA]

INV.260

symb

$0(x)=\int[D^2\phi\otimes h_{\mu\nu}]\,dm\,\,/\,\,e^{\text{}}(x\log x)\,;\,\,0(x)=V(x)\,\,\,/\,\,\int[D^2\phi\otimes h_{\mu\nu}]\,dm$

[MANIFOLD]

INV.261

symb

$M3=e^{\text{}}(x\log x)\,,\,\,x^{\text{}}(1/2+iy)=e^{\text{}}(x\log x)\,,\,\,0(x)=M3\,\,/\,\,e^{\text{}}(x\log x)=n\,E_x$

[ZETA, QUANTUM]

INV.262

differs

$V/S^2=p/c^3\,,\,\,h\nu\neq p/(mv)\,,\,\,E1=h\nu\,,\,\,E2=mc^2\,,\,\,E1\equiv E2\,\,\,\,\,=4.13567\,\text{vs}\,\,510999$

[SR, QUANTUM]

INV.263

calc

$l=\sqrt{\hbar G/c^3}\,\,\,\,\,\,=1.61622e-35$

[SR, QUANTUM]

INV.264

symb

$\text{ff}_{\text{1}}\,(y\log y)^{\text{}}(1/2)\,dy^{\text{}}m\,\,/\,\,\text{ff}_{\text{1}}\,(x\log x)^2\,dx^{\text{}}m=1/i$

[ENTROPY]

INV.265

symb

$i\,\,h\,c=G\,,\,\,h\,c=G/i\,,\,\,(1/2)/(1/(2i))=\sqrt[3]{1}\sqrt[3]{2}\leq 1$

[QUANTUM]

INV.266

symb

$A=BQ+R$

[OTHER]

INV.267

symb

$ds^2=g_{\mu\nu}\,dx^{\text{}}_{\mu}\,dx^{\text{}}_{\nu}+\kappa^2\,(A_{\mu\nu})^2$

[MANIFOLD]

INV.268

differs

$\Gamma(x)=\int\,e^{\text{}}(-x)\,x^{\text{}}(1-t)\,dx=\delta(x)\,\,\pi(x)\,f_{\text{n}}(x)\,\,\,\,\,=1.77245\,\text{vs}\,\,0.886019$

[GAMMA]

INV.269

symb

$ds^2=[T^2\,d^2\phi]\,,\,\,0(x)=[x]$

[MANIFOLD]

INV.270

symb

$\nabla\phi^2=8\pi G(\,\,p/c^3\,\,\circ\,\,V/S\,\,)$

[SR]

INV.271

symb

$p\,V\,\,/\,\,S=h$

[QUANTUM]

INV.272

differs

$\beta(p,q)=\Gamma(p)\Gamma(q)/\Gamma(p+q)\cong\Gamma(p+q)/(\Gamma(p)\Gamma(q))\,\,\,\,\,=0.0833333\,\text{vs}\,\,12$

[GAMMA, BETA]

INV.273

symb

$\ker f\,\,/\,\,\text{im}\,f\cong\text{im}\,f\,\,/\,\,\ker f$

[MANIFOLD]

INV.274

symb

$\otimes\nabla g(x)=[\,\,\nabla_{\text{i}}\nabla_{\text{j}}\int\,\,\nabla f(x)\,d\eta\,\,],\,\,\bigcup_{\{k=0\}}^{\text{}}\infty\,(\,\,\otimes\nabla f(x)\,\,)=\square\,\,\text{fff}\,\,\nabla g(x)\,d\eta$

[MANIFOLD]

INV.275

calc

$a^{\text{'}}=\sqrt{(v\,\,/\,\,(1-(v/c)^2))}\,,\,\,F=m\,a^{\text{'}}\,\,\,\,\,=0.968246$

[SR]

INV.276

symb

$\nabla f(x)=\int_{\text{M}}\square\otimes\nabla f(x)^{\text{}}n\,dm$

[MANIFOLD]

INV.277

symb

$\square=2(T-t)\,|\,R_{\text{ij}}+\nabla\nabla f-(1/(2(T-t)))\,g^2_{\text{ij}}$

[MANIFOLD]

INV.278

symb

$(\square+m)\cdot\phi=0$

[SR, QUANTUM]

INV.279

symb

$\square\times\square=(\square+m^2)\cdot\phi\,,\,\,(\partial\,\gamma_{\text{n}}+\delta\phi)\cdot\phi=0$

[SR, QUANTUM]

INV.280

symb

$\nabla_{\text{i}}\nabla_{\text{j}}\,\,\text{ff}\,\,\otimes\nabla f(t)\,dt=\square\,(\,\,\bigcup_{\{k=0\}}^{\text{}}\infty\otimes\nabla g(x)\,d\eta\,\,)$

[MANIFOLD]

INV.281

symb

$\int\otimes(l\times l)\,dm=\Sigma\,l\otimes l\,d\eta$

[OTHER]

INV.282

symb

$=[D^2\phi\otimes h_{\mu\nu}]^{\text{}}(1/2+iy)=H3(M1)$

[ZETA, MANIFOLD]

INV.283

symb

$y=\nabla_{\text{i}}\nabla_{\text{j}}\int\,\,\nabla g(x)\,dx\,,\,\,y^{\text{'}}-y+(d/dx)\,z+[n]=0$

[OTHER]

INV.284

holds

$z=\cos x+i\sin x=e^{\text{}}(i\theta)\,\,\,\,\,=0.764842+0.644218i\,\text{vs}\,\,0.764842+0.644218i$

[ROT]

INV.285

symb

$T_{\mu\nu}=-2R_{\text{ij}}\,,\,\,(d/dt)\,g_{\text{ij}}(t)=-2R_{\text{ij}}$

[MANIFOLD]

INV.286

symb

$G_{\mu\nu}=R_{\mu\nu}\,T_{\mu\nu}\,,\,\,\sigma(x)\otimes\delta(x)=(E_{\text{n}}\times H_{\text{m}})$

[MANIFOLD]

INV.287

symb

$G_{\mu\nu}=[\,\,\partial R_{\text{ij}}/\partial f\,\,]^2\,,\,\,\delta(x)\cdot V(x)=\lim_{\{n\rightarrow 1\}}\,\,\delta(x)$

[MANIFOLD]

INV.288

symb

$\lim_{\{n\rightarrow\infty\}}\,\,\text{mesh}\,V(x)=m/(m+1)\,,\,\,V(x)=\sigma\cdot S^2(x)\,,\,\,\text{fff}\,V(x)/S^2(x)\,dm=[D^2\phi\otimes h_{\mu\nu}]$

[MANIFOLD]

INV.289

symb

$g(x)|\,\delta(x,y)=(d/dt)\,g_{\text{ij}}(t)\,,\,\,\sigma(x,y)\cdot g(x)|\,\delta(x,y)=R_{\text{ij}}|\,\sigma(x,y)=\int\,R^{\text{}}\{a(x-y)^{\text{}}n+r^{\text{}}n\}_{\text{ij}}\,(ux+vy+wz)/\Gamma=\int\,R^{\text{}}\{(x-u)(y-v)(z-w)\}_{\text{ij}}\,dV$

[GAMMA, MANIFOLD]

INV.290

symb

$(\square+m)\cdot\phi=0\,,\,\,E=mc^2\,,\,\,(\partial/\partial f)\,\square\phi=4\pi G\rho$

[SR, MANIFOLD]

INV.291

differs

$(\partial\,\gamma_{\text{n}}+m)\cdot\phi=0\,,\,\,E=mc^2-(1/2)\,m\,v^2=(\,-(1/2)(v/c)^2+m\,\,)\cdot c^2=(\,-(1/2)\,a^2+m\,\,)\cdot c^2\,\,\,\,\,=1.64\,\text{vs}\,\,1.82$

[SR]

INV.292

symb

$F=m\,a\,,\,\,\int\,a\,dx=(1/2)\,a^2+C\,,\,\,T_{\mu\nu}=-(1/2)\,a^2\,,\,\,(e^{\text{}}(i\theta))^{\text{'}}=i\,e^{\text{}}(i\theta)$

[ROT, OTHER]

INV.293

symb

$ds^2=[\,g^2_{\text{}}\mu\nu\,,\,dx\,]$

[MANIFOLD]

INV.294

symb

$ds^2=g^{\text{}}(-1)_{\mu\nu}\,(\,g^2_{\text{}}\mu\nu(x)-dx\,g^2_{\text{}}\mu\nu\,)=h(x)\otimes g_{\mu\nu}\,d^2x-h(x)\otimes dx\,g_{\mu\nu}(x)\,,\,h(x)=(\,f^2(\mathfrak{X})-\vec{E}^{\text{}}+)$

[MANIFOLD]

INV.295

symb

$G_{\mu\nu}=R_{\mu\nu}\,T_{\mu\nu}\,,\,\,\partial M2=\otimes\nabla C_{+-}$

[MANIFOLD]

INV.296

holds

$r=2\,f^{\text{}}(1/2)(x)\,,\,\,f(x)=(1/4)\,||r||^2\,\,\,\,\,=3\,\text{vs}\,\,3$

[OTHER]

INV.297

symb

$E_{\text{+}}=f^{\text{}}(-1)\times f(x)\,,\,\,h(x)\otimes g(\vec{x})\cong V/S\,,\,\,R/M2=E_{\text{+}}-\phi=M3\supset R$

[MANIFOLD]

INV.298

symb

$M_{\text{+}2}=E_{\text{+}1}\cup E_{\text{+}2}\rightarrow E_{\text{+}1}\otimes E_{\text{+}2}=M1\otimes\nabla C_{+-}\,,\,\,(E_{\text{+}1}\otimes E_{\text{+}2})\cdot(R\subset C_{\text{+}})$

[MANIFOLD]

INV.299

symb

$R/M2=E_{\text{+}}-\{\phi\}=M3\supset R\,,\,\,M_{\text{+}3}\cong h(x)\cdot R_{\text{+}3}=\otimes\nabla C_{+-}\,,\,\,R=E_{\text{+}}\otimes M2-(E_{\text{+}}\cap M2)$

[MANIFOLD]

INV.300

symb

$E_{\text{+}}=g_{\mu\nu}\,dx\,g_{\mu\nu}\,,\,\,M2=g_{\mu\nu}\,d^2x\,,\,\,F=\rho\,g\,l\rightarrow V/S$

[MANIFOLD]

INV.301

calc

$0(x)=\delta(x)[\,f(x)+g(\vec{x})\,]+\rho\,g\,l\,,\,\,F=(1/2)\,m\,v^2-(1/2)\,k\,x^2\,,\,\,M2=P^2n\,\,\,\,\,=7$

[OTHER]

INV.302

symb

$V=R_{\text{+}}\,\Sigma\,K^{\text{}}m\,,\,\,W=C_{\text{+}}\,\Sigma_{\{k=0\}}^{\text{}}\infty\,K^{\text{}}(n2)\,,\,\,V/W=R_{\text{+}}\,\Sigma\,K^{\text{}}m\,\,/\,\,C_{\text{+}}\,\Sigma\,K^{\text{}}(n2)=R_{\text{+}}/C_{\text{+}}\,\Sigma\,x^{\text{}}k/(a_k\,f_k(x))=M_{\text{+}}-$

[MANIFOLD]

INV.303

symb

$(d/df)\,F=m(x)\rightarrow M_{\text{+}}-\,,\,\,\Sigma_{\{k=0\}}^{\text{}}\infty\,x^{\text{}}k/(a_k\,f_k(x))=a_k\,x^{\text{}}k\,\,/\,\,\zeta(x)$

[ZETA]

INV.304

symb

$\{f,g\}/[f,g]=(fg+gf)/(fg-gf)\,,\,\,\nabla f=2\,,\,\,\partial H3=2\,,\,\,(1+f)/(1-f)=1\,,\,\,(d/df)F=\otimes\nabla C_{+-}\,,\,\,\vec{F}=1/2$

[QUANTUM]

INV.305

symb

$H1\cong H3=M3\,;\,H3\rightarrow H1\rightarrow\pi(\chi,x)\,,\,\,H_{\text{n}}\,,\,H_{\text{m}}=\text{rank}(\mathfrak{m},n)\,,\,\,\lim\,\text{mesh}(\text{rank}(\mathfrak{m},n))\rightarrow 0$

[MANIFOLD]

INV.306

symb

$(f\mathfrak{g})^{\text{'}}=f\,g^{\text{'}}+g\,f^{\text{'}}\,,\,\,(f/g)^{\text{'}}=(f'g-g'f)/g^2\,,\,\,\{f,g\}/[f,g]=(fg)^{\text{'}}\otimes d_x\,fg$

[OTHER]

INV.307

symb

$(f/g)^{\text{'}}\otimes g^{\text{}}(-2)\,d_x\,fg=(fg)^{\text{'}}\otimes d_x\,fg\,\,/\,\,((f/g)^{\text{'}}\otimes g^{\text{}}(-2)\,d_x\,fg)=(d/df)\,F$

[OTHER]

4. 全方程式レジストリ (2111 本) p.24/33

INV.308	symb	$\hbar\phi=(1/i)\,H\Psi\,,\;i[H,\,\phi]=-H\Psi\,,\;(\{f,g\}/[f,g])'=(i)^2$	[QUANTUM]
INV.309	symb	$[\nabla_{\cdot i}\nabla_{\cdot j}\,f(x),\,\delta(x)]=\nabla_{\cdot i}\nabla_{\cdot j}\,\int f(x,y)\,dm_{xy}\,,\;f(x,y)=[f(x),\,h(x)]\times[g(x),\,h^{(-1)}(x)]$	[MANIFOLD, QUANTUM]
INV.310	symb	$\delta(x)=1/f'(x)\,,\;[H,\,\phi]=\Delta f(x)\,,\;0(x)=\nabla_{\cdot i}\nabla_{\cdot j}\,\int\delta(x)\,f(x)\,dx\,,\;0(x)=\int\delta(x)\,f(x)\,dx$	[QUANTUM]
INV.311	symb	$R+\cap E+\supset x\,,\;M\times R+\supset M3\,,\;Q\supset C+-\,,\;Z\in Q\nabla f\,,\;f\equiv\oplus_{\{k=0\}}^n\nabla C+-$	[MANIFOLD]
INV.312	symb	$\oplus_{\{k=0\}}^{\infty}\nabla C+-=M1\,,\;\oplus_{\{k=0\}}^{\infty}\nabla M+-\equiv E+-\,,\;M3\equiv M1\,,\;\oplus_{\{k=0\}}^{\infty}\nabla V+-/S$	[MANIFOLD]
INV.313	symb	$P^2n\,/\,M2\equiv\oplus_{\{k=0\}}^{\infty}\nabla C+-\,,\;E+-\times R+-\equiv M2$	[MANIFOLD]
INV.314	symb	$\zeta(x)=P^2n\times\Sigma_{\{k=0\}}^{\infty}a_kx^k\,,\;M2\equiv P^2n\,/\,ker\,f\rightarrow\oplus\nabla C+-$	[ZETA, MANIFOLD]
INV.315	symb	$S+-\times V+-\equiv(V/S)\,\oplus_{\{k=0\}}^{\infty}\nabla C+-\,,\;V+-=M+-\otimes S+-\,,\;Q\times M1\subset\oplus\nabla C+-$	[MANIFOLD]
INV.316	symb	$\Sigma_{\{k=0\}}^{\infty}Z\otimes Q+-=\oplus_{\{k=0\}}^{\infty}\nabla M1=\oplus_{\{k=0\}}^{\infty}\nabla C+-\times\Sigma_{\{k=0\}}^{\infty}M1\,,\;x\in R+\times C+-\supset M1\,,\;M1\subset M2\subset M3$	[MANIFOLD]
INV.317	symb	Thurston geometries: $S^3,\,H^1\times E^1,\,E^1,\,S^1\times E^1,\,S^2\times E^1,\,H^1\times S^1,\,H^1,\,S^2\times E\,(X2)$	[MANIFOLD]
INV.318	symb	$\otimes\nabla C+-\equiv M3\,,\;R\supset Q\,,\;R\cap Q\,,\;R\subset M3\,,\;C+\otimes M_{\cdot n},\,E+\cap R+\,,\,E2\otimes E1\,,\;R-\subset C+\,,\;M+-,C+-\,,\;M+-\nabla C+-\,,\;C+\nabla H_{\cdot m},\,E+\nabla R+-\,,\,E2\nabla E1\,,\;R-\nabla C+-\,(X3)$	[MANIFOLD]
INV.319	symb	$\nabla/\Delta\,\int x\,f(x)\,dx\,,\;\nabla R\,/\,\Delta f\,,\;\square=2\,\int(R+\nabla_{\cdot i}\nabla_{\cdot j}\,f)^2\,/\,-(R+\Delta f)\,e^{(-f)}\,dV$	[MANIFOLD, ENTROPY]
INV.320	symb	$\square=\nabla R/\Delta f\,,\;(d/dt)\,g_{ij}=\square\rightarrow\nabla f/\Delta x\,,\;(R+ \nabla f ^2)\,dm\rightarrow-2\,(R+\nabla_{\cdot i}\nabla_{\cdot j}\,f)^2\,e^{(-f)}\,dV$	[MANIFOLD, ENTROPY]
INV.321	symb	$x^{\wedge}n+y^{\wedge}n=z^{\wedge}n\rightarrow\nabla\phi^2=8\pi G\,T_{\cdot\mu\nu}\,,\;f(x+y)\geq f(x)\circ f(y)$	[SR, MANIFOLD]
INV.322	symb	$im\,f\,/\,ker\,f=\partial f\,,\;ker\,f=\partial f\,,\;ker\,f\,/\,im\,f\equiv\partial f\,,\;ker\,f=f^{(-1)}(x)\,x\,f(x)$	[MANIFOLD]
INV.323	holds	$f^{(-1)}(x)\,x\,f(x)=\int\partial f(x)\,d(ker\,f)\rightarrow\nabla f=2\,;\,nC_r=nC(n-r)\rightarrow im\,f/ker\,f\equiv ker\,f/im\,f\quad=35\,vs\,35$	[OTHER]
INV.324	symb	$\Sigma_{\{k=0\}}^{\infty}a_kf_k=T^2\,d^2\phi\,,\;a_k\equiv\Sigma_{\{r=0\}}^{\infty}nC_r$	[MANIFOLD]
INV.325	symb	$V/W=R/C\,\Sigma_{\{k=0\}}^{\infty}x^k/(a_kf_k)\,,\;W/V=C/R\,\Sigma_{\{k=0\}}^{\infty}a_kf_k\,/\,x^k\,,\;V/W\equiv W/V\equiv R/C\,(\Sigma_{\{r=0\}}^{\infty}nC_r)^{(-1)}\,\Sigma_{\{k=0\}}^{\infty}x^k$	[OTHER]
INV.326	symb	$W/V=x\,F(x)\,,\;\chi(x)=(-1)^k a_k\,,\;\Gamma(x)=\int e^{(-x)}\,x^{(1-t)}\,dx$	[GAMMA, MANIFOLD]
INV.327	symb	$\Sigma_{\{k=0\}}^n a_kf_k=(f_k)'\,,\;\Sigma_{\{k=0\}}^n a_kf_k=\Sigma_k nC_r f_k=(f_k)'\,,\;\Sigma_{\{k=0\}}^{\infty} a_kf_k=[f(x)]\,,\;\Sigma a_kf_k=\alpha$	[OTHER]
INV.328	symb	$\Sigma_{\{k=0\}}^{\infty}1/(a_kf_k)\,,\;\Sigma_{\{k=0\}}^{\infty}(a_kf_k)^{(-1)}=1/(1-z)$	[OTHER]
INV.329	differs	$\mathcal{J}\mathcal{J}_1\,(x\log x)(y\log y)\,dxy=nC_rxy\,/\,(nC(n-r)\,(x\log x)(y\log y))^{(-1)}=(nC(n-r))^2\quad=0.0625002\,vs\,4$	[ENTROPY]
INV.330	symb	$\Sigma_{\{k=0\}}^{\infty}(\,1/(x\log x)-1/(y\log y)\,)\,d(1/(nxy))\times xy=\Sigma_{\{k=0\}}^{\infty}a_kf_k=\alpha$	[ENTROPY]
INV.331	symb	$Z\supset C\otimes\nabla R+\,,\;\nabla(R+\cap E+)\supset x\,,\;\Delta(C\subset R)\supset x\,,\;M+-\otimes R+\,,\;E+\in\otimes\nabla R+\,,\;S+-\subset R+_2\,,\;V+-\times R+-\equiv V/S$	[MANIFOLD]
INV.332	symb	$C\cup V+-\supset M1\otimes\nabla C+-\,,\;Q\,\overline{\mathbb{R}}\,R+-\,,\;Q\subset\otimes M+-\,,\;\otimes Q\subset\zeta(x)\,,\;\otimes\nabla C+-\equiv M3$	[ZETA, MANIFOLD]
INV.333	symb	$[-\Delta v+\nabla_{\cdot i}\nabla_{\cdot j}\,v_{\cdot ij}-R_{ij}\,v_{\cdot ij}-v_{\cdot ij}\,\nabla_{\cdot i}\nabla_{\cdot j}+2<\nabla f,\,\nabla h>+(R+\nabla f^2)(v/2-h)\,]$	[MANIFOLD, ENTROPY]
INV.334	symb	$x^{(1/2+iy)}=[f(x)\circ g(x),\,h(x)]\,/\,\partial f\partial g\partial h$	[ZETA]
INV.335	symb	$x^{(1/2+iy)}=\exp[\,\int\,\nabla_{\cdot i}\nabla_{\cdot j}\,f(g(x))\,g'(x)\,/\,\partial f\partial g\,]$	[ZETA]
INV.336	symb	$0(x)=\{\,[f(x)\circ g(x),\,\bar{h}(x)],\,g^{(-1)}(x)\,\}$	[QUANTUM]
INV.337	symb	$\exists[\,\nabla_{\cdot i}\nabla_{\cdot j}(R+\Delta f),\,g(x)\,]=\oplus_{\{k=0\}}^{\infty}\nabla\,\int\,\nabla_{\cdot i}\nabla_{\cdot j}\,f(x)\,dm\,,\;\vee(\nabla_{\cdot i}\nabla_{\cdot j}\,f)=\otimes\nabla E+$	[MANIFOLD]
INV.338	symb	$g(x,y)=0(x)[\,f(x)+\bar{h}(x)\,]+T^2\,d^2\phi$	[MANIFOLD]
INV.339	symb	$0(x)=\int\,[g(x)]\,e^{(-f)}\,dV'-\Sigma\,\delta(x)$	[MANIFOLD]
INV.340	symb	$0(x)=[\nabla_{\cdot i}\nabla_{\cdot j}\,f(x)]'\equiv nC_r\,f(x)^n\,f(y)^{(n-r)}\,\delta(x,y)\,,\;V(\varepsilon)=\int[f(x)]\,dm\,/\,\partial f_{\cdot xy}$	[MANIFOLD]
INV.341	symb	$\square\phi=8\pi G\,T_{\cdot\mu\nu}\,,\;(\square\phi)'=\nabla_{\cdot i}\nabla_{\cdot j}\,(\delta(x)\circ G(x))_{\cdot\mu\nu}\,(p/c^2\circ V/S)\,,\;x^{(1/2+iy)}=e^{(x\log x)}$	[SR, ZETA, MANIFOLD]
INV.342	symb	$\delta(x)\,\phi=\vee[\,\nabla_{\cdot i}\nabla_{\cdot j}\,f\circ g(x)\,]\,,\;\exists(R+\Delta f)$	[MANIFOLD]
INV.343	symb	$-nC_r=(1/i)\,H\phi\,C\,\hbar\phi+[H,\,\phi]\,C^{(-n-r)}\,;\,nC_r=nC(n-r)$	[QUANTUM]
INV.344	symb	$\mathcal{J}\mathcal{J}_1\,(x\log x)^2\,dx^m\rightarrow 0(x)=[\nabla_{\cdot i}\nabla_{\cdot j}\,f]'\,/\,\partial f_{\cdot xy}$	[ENTROPY]
INV.345	symb	$\bigcup_{\{x=0\}}^{\infty}f(x)=\nabla_{\cdot i}\nabla_{\cdot j}\,f(x)\otimes\Sigma\,f(x)=\otimes\nabla f(x)\,;\,\nabla_{\cdot i}\nabla_{\cdot j}\,f\equiv\partial x\partial y\,\int\,\nabla_{\cdot i}\nabla_{\cdot j}\,f\,dm\equiv\int\,[f(x)]\,dm\equiv\{[f(x),\,g(x)],\,g^{(-1)}(x)\}\equiv\square\phi\equiv\nabla\phi^2\equiv f(x\circ y)\leq f(x)\circ g(x)\equiv f(x) + g(x) $	[MANIFOLD]
INV.346	symb	$\delta(x)\,\phi=<f,\,g>\circ[h^{(-1)}(x)]\,,\;\partial f_{\cdot x}\cdot\delta(x)\,\phi=x\,,\;x\in 0(x)\,,\;0(x)=\{\,[f\circ g,\,h^{(-1)}(x)],\,g(x)\,\}$	[QUANTUM]
INV.347	symb	$\lim_{\{n\rightarrow\infty\}}\Sigma_{\{k=n\}}^{\infty}\nabla f=[\,\nabla\,\int\,\nabla_{\cdot i}\nabla_{\cdot j}\,f(x)\,dx^m,\,g^{(-1)}(x)\,]\rightarrow\oplus_{\{k=0\}}^{\infty}\nabla E+-=M3=\oplus_{\{k=0\}}^{\infty}E+-$	[MANIFOLD]
INV.348	symb	$dx^2=[g^2\,_{\mu\nu}\,,\,dx]\,,\;g^{(-1)}=dx\,\int\,\delta(x)\,f(x)\,dx\,,\;f(x)=\exp[\,\nabla_{\cdot i}\nabla_{\cdot j}\,f(x),\,g^{(-1)}(x)\,]$	[MANIFOLD]
INV.349	symb	$(\,g(x)/f(x)\,)'=\lim_{\{n\rightarrow\infty\}}g(x)/f(x)=g'(x)/f'(x)$	[OTHER]
INV.350	symb	$\nabla F=f\cdot(1/4) r ^2\,,\;\nabla_{\cdot i}\nabla_{\cdot j}\,f=(d/dx_{\cdot i})(d/dx_{\cdot j})\,f(x)\,g(x)$	[MANIFOLD]
INV.351	symb	$\mathcal{J}\mathcal{J}_1\,(x\log x)^2\,dx^m=[\,\nabla_{\cdot i}\nabla_{\cdot j}\,\int\,\nabla f(x)\,d\eta\,]\times U(r)$	[ENTROPY]
INV.352	symb	$S^{\sim mn}_1\otimes S^{\sim mn}_2=\int\,[D^2\phi\otimes h_{\cdot\mu\nu}]\,dm$	[MANIFOLD]
INV.353	symb	$U(r)=(1/2)\,\sqrt[4]{(1+f'(r))}\,/\,f(r)+m\,g\,r\,,\;F_{\cdot m}^{\sim}t=(1/4) r ^2$	[OTHER]
INV.354	symb	$\mathcal{J}\mathcal{J}_1\,(y\log y)^{(1/2)}\,dy^m=[\,\nabla_{\cdot i}\nabla_{\cdot j}\,\int\,\nabla f(x)\,d\eta\,]\times E+-\,,\;E+-=\exp[L(x)]\,dm\,d\phi+0(N^{-})$	[QUANTUM, ENTROPY]
INV.355	calc	$(d/df)\,F=[\,\nabla_{\cdot i}\nabla_{\cdot j}\,\int\,\nabla f(x)\,d\eta\,]\,(\,U(r)+E+-\,)=(1/2)\,m\,v^2+m\,c^2\quad=8.99255e+16$	[SR]
INV.356	symb	$\nabla_{\cdot i}\nabla_{\cdot j}\,\int\,\nabla f(x)\,d\eta\,/\,\nabla_{\cdot i}\nabla_{\cdot j}\,\otimes M3=M3/P^2n\equiv P^2n/M3\equiv M1=[M1]$	[MANIFOLD]
INV.357	symb	$[f(x)]=\Sigma_{\{k=0\}}^{\infty}a_kf_k\,,\;\lim_{\{n\rightarrow1\}}[f(x)]=\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}}^{\infty}a_kf_k=\alpha\,,\;e^{(i\theta)}=\cos\theta+i\sin\theta\,,\;H3(M1)=0$	[ROT, ZETA]
INV.358	symb	$x^2\cos\theta\,/\,a+y^2\sin\theta\,/\,b=r^2$	[ROT]
INV.359	differs	$[[\cos\theta,\,-\sin\theta],[\sin\theta,\,\cos\theta]]\,[x;\,y]=[[1,0],[0,-1]]\quad=1\,vs\,-1$	[ROT, QUANTUM]
INV.360	differs	$\chi(x)=\Sigma_{\{k=0\}}^{\infty}(-1)^n\,r^{\sim}n\,,\;(d/df)F=(d/df)\,\Sigma\Sigma a_kf_k= a_1a_2\dots a_n - a_1\dots a_{n-1} - a_n\dots a_1 \,,\,\lim_{\{n\rightarrow0\}}\chi(x)=2\quad=0.666667\,vs\,2$	[MANIFOLD]
INV.361	symb	$\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}}^{\infty}a_kf_k=nC_r\,f(x)^n\,f(y)^{(n-r)}\,\delta(x,y)\,;\,\lim_{\{n\rightarrow\infty\}}nC_r\,f(x)^n\,f(y)^{(n-r)}\,\delta(x,y)$	[OTHER]
INV.362	holds	$\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}}^{\infty}(\,1/(n+1)\,)^{\sim}s=\lim_{\{n\rightarrow1\}}Z^{\sim}r=1/z\quad=1.64493\,vs\,1.64493$	[ZETA]
INV.363	differs	$\beta(p,q)=\Gamma(p)\Gamma(q)/\Gamma(p+q)\equiv\Gamma(p+q)/(\Gamma(p)\Gamma(q))\,,\;\lim_{\{n\rightarrow1\}}a_kf_k\equiv\lim_{\{n\rightarrow\infty\}}\zeta(s)/(a_kf_k)\,,\;ker\,f/im\,f\equiv im\,f/ker\,f\quad=0.0833333\,vs\,12$	[GAMMA, ZETA, BETA]
INV.364	differs	$\lim_{\{n\rightarrow1\}}\zeta(s)=0\,,\;0(x)=\zeta(s)\,\Sigma_{\{x=0\}}^{\infty}f(x)\rightarrow\oplus_{\{k=0\}}^{\infty}\nabla f(x)=\int_{\cdot M}\delta(x)\,f(x)\,dx\quad=1000.58\,vs\,0$	[ZETA]
INV.365	symb	$\mathcal{J}\mathcal{J}\mathcal{J}_{\cdot M}\,V/S^2\,e^{(-f)}\,dV=\mathcal{J}\mathcal{J}_{\cdot D}\,-(f(x,y)^2,\,g(x,y)^2)\,-\mathcal{J}\mathcal{J}_{\cdot D}\,(\,g(x,y)^2,\,f(x,y)^2\,)$	[MANIFOLD]
INV.366	symb	$\lim_{\{n\rightarrow1\}}\Sigma_{\{k=0\}}^{\infty}a_kf_k=\int\,[D^2\phi\otimes h_{\cdot\mu\nu}]\,dm=\int\,\exp[L(x)]\,d\phi\,dm\times E+-=S^{\sim mn}_1\otimes S^{\sim mn}_1=Z1\otimes Z1=M1$	[ZETA, MANIFOLD, QUANTUM]
INV.367	symb	$H^{\sim mn}_n(\chi,\,h)=\mathcal{J}\mathcal{J}_{\cdot M}\,V\,(R+\Delta f)\,e^{(-f)}\,dV\,,\;V/S^2=\mathcal{J}\mathcal{J}\mathcal{J}_{\cdot M}\,[D\phi\otimes h_{\cdot\mu\nu}]\,dm$	[MANIFOLD, ENTROPY]
INV.368	symb	$\mathcal{J}\mathcal{J}\mathcal{J}_{\cdot M}\,V/S^2\,dm=\int_{\cdot D}\,(l\times l)\,dm$	[MANIFOLD]
INV.369	symb	$\mathcal{J}\mathcal{J}_{\cdot D}\,-g(x,y)^2\,dm-\mathcal{J}\mathcal{J}_{\cdot D}\,-f(x,y)^2\,dm=-2R_{\cdot ij}\,,\,[f(x),\,g(x)]\times[h(x),\,g^{(-1)}(x)]$	[MANIFOLD]

INV.370	symb	$[[D_m, d_x],[d_x, \partial_m]] [[\cos \theta, -\sin \theta],[\sin \theta, \cos \theta]] [x; y] = [[1,0],[0,-1]]^{(1/2)} = (D_m, d_x) \cdot (\cos \theta, \sin \theta) \times (d_x, \partial_m) \cdot (\cos \theta, \sin \theta) (\times 2)$	[ROT, QUANTUM]
INV.371	symb	$[[1,0],[0,i]] \beta(x,\theta) [x; y] = [[i,0],[0,-1]]$	[BETA, QUANTUM]
INV.372	calc	$l = \sqrt{(\hbar G/c^2)} \quad , \quad \sigma_m \cdot [[\delta(x), -1],[1, \varepsilon(x)]]^{(1/2)} = [[i,0],[0,-i]] \quad = 1.61622e-35$	[QUANTUM]
INV.373	symb	$(\partial/\partial \tau f(x,y,z))^{(3')} = A_{\mu \nu} \quad , \quad (d/dt) g_{ij}(t) = -2R_{ij}$	[MANIFOLD]
INV.374	symb	$\lim_{\{n \rightarrow 1\}} \sum_{\{k=0\}}^{\infty} a_k f_k = \lim_{\{n \rightarrow 1\}} a_n / a_{\{n-1\}} \equiv \alpha \quad ; \quad \lim_{\{n \rightarrow 1\}} \sum_{\{k=0\}}^{\infty} a_k/a_{\{k+1\}} = \sum_{\{k=0\}}^{\infty} a_k f_k$	[OTHER]
INV.375	calc	$\square = (8\pi G/c^4) T_{\mu \nu} \quad = 2.07655e-43$	[SR, MANIFOLD]
INV.376	symb	$\int x \log x \, dx = \iint_M (e^{(x \log x) \sin \theta} d\theta) \, dx \, d\theta = (\log \sin \theta \, d\theta)^{(1/2)} - (\delta(x) \cdot \varepsilon(x))^{(1/2)}$	[ROT, ENTROPY]
INV.377	symb	$T_{\mu \nu} = \iint_M [\nabla_i \nabla_j e^{(\int x \log x \, dx) + 0(N^{(-1)})}] \, dm \, d\phi \, $	[MANIFOLD, ENTROPY]
INV.378	symb	$[[d_x, \delta(x)],[\varepsilon(x), \partial_m(x)]] (f_{mn}(x), g_{\mu \nu}(x)) = [[1,0],[0,-1]]^{(1/2)}$	[QUANTUM]
INV.379	symb	$ ds^2 = H(x)_{mn} \otimes K(y)_{mn}$	[MANIFOLD]
INV.380	symb	$ H(x) ^{(-1/2+iy)} \subset R+ \cup C+- \cong M3 \quad ; \quad H(x) ^{(1/2+iy)} \subseteq M3$	[ZETA, MANIFOLD]
INV.381	symb	$K(y) = x \, y \, z; a \, b \, c ^2 \, / \, g_{\mu \nu}(x) \equiv f(x,y,z) \, / \, (g(a,b,c) \, h^{(-1)}(u,v,w))$	[MANIFOLD]
INV.382	symb	$V = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm \quad , \quad S^2 = \int e^{(-2 \sin \theta \cos \theta)} \cdot \log \sin \theta \, dx_{\theta} + 0(N^{(-1)}) \quad , \quad 0(x) = \zeta(s) \, / \, \lim_{\{x \rightarrow 1\}} \sum a_k f_k = \alpha$	[ROT, ZETA]
INV.383	symb	$0(x) = T_{\mu \nu} \quad , \quad \lim_{\{x \rightarrow 1\}} \sum_{\{k=0\}}^{\infty} a_k f_k = T_{\mu \nu}$	[ZETA]
INV.384	symb	$G_{\mu \nu} = R_{\mu \nu} T_{\mu \nu} \quad , \quad M3 = \iiint V/S^2 \, dm \quad , \quad -(1/(2(T-t))) \mid R_{ij} = \square \phi$	[MANIFOLD]
INV.385	symb	$m(x) = [f(x)] \quad , \quad f(x) = \iint e^{(\int x \log x \, dx) + 0(N^{(-1)})} + T^2 \, d^2 \phi$	[ENTROPY]
INV.386	symb	$F(x) = \nabla_i \nabla_j \int \nabla f(x,y) \, dm \, ^{(1/2+iy)}$	[ZETA]
INV.387	symb	$G_{\mu \nu} = [\nabla_i \nabla_j \int \nabla g(x,y,z) \, dx \, dy \, dz] \, ^m / (2 \sin \theta \cos \theta \log \sin \theta \, d\theta \, d\phi)$	[ROT, MANIFOLD]
INV.388	symb	$T_{\mu \nu} = F(x) \quad , \quad 2(T-t) \mid g^2_{ij} = \iint 1 \, (x \log x)^2 \, dx^m$	[ENTROPY]
INV.389	symb	$\phi \delta(x) = [m(x)] \quad , \quad \nabla(\square \phi) = \nabla_i \nabla_j \int \nabla g(x,y) \, d\eta \quad , \quad \nabla \cdot (\square \phi) = (1/4) \, g^2_{ij} \quad , \quad \square \phi = (8\pi G/c^4) T_{\mu \nu}$	[SR, MANIFOLD]
INV.390	symb	$\rho \, V \, / \, c^2 = S \circ h_{\mu \nu} = h$	[QUANTUM]
INV.391	symb	$T_{\mu \nu} = \hbar \nu \, / \, S \quad , \quad T_{\mu \nu} = \iiint V/S^2 \, dm \quad , \quad (d/df) \, m(x) = V(x)/F(x)$	[QUANTUM]
INV.392	symb	$y = x \quad , \quad (d/df)F = m(x) \quad , \quad R_{ij} \mid g_{\mu \nu}(x) = [\nabla_i \nabla_j g(x,y)]^{(1/2+iy)}$	[ZETA]
INV.393	symb	$\nabla \circ (\square \phi) = (\partial/\partial f) F = \iint \nabla_i \nabla_j f(x) \, d\eta_{\mu \nu} \quad , \quad \int [\nabla_i \nabla_j g(x,y)] \, dm = (\partial/\partial f) R_{ij} \mid g_{\mu \nu}(x)$	[MANIFOLD]
INV.394	symb	$G(x) = \nabla_i \nabla_j f + R_{ij} \mid g_{\mu \nu}(x) + \nabla(\square \phi) + (\square \phi)^2$	[MANIFOLD]
INV.395	symb	$G_{\mu \nu} + \Lambda g_{ij} = T_{\mu \nu} \quad , \quad T_{\mu \nu} = (d/dx^{\mu})(d/dx^{\nu}) f_{\mu \nu} + -2(T-t) \mid R_{ij} + f'' + (f')^2 = \int \exp[L(x)] \, dm + 0(N^{(-1)}) = \int e^{((2/m) \sin \theta \cos \theta \cdot \log(\sin \theta))} \, dx + 0(N^{(-1)})$	[ROT, MANIFOLD, QUANTUM]
INV.396	symb	$(\partial/\partial f) F = (\nabla_i \nabla_j)^{(-1)} \circ F(x)$	[MANIFOLD]
INV.397	symb	$0(x) = \int [\nabla_i \nabla_j \int \nabla f(x) \, dm] \, d\phi = \int [\nabla_i \nabla_j f(x) \, d\eta_{\mu \nu}] \, d\phi$	[MANIFOLD]
INV.398	symb	$\nabla f = \int \nabla_i \nabla_j [\int^{\{S^{(-3)}\}} \delta(x) \, dV] \, dm$	[MANIFOLD]
INV.399	symb	$ \int [\nabla_i \nabla_j f] \, dm \, ^{(1/2+iy)} = \text{rot}(\text{div}, E, E1) = 2\langle f, h \rangle \quad ; \quad V(x)/f(x) = \rho(x)$	[ROT, ZETA]
INV.400	symb	$\int_M \rho(x) \, dx = \square \phi \quad , \quad -2\langle g, h \rangle = \text{div}(\text{rot} \, E, E1) = -2R_{ij}$	[MANIFOLD]
INV.401	symb	$0(x) = \nabla_i \nabla_j \, / \, S^2 \int [\nabla_i \nabla_j f \cdot g(x) \, dx \, dy] \, d\phi \, = \int (\delta(x))^2 \sin \theta \cos \theta \log \sin \theta \, d\theta \, d\phi$	[ROT]
INV.402	symb	$\delta \cdot 0(x) = [\nabla_i \nabla_j f \, d\eta \, \, / \, \int e^{(2 \sin \theta \cos \theta)} \cdot \log \sin \theta \, d\theta]$	[ROT]
INV.403	symb	$i\hbar \phi = \int (\partial/\partial z) [i(xy + yx) / (z - \bar{z})] \, dm \, d\phi \, $	[QUANTUM]
INV.404	symb	$\delta(x) \int_C R^2 \sin \theta \cos \theta = \Gamma(p+q)$	[ROT, GAMMA]
INV.405	symb	$(\square + m) \cdot \phi = (\nabla_i \nabla_j f \mid g_{\mu \nu}(x) + v \nabla_i \nabla_j) \int [m(x) (\text{rot} \cdot \text{div}(E, E1))] \, dm \, d\phi$	[ROT, QUANTUM]
INV.406	symb	$G_{\mu \nu} = \square \iiint (x,y,z)^3 \, dx \, dy \, dz = (8\pi G/c^4) T_{\mu \nu}$	[SR, MANIFOLD]
INV.407	symb	$(d/dV) F = \delta(x) \int \nabla_i \nabla_j f \, d\eta_{\mu \nu}$	[MANIFOLD]
INV.408	symb	$x^{\perp n} + y^{\perp n} = z^{\perp n} \quad , \quad \delta(x) \int z^{\perp n} = (d/dV) z^2 \quad , \quad (x,y) \cdot (\delta m, \partial m) = (x,y) \cdot (z^{\perp n}, f) \quad , \quad n \perp x \quad , \quad n \perp y = 0$	[OTHER]
INV.409	symb	$V(\nabla_i \nabla_j f) \cdot XOR(\square \phi) = (d/df) \int_M F \, dV$	[OTHER]
INV.410	holds	$\partial(x) \int z^3 = (d/dV) z^2 \quad , \quad \sum_{\{k=0\}}^{\infty} 1/(n+1)^s = 0(x) \quad = 1.20206 \text{ vs } 1.20206$	[ZETA]
INV.411	symb	$\int 0(x) \, dx = \delta(x) \pi(x) f(x) \quad , \quad 0(x) = \int \sigma(x)^2 \sin \theta \cos \theta \log \sin \theta \, d\theta \, d\phi$	[ROT, ZETA]
INV.412	symb	$y = x \quad , \quad 0(x) = \nabla_i \nabla_j f \, / \, S^2 \int [\nabla_i \nabla_j f \circ g(x) \, dx \, dy] \, $	[MANIFOLD]
INV.413	holds	$\lim_{\{n \rightarrow 1\}} \sum_{\{k=0\}}^{\infty} a_k/a_{\{k+1\}} = (\log \sin \theta \, dx)' = \cos \theta \, / \, \sin \theta = x/y \quad = 1.73205 \text{ vs } 1.73205$	[ROT]
INV.414	symb	$\lim_{\{n \rightarrow 1\}} \sum_{\{k=0\}}^{\infty} a_k f_k = 1/(1-z)$	[OTHER]
INV.415	symb	$\square \phi = (8\pi G/c^4) T_{\mu \nu} \quad , \quad (\partial/\partial f) \square \phi = 4\pi G \rho \quad , \quad \int \rho(x) = \square \phi \quad , \quad V(x)/f(x) = \rho(x)$	[SR, MANIFOLD]
INV.416	symb	$(\partial^{\perp}/\partial f^{(n-1)}) F = \int [D^2 \phi \otimes h_{\mu \nu}] \, dm = (P1 \, P3 \dots P_{\{2n-1\}}) \, / \, (P0 \, P2 \dots P_{\{2n+2\}})$	[ZETA]
INV.417	symb	$\otimes \nabla M1 \otimes \nabla M1 = \sigma_n(x,x) \otimes \sigma_{(n-1)}(x,x) = \{f,h\} \circ [f,h]^{(-1)} = g^{(-1)}(x)_{\mu \nu} \, d_x g_{\mu \nu}(x)$	[MANIFOLD, QUANTUM]
INV.418	symb	$\sum_{\{k=0\}}^{\infty} \nabla^{\perp n} nCr f^{\perp n}(x) \equiv \sum_{\{k=0\}}^{\infty} \nabla^{\perp n} \nabla^{(n-1)} nCr f^{\perp n}(x) g^{\perp(n-r)}(x)$	[OTHER]
INV.419	symb	$\sum_{\{k=0\}}^{\infty} (\partial^{\perp n}/\partial^{(n-1)} f) \circ \zeta'(x)/n! = \lim_{\{n \rightarrow 1\}} \sum a_k f_k \quad ; \quad (f)^{\perp n} = nCr f^{\perp n}(x) g^{\perp(n-r)}(x) \cdot \delta(x,y)$	[ZETA]
INV.420	holds	$(e^{(i \, \theta)})' = i \, e^{(i \, \theta)} \quad , \quad \int e^{(i \, \theta)} = (1/i) \, e^{(i \, \theta)} \quad , \quad i \, h \, c = G \quad , \quad h \, c = (1/i) \, G \quad = -0.644218 + 0.764842i \text{ vs } -0.644218 + 0.764842i$	[ROT]
INV.421	symb	$(\square \phi, \nabla f^2) \cdot ((8\pi G/c^4) T_{\mu \nu}, 4\pi G \rho) = (-1/2) m \, v^2 + m \, c^2, (1/2) k \, T^2 + (1/2) m \, v^2) \cdot [[\cos \theta, -\sin \theta],[\sin \theta, \cos \theta]] = [[1,0],[0,i]]$	[ROT, SR]
INV.422	symb	$(\{f,g\}/[f,g])' = i^2 \quad , \quad \nabla f^2 \, / \, \square \phi = (1/2) \iint \{f,g\}/[f,g] = 1/(2i)$	[QUANTUM]
INV.423	symb	$\iint 1 \, (y \log y)^{(1/2)} \, dy^m = (1/2) \iint 1 \, (x \log x)^2 \, dx^m = 1/(2i) \quad , \quad (d/dt) g_{ij} = -2R_{ij}$	[ENTROPY]
INV.424	symb	$(\partial/\partial x)(f(x) g(x))' = \otimes \nabla_i \nabla_j f(x) g(x) \quad ; \quad \int f'(x) g(x) \, dx = [f(x) g(x)] - \int f(x) g'(x) \, dx$	[OTHER]
INV.425	symb	$f = n \, \nu \, \lambda \quad , \quad \lambda = x/L \quad , \quad \int d n \, \nu \, \lambda = f(x) \quad , \quad x \, f(x) = F(x) \quad , \quad [f(x)] = \nu \, h \text{ (see INV.200)}$	[QUANTUM]
INV.426	symb	$p_x = m \, v \quad , \quad \lambda(x) = \text{esperial } f(x) \quad , \quad [F(x)] = 1/(1-z) \quad , \quad 0 < \tan \theta < i \quad , \quad -1 < \tan \theta < i \quad , \quad -1 \leq \sin \theta \leq 1 \quad , \quad -1 \leq \cos \theta \leq 1$	[ROT, QUANTUM]
INV.427	symb	$R \nabla E+ = f(x) \nabla e^{(x \log x)} \quad , \quad Q \nabla C+ = (d/df)F(x) \nabla \int \delta(x) f(x) \, dx \quad , \quad E+ \nabla f = e^{(x \log x)} \nabla n! f(x) \, / \, E(x) \text{ (see INV.155-158)} (\times 2)$	[ENTROPY]
INV.428	symb	$(u + v + w)(x + y + z) \, / \, \Gamma$	[GAMMA]
INV.429	symb	$\exp(\nabla(R \cap E+), \Delta(C \supset R)) = \pi(R, C \nabla E+) = \text{rot}(E1, \text{div } E2) \text{ (see INV.153)} (\times 3)$	[ROT]
INV.430	symb	$d(R \nabla E+) = \Delta f(x) \circ E+(x) \quad ; \quad \square x = \int^{\{\nabla_i \nabla_j (R+E+)\}} \nabla(R \cap E+) \, d \square x \quad ; \quad x^{\perp n} + y^{\perp n} = z^{\perp n} \rightarrow \square x = \dots$	[MANIFOLD]
INV.431	symb	$\square = -2(T-t) \mid R_{ij} + \nabla \nabla f + (1/(-2(T-t))) \, g^2_{ij}$	[MANIFOLD, ENTROPY]
INV.432	symb	$\square = -2 \int (R + \nabla_i \nabla_j f) \, / \, -(R + \Delta f) \, e^{(-f)} \, dV$	[MANIFOLD, ENTROPY]

4. 全方程式レジストリ (2111 本) p.26/33

INV. 433	symb	$R\nabla E^+ = \Delta f(x) \circ E^+(x)$, $d(R\nabla E^+) = \nabla_{-i}\nabla_{-j}(R + E^+)$, $R\nabla E^+ = f(x)$, $f(x) = M1$	[MANIFOLD]
INV. 434	symb	$-2(T-t) \mid R_{ij} + \nabla \nabla f = \int_{-1}^1 (y \log y)^{\wedge (1/2)} dy^{\wedge} m ; 1/(-2(T-t)) \mid g^2_{-ij} = \int_{-1}^1 (x \log x)^2 dx^{\wedge} m$	[MANIFOLD, ENTROPY]
INV. 435	symb	$(\Box + m^2) \phi = 0$, $(\gamma^{\wedge} n \partial_{-n} + m^2) \phi = 0$	[SR, QUANTUM]
INV. 436	symb	$\Box = (\delta \phi + m^2) \phi$, $(\delta \phi + m^2 c) \phi = 0$, $E = mc^2$, $E = -(1/2) m v^2 + m c^2$, $\Box \phi^2 = (\partial \phi + m^2) \phi$	[SR, QUANTUM]
INV. 437	symb	$\Box \phi^2 = (8 \pi G/c^4) T_{\mu \nu}$, $\nabla \phi^2 = 8 \pi G(p/c^3 + V/S)$, $(d/dt) g_{ij} = -2R_{ij}$, $f(x) + g(x) \geq f(x) \circ g(x)$	[SR, MANIFOLD]
INV. 438	symb	$\int_{- \beta}^{\wedge} \alpha a(x-1)(y-1) \geq 2 \int f(x) g(x) dx$	[OTHER]
INV. 439	calc	$m^2 = 2 \pi T \left((26 - D_n)/24 \right)$, $r^{\wedge} n = 1/(1-z) \quad = 5.75959$	[QUANTUM]
INV. 440	symb	$(\partial \phi + m^2 c) \phi = 0$, $T_{\mu \nu} = (1/2) m v^2 - (1/2) k x^2$	[QUANTUM]
INV. 441	symb	$\int [f(x)] dx = \mid \mid \int f(x) dx \mid \mid$, $\int \nabla x dx = \Delta \Sigma f(x) dx$, $(e^{\wedge}(i \theta))^{\wedge} = i e^{\wedge}(i \theta)$	[ROT]
INV. 442	symb	$T_{\mu \nu} = n h \nu$; $T_{\mu \nu} = (1/2) m v^2 - (1/2) k x^2 \geq m c^2 - (1/2) m v^2$	[SR, QUANTUM]
INV. 443	symb	$Z \in R \cap Q$, $R \subset M^+$, $C^+ \otimes_{\{k=0\}^{\wedge} n} H_m$, $E^+ \cap R^+$, $M^+ = \Sigma_{\{k=0\}^{\wedge} n} C^+ \otimes H_m$, $M^+ = \Sigma_{\{k=0\}^{\wedge} n} C^+ \cup H^+$, $E2 \otimes E1$, ... , $(d/df)F = \otimes \nabla M^+ , \otimes \nabla C^+$	[MANIFOLD]
INV. 444	symb	$\nabla \circ \Delta = \nabla \Sigma f(x) / \Delta \rightarrow mesh f(x) dx$, $\partial x/\nabla \rightarrow \partial xy$, $(d/dx) f(x) (d/dx) g(x) / \Box x = -[f, g]$	[QUANTUM]
INV. 445	symb	$\lim_{\{x \rightarrow \infty\}} \Sigma f(x) dx = \int dx$, $\nabla \int dx$	[OTHER]
INV. 446	symb	$(E2 \otimes_{\{k=0\}^{\wedge} n} E1) \cdot (R - C^+) = \otimes_{\{k=0\}^{\wedge} n} \nabla C^+ , \vee \int^{\wedge} \{C^+\} \nabla H_m$	[MANIFOLD]
INV. 447	symb	$\Delta (M^+ - \nabla C^+) = \wedge M^+ - \otimes_{\{k=0\}^{\wedge} n} C^+ , \exists (M^+ - \nabla C^+) = XOR(\otimes_{\{k=0\}^{\wedge} n} \nabla M^+)$, $-[E^+ \nabla R^+] = \nabla + \nabla - C^+$	[MANIFOLD]
INV. 448	symb	$\int dx$, ∂x , $\nabla_{-i}\nabla_{-j}$, $\Delta x \rightarrow E^+ \nabla M1$, $E^+ \cap R \in M1$, $R \nabla C^+$	[MANIFOLD]
INV. 449	holds	$[[\cos x, \sin x], [\sin x, -\cos x]] [x; y] = [[1, 0], [0, -1]] \quad = 0$ vs 0	[ROT, QUANTUM]
INV. 450	differs	$[[\cos x, -1], [1, -\sin x]] [\cos x; \sin x] = [[1, 0], [0, -1]] (\times 3) \quad = 0.25 + 0.616025i$ vs $1 - 1i$	[ROT, QUANTUM]
INV. 451	holds	$\Sigma_{\{k=0\}^{\wedge} n} \cos k \theta = \sin((n+1) \theta/2) / \sin(\theta/2) \cdot \cos(n \theta/2) \quad = 3.83538$ vs 3.83538	[ROT]
INV. 452	holds	$\Sigma_{\{k=0\}^{\wedge} n} \sin k \theta = \sin((n+1) \theta/2) / \sin(\theta/2) \cdot \sin(n \theta/2) \quad = 3.57302$ vs 3.57302	[ROT]
INV. 453	holds	$\sin \theta = (e^{\wedge}(i \theta) - e^{\wedge}(-i \theta))/(2i)$, $\cos \theta = (e^{\wedge}(i \theta) + e^{\wedge}(-i \theta))/2$, $\tan \theta = (e^{\wedge}(i \theta) - e^{\wedge}(-i \theta)) / (i(e^{\wedge}(i \theta) + e^{\wedge}(-i \theta))) \quad =$ $2.25135 + 0i$ vs $2.25135 + 0i$	[ROT]
INV. 454	differs	$\lim_{\{\theta \rightarrow 0\}} \sin \theta / \theta \rightarrow 1$, $\lim_{\{\theta \rightarrow 0\}} \cos \theta / \theta \rightarrow 1$ (sic; diverges) $\quad = 10001$ vs 2	[ROT]
INV. 455	symb	$(e^{\wedge}(i \theta))^{\wedge} = 2$, $\nabla f = 2$, $e^{\wedge}(i \theta) = \cos \theta + i \sin \theta$	[ROT]
INV. 456	symb	$\rightarrow [\cos^2 \theta + \sin \theta + \cos \theta - 2 \sin^2 \theta] [[1, 0], [0, -1]]$	[ROT]
INV. 457	differs	$2 \sin \theta \cos \theta = 2 n \lambda \sin \theta \quad = 0.866025$ vs 0.5	[ROT, QUANTUM]
INV. 458	symb	$\lim_{\{\theta \rightarrow 0\}} (1/\theta)[\sin \theta; \cos \theta][[\theta, 1], [1, \theta]][\cos \theta; \sin \theta] = [[1, 0], [0, -1]]$, $f^{\wedge}(-1)(x) \times f(x) = 1$ (see INV.218)	[ROT, QUANTUM]
INV. 459	calc	$\sin \theta = \lambda/d$, $-p y1 \sin 90^{\circ} \leq \sin \theta \leq p y2 \sin 90^{\circ}$, $\lambda = h/(mv) \quad = 7.2739e-10$	[ROT, QUANTUM]
INV. 460	symb	$\lambda 2/\lambda 1 = \sin \theta / \sin(\theta/2)$, h , $\lambda^{\wedge} \geq 2h$, $\int \sin 2 \theta = \mid \mid x - y \mid \mid$	[ROT, QUANTUM]
INV. 461	symb	$D^2 \phi = \nabla \int (\nabla_{-i}\nabla_{-j} f)^2 d \eta$	[MANIFOLD]
INV. 462	calc	$E = mc^2$, $E = (1/2) m v^2 - (1/2) k x^2$, $G_{\mu \nu} = (1/2) \wedge g_{ij}$, $\Box = (1/2) k T^2 \quad = 8.98755e+16$	[SR]
INV. 463	symb	$\ker f / \im f \cong S^{\wedge} \mu \nu_m$, $S^{\wedge} \mu \nu_m = \pi(\chi, \chi) \otimes h_{\mu \nu}$	[MANIFOLD]
INV. 464	symb	$D^2 \phi = 0(x) (p/c^3 + V/S)$, $V(x) = D^2 \phi \otimes M^+_3$, $S^{\wedge} \mu \nu_m \otimes S^{\wedge} \mu \nu_n = -2R_{ij} / V(\tau) [D^2 \phi]$	[MANIFOLD]
INV. 465	symb	$\nabla_{-i}\nabla_{-j} [S^{\wedge} mn_1 \otimes S^{\wedge} mn_2] = \int^{\wedge} \{V(\tau)\} f(x) [D^2 \phi] = \int^{\wedge} \{V(\tau)\} f(x) 0(x)$	[MANIFOLD]
INV. 466	symb	$z(x) = g(cx + d) / (f(ax + b) h(ex + l)) = \int^{\wedge} \{V(\tau)\} f(x) 0(x)$	[OTHER]
INV. 467	symb	$V(x)/f(x) = m(x)$, $0(x) = m(x)[D^2 \phi(x)]$, $(d/df)F = m(x)$, $\int F dx^{\wedge} m = \Sigma_{\{k=0\}^{\infty}} m(x)$	[MANIFOLD]
INV. 468	symb	$\int \nabla \phi^2 d \nabla \phi = \Box \phi$, $\Box \phi = \int [D^2 \phi \otimes h_{\mu \nu}] dm$	[MANIFOLD]
INV. 469	symb	$0(x) = [\nabla_{-i}\nabla_{-j} \int \nabla f(x) d \eta]^{\wedge} (1/2)$, $\delta \cdot 0(x) = [\nabla_{-i}\nabla_{-j} \int \nabla g(x) dx_{ij}]^{\wedge} (iy)$	[ZETA]
INV. 470	symb	$\mid \mid ds^2 \mid \mid = e^{\wedge}(-2 \pi T \mid \phi \mid) [0(x) + \delta 0(x)] dx^{\wedge} \nu dx^{\wedge} \nu + \lim_{\{n \rightarrow 1\}} \Sigma_{\{k=0\}^{\infty}} a_k f_k$	[ZETA, MANIFOLD]
INV. 471	calc	$\log(x \log x) \geq 2 (y \log y)^{\wedge} (1/2)$, $(y \log y)^{\wedge} (1/2) / \log(x \log x) \leq 1/2 \quad = 1.52218$	[ENTROPY]
INV. 472	symb	$\delta(x) = (\text{reality of value})/(\text{exist of value}) \leq 1$; $\log(x \log x) = \Box \phi$; $(y \log y)^{\wedge} (1/2) = \nabla \phi^2$	[ENTROPY]
INV. 473	symb	$l(x) = 2x^2 + qx + r = (ax + b)(cx + d) + r$, $e^{\wedge}(l(x)) = (d/df) L(x)$, $G_{\mu \nu} = g(x) \wedge f(x)$	[OTHER]
INV. 474	symb	$V(\tau) = \int \exp[L(x)] dx^{\wedge} m + 0(N^{\wedge}(-1))$, $g(x) = L(x) \circ \int e^{\wedge}(2x^2 + 2x + r) dx^{\wedge} m$	[QUANTUM]
INV. 475	symb	$\mid \mid ds^2 \mid \mid = \mid \mid (d/df) L(x) \mid \mid$, $\eta = [\nabla_{-i}\nabla_{-j} \int \nabla f(x) d \eta]^{\wedge} (1/2)$, $\bar{h} = [\nabla_{-i}\nabla_{-j} \int \nabla g(x) d_{ij}]^{\wedge} (iy)$	[ZETA, MANIFOLD]
INV. 476	symb	$(1/\tau) (N/2 + r(2 \Delta F - \mid \nabla f \mid^2 + R) + f) \bmod N^{\wedge}(-1)$	[ENTROPY]
INV. 477	symb	$(x^2/a) \cos x + (y^2/b) \sin x = r^2$	[ROT]
INV. 478	calc	$S^2_m = \mid \mid \int \pi r^2 dr \mid \mid^2$, $V(\tau) = r^2 \times S^2_m$, $S^{\wedge} mn_1 \otimes S^{\wedge} mn_2 = \int [D^2 \phi \otimes h_{\mu \nu}] dm \quad = 1.09662$	[MANIFOLD]
INV. 479	symb	$V(x) = \int (1/\sqrt{(2 \pi q)}) (\exp L(x) dx) + 0(N^{\wedge}(-1))$	[MANIFOLD]
INV. 480	symb	$V(x) = 2 \int (R + \nabla_{-i}\nabla_{-j} f) / -(R + \Delta f) e^{\wedge}(-f) dV$, $V(\tau) = \int \tau(p)^{\wedge}(-n/2) \exp(-(1/\sqrt{(2 \pi q)}) L(x) dx) + 0(N^{\wedge}(-1))$	[MANIFOLD, ENTROPY]
INV. 481	symb	$Zeta(x, h) = \exp(q f(x))^{\wedge} m / m$	[ZETA]
INV. 482	symb	$\mid x y z ; u v w \mid^2 / g_{\mu \nu}(x) = (f(x) dx^{\wedge} \mu dx^{\wedge} \nu , f^{\wedge}(y) dy^{\wedge} \mu dy^{\wedge} \nu , f^{\wedge}(z) dz^{\wedge} \mu dz^{\wedge} \nu) \cdot (u, v, w) =$ $[[1, 0, 0], [0, 1, 0], [0, 0, i]] \cong g(x, y, z) / (f(a, b, c) \cdot h^{\wedge}(-1)(u, v, w))$	[MANIFOLD, QUANTUM]
INV. 483	symb	$(d/df) \iiint \Box \phi d \phi_{xy} = V(\Box \phi)$, $\lim_{\{n \rightarrow \infty\}} \Sigma_{\{k=0\}^{\infty}} V_k(\Box \phi) = (\partial / \partial f) i h c$	[QUANTUM]
INV. 484	symb	$\lim_{\{n \rightarrow \infty\}} \Sigma_{\{k=0\}^{\infty}} G_{\mu \nu} = f(x) \circ m(x)$, $V(x)/f(x) = m(x)$	[MANIFOLD]
INV. 485	symb	$(a_k f_k)^{\wedge} = nC0 a_0 f^{\wedge} n + nC1 a_1 f^{\wedge} (n-1) ... nC(r-1) a_n f^{\wedge} (n-1)$	[OTHER]
INV. 486	symb	$\int a_k f_k dx_k = (a_{(n+1)/(k+1)}) f^{\wedge}(n+1) + (a_{(k+2)/(k+2)}) f^{\wedge}(k+2) + ... + (a_0/k) f_k$	[OTHER]
INV. 487	symb	$(\partial / \partial f) \Box \phi = (1/4) g^2_{-ij} (\times 2)$, $(\nabla \phi^2 / \Box \phi)^{\wedge} = 0 (\times 2)$	[MANIFOLD]
INV. 488	symb	$(y \log y)^{\wedge} (1/2) / \log(x \log x) = (1/2)/(1/(2i))$	[ENTROPY]
INV. 489	symb	$\{f, g\}/[f, g] = 1/i$, $(\{f, g\}/[f, g])^{\wedge} = i^2$, $(i)^2 \rightarrow (1/4) g_{ij}$, $F_m^{\wedge} t = (1/4) g^2_{-ij}$, $f(r) = (1/4) \mid r \mid^2$, $4 f(r) = g^2_{-ij}$	[QUANTUM]
INV. 490	symb	$(1/y) \cdot (1/y^{\wedge}) \cdot (y^{\wedge})^{\wedge} / y^{\wedge} \cdot (y^{\wedge})^{\wedge} / y^{\wedge} \cdot \dots = nCr y2 \cdot y3 \cdot \dots / (nCr y1 y2 \cdot \dots)$	[OTHER]
INV. 491	symb	$(\partial y / \partial x) \cdot (\partial / \partial y) f(y) = y^{\wedge} \cdot f^{\wedge}(y)$	[OTHER]
INV. 492	symb	$\int l \times l dm = (l \otimes l) m$, $= (d/dx^{\wedge} \mu)(d/dx^{\wedge} \nu) f_{\mu \nu} \cdot \nabla \phi^2 = \Box \phi$	[OTHER]
INV. 493	differs	$\nabla \phi^2 / \Box \phi = 1/2$, $l = 2 \pi r$, $V/S = (4 \pi r^2)/(4 \pi r^3)$ (sic) , $2 \pi r = 2 \cdot (\pi r^2) = \pi r^2$, $H3 = 2$, $\pi(H3) = 0 \quad = 0.666667$ vs 1	[OTHER]
INV. 494	symb	$f(r) = (1/2) \sqrt{(1 + f^{\wedge}(r))} / f(r) + m g f(r)$	[OTHER]
INV. 495	symb	$S^{\wedge} m_n = \mid S2 S1 - S1 S2 \mid$, $\Box \phi = V \cdot S^{\wedge} m_n$, $G_{\mu \nu} \cong R_{\mu \nu}$	[MANIFOLD]
INV. 496	symb	$\nabla_{-i}\nabla_{-j}(\Box \phi) d \phi_{xy} = (\partial / \partial f) \Box \phi \cdot T_{\mu \nu} = (8 \pi G/c^4)(T_{\mu \nu})^2$, $(\chi \otimes \pi) = \iiint \Box \phi d^3 \phi = \text{div}(\text{rot } E, E1) \cdot e^{\wedge}(-i \times \log x)$	[ROT, SR, ENTROPY]
INV. 497	symb	$\sigma(H_n \otimes K_m) = E_n \times H_m$, $H_n = [\nabla_{-i}\nabla_{-j} \int \nabla g(x) dx_{-i} dx_{-j}]^{\wedge} (iy)$, $K_m = [\nabla_{-i}\nabla_{-j} \int \nabla f(x) d \eta]^{\wedge} (1/2)$	[ZETA, MANIFOLD]

4. 全方程式レジストリ (2111 本) p.27/33

INV. 498	symb	$ ds^2 = \sigma(H_n \times K_m), \sigma(\chi, x) \times \pi(\chi, x) = (\partial/\partial f) L(x), V'_-\tau(x) = (\partial/\partial V) L(x) (\times 2)$	[MANIFOLD]
INV. 499	symb	$(\Box \phi)' = (\partial/\partial f) \circ (\iiint f(x,y,z) dx dy dz)' d\phi ; V'_-\tau(x) = \text{same} (\times 3)$	[MANIFOLD]
INV. 500	symb	$(\partial/\partial V) L(x) = V(\tau) \int e^{-(\int x \log x dx + O(N^*(-1)))} d\phi$	[ENTROPY]
INV. 501	symb	$(d/df) \sum_{\{k=1\}^n} \sum_{\{k=0\}^\infty} a_k f_k = (d/df) m(x), V(x)/f(x) = m(x)$	[OTHER]
INV. 502	symb	$4 V'_-\tau(x) = g^2_{ij}, (d/dl) L(x) = \sigma(\chi, x) \times V_-\tau(x)$	[MANIFOLD]
INV. 503	symb	$ ds^2 = e^{(-2\pi T \phi)}[\eta_{\mu\nu} + \hat{h}_{\mu\nu}(x)] dx^\mu dx^\nu + T^2 d\phi^2 (\times 3 \text{ more})$	[MANIFOLD]
INV. 504	symb	$f^-(2)(x) = [\nabla_i \nabla_j \int \nabla f^-(5) d\eta]^{-(1/2)} = [f^-(2)(x) d\eta]^{-(1/2)}$	[MANIFOLD]
INV. 505	symb	$\nabla_i \nabla_j \int F(x) d\eta = (\partial/\partial f) F, \nabla f = (d/dx) f, \nabla_i \nabla_j \int \nabla f d\eta = (\partial/\partial x_i)(\partial/\partial x_j)((d/dx) f)$	[MANIFOLD]
INV. 506	symb	$(z_3 z_2 - z_2 z_3)/(z_2 z_1 - z_1 z_2) = \omega, (\bar{z}_3 z_2 - \bar{z}_2 z_3)/(\bar{z}_2 z_1 - \bar{z}_1 z_2) = \bar{\omega}, \omega \cdot \bar{\omega} = 0, z_n = \omega - \{x\}, z_n \cdot \bar{z}_n = 0, \bar{z}_n \cdot \bar{z}_n = 0$	[ROT]
INV. 507	symb	$[f, g] \times [g, f] = fg + gf = \{f, g\}$	[QUANTUM]
INV. 508	symb	$V(\tau) = \int e^{-(\int x \log x + O(N^*(-1)))} d\phi$	[ENTROPY]
INV. 509	symb	$(\Box \phi)' = 4 \tilde{V}(x), (\partial/\partial V) L(x) = m(x), V(\tau) = \int (1/\sqrt{2\tau q}) \exp[L(x)] d\phi + O(N^*(-1))$	[MANIFOLD]
INV. 510	symb	$V(\tau) = \iiint V/S^2 dm, f(r) = \dots, \log(x \log x) \geq 2(y \log y)^{1/2}, F_m{}^t = (1/4) g^2_{ij}, (d/dt) g_{ij}(t) = -2R_{ij}$	[MANIFOLD, ENTROPY]
INV. 511	calc	$\nabla_i \nabla_j v = (1/2) m v^2 + m c^2, \int \nabla_i \nabla_j v dv = (\partial/\partial f) L(x) = 8.99255e+16$	[SR]
INV. 512	symb	$(\Box \phi)^2 = -2 \int \nabla_i \nabla_j v d^2 v, (\Box \phi)^2 = (\nabla \phi^2/\Box \phi)' = (d/df) \int_1 (x \log x)^2 dm$	[ENTROPY]
INV. 513	symb	$\otimes \nabla M+_3 = \int \sqrt{(R + \nabla_i \nabla_j f)^2} / \exists (R + \Delta f) dV = (x,y,z) \cdot (u,v,w)/\Gamma$	[GAMMA, MANIFOLD]
INV. 514	symb	$\otimes C+- = \int \exp[\int \nabla_i \nabla_j f d\eta] d\phi = L(x) \cdot (\partial/\partial l) F(x) = (\Box \phi)^2$	[QUANTUM]
INV. 515	calc	$l = \sqrt{(hG/c^2)}, T_{\mu\nu} = (1/2) k l^2 + (1/2) m v^2, E = mc^2 - (1/2) m v^2 = 4.05126e-35$	[SR, QUANTUM]
INV. 516	holds	$e^{(x \log x)} = x^x, x = \log x^x / \log x, y = x, x = e = 9.88212 \text{ vs } 9.88212$	[ENTROPY]
INV. 517	symb	$\int_1 (x \log x) dx = i \int x \log x dx + \int_1 (x \log x) dx$	[ENTROPY]
INV. 518	symb	$\int_1 (x \log x)^2 dx^m = i (1/2) x^2; \int_1 (x \log x)^2 dx^m = i \int_1 M dx^m \leq 1/(2i) + x^2$	[ENTROPY]
INV. 519	calc	$E = -(1/2) m v^2 + m c^2 (\times 4 \text{ overall}) = 8.98255e+16$	[SR]
INV. 520	symb	$\lim_{\{x \rightarrow \infty\}} \int_1 (x \log x)^2 dx^m \geq 1/(2i), (d/df) \int_1 (x \log x)^2 dx^m = 1/(2i)$	[ENTROPY]
INV. 521	calc	$\lim_{\{x \rightarrow \infty\}} x^2 / e^{(x \log x)} = 0 = 1e-08$	[ENTROPY]
INV. 522	symb	$\int dx \rightarrow \partial f \rightarrow dx \rightarrow \text{cons}$	[OTHER]
INV. 523	symb	$(\Box \phi)' = (\exists \int \sqrt{(R + \nabla_i \nabla_j f)} e^{(-f)} dV)' d\phi$	[MANIFOLD, ENTROPY]
INV. 524	symb	$\otimes M+_3 = (\partial/\partial f) L(x), (\partial/\partial l) L(x) = \nabla_i \nabla_j \int \nabla f(x) d\eta, L(x) = V(x)/f(x), l(x) = L'(x)$	[MANIFOLD]
INV. 525	symb	$V'(\tau) = \int e^{-(\int x \log x dx + O(N^*(-1)))} d\phi; F = \text{same}$	[ENTROPY]
INV. 526	symb	$T_{\mu\nu} = \iiint V(x)/S^2 dm, S^2 = \pi r^2 \cdot S(x) = 4\pi r^3 / \tau(x)$	[MANIFOLD]
INV. 527	symb	$\eta = \nabla_i \nabla_j \int \nabla f(x) d\eta, \hat{h} = \nabla_i \nabla_j \int \nabla g(x) dx_i dx_j, \delta 0(x) = [\nabla_i \nabla_j \int f(x) d\eta]^{-(1/2+iy)}$	[ZETA, MANIFOLD]
INV. 528	symb	$Z(x, h) = \lim_{\{x \rightarrow \infty\}} \sum_{\{k=0\}^\infty} q T^m / m = \delta(x)$	[ZETA]
INV. 529	symb	$l(x) = 2x^2 + p x + q, m(x) = \lim_{\{x \rightarrow \infty\}} \sum_{\{k=0\}^\infty} (q x^m)' / f(x)$	[OTHER]
INV. 530	holds	$Z(T, X) = \exp(\sum_{\{m=1\}^\infty} q^k T^m / m), Z(x, h) = \exp(q f(x))^m / m = 1.38889 \text{ vs } 1.38889$	[ZETA]
INV. 531	symb	$\lim_{\{x \rightarrow 1\}} \text{mesh } m/(m+1) = 0, \int x^m = x^m/(m+1) \text{ (sic; } x^{\{m+1\}/(m+1)})$	[OTHER]
INV. 532	calc	$(d/df) \int x^m = m x^m, \lim_{\{x \rightarrow 1\}} \text{mesh}(x) = \lim_{\{m \rightarrow \infty\}} m/(m+1), \lim_{\{x \rightarrow 1\}} \sum a_k f_k = \alpha = 0.999001$	[OTHER]
INV. 533	symb	$(\partial/\partial V) ds^2 = T_{\mu\nu}, V(\tau) = \int e^{(x \log x)} d\phi = l(x)$	[MANIFOLD]
INV. 534	symb	$R_{ij} = (\partial/\partial x_i)(\partial/\partial x_j) T_{\mu\nu}, G_{\mu\nu} = R_{\mu\nu} - T_{\mu\nu}$	[MANIFOLD]
INV. 535	symb	$F(x) = \int e^{-(\int x \log x dx + O(N^*(-1)))} d\phi, (d/dV) F(x) = V'(x)$	[ENTROPY]
INV. 536	symb	$T_{\mu\nu} = R_{\mu\nu}, T_{\mu\nu} = \iiint V/S^2 dm, \delta 0(x) = [D^2 \phi \otimes h_{\mu\nu}] dm$	[MANIFOLD]
INV. 537	symb	$\nabla(\Box \phi)' = [\nabla_i \nabla_j \int \nabla f(x) d\eta]^{-(1/2+iy)}$	[ZETA]
INV. 538	symb	$(f(x), g(x))' = (A_{\mu\nu})' [[d_x, \delta(x)], [\varepsilon(x), \partial_x]] \cdot (f(x,y), g(x,y)) = [[1,0],[0,-1]]$	[QUANTUM]
INV. 539	calc	$\delta(x) \cdot 0(x) = [[1,0],[0,-1]]^{(1/2)} = 6.12323e-17 + 1i$	[QUANTUM]
INV. 540	symb	$p = e^{(x \log x)}, e^{(-x \log x)}; p = \otimes (i\hbar \nabla) \otimes L$	[ZETA, QUANTUM]
INV. 541	symb	$ ds^2 = 0, 1 \text{ (Hilbert manifold of genus 3)}$	[MANIFOLD]
INV. 542	symb	$ ds^2 = e^{(-2\pi T \phi)}[\eta_{\mu\nu} + \hat{h}(x)] dx^\mu dx^\nu + T^2 d^2 \phi = [\infty]/e^{(-2\pi T \phi)} + T^2 d^2 \phi \geq [\infty]/e^{(-2\pi T \phi)} \cdot T^2 d^2 \phi = (n/(n+1)) \Gamma_n = \int e^{(-x)} x^{(1-t)} dx$	[GAMMA, MANIFOLD]
INV. 543	symb	$\lim_{\{x \rightarrow \infty\}} \sum_{\{x=0\}^\infty} n/(n+1) = a_k f_k$	[OTHER]
INV. 544	calc	$\beta(p,q) = \int e^{(-\sin \theta \cos \theta)} \int \sin \theta \cos \theta d\theta = \int \Gamma(\gamma)' dx^m = 0.578244$	[ROT, GAMMA, BETA]
INV. 545	symb	$\int \Gamma(\gamma)' dx^m = \int \Gamma dx^m \cdot (d/d\gamma) \Gamma \leq \int \Gamma dx^m + (d/d\gamma) \Gamma = \otimes (i\hbar \nabla) \otimes L' \int \otimes (i\hbar \nabla) \otimes L dx^m, \otimes (i\hbar \nabla) \otimes L = n$	[GAMMA, QUANTUM]
INV. 546	differs	$\beta(p,q)^{-(-1)} = \Gamma(p) \Gamma(q) / \Gamma(p+q) \text{ (sic; this is } \beta \text{ itself)} = 12 \text{ vs } 0.0833333$	[GAMMA, BETA]
INV. 547	differs	$\sqrt{g} = 1; \Gamma(p+q)/(\Gamma(p) \Gamma(q)) = \beta(p,q) \text{ (sic; } = 1/B) = 12 \text{ vs } 0.0833333$	[GAMMA, BETA]
INV. 548	symb	$x^n + y^n \geq z^n$	[OTHER]
INV. 549	symb	$(\Gamma(p) \Gamma(q))^2 - \Gamma(p,q)^2 = 0; (\Gamma(p) \Gamma(q) - \Gamma(p,q))(\Gamma(p) \Gamma(q) + \Gamma(p,q)) = 0$	[GAMMA]
INV. 550	calc	$((d/d\gamma) \Gamma'(\gamma)) (\int \Gamma'(\gamma) dx^m) (e^\pi - \pi^e) = (\Box - \Box')(\Box + \Box') = 0.681535$	[GAMMA, OTHER]
INV. 551	differs	$(2(\sin(i x \log x) + \cos(i x \log x))(\cos(i x \log x) - i \sin(i x \log x)) / (2(\sin(i x \log x) + \cos(i x \log x))(\cos(i x \log x) + i \sin(i x \log x))) = 0 = 16 + 0i \text{ vs } 0 + 0i$	[ROT, ENTROPY]
INV. 552	symb	$= (99 - 96)(94 + 92)(90 - 87)(85 + 82)(80 - 78) \cdots = 0$	[ZETA]
INV. 553	holds	$\beta(p,q)^{-(-1)} = 1 / ((5-3)(7+13)(17-19)(23+29)(31-37)(41+47)(51-53) \cdots) = 1 / (2 \cdot 20 \cdot (-2) \cdot 52 \cdot (-6) \cdot 88 \cdot (-2) \cdots) = \int_1 \beta(p,q) dx = \int_1 t^2 dt = -2.27637e-07 \text{ vs } -2.27637e-07$	[ZETA, BETA]
INV. 554	differs	$(\beta(p,q) - \beta(p,q)^{-(-1)})(\beta(p,q) + \beta(p,q)^{-(-1)}) = 0 = -143.993 \text{ vs } 0$	[BETA]
INV. 555	symb	$\Gamma(2) = \beta(5, -3) = \Gamma(5) \Gamma(-3) / \Gamma(5-3)$	[GAMMA, BETA]
INV. 556	differs	$\Gamma(2) = \int e^{(-2) 2^{-(t-1)} dx} = \sqrt{e} = \zeta(s) = 1 \text{ vs } 1.64872$	[GAMMA, ZETA]
INV. 557	calc	$\beta(p,q) = \Gamma(-1) = -1/12 = -0.0833333$	[GAMMA, ZETA, BETA]
INV. 558	calc	$x = (1/(2\pi i)) \log(\sqrt{(1 - 1/\Box^2)}) = 0 + 0.0355144i$	[ROT, SR]
INV. 559	symb	$\Box = 2(\sin(i x \log x) + \cos(i x \log x))$	[ROT]
INV. 560	differs	$\Gamma(5) = 3 = \Box = 3; \Gamma(3) = 2 = \Box = 2; \Gamma(2) = 1 = \Box = 1 = 24 \text{ vs } 3$	[GAMMA, ZETA]
INV. 561	differs	$e^\pi = \pi^e, x = \sqrt{g} = 23.1407 \text{ vs } 22.4592$	[OTHER]

4. 全方程式レジストリ (2111 本) p.28/33

INV. 562	calc	$x = 2\,e^{(-1)}\,e^{(-1)},\,\sqrt{g} = 1,\,\sqrt{g} = \sqrt{e} = 0.270671$	[OTHER]
INV. 563	holds	$1/(x\log x) = \sqrt{g};\,e^{(x\log x)},\,x = 2,\,2^2 = 4,\,2^2 = e^{(2\log 2)},\,4 = e^{(\log 4)},\,\log 4 = \log \log 4 = \sqrt{(4\log 4)}$ $= 1 - 2 = 1\text{ (sic)} = 4\text{ vs }4$	[ENTROPY]
INV. 564	symb	$\Sigma_{(k=0)}^\infty a_k\,f(x,y)^{(a_k)} = \pi(\chi,x) = \int x\log x\,dx$	[ENTROPY]
INV. 565	symb	$\beta(p,q)/(x\log x) = \zeta(s)$	[ZETA, BETA, ENTROPY]
INV. 566	symb	$\pi(\chi,x) = i\,\pi(\chi,x) \circ f(x) - f(x) \circ \pi(\chi,x)$	[MANIFOLD, QUANTUM]
INV. 567	symb	$\iint_1(x\log x)^2\,dx^{\sim}m \geq 1/(2i)(\times 3);y = x \Rightarrow \iint_1(y\log y)^{(1/2)}\,dy^{\sim}m \geq 1/2$	[ENTROPY]
INV. 568	symb	$F_m^{\sim}t \geq \int_M(R + \nabla_i\nabla_j f)\,e^{(-f)}\,dV$	[MANIFOLD, ENTROPY]
INV. 569	symb	$F_t \geq (2/n)\,f^2$	[MANIFOLD, ENTROPY]
INV. 570	symb	$(d/df)\,F = 2\int(R + \nabla_i\nabla_j f)^2/-(R + \Delta f)\,dm(\times 3)$	[MANIFOLD, ENTROPY]
INV. 571	symb	$(d/df)\,F_t = (1/4)\,g^2_{ij}$	[MANIFOLD]
INV. 572	differs	$f(r) = (1/(2m))\sqrt{(1 + f'(r))}/f(r) - m\,g\,f(r);m = 1,\,g = 1 \Rightarrow f(r) = (1/4) r ^2 = 1\text{ vs }-0.292893$	[OTHER]
INV. 573	symb	$\iint_1(x\log x)^2\,dx^{\sim}m + \iint_1(y\log y)^{(1/2)}\,dy^{\sim}m = 0$	[ENTROPY]
INV. 574	symb	$x^{(1/2 + iy)} = e^{(x\log x)}$	[ZETA, ENTROPY]
INV. 575	symb	$(d/df)\,F(v_{ij},h) = \int e^{(-f)}[-\Delta v + \nabla_i\nabla_j v_{ij} - R_{ij}v_{ij} - v_{ij}\nabla_i\nabla_j + 2\langle \nabla f,\nabla h \rangle + (R + \nabla^2 f)(v/2 - h)]$	[MANIFOLD, ENTROPY]
INV. 576	symb	$F \geq (d/df)\,\iint_1(x\log x)^2\,dx^{\sim}m + (d/df)\,\iint_1(y\log y)^{(1/2)}\,dy^{\sim}m$	[ENTROPY]
INV. 577	differs	$\zeta(2) = 1/4 = \pi^2/6\text{ (sic)} = 1.64493\text{ vs }0.25$	[ZETA]
INV. 578	calc	$\Gamma(-1) = -1/12 = -0.0833333$	[GAMMA, ZETA]
INV. 579	calc	$\beta(2,-3) = \Gamma(2)\Gamma(-3)/\Gamma(2-3); \Gamma(2)\Gamma(-3)/(-1/12) = \Gamma(2) = \int e^{(-2)}(-2)^{(t-1)}\,dx,\,\Gamma(-3) = \int e^{-3}(-3)^{(t-1)}\,dx = 0.166667$	[GAMMA, BETA]
INV. 580	symb	$\int e^{(-2)}e^{-3}(-2)^{(t-1)}(-3)^{(t-1)}\,dx = \int e^{-1}(-2)^u(-3)^u\,dx; \int e^{-1}(-1)z^{\sim}n\,dx = \int w\,e\,dx = -(\log x)' = -1/x$	[OTHER]
INV. 581	differs	$1 + 1/2 + 1/3 + 1/4 + 1/5 + \cdots = -1/12\text{ (sic; harmonic series diverges)} = 9.78761\text{ vs }-0.0833333$	[ZETA]
INV. 582	differs	$\beta(p,q) = (\beta(p,q))^{(-1)} = 0.0833333\text{ vs }12$	[BETA]
INV. 583	differs	$\Gamma(-1) = 1 + 2 - 3 + 4 - 5 + \cdots = -1/12 = 1.74\text{ vs }-0.0833333$	[GAMMA, ZETA]
INV. 584	symb	$\beta\Box\beta = \beta(p,q)/\log x$	[ZETA, BETA]
EXP. 1	calc	$f(x) = x^2 + 1,\,f : x \rightarrow y = 5$	[OTHER]
EXP. 2	symb	$T' = \int \Gamma(\gamma)' \,dx^{\sim}m$	[GAMMA]
EXP. 3	symb	<chem>CH2(OH)-CH(OH)-C(=O)-OH + 2NH4+ -> Me/Me-C-CH3/CH2-OH-COOH</chem> (garbled organic scheme)	[OTHER]
EXP. 4	symb	<chem>NH2 / NH2-H2N ; NH-CH2-CH(NH2)-COOH</chem>	[OTHER]
EXP. 5	symb	<chem>H2C-CH-CHO-C(=O)-H0 + 2NH4+ -> Me Me HCHO C(=O) H0</chem>	[OTHER]
EXP. 6	symb	<chem>[CH2-C-N-CH2] Cl + 0Na + heat -> C02 + H20 + NaCl</chem>	[OTHER]
EXP. 7	symb	<chem>[CH2-C-N-CH2] Cl + Na0-CH2-C(=O)-CH2-CH3 + heat -> C02 + H20 + NaCl</chem>	[OTHER]
EXP. 8	symb	<chem>R1[CH2-C(=O)-N(H)]R2 + Na0-CH2-C(=O)-CH2-CH3 + heat -> C02 + H20 + NaCl + R1-C(=O)-OH + R2-NH4+</chem>	[OTHER]
EXP. 9	symb	<chem>R1[CH2-C(=O)-N(H)]R2 + K0-CH2-C(=O)-CH2-CH3 + heat -> C02 + H20 + KCl + R1-C(=O)-OH + R2-NH4+</chem>	[OTHER]
EXP. 10	calc	$Na : 10/11\text{ mol},\,K : 25/19\text{ mol (per }1\text{ mL}^3) = 2.22488$	[OTHER]
EXP. 11	symb	$\nabla_i\nabla_j(\Box\phi) = \Box + \Box'$	[MANIFOLD]
EXP. 12	symb	$\Box' = (8\pi G/c^4)\,T_{\mu\nu}/\log x (=INV.119)(\times 2)$	[SR, MANIFOLD]
EXP. 13	symb	$(R_{\mu\nu} + (1/2)\wedge g_{ij})_{\mu\nu} = \kappa(T^{\mu\nu}_e + T^{\mu\nu}_y + T^{\mu\nu}_g + T^{\mu\nu}_h + \cdots)$	[SR, MANIFOLD]
EXP. 14	symb	$(d/dM)(\kappa\,T_{\mu\nu}) \rightarrow \otimes (i\hbar\nabla) \otimes L,\,H\Psi = i\hbar\phi$	[QUANTUM]
EXP. 15	symb	$\otimes (i\hbar\nabla) \otimes L \rightarrow (d/dM)(\kappa\,T_{\mu\nu})$	[QUANTUM]
EXP. 16	symb	$ ds^2 = \lim_{\{x\rightarrow\infty}\left[\delta(x)\int\int\pi(\sum_{k=0}^\infty(n\sqrt{p},x/n))^{(1/2)}d\tau\right]_{\mu\nu} (=INV.106)(\times 4)$	[ZETA, MANIFOLD]
EXP. 17	symb	$ ds^2 = \int [D^2\phi \otimes h\nu]\,d\tau = (d/dM)\,F$	[MANIFOLD, QUANTUM]
EXP. 18	calc	$E = mc^2 - (1/2)m\,v^2 = ds^2 = \int \Gamma(\gamma)' \,dx^{\sim}m = \int e^{(-f)}\,dV = \int dvol = e^{(-f)} + e^f \geq e^f - e^{(-f)} \geq e^{(-f)} - e^f = 0.82$	[SR, GAMMA]
EXP. 19	calc	$= R'_{\mu\nu} + (1/2)\wedge g'_{ij} = \int(i\,v/\sqrt{(1-(v/t)^2)} + v/\sqrt{(1-(v/t)^2)})\,dvol (=INV.117) = 0.4 + 0.4i$	[SR]
EXP. 20	symb	$= (d/df)\,\iint_1(x\log x)^2\,dx^{\sim}m + (d/df)\,\iint_1(y\log y)^{(1/2)}\,dy^{\sim}m (=INV.31)$	[ENTROPY]
EXP. 21	symb	$= \int C\,dx^{\sim}m = \int \kappa\,T_{\mu\nu}\,dx^{\sim}m = T_{\mu\nu}\,T^{\mu\nu}' (=INV.118)$	[MANIFOLD]
EXP. 22	symb	$(d/dM)\,F = (d/df)\,F = -2\int(R + \nabla_i\nabla_j)/(\Delta + R)\,e^{(-f)}\,dV$	[MANIFOLD, ENTROPY]
EXP. 23	symb	Seifert manifold = (x, y, z)/Γ	[GAMMA, MANIFOLD]
EXP. 24	symb	$\int(\kappa\,T_{\mu\nu})\,\nabla\,dx^{\sim}m = \int e^{(x\log x)}\text{div}(\text{rot }E)\,dx^{\sim}m = e^{(-x\log x)}$	[ROT, ENTROPY]
EXP. 25	symb	$\int f(x)\,dx = \int \Gamma(\gamma)' \,dx^{\sim}m = 2(\cos(i\,x\log x) - i\sin(i\,x\log x)) (=INV.94)$	[ROT, GAMMA]
EXP. 26	symb	$(\int f(x)dx/\log x) = \lim_{\{\theta\rightarrow\infty}\}(\int f(x)dx/\theta) = 0,1;e^{(i\theta)} = \cos\theta + i\sin\theta (=INV.95/96)$	[ROT]
EXP. 27	symb	$(\int f(x)dx)' = 2(i\sin(ix\log x) - \cos(ix\log x)) = 2(-\cos(ix\log x) + i\sin(ix\log x)) (=INV.97)$	[ROT]
EXP. 28	symb	$(\cos(ix\log x) - i\sin(ix\log x))' = (d/de^{(i\theta)})(\cos,-\sin)\cdot(\sin,\cos) (=INV.98)$	[ROT]
EXP. 29	symb	$\int \Gamma(\gamma)'' \,dx^{\sim}m = \int \Gamma(\gamma)' \,dx^{\sim}m/\nabla L = \int \Gamma\,dx^{\sim}m \cdot (d/d\gamma)\Gamma/\nabla L \leq (\int \Gamma\,dx^{\sim}m + (d/d\gamma)\Gamma)/\nabla L \leq e^f - e^{(-f)} \leq e^{(-f)} + e^f = 0,1 (=INV.99/100)$	[GAMMA]
EXP. 30	holds	$\log x \nabla L;g_{ij} = f\,f',\,F\,f \nabla L;g_{ij} : x \rightarrow y,\,x^{\sim}p \rightarrow y;f(x) = \log x^{\sim}p = p\log x,\,f(y) = p\log x = 2.74653\text{ vs }2.74653$	[MANIFOLD]
EXP. 31	symb	$(d^2/df^2)\,F = F\,f' \cdot f\,f',\,\iint F\,dx^{\sim}m = F\,f \cdot F(f)'$	[OTHER]
EXP. 32	symb	$(d/(df\,dg))(f,g) = (f\cdot g)\,f' + g'$	[OTHER]
EXP. 33	symb	$(F\,f \cdot G\,g) = \int (f\cdot g)\,f' + g';\iint F \cdot G\,dx^{\sim}m = [F\,f \cdot G\,g] - \int (f\cdot g)\,f' + g'$	[OTHER]
EXP. 34	symb	$\int (d/(df\,dg))\,FG = \int F\,f'\,G + \int F\,G\,g';\int F\,f'\,G\,dx^{\sim}m = [F\,f\,G\,g] - \int F\,G\,g'\,dx^{\sim}m$	[OTHER]
EXP. 35	symb	$(F(x)/G(x))(fg)' = (F\,f'\,G - F\,G\,g')/G\,g$	[OTHER]
EXP. 36	symb	$(d/(df\,dg))\,FG = (d/df^{\sim}m)\,FG,(\partial/\partial f^{\sim}m)\,FG = F\,f^{\sim}\mu\nu \cdot G + F \cdot G\,g^{\sim}\mu\nu,\int F\,dx^{\sim}m = F\,f$	[OTHER]
EXP. 37	holds	$\beta(p,q) = \Gamma(p)\Gamma(q)/\Gamma(p+q) = 0.19635\text{ vs }0.19635$	[GAMMA, BETA]
EXP. 38	calc	$t = \Gamma(x),\,T = \int \Gamma(x)\,dx = 0.922746$	[GAMMA]
EXP. 39	symb	$\beta(p,q) = -\int_1 t^2\,dt$	[BETA]
EXP. 40	symb	$T_m = \int \Gamma(x)\,dx^{\sim}m$	[GAMMA]

4. 全方程式レジストリ (2111 本) p.29/33

EXP. 41	symb	$T' = t' \log t \, dt + C$ (C integration constant)	[OTHER]
EXP. 42	symb	$dx^{\sim}m = (1/\log x) \, dx$, $dx^{\sim}m = (\log x^{(-)})'$	[ZETA]
EXP. 43	symb	$T' = \int \Gamma(\gamma)' \, dx^{\sim}m$	[GAMMA]
EXP. 44	symb	$\otimes T \nabla = \int T \, dx^{\sim}m$, $\delta(t) = t \, dx^{\sim}m$	[OTHER]
EXP. 45	differs	$i^{\sim}i = (\sqrt{i})^{\sim}(\sqrt{i}) = e^{\sim}(x \log x) = 0.20788 + 0i$ vs $0.487619 + 0.30257i$	[ROT, ENTROPY]
EXP. 46	holds	$e^{\sim}x = i$, $e^{\sim}(2\pi m) = i^{\sim}n$, $(d/dx) \, e^{\sim}(2\pi m) = i^{\sim}n = 6.12323e-17 + 1i$ vs $0 + 1i$	[ROT]
EXP. 47	symb	$e^{\sim}(i\theta) = \cos\theta + i \sin\theta$, $e^{\sim}(2\pi m) = i^{\sim}n$, $2\pi m = n$, $e = i$ (sic)	[ROT]
EXP. 48	holds	$(d/di) \, i^{\sim}i = i^{\sim}i = e^{\sim}(x \log x)$, $m = 1/(2\pi)$, $l = 2\pi r$, $\pi = l/(2r) = 3,14159$ vs $3,14159$	[ROT]
EXP. 49	symb	$ ds^2 = e^{\sim}(-2\pi T \phi)[\eta_{\sim}\mu\nu + \bar{h}(x)] \, dx^{\sim}\mu \, dx^{\sim}\nu + T^2 \, d^2\phi$ (=INV.143)	[MANIFOLD]
EXP. 50	symb	$[\eta_{\sim}\mu\nu + \bar{h}(x)] \, dx^{\sim}\mu \, dx^{\sim}\nu / i = H\Psi = i\hbar\phi$, $H\Psi = (1/i)[H, \Psi]$	[QUANTUM]
EXP. 51	symb	$\otimes (i\hbar\nabla) \otimes L = e^{\sim}(i \, x \log x) \, e^{\sim}(i \, x \log x)'$, $H\Psi = \otimes H\Psi / \nabla L$	[QUANTUM, ENTROPY]
EXP. 52	symb	$n!^{(-)}(-1) = \otimes \nabla L$, $n!^{(-)}(-1) = \int T'_{\sim}\mu\nu \, dV - \int T_{\sim}\mu\nu \, dV = \int f(x)^{\sim}(-1) \, x \, f(x) - f(x)$	[GAMMA]
EXP. 53	symb	$\exists \, x = a$, $1 - \exists \int x \, dx$, $\Lambda = \int A \, dx$	[OTHER]
EXP. 54	symb	$\nabla f(x) = (1/n!) \int \int \cdots \int dx \, \Sigma \int (\int \cdots A \int dz) / n!$	[OTHER]
EXP. 55	symb	$M^2 \, a(m) = k \, a(m)$	[QUANTUM]
EXP. 56	symb	$Z_{M3} = \int dA \, e^{\sim}(k \, i / (4\pi) \, L_{M3})$	[JONES, QUANTUM]
EXP. 57	symb	$Z_M = \{M1 \mid M2\}$	[JONES, QUANTUM]
EXP. 58	holds	$e^{\sim}x = \Sigma \, x^{\sim}n / n!$ (r = ∞) = 4.48169 vs 4.48169	[OTHER]
EXP. 59	symb	$(d/df) \int \, dx^{\sim}m = (1/n!) \Sigma \, x^{\sim}n$	[OTHER]
EXP. 60	calc	$\int_{\sim}n^{(n+1)} \, dx/x < 1/n$ (n ≥ 1) = 0.0456513	[ZETA]
EXP. 61	calc	$1 + 1/2 + \cdots + 1/n > \int_{\sim}1^{(n+1)} \, dx/x = \log(n+1) = 0.531073$	[ZETA]
EXP. 62	symb	$1 + 1/2 + \cdots + 1/n - \log x > \log(n+1) - \log n > 0$; $1/(n+1) < \int_{\sim}0^{(n+1)} \, dx/x = \log(n+1) - \log n$ (sic lower limit n)	[ZETA]
EXP. 63	holds	$\lim_{\{n \rightarrow \infty\}} (1 + 1/2 + \cdots + 1/n - \log n) = C = \int (\int_{\sim}1 \, x^{\sim}s \, dx - \log x) \, dvol = \int C \, dx^{\sim}m$, $C = 0.5772156\cdots = 0.577266$ vs 0.577216	[GAMMA, ZETA]
EXP. 64	symb	$C = \int_{\sim}1 \, x^{\sim}s \, dx - \log x$; $= \int (\int_{\sim}1 \, x^{\sim}s \, dx - \log x) \, dvol = \int\!\!\int_{\sim}1 \, x^{\sim}s \, dx - \int \log x \, dvol$	[ZETA]
EXP. 65	symb	$F = \Gamma = \int e^{\sim}(-x) \, x^{\sim}(1-t) \, dx = (d/df) \int\!\!\int_{\sim}1 (y \log y)^{\sim}(1/2) \, dx^{\sim}m - \int e^{\sim}(-x) \, x^{\sim}(1-t) \log x \, dx^{\sim}m$	[GAMMA, ENTROPY]
EXP. 66	symb	$\int \, x^{\sim}(1-t) \, e^{\sim}(-x) \, dV = \int \, x^{\sim}(1-t) \, dm$; $\int \, x^{\sim}(1-t) \, e^{\sim}(-x) \, dV = \int \, x^{\sim}(1-t) \, dvol$	[GAMMA, MANIFOLD]
EXP. 67	differs	$f = \gamma = \Gamma' = \int e^{\sim}(-x) \, x^{\sim}(1-t) \log x \, dx = (d/d\gamma) \Gamma^{\sim}(-1) - \int e^{\sim}(-x) \, x^{\sim}(1-t) \log x \, dx^{\sim}m = (d/d\gamma) \Gamma^{\sim}(-1) - (\gamma) \gamma' = e^{\sim}(-f) - e^{\sim}f = 2 \cos(i \, x \log x)$ (sic) = -0.577212 vs 0.577216	[GAMMA, ZETA]
EXP. 68	symb	$ds^2 = g_{\sim}\mu\nu(x) \, dx^{\sim}\mu \, dx^{\sim}\nu + \phi^2(x)(\, dx^2 + \kappa^2 \, A_{\sim}\mu(x) \, dx^{\sim}\nu)^2$	[SR, MANIFOLD]
EXP. 69	symb	$ds^2 = -N(r)^2 \, dt^2 + \phi^2(x)(\, dr^2 + r^2 \, d\theta^2)$	[SR, MANIFOLD]
EXP. 70	symb	$ds^2 = -dt^2 + r^{\sim}(-8\pi Gm) (\, dr^2 + r^2 \, d\theta^2)$	[SR, MANIFOLD]
EXP. 71	symb	$dx^2 = g_{\sim}\mu\nu(x)(\, g_{\sim}\mu\nu(x) \, dx^2 - dx \, g_{\sim}\mu\nu(x))$; $dx = (\, g_{\sim}\mu\nu(x)^2 \, dx^2 - g_{\sim}\mu\nu(x) \, dx \, g_{\sim}\mu\nu(x))^{\sim}(1/2)$ (=INV.197) ($\times 2$)	[MANIFOLD]
EXP. 72	symb	$\pi(\chi, x) = i\pi(\chi, x) \, f(x) - f(x) \, \pi(\chi, x)$ (=INV.566) ($\times 2$)	[MANIFOLD, QUANTUM]
EXP. 73	calc	$ds^2 = -(1 - r_{\sim}s/r) \, c^2 \, dt^2 + dr^2/(1 - r_{\sim}s/r) + r^2 \, d\theta^2 + r^2 \sin^2\theta \, d\phi^2 = 6.96077e-10$	[SR, MANIFOLD]
EXP. 74	calc	$R_{\sim}\mu\nu + (1/2) \, g_{ij} \, \Lambda = \kappa \, T_{\sim}\mu\nu = 2.07655e-43$	[SR, MANIFOLD]
EXP. 75	symb	$\int \kappa \, T_{\sim}\mu\nu \, dvol = \int (\, R_{\sim}\mu\nu + (1/2) \, g_{ij} \, \Lambda) \, dvol$	[MANIFOLD]
EXP. 76	symb	$\int \Gamma \cdot (d/d\gamma) \Gamma \, dx^{\sim}m \leq \int \Gamma \, dx^{\sim}m + (d/d\gamma) \Gamma = \int \Gamma(\gamma)' \, dx^{\sim}m$ ($\times 2$)	[GAMMA]
EXP. 77	symb	$\int\!\!\int (\, \int_{\sim}1 \, x^{\sim}s \, dx - \log x) \, dvol = e^{\sim}f - e^{\sim}(-f) \leq e^{\sim}f + e^{\sim}(-f)$ ($\times 2$)	[ZETA]
EXP. 78	symb	$(d/df) \, F + \int C \, dx^{\sim}m = 2(\cos(i \, x \log x) - i \sin(i \, x \log x))$ ($\times 2$)	[ROT, ZETA]
EXP. 79	symb	$(d/df) \, F \geq (d/df) \int\!\!\int_{\sim}1 (x \log x)^2 \, dx^{\sim}m + (d/df) \int\!\!\int_{\sim}1 (y \log y)^{\sim}(1/2) \, dy^{\sim}m$ ($\times 2$; cf INV.576)	[ENTROPY]
EXP. 80	symb	$H\Psi = \otimes (\, i\hbar\nabla \otimes L) = (1/(2i)) \, e^{\sim}(i\hbar)$, $(i\hbar)' = (-e^{\sim}(i\hbar))' = -i \, e^{\sim}(i\hbar)$ ($\times 2$)	[QUANTUM]
EXP. 81	symb	$\phi(x) = e^{\sim}(-i\hbar t)$, $\otimes (i\hbar\nabla) \otimes L = (1/2) \, e^{\sim}(i\hbar) (-i \, e^{\sim}(i\hbar)) = (1/(2f))^{\sim}(-if) = (1/2)^{\sim}(-if) \cdot e^{\sim}(-x \log x) \cdot (f)^{\sim}i = \int e^{\sim}(-x) \, x^{\sim}(t-1) \, dx$ ($\times 2$)	[GAMMA, QUANTUM]
EXP. 82	holds	$(d/d\gamma) \, \Gamma = e^{\sim}(-x \log x)$; $f = x$, $i = t$, $1/2 = a$; $\otimes a^{\sim}(-t) \, x \, x^{\sim}t \, [Im] \equiv \int e^{\sim}(-x) \, x^{\sim}(t-1) \, dx$ ($\times 2$) = 1.32934 vs 1.32934	[GAMMA]
EXP. 83	symb	$ \phi(t)_{\sim}s = e^{\sim}(-i\hbar t) \, \Psi\rangle_H$, $\hat{A}_s = \hat{A}_H(\theta)$ ($\times 2$)	[QUANTUM]
EXP. 84	holds	$ \Psi(t)_{\sim}s \rightarrow (d/dt) : i \, (d/dt) \, \phi(t)_{\sim}s = \hat{H} \, \phi(t)_{\sim}s$ ($\times 2$) = 0 - 0.644218i vs 0 - 0.644218i	[QUANTUM]
EXP. 85	symb	$\langle \hat{A}(t) \rangle = \langle \Psi(t) \hat{A}(0) \Psi(t) \rangle$ ($\times 2$)	[QUANTUM]
EXP. 86	symb	$(d/dt) \, \hat{A} = (1/i) \, [\hat{A}, H]$ ($\times 2$)	[QUANTUM]
EXP. 87	calc	$\hat{A}(t) = e^{\sim}(i\hbar t) \, \hat{A}(0) \, e^{\sim}(-i\hbar t)$ ($\times 2$) = 0.169967 + 0.98545i	[QUANTUM]
EXP. 88	symb	$\lim_{\{\theta \rightarrow 0\}} (1/\theta) \, [\sin\theta; \cos\theta][[\theta, 1], [1, \theta]][\cos\theta; \sin\theta] = [[1, 0], [0, -1]]$; $f^{(-)}(-1)(x) \, x \, f(x) = I'_{\sim}m$, $I'_{\sim}m = [1, 0] \times [0, 1]$ (=INV.218/219) ($\times 2$)	[ROT, QUANTUM]
EXP. 89	symb	$x + y \geq \sqrt{(xy)}$	[OTHER]
EXP. 90	symb	$x^{\sim}(1/2 + iy) / e^{\sim}(x \log x) = 1$	[ZETA, ENTROPY]
EXP. 91	symb	$0(x) = \nabla_{\sim}i \nabla_{\sim}j \int e^{\sim}((2/m) \sin\theta \cos\theta) \times N \bmod(e^{\sim}(x \log x))$; $0(x)(x + \Delta f ^2)^{\sim}(1/2)$ (=INV.215/216) ($\times 2$)	[ROT]
EXP. 92	symb	$x \, \Gamma(x) = 2 \int \sin 2\theta ^2 \, d\theta$, $0(x) = m(x)[D^2\phi]$ (=INV.217) ($\times 2$)	[ROT, GAMMA]
EXP. 93	symb	$i^2 = (0, 1) \cdot (0, 1)$, $ a b \cos\theta = -1$, $E = \text{div}(E, E1)$, $\{f, g\}/[f, g] = i^2$, $E = mc^2$, $I' = i^2$ (=INV.220/221) ($\times 2$)	[ROT, SR, QUANTUM]
EXP. 94	symb	$H\Psi = \otimes (i\hbar\nabla) \otimes L = \otimes H\Psi / \nabla L = e^{\sim}(x \log x) = x^{\sim}(x)'$; $(d/df) \, F = m(x)$	[QUANTUM, ENTROPY]
EXP. 95	symb	$(d/dt) \, \phi(t) = \hbar = (1/(2i)) \, e^{\sim}(i\hbar)$, $(i\hbar)' = -i \, e^{\sim}(i\hbar)$	[QUANTUM]
EXP. 96	symb	$(d/d\gamma) \, \Gamma = m(x) = e^{\sim}(-x \log x)$	[GAMMA, ENTROPY]
EXP. 97	differs	$\sin(ix) = (e^{\sim}(-x) + e^{\sim}x)/(2i)$ (sic; correct is $(e^{\sim}\{-x\} - e^{\sim}\{x\})/(2i) = i \sinh x$) = 0 + 1.1752i vs 0 - 1.54308i	[ROT]
EXP. 98	symb	$(d/df) \, F = m(x) = e^{\sim}f + e^{\sim}(-f) = 2i \sin(i \, x \log x)$ (sic)	[ROT, ENTROPY]
EXP. 99	symb	$\beta(p, q) = \int x^{\sim}(1-t)(1-x)^{\sim}t \, dx = \int t^{\sim}x \, (1-t)^{\sim}(x-1) \, dt$ (=INV.160)	[BETA]
EXP. 100	holds	$\int_{\sim}0^1 x^{\sim}10 \, (1-x)^{\sim}20 \, dx = B(11, 21) = \Gamma(11) \Gamma(21)/\Gamma(32) = 10!20!/31! = 1/931395465$ (=INV.161) = 1 vs 1	[GAMMA, BETA]
EXP. 101	symb	$1/931395465 = 1/9 = 1/(1-x)$; $1/(1-z) = \Sigma \, z^{\sim}k$; $1/(1+z^2) = \Sigma (-1)^{\sim}k \, z^{\sim}2k$; $f(x) = \Sigma \, a_k \, z^{\sim}k$ (=INV.162-164)	[OTHER]
EXP. 102	symb	$d^{\sim}n \, y/dx^{\sim}n = n!$ $y^{\sim}(n+1)$, $f^{\sim}(0)(0) = n!$ $f(0)^{\sim}(n+1) = n!$, $f(x) \equiv \Sigma \, x^{\sim}n = 1/(1-x)$ (=INV.165/166)	[OTHER]
EXP. 103	holds	$dy/dx = y^2 \Rightarrow -1/y = x - C \Rightarrow y = 1/(C - x)$; $x=0$, $y=1 \Rightarrow y = 1/(1-x)$ (=INV.167/168) = 4 vs 4	[OTHER]

4. 全方程式レジストリ (2111 本) p.30/33

EXP. 104	holds	$H\Psi = \otimes H\Psi \ / \ \nabla L \ , \ \Gamma = \int e^{\sim}(-x) \ x^{\sim}(1-t) \ dx \ , \ \gamma = \int e^{\sim}(-x) \ x^{\sim}(1-t) \ \log x \ dx = \int \Gamma(\gamma)' \ dx^{\sim}m = (d/d\gamma) \ \Gamma \quad = 0.422784 \ \text{vs} \quad 0.422784$	[GAMMA]
EXP. 105	differs	$E = mc^2 \ , \ E = m^2 \ c^2 \quad = 1.79751e+17 \ \text{vs} \ 3.59502e+17$	[SR]
EXP. 106	symb	$\nabla(i\hbar\nabla)\otimes L \ , \ \otimes(i\hbar\nabla)\otimes L \ , \ \square(i\hbar\nabla)\otimes L \ (=INV.93) \ (\times 2)$	[JONES, QUANTUM]
EXP. 107	symb	$(d/d\gamma) \ \Gamma = e^{\sim}f + e^{\sim}(-f) \geq e^{\sim}f - e^{\sim}f$	[GAMMA, JONES]
EXP. 108	symb	$(d/d\gamma) \ \Gamma + \int \mathbb{C} \ dx^{\sim}m = 2(\cos(i \ x \ \log x) - i \ \sin(i \ x \ \log x) \)$	[ROT, GAMMA]
EXP. 109	symb	$e^{\sim}(-\theta) = \cos(i \ x \ \log x) - i \ \sin(i \ x \ \log x)$	[ROT, JONES]
EXP. 110	symb	$\int \Gamma(\gamma)' \ dx^{\sim}m = 2(\cos(i \ x \ \log x) - i \ \sin(i \ x \ \log x) \)$	[ROT, GAMMA]
EXP. 111	calc	$\square = -(16\pi G/c^4) \ T_{\mu \nu} \quad = -4.15311e-43$	[SR, MANIFOLD]
EXP. 112	symb	$\hat{t}\!\!\!\int\!\!\!\int\!\!\!\int \text{cohom} \ D_{\chi} \ [Im] = \oint (p \ x^{\sim}n + q \ x + r) \ \nabla_{\perp} l \ ; \ (d/dl) \ L(x,y) = 2 \ \int \sin 2x ^2 \ d\tau \ , \ (d/d\gamma) \Gamma \ (=INV.120/121)$	[ROT, MANIFOLD]
EXP. 113	symb	$= [\ i \ \pi(\chi,x), f(x) \] \ (=INV.25)$	[QUANTUM]
EXP. 114	calc	$(d/d\gamma) \ \Gamma = \int \mathbb{C} \ dx^{\sim}m = \int (\ \int_{\perp} 1 \ x^{\sim}s \ dx - \log x \) \ dvol \quad = 0.577212$	[GAMMA, ZETA]
EXP. 115	symb	$m \ \otimes(i\hbar\nabla)\otimes L + n \ \otimes(i\hbar\nabla)\otimes L = (m+n)\otimes(i\hbar\nabla)\otimes L \ ; \ (m-n) \ ; \ \otimes L^{\sim}m \times \otimes L^{\sim}n = \otimes L^{\sim}(m+n) \ ; \ \otimes L^{\sim}m \ / \ \otimes L^{\sim}n = \otimes L^{\sim}(m/n) \ (=INV.85-88)$	[QUANTUM]
EXP. 116	symb	$d/df \ \otimes(i\hbar\nabla)\otimes L = \otimes(i\hbar\nabla)\otimes L' \ , \ \int \otimes(i\hbar\nabla)\otimes L \ dx^{\sim}m \ , \ \otimes(i\hbar\nabla)\otimes L = n \ , \ n^{\sim}(L+1)/(n+1) = \int e^{\sim}x \ x^{\sim}(1-t) \ dx \ (=INV.89-91)$	[GAMMA, QUANTUM]
EXP. 117	symb	$(\ \otimes(i\hbar\nabla)\otimes L + m \ \otimes(i\hbar\nabla)\otimes L + n \) = n^{\sim}(L+1)/(n+1) = t \ \int (1-x)^{\sim}n \ x^{\sim}(m-1) \ dx \ ; \ \int x^{\sim}(m-1)(1-x)^{\sim}(n-1) \ dx \ (=INV.92)$	[GAMMA, BETA]
EXP. 118	symb	$\boxtimes(i\hbar\nabla) L \ dx^{\sim}m \ , \ \boxplus(i\hbar\nabla L \ dx^{\sim}m) \ , \ x^{\sim}(1/2+iy) = e^{\sim}(x \ \log x) \ (=INV.93/47) \ (\times 2)$	[ZETA, QUANTUM]
EXP. 119	holds	$C = b0 + c1/(b1 + c2/(b2 + c3/(b3 + c4/(b4 + \cdots)))) \quad = 0.577216 \ \text{vs} \ 0.577216$	[ZETA]
EXP. 120	calc	$(2.71828)^2.828 = a^{\sim}a + b^{\sim}b + c^{\sim}c + d^{\sim}d \ \cdots = \int e^{\sim}f \cdot x^{\sim}(1-t) \ dx \quad = 16.9116$	[GAMMA, ENTROPY]
EXP. 121	symb	$(d/df) F(v_{\perp ij} \ , \ h) = \int e^{\sim}(-f) \ dV \ [-\Delta v + \nabla_{\perp} i \nabla_{\perp} j \ v_{\perp ij} - R_{\perp ij} \ v_{\perp ij} - v_{\perp ij} \ \nabla_{\perp} i \nabla_{\perp} j + 2\langle \nabla f, \nabla h \rangle + (R + \nabla^2 f)(v/2 - h) \] = (d/df)F = 2 \ \int (R + \nabla_{\perp} i \nabla_{\perp} j \ f)^2 \ / \ -(R + \Delta f) \ dm \ (=INV.575/570)$	[MANIFOLD, ENTROPY]
EXP. 122	symb	$F \geq (d/df) \ \hat{\int\!\!\!\int}_1 (x \ \log x)^2 \ dx^{\sim}m + (d/df) \ \hat{\int\!\!\!\int}_1 (y \ \log y)^{\sim}(1/2) \ dy^{\sim}m \ (=INV.576)$	[ENTROPY]
EXP. 123	symb	$V(\tau) = [f(x), g(x)] \times [f^{\sim}(-1)(x), h(x)]$	[MANIFOLD, QUANTUM]
EXP. 124	symb	$\Gamma(p,q) = \int e^{\sim}(-x) \ x^{\sim}(1-t) \ dx = \beta(p,q) = \pi(\ f(\chi,x), \ x \)$	[GAMMA, BETA, MANIFOLD]
EXP. 125	symb	$ ds^2 = 0(x)[\ f(x) \circ g(x)_{\perp} \mu \nu \] \ dx^{\sim}\mu \ dx^{\sim}\nu = \lim_{\{x \rightarrow \infty\}} \ \Sigma_{\{k=0\}}^{\sim} \infty \ a_k \ f_k$	[MANIFOLD]
EXP. 126	symb	$G_{\mu \nu} = (\partial/\partial f) \ \int [\ f(x)_{\perp} \mu \nu \circ G(x)_{\perp} \mu \nu \ dx^{\sim}\mu \ dx^{\sim}\nu \]_{\perp} \mu \nu \ dm = g_{\mu \nu}(x) \ dx^{\sim}\mu \ dx^{\sim}\nu - f(x)_{\perp} \mu \nu \ dx^{\sim}\mu \ dx^{\sim}\nu$	[MANIFOLD]
EXP. 127	symb	$[\ i \ \pi(\chi,x), f(x) \] = i \ \pi \ f(x) - f(x) \ \pi(\chi,x)$	[QUANTUM]
EXP. 128	symb	$T_{\mu \nu} = (\ \lim_{\{x \rightarrow \infty\}} \ \Sigma_{\{k=0\}}^{\sim} \infty \ \hat{\int\!\!\!\int} [\ V(\tau) \circ S_{\perp} \mu \nu(\chi,x) \] \ dm)_{\perp} \mu \nu \ dx^{\sim}\mu \ dx^{\sim}\nu \ , \ G_{\mu \nu} = R_{\mu \nu} \ T_{\mu \nu}$	[MANIFOLD]
EXP. 129	symb	$[[D_m, \ d_x],[d_x, \ \sigma_m]] \ [[\cos\theta, \ -\sin\theta],[\sin\theta, \ \cos\theta]] \ [x; y] = [[1,0],[0,-1]]^{\sim}(1/2) \ (\text{cf} \ INV.370) \ (\times 2)$	[ROT, QUANTUM]
EXP. 130	symb	$\sigma_m \ll [[\delta(x), \ -1],[1, \ \varepsilon(x)]]^{\sim}(1/2) = [[i,0],[0,-i]] \ (=INV.372) \ (\times 3)$	[QUANTUM]
EXP. 131	symb	$V(M) = (\partial/\partial f) \ (\ N \ \int [f \setminus M] \otimes N \)_{\perp} \mu \nu \ dx^{\sim}\mu \ dx^{\sim}\nu$	[MANIFOLD]
EXP. 132	calc	$V(M) = \pi(\ 2 \ \int \sin^2 dx \) \otimes (d/df) \ F_M \ dx^{\sim}m \quad = 9.8696$	[ROT, MANIFOLD]
EXP. 133	symb	$\lim_{\{x \rightarrow \infty\}} \ \Sigma_{\{k=0\}}^{\sim} \infty \ a_k \ f_k = \int (\ F(V) \ dx^{\sim}m)_{\perp} \mu \nu \ dx^{\sim}\mu \ dx^{\sim}\nu \ , \ \otimes_{\{k=0\}}^{\sim} \infty [f \setminus g] = \vee(M \wedge N)$	[MANIFOLD]
EXP. 134	symb	$\pi 1(M) = e^{\sim}(-f^2 \ \int \sin^2 x \ dm + 0(N^{\sim}(-1))) = [\ i \ \pi(\chi,x), \ f(x) \]$	[ROT, MANIFOLD]
EXP. 135	symb	$M \circ f(x) = e^{\sim}(-f \ \int \sin x \cos x \ dx^{\sim}m) + \log(\ 0(N^{\sim}(-1)) \)$	[ROT]
EXP. 136	symb	$(d/dL) \ V(\tau) = (d/df) \ \hat{\int\!\!\!\int}_M 1/(x \ \log x)^2 \ dx^{\sim}m + (d/df) \ \hat{\int\!\!\!\int}_M 1/(y \ \log y)^{\sim}(1/2) \ dy^{\sim}m \ (\times 2)$	[MANIFOLD, ENTROPY]
EXP. 137	symb	$\varepsilon_S(\nu) = \square v \cdot (\partial/\partial \chi)(\ ^5\!\!\sqrt{p} \wedge g^2 \) \ d\chi$	[MANIFOLD]
EXP. 138	symb	$\wedge(F_m^{\sim}t)'^{\sim} = (1/12) \ g^2_{\perp ij}$	[MANIFOLD]
EXP. 139	symb	$\pi(V_{\perp} \tau) = e^{\sim}(-\sqrt{\pi/16} \ \log x)^{\sim}\delta \times 1/(x \ \log x)$	[ENTROPY]
EXP. 140	symb	$(d/dt)(g_{\perp ij})^2 = (1/24)(F_m^{\sim}t)^2$	[MANIFOLD]
EXP. 141	calc	$m^2 = 2\pi T \ (\ (26 - D_n)/24 \) \ (=INV.439) \quad = 0$	[QUANTUM]
EXP. 142	symb	$g_{\perp ij} \wedge \pi(\nu_{\perp} \tau) = e^{\sim}(-2\pi T \phi)[\eta_{\perp} \mu \nu + \hbar_{\perp} \mu \nu(x)] \ dx^{\sim}\mu \ dx^{\sim}\nu + T^2 \ d\phi^2 \ ; \ ds^2 = g_{\perp ij} \wedge \pi(\nu_{\perp} \nu)$	[MANIFOLD]
EXP. 143	differs	$e^{\sim}(i\theta) = \cos\theta + i \ \sin\theta \ , \ e^{\sim}(x \ \log x) = x^{\sim}(1/2+iy) \ , \ x \ \log x = \log(\cos\theta + i \ \sin\theta) = \log \cos\theta + i \ \log \sin\theta \ (\text{sic}) \quad = 0 + 0.523599i \ \text{vs} \ -0.143841 - 0.693147i$	[ROT, ZETA]
EXP. 144	symb	$\log(\sin\theta + i \ \cos\theta) = \log(\sin\theta - i \ \cos\theta) \ ; \ \log(\ \sin\theta \ / \ (i \ \cos\theta) \) = -2R_{\perp ij} \ , \ (d/dt) \ g_{\perp ij}(t) = -2R_{\perp ij}$	[ROT, MANIFOLD]
EXP. 145	symb	$0(x) = \zeta(s) \ / \ \Sigma_{\{k=0\}}^{\sim} \infty \ a_k \ f_k \ , \ \text{Im} \ f = \ker f \ , \ \chi(x) = \ker f \ / \ \text{Im} \ f \ , \ H(3) = 2 \ , \ \nabla H(x) = 2 \ , \ \pi(x) = 0$	[ZETA, MANIFOLD]
EXP. 146	symb	$[f(x)] = \infty \ , \ ds^2 = 0(x)[\eta_{\perp} \mu \nu + \hbar_{\perp} \mu \nu(x)] \ dx^{\sim}\mu \ dx^{\sim}\nu + T^2 \ d^2 \phi \ , \ 0(x) = e^{\sim}(-2\pi T \phi) \ (\times 2)$	[MANIFOLD]
EXP. 147	symb	$T^2 \ d^2 \phi = [f(x)] \ , \ T^2 \ d^2 \phi = \lim_{\{x \rightarrow 1\}} \ \Sigma_{\{k=0\}}^{\sim} \infty \ a_k \ f_k = [T^2 \ d^2 \phi] \ (\times 2)$	[ZETA, MANIFOLD]
EXP. 148	symb	$(d/dL) \ V(\tau) = (d/df) \ \hat{\int\!\!\!\int}_M (\ ^5\!\!\sqrt{x^2} \) \ d\Lambda + (d/df) \ \hat{\int\!\!\!\int}_M N(\ ^3\!\!\sqrt{x} \) \otimes N \ d\Lambda$	[MANIFOLD]
EXP. 149	symb	$M(\ \vee(\wedge \ f \circ g) \ N \)^{\sim}(1/2) = (d/df) \ \hat{\int\!\!\!\int}_M 1/(x \ \log x)^2 \ dx^{\sim}m + (d/df) \ \hat{\int\!\!\!\int}_M 1/(y \ \log y)^{\sim}(1/2) \ dy^{\sim}m$	[ENTROPY]
EXP. 150	symb	$G_{\perp} \mu \nu = R_{\perp} \mu \nu \ T_{\perp} \mu \nu = -(1/2) \ \wedge \ g_{\perp ij}(x) + T_{\perp} \mu \nu$	[MANIFOLD]
EXP. 151	differs	$C = \hat{\int\!\!\!\int}_1 (x \ \log x)^2 \ dx^{\sim}m \quad = 0.0740741 \ \text{vs} \ 0.577216$	[ZETA, ENTROPY]
EXP. 152	calc	$(D_m, \ d_x) \cdot (\cos\theta, \ \sin\theta) \times (d_x, \ \partial_m) \cdot (\cos\theta, \ \sin\theta) \ ; \ [[1,0],[0,i]] \ \beta(x, \theta) \ [x; y] = [[1,0],[0,-1]] \ ; \ l = \sqrt{(\hbar G/c^3)} \ (=INV.370-372) \ (\times 2) \quad = 1.61622e-35$	[ROT, BETA, QUANTUM]
EXP. 153	symb	$(\ \partial/\partial \tau \ f(x,y,z) \)^{\sim}(3') = A_{\perp} \mu \nu \ , \ (d/dt) \ g_{\perp ij}(t) = -2R_{\perp ij} \ (=INV.373)$	[MANIFOLD]
EXP. 154	symb	$x^{\sim}y = 1/y^{\sim}x \rightarrow$	[OTHER]
EXP. 155	symb	$\pi(\chi,x) = [\ i \ \pi(\chi,x), f(x) \] \ , \ (d/df) \ F = F \ f'$	[QUANTUM]
EXP. 156	symb	$\text{SiO2 prism; (Al, Mg)} \rightarrow \text{S; material list: Cs, Fe}\cdot\text{Co60}\cdot\text{Pt}\cdot\text{Al alloy, H2S04, Al203, He, H2, Mg, Al (heat), CnH2n}$	[OTHER]
EXP. 157	symb	$(\partial/\partial f) F(x) = \hat{\int\!\!\!\int} \text{cohom} \ D.k(x) \ [Im] \ ; \ (\partial/\partial f)F = \hat{t}\!\!\!\int\!\!\!\int \text{cohom} \ D.k(x) \ \ll p = \square' \ (=INV.114/115) \ (\times 6)$	[MANIFOLD]
EXP. 158	symb	$\nabla_{\perp} i \nabla_{\perp} j \ f(x) \ d\eta = \partial^2/(\partial x \partial y) \ \int \square' \ d\eta \ (=INV.116)$	[MANIFOLD]
EXP. 159	calc	$R'_{\perp} \mu \nu + (1/2) \ \wedge \ g'_{\perp ij} = \int (\ i \ v/\sqrt{1-(v/t)^2} + v/\sqrt{1-(v/t)^2} \) \ dvol = (d/df)\hat{\int\!\!\!\int}(x \ \log x)^2 \ dx^{\sim}m + (d/df)\hat{\int\!\!\!\int}(y \ \log y)^{\sim}(1/2) \ dy^{\sim}m = \int \mathbb{C} \ dx^{\sim}m = \int \kappa \ T_{\perp} \mu \nu \ dx^{\sim}m = T_{\perp} \mu \nu \ T^{\sim} \mu \nu' \ (=INV.117/118, EXP.19) \quad = 1.33333 + 1.33333i$	[SR, ENTROPY]
EXP. 160	symb	$\square' = (8\pi G/c^4) \ T_{\perp} \mu \nu \ / \ \log x \ ; \ \hat{t}\!\!\!\int\!\!\!\int\!\!\!\int \text{cohom} \ D_{\chi} [Im] = \oint (p \ x^{\sim}n + q \ x + r) \ \nabla_{\perp} l \ ; \ (d/dl) \ L(x,y) = 2 \int \sin 2x ^2 \ d\tau \ (=INV.119-121)$	[ROT, SR]
EXP. 161	symb	$\oint \! \! \! \cong ds^2 = \lim_{\{x \rightarrow \infty\}} [\ \delta(x) \ \hat{\int\!\!\!\int} \ \pi(\Sigma \ ^{\sim}\!\!\sqrt{p}, \ x/n)^{\sim}(1/2) \ d\tau \]_{\perp} \mu \nu \ ; \ e^{\sim}(-2\pi T \phi) [\eta + \hbar] \ dx^{\sim}\mu \ dx^{\sim}\nu + T^2 \ d^2 \phi \ (=INV.122/123)$	[ZETA, MANIFOLD]
EXP. 162	symb	$-(16\pi G/c^4) \ T_{\perp} \mu \nu \ / \ \log x = \square' \ ; \ -(16\pi G/c^4) \ T_{\perp} \mu \nu \ / \ e^{\sim}(-2\pi T \phi)) = 4\pi G\rho \ \ (=INV.124)$	[SR]
EXP. 163	symb	$\partial/(\partial x \partial y) = \nabla_{\perp} i \nabla_{\perp} j \ ; \ \square \hat{\int\!\!\!\int} = \hat{t}\!\!\!\int\!\!\!\int \ ; \ (\partial/\partial x) \hat{\int\!\!\!\int} = \nabla_{\perp} i \nabla_{\perp} j \ \int \nabla f(x) \ d\eta \ ; \ (\partial/\partial f) \hat{\int\!\!\!\int} = \square \hat{\int\!\!\!\int} \ ; \ \Gamma \Gamma, B B, E E, C C, F F, \beta \beta, D D, \hat{t}\!\!\!\int\!\!\!\int \text{cohom} \ D.k(x) \ [Im] \ , \ D \otimes L \ (=INV.125/126)$	[GAMMA, BETA, MANIFOLD]
EXP. 164	symb	$\square + \square' = \emptyset \ , \ \square' \setminus \emptyset = \square \ , \ [[1,0],[0,-i]]^{\sim}(1/2) \ [[1,0],[0,-i]] \ , \ x^{\sim}(1/2) \cong x \ , \ \emptyset^{\sim}(1/2) = \emptyset \ (=INV.127-129)$	[QUANTUM]

4. 全方程式レジストリ (2111 本) p. 31/33

EXP. 165	symb	$\int^\wedge\langle\llcorner n\rangle ds^2 =\iiint8\pi G\left(\rho/c^3+V/S\right)dx\;dy\;dz=\left(\partial/\partial f^n\right) ds^2 $	[SR, ENTROPY]
EXP. 166	symb	$f_r\;dx^m=0(x,y);\;{}^\frown t\iiint D(\chi,x)\;Hom[D^2\phi]\;\llcorner p\equiv vol(V/S)\;(=INV.130/131)\;(\times 2)$	[MANIFOLD]
EXP. 167	symb	$(d/df)F=(F)f',\;\int F\;dx^m=(F)f,\;(d/df)\int_1(x\log x)^2\;dx^m=\int_1(x\log x)^2\;dx^m,\;f'/f'=2x(\log x+\log(x+1)),\;\int\!\!\int dx^m=(1/(x\log x)^2)(\int 2x(\log x+\log(x+1))dx)=e^{\frown}(-f),\;(d/df)F=(1/(x\log x)^2)(2x(\log x+\log(x+1)))=e^f$ $(=INV.132-136)\;(\times 2)$	[ENTROPY]
EXP. 168	symb	$\log(x\log y)\geq 2\;(y\log y)^{\frown}(1/2)=2\sqrt{\frown}(y\log y)\;(=INV.137)\;(\times 3)$	[ENTROPY]
EXP. 169	symb	$ ds^2 =8\pi G(\rho/c^3+V/S);\;\int ds^2 \;dx^m=\int 8\pi G(\rho/c^3+V/S)\;dvol=\dots\;dy^m=\int_1\dots\;dx^m\;(=INV.45/138)$	[SR]
EXP. 170	symb	$(d/df)F=m(x),\;\otimes i\hbar\nabla\odot L=\nabla_i\nabla_j\int\;\nabla f(x)\;d\eta,\;dx^m=y/\log x,\;dy^m=x/\log x,\;e^{\frown}(-f)\;dV=dy^m=dvol$ $(=INV.139-141)$	[MANIFOLD]
EXP. 171	symb	$\beta(p,q)=(number)+Abel\;manifold=AdS5\;manifold=(d/df)F+\int C\;dx^m=\int\;\Gamma(\gamma)'\;dx^m\;(=INV.142)\;(\times 2)$	[GAMMA, BETA]
EXP. 172	symb	$ ds^2 =e^{\frown}(-2\pi G \phi)[\eta+\hbar(x)]\;dx^\mu\;dx^\nu+T^2\;d^2\phi\;(=INV.143)$	[MANIFOLD]
EXP. 173	symb	$\int\;dvol=\square\phi,\;\int\;\nabla\phi^2\;d\nabla\phi=\square\phi\;(\times 2);\;expanding=\log(x\log x)=\square\phi;\;freeze-out=(y\log y)^{\frown}(1/2)=\nabla\phi\;(=INV.144-146)$	[ENTROPY]
EXP. 174	symb	$x^{\frown}n+y^{\frown}n=z^{\frown}n,\;\beta(p,q)=x^{\frown}n+y^{\frown}n-\delta(x)=z^{\frown}n-\delta(x);\;(d/dt)g_{ij}=-2R_{ij};\;\cos x/((\cos x)'\cdot(\sin x)')=z^{\frown}n,\;z^{\frown}n$ $=-2e^{\frown}(x\log x);\;z=e^{\frown}(-f)+e^f-y;\;\beta(p,q)=e^{\frown}(-f)+e^f\;(=INV.147-150)\;(\times 2)$	[ROT, BETA, MANIFOLD]
EXP. 175	symb	$\square x=\int^\frown\{f(x)\}\;\nabla(R+ E+) \;d\square x=\int^\frown\{\Delta f(x)\circ E+\}\;\nabla(R+ E+) \;\square x;\;\square x=\int^\frown\{d(R\nabla E+)\}\dots=\int^\frown\{\nabla_i\nabla_j(R+E+)\}\dots;\exp(\nabla(R+ E+),\Delta(C\supset R))=\pi(RI\subset\nabla E+)=rot(E1,\;div\;E2)\;(=INV.151-153)$	[ROT, MANIFOLD]
EXP. 176	holds	$x\;f(x)=F(x),\;s\;\Gamma(s)=\Gamma(s+1);\;Q\nabla C+=(d/df)F(x)\;\nabla\int\;\delta(s)\;f(x)\;dx;\;E+\nabla f=e^{\frown}(x\log x)\;\nabla\;n!\;f(x)/E(x);\;(d/df)F\;F$ $f'=e^{\frown}(x\log x);\;(C\nabla)\equiv Q=e^{\frown}(x\log x);\;R\nabla E+=f(x)\nabla e^{\frown}(x\log x)\;(=INV.154-158)\quad=8.85534\;vs\;8.85534$	[GAMMA, ENTROPY]
EXP. 177	symb	$\int\;\delta(x)\;f(x)\;dx;\;\int\;\nabla\phi^2\;d\nabla\phi=\square\phi;\;\int\;dx=x+C\;(=INV.159)$	[OTHER]
EXP. 178	holds	$\beta(p,q)=\int\;x^{\frown}(1-t)(1-x)^{\frown}t\;dx=\int\;t^{\frown}x(1-t)^{\frown}(x-1)\;dt;\;\int_0^1x^{\frown}10x^{\frown}10(1-x)^{\frown}20\;dx=B(11,21)=10!20!/31!=1/931395465;$ $series\;1/(1-z),\;1/(1+z^2);\;d^{\frown}ny/dx^{\frown}n=n!\;y^{\frown}(n+1);\;dy/dx=y^2\rightarrow y=1/(1-x)\;(=INV.160-168,\;EXP.99-103)\;(\times 3\;overall)$ $=1\;vs\;1$	[GAMMA, BETA]
EXP. 179	symb	$(dx,\;\partial x)\cdot(\varepsilon x,\;\delta x)=[[1,0],[0,-1]]^{\frown}(1/2),\;(d/df)F=m(x)\;(=INV.169)$	[QUANTUM]
EXP. 180	symb	$\square((\sigma1+\sigma2)/2)=[\;3\pi(\chi,x)\circ f(x),\;\sigma(x)\;]$	[ROT, QUANTUM]
EXP. 181	symb	$\square\phi=8\pi G\;T_{\mu\nu},\;(d/dt)\;g_{ij}(t)=-2R_{ij},\;\nabla\phi^2=4\pi G\rho$	[SR, MANIFOLD]
EXP. 182	symb	$\square(\sigma1+\sigma2)=[\;6\pi(\chi,x)\circ f(x),\;\sigma(x)\;]=[\;i\pi(\chi,x),\;f(x)\;];\;\square\phi=[\;12\pi(\chi,x)\circ f(x),\;\sigma(x)\;]$	[QUANTUM]
EXP. 183	symb	$(d/df)\;F=\int\;e^{\frown}(-f)\;[-\Delta v+R_{ij}\;v_{ij}+\nabla_i\nabla_j\;f+v\;\nabla_i\nabla_j+2<f,h>+(R+\nabla f)(v/2-h)]=[\;i\pi(\chi,x),\;f(x)\;]$	[QUANTUM, ENTROPY]
EXP. 184	symb	$f^{\frown}(-1)(x)\;x\;f(x)=1,\;H_m=E_m\times K_m$	[QUANTUM]
EXP. 185	differs	$i=(1,0)\cdot(1,0)\;(sic),\;e^{\frown}(i\theta)=\cos\theta+i\sin\theta,\;e^{\frown}(-i\theta)=\cos\theta-i\sin\theta\quad=1.76484-0.644218i\;vs\;0.764842+0.355782i$	[ROT]
EXP. 186	differs	$\sin\theta=(e^{\frown}(i\theta)-e^{\frown}(-i\theta))/(2i),\;\sin(i\theta)=(e^{\frown}(-\theta)-e^{\frown}\theta)/2\;(sic;\;=i\sinh\theta)\quad=0+1.1752i\;vs\;-1.1752+0i$	[ROT]
EXP. 187	symb	$\pi(\chi,x)=\cos\theta+i\sin\theta$	[ROT, MANIFOLD]
EXP. 188	holds	$R(-\alpha)=[[cos\alpha,\;sin\alpha],[-sin\alpha,\;cos\alpha]],\;R(\alpha)=[[cos\alpha,\;-sin\alpha],[sin\alpha,\;cos\alpha]]\quad=1\;vs\;1$	[ROT]
EXP. 189	holds	$R(\alpha)\;M\;R(-\alpha)=R(\alpha)[[1,0],[0,-1]]R(-\alpha)=[[cos^2\alpha-sin^2\alpha,\;2\sin\alpha\cos\alpha],[2\sin\alpha\cos\alpha,\;-cos^2\alpha+sin^2\alpha]]=[[cos\;2\alpha,\;\sin\;2\alpha],[sin\;2\alpha,\;-cos\;2\alpha]]\quad=4.44089e-16\;vs\;0$	[ROT, QUANTUM]
EXP. 190	symb	$[[cos\alpha,\;-1],[1,\;-sin\alpha]]\;[cos\alpha;\;sin\alpha]=[[1,0],[0,-1]]\;(=INV.450)$	[ROT]
EXP. 191	symb	$[[cos\alpha,\;sin\alpha],[sin\alpha,\;-cos\alpha]][x;\;y]=[[1,0],[0,-1]]\;(=INV.449)$	[ROT]
EXP. 192	symb	$\lim_{\theta\rightarrow 0}[\theta\rightarrow 0](1/\theta)[sin\theta;\cos\theta][[0,1],[1,0]][cos\theta;\sin\theta]=[[1,0],[0,-1]];\;f^{\frown}(-1)\;x\;f(x)=1\;(=INV.218)$	[ROT]
EXP. 193	differs	$(\log x)'=1/x,\;x^{\frown}n+y^{\frown}n=z^{\frown}n,\;x^{\frown}n=-y^{\frown}n+c,\;n\;x^{\frown}(n-1)=-n\;y^{\frown}(n-1)\;y',\;y'=n\;x^{\frown}(n-1)/(n\;y^{\frown}(n-1))=x^{\frown}(n-1)/y^{\frown}(n-1)=-(y/x)\cdot(x/y)^{\frown}n\quad=0.25\;vs\;-0.25$	[OTHER]
EXP. 194	differs	$-\cos x/((\cos x)'\;(\sin x)')=z^{\frown}n,\;z^{\frown}n=-2\;e^{\frown}(x\log x)\quad=1.1884\;vs\;-2$	[ROT]
EXP. 195	symb	$\lim_{x\rightarrow\infty}f(x)=a,\;\lim_{y\rightarrow\infty}f(y)=b,\;\lim_{x,y\rightarrow\infty}\{f(x)+f(y)\}=a+b$	[OTHER]
EXP. 196	symb	$\lim_{z\rightarrow 0}f(z)=c,\;\delta\int\;z^{\frown}n=(d/dV)\;x^3;\;\lim_{x\rightarrow\infty}=c-\lim f(y)\cdot\lim f(x)+\lim f(y)=\lim_{z\rightarrow\infty}f(z)$	[OTHER]
EXP. 197	holds	$z^{\frown}n=\cos n\theta+i\sin n\theta=-2\;e^{\frown}(x\log x)\quad=0.362358+0.932039i\;vs\;0.362358+0.932039i$	[ROT]
EXP. 198	symb	$(d/d\sigma)[\;(\sigma1+\sigma2)/2\;]=\sigma(1\downarrow)+\sigma(\uparrow\uparrow)+\sigma(\mathbb{L})+\sigma(\Leftrightarrow)+\sigma(\Rightarrow)$	[QUANTUM]
EXP. 199	symb	$(d/df)\;\int_1(x\log x)^2\;dx^m=\sigma(t);\;(d/df)\;\int_1(y\log y)^{\frown}(1/2)\;dy^m=\sigma(l)$	[QUANTUM, ENTROPY]
EXP. 200	symb	$\sigma(\leftrightarrow)+\sigma(\rightharpoonup)=\int\;e^{\frown}(-f)\;[-\Delta v+R_{ij}\;v_{ij}+\nabla_i\nabla_j\;v+v\nabla_i\nabla_j+2<f,h>+(R+\nabla f)(v-h/2)\;]$	[QUANTUM, ENTROPY]
EXP. 201	symb	$\sigma(\downarrow)=\sigma(t\downarrow+\uparrow\uparrow+\Rightarrow),\;\sigma(1)=\sigma(t\downarrow+\mathbb{L}+\Leftrightarrow),\;weak\;electric\;theorem=\sigma(1),\;strong\;electric\;theorem=\sigma(\downarrow)$	[QUANTUM]
EXP. 202	symb	$C=\int_1x^s\;dx-\log x=\int\!(\int_1x^s\;dx-\log x)\;dvol=\int_1x^s\;dvol-\int\log x\;dvol$	[ZETA]
EXP. 203	symb	$=(d/df)\;\int_1(y\log y)^{\frown}(1/2)\;dy^m-\int\;e^{\frown}(-x)\;x^{\frown}(1-t)\log x\;dx^m;\;\int\;x^{\frown}(1-t)\;e^{\frown}(-x)\;dV=\int\;x^{\frown}(1-t)\;dm=\int\;x^{\frown}(1-t)\;dvol$ $;f=\gamma=\Gamma'=\int\;e^{\frown}(-x)\;x^{\frown}(1-t)\log x\;dx=(d/d\gamma)\Gamma^{\frown}(-1)-(\gamma)\gamma'=e^{\frown}(-f)-e^{\frown}f\;(=EXP.65-67)$	[GAMMA]
EXP. 204	symb	$e^{\frown}(-f)-e^{\frown}f=(d/df)\int_1(y\log y)^{\frown}(1/2)\;dy^m-(d/df)\int_1(x\log x)^2\;dx^m=2\cos(i\;x\log x)$	[ROT, ENTROPY]
EXP. 205	symb	$(d/df)\int_1(y\log y)^{\frown}(1/2)\;dy^m+(d/df)\int_1(x\log x)^2\;dx^m=(d/df)F=2i\sin(i\;x\log x)$	[ROT, ENTROPY]
EXP. 206	symb	$(d/df)\;F+\int C\;dx^m=2(\cos(i\;x\log x)+i\sin(i\;x\log x))=2\;e^{\frown}(-\theta)=2\;e^{\frown}(-i\;x\log x)=(d/dt)\;g_{ij}(t)=-2R_{ij}$	[ROT, MANIFOLD, ENTROPY]
EXP. 207	holds	$\log(x\log y)\geq 2\sqrt{\frown}(y\log y),\;\log(x\log x)=\log x+\log\log x\quad=1.19266\;vs\;1.19266$	[ENTROPY]
EXP. 208	symb	$\nabla\phi^2=8\pi G(\;\rho/c^3+V/S\;);\;\nabla\phi^2=8\pi G\hbar+8\pi\;V/S$	[SR, QUANTUM]
EXP. 209	calc	$\square v=2\sqrt{\frown}(2\pi G\hbar)+2\sqrt{\frown}(2\pi\;V/S)\quad=2.89441$	[QUANTUM]
EXP. 210	symb	$\sqrt{\frown}(2\pi T)=2\sqrt{\frown}(2\pi\;V/S)$	[OTHER]
EXP. 211	holds	$2\cos(i\;x\log x)+2i\sin(i\;x\log x)=2\;e^{\frown}(-f);\;2\sqrt{\frown}(2\pi\;V/S)=(d/df)\;\int_1(x\log x)\;dx^m\quad=0.5+0i\;vs\;0.5$	[ROT, ENTROPY]
EXP. 212	symb	$e^{\frown}(-2\pi \phi)[\eta_{\mu\nu}+\hbar x]dx^\mu\;dx^\nu+T^2\;d^2\phi=e^{\frown}(-f)+e^{\frown}f=(u+d)+c,\;(w+s)+b=e^{\frown}(-f)+e^{\frown}f$	[MANIFOLD, QUANTUM]
EXP. 213	symb	$\square=(atom)\rightarrow[\eta_{\mu\nu}+\hbar(x)]\;dx^\mu\;dx^\nu/\;e^{\frown}(2\pi T \phi);\; ds^2 =e^{\frown}(2\pi T \phi)([\eta_{\mu\nu}+\hbar(x)]\;dx^\mu\;dx^\nu)^{\frown}(-1)+(T^2$ $d^2\phi)^{\frown}(-1);\; ds^2 =e^{\frown}(-2\pi T \phi)[\eta_{\mu\nu}+\hbar(x)]\;dx^\mu\;dx^\nu+T^2\;d^2\phi$	[MANIFOLD]
EXP. 214	symb	$(u+d)+c=e^{\frown}(-f)+e^{\frown}f,\;(w+s)+b=e^{\frown}(-f)+e^{\frown}f;\;R'_{ij}=-R_{ij}\;(\times 2)$	[ROT, QUANTUM]
EXP. 215	symb	$ ds^2 =\square+\rho,\; ds^2 =\nabla\Psi^2+\phi/\;(\eta_{\mu\nu}+\hbar(x))$	[MANIFOLD]
EXP. 216	symb	$ploximetry=\Delta x\;\Delta p-\Delta p\;\Delta x+\delta(p,\;x)$	[QUANTUM]
EXP. 217	symb	$ ds^2 =e^{\frown}(-2\pi T \phi)+[\eta_{\mu\nu}+\hbar(x)]\;dx^\mu\;dx^\nu+T^2\;d^2\phi$	[MANIFOLD]
EXP. 218	calc	$(d/df)\;F=e^{\frown}f+e^{\frown}(-f),\;(d/df)\;\int C\;dx^m=e^{\frown}f-e^{\frown}(-f)\quad=3.08616$	[ZETA]
EXP. 219	symb	$(u+d)+c=e^{\frown}(-f)-e^{\frown}f,\;b+(w+s)=e^{\frown}(-f)+e^{\frown}f$	[QUANTUM]

4. 全方程式レジストリ (2111 本) p.32/33

EXP.220	symb	$\int\!\! \int_{-1} x^s \, \mathrm{dvol} - \int \log x \, \mathrm{dvol} = (d/d\mathfrak{f}) \int\!\! \int_{-1} (y \log y)^{\wedge(1/2)} \, \mathrm{d}\mathfrak{y}^{\mathfrak{m}} - \int \mathbb{C} \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = \mathfrak{e}^{\wedge}(-\mathfrak{f}) - \mathfrak{e}^{\mathfrak{f}}$	[ZETA, ENTROPY]
EXP.221	symb	$(d/d\mathfrak{f}) \, \mathfrak{F} = (d/d\mathfrak{f}) \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} + (d/d\mathfrak{f}) \int\!\! \int_{-1} (y \log y)^{\wedge(1/2)} \, \mathrm{d}\mathfrak{y}^{\mathfrak{m}} = \mathfrak{e}^{\mathfrak{f}} + \mathfrak{e}^{\wedge}(-\mathfrak{f})$ (Jones; weak electric)	[JONES, ENTROPY]
EXP.222	symb	$ \mathrm{ds}^2 = \mathfrak{e}^{\wedge}(-2\pi\mathbb{T} \phi)[\eta + \mathfrak{h}(x)] \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \, \mathrm{d}\mathfrak{x}^{\mathfrak{v}} + \mathbb{T}^2 \, \mathrm{d}^2 \phi$	[MANIFOLD]
EXP.223	calc	$\square = 2(\cos(\mathfrak{i} \, x \log x) + \mathfrak{i} \sin(\mathfrak{i} \, x \log x)) = 0.5 + 0\mathfrak{i}$	[ROT]
EXP.224	symb	$\pi(\chi, x) = [\mathfrak{i} \pi(\chi, x), \mathfrak{f}(x)]$; $\int_{-1} (x \log x) \, \mathrm{d}\mathfrak{x} = \mathfrak{i} \int_{-1} (x \log x) \, \mathrm{d}\mathfrak{x} \, \mathbb{N} + \mathbb{N} \mathfrak{i} \int_{-1} (x \log x) \, \mathrm{d}\mathfrak{x}$	[QUANTUM, ENTROPY]
EXP.225	symb	$\int_{-1} (x \log x) \, \mathrm{d}\mathfrak{x} = \mathfrak{i} \int x \log x \, \mathrm{d}\mathfrak{x} - \int_{-1} (x \log x) \, \mathrm{d}\mathfrak{x} \, (=INV.29)$; $\int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = \mathfrak{i} \, (1/2) \, x^2 = \mathfrak{i} \int\!\! \int_{-M} \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \leq 1/(2\mathfrak{i}) + x^2 \, (=INV.518)$	[ENTROPY]
EXP.226	calc	$\mathfrak{E} = -(1/2) \, \mathfrak{m} \, \mathfrak{v}^2 + \mathfrak{m} \, \mathfrak{c}^2$; $\lim_{\{x \rightarrow \infty\}} \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \geq 1/(2\mathfrak{i})$; $(d/d\mathfrak{f}) \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = 1/(2\mathfrak{i}) = \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \, (=INV.519/520) \, (\times 2) = 8.98255\mathfrak{e}+16$	[SR, ENTROPY]
EXP.227	differs	$(\mathfrak{f})' / ((\mathfrak{f})(\mathfrak{f})') = (\mathfrak{f} \, \mathfrak{f})' = \mathfrak{e}^{\wedge}(x \log x) = 6.77259 \, \mathrm{vs} \, 4$	[ENTROPY]
EXP.228	symb	$\Gamma = \int \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(1-t) \, \mathrm{d}\mathfrak{x}$, $(d/d\mathfrak{f}) \, \mathfrak{F} = \mathfrak{e}^{\mathfrak{f}}$, $\int \mathfrak{F} \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = \mathfrak{e}^{\wedge}(-\mathfrak{f})$, $\int x^{\wedge}(1-t) \, \mathrm{d}\mathfrak{x} = (d/d\mathfrak{f}) \, \mathfrak{F}$, $\int \mathfrak{e}^{\wedge}(-x) \, \mathrm{d}\mathfrak{x} = \int \mathfrak{F} \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}}$	[GAMMA]
EXP.229	symb	$x \perp y \rightarrow x \cdot y = \vec{0}$; $(d/d\mathfrak{f}) \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = 1/(2\mathfrak{i})$; $(\mathfrak{f})'$	[OTHER]
EXP.230	differs	$1/2 + 1/(2\mathfrak{i}) = \vec{0}$, $(1/2) \cdot (1/(2\mathfrak{i}))$; $A + B = \vec{0}$, $A \cdot B = 1/(4\mathfrak{i})$, $\tan 90 \neq 0$, $ \mathrm{ds}^2 = A + B = 0.5 - 0.5\mathfrak{i} \, \mathrm{vs} \, 0 + 0\mathfrak{i}$	[ZETA]
EXP.231	symb	$\sin 0 = 0$, $\mathfrak{e}^{\wedge}(\mathfrak{i} \, \theta) = \cos \theta + \mathfrak{i} \sin \theta$	[ROT]
EXP.232	symb	$(d/d\mathfrak{f}) \, \mathfrak{F} = (d/d\mathfrak{f}) \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} + (d/d\mathfrak{f}) \int\!\! \int_{-1} (y \log y)^{\wedge(1/2)} \, \mathrm{d}\mathfrak{y}^{\mathfrak{m}} = \Gamma(x, y)$	[GAMMA, ENTROPY]
EXP.233	symb	$(d/d\gamma) \, \Gamma = (\mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(1-t)) \, \gamma'$ $\rightarrow (d/d\gamma) \, \Gamma = \Gamma(\gamma)'$	[GAMMA]
EXP.234	holds	$\Gamma(\mathfrak{s}) = \int_{0^{\wedge}\infty} \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(\mathfrak{s}-1) \, \mathrm{d}\mathfrak{x}$, $\Gamma'(\mathfrak{s}) = \int_{0^{\wedge}\infty} \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(\mathfrak{s}-1) \log x \, \mathrm{d}\mathfrak{x} \, (\times 2) = -0.577212 \, \mathrm{vs} \, -0.577216$	[GAMMA]
EXP.235	holds	$x^{\wedge}(\mathfrak{s}-1) = \mathfrak{e}^{\wedge}((\mathfrak{s}-1) \log x)$; $(\partial/\partial \mathfrak{s}) \, x^{\wedge}(\mathfrak{s}-1) = (\partial/\partial \mathfrak{s}) \, \mathfrak{e}^{\wedge}(-x) \cdot (\mathfrak{e}^{\wedge}((\mathfrak{s}-1) \log x)) = \mathfrak{e}^{\wedge}(-x) \cdot \log x \cdot \mathfrak{e}^{\wedge}((\mathfrak{s}-1) \log x) = \mathfrak{e}^{\wedge}(-x) \cdot \log x \cdot x^{\wedge}(\mathfrak{s}-1) = 0.375229 \, \mathrm{vs} \, 0.375229$	[GAMMA]
EXP.236	symb	$(d/d\gamma) \, \Gamma = \Gamma(\gamma)'$; $(d/d\gamma) \, \Gamma(\mathfrak{s}) = \int_{0^{\wedge}\infty} \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(\mathfrak{s}-1) \, \mathrm{d}\mathfrak{x} / ((\int_{0^{\wedge}\infty} \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(\mathfrak{s}-1) \log x \, \mathrm{d}\mathfrak{x})' = \Gamma((\int \log x \, \mathrm{d}\mathfrak{x})' = \mathfrak{e}^{\wedge}(-x \log x)$	[GAMMA]
EXP.237	symb	$(d/d\mathfrak{f}) \, \mathfrak{F} = \int x^{\wedge}(\mathfrak{s}-1) \, \mathrm{d}\mathfrak{x}$, $\int \mathfrak{F} \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = \int \mathfrak{e}^{\wedge}(-x) \, \mathrm{d}\mathfrak{x}$, $(d/d\mathfrak{f}) \, \mathfrak{F} = \mathfrak{F}(\mathfrak{f})'$, $\int \mathfrak{F} \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} = \mathfrak{F}(\mathfrak{f})$	[GAMMA]
EXP.238	symb	$\mathbb{H}\Psi = \otimes (\mathfrak{i} \hbar \nabla) \otimes \mathbb{L} = \otimes \mathbb{H}\Psi/\nabla \mathbb{L} = \mathfrak{e}^{\wedge}(x \log x) = x^{\wedge}(x)'$; $(d/d\mathfrak{f}) \mathfrak{F} = \mathfrak{m}(x)$ (=EXP.94)	[QUANTUM]
EXP.239	symb	$(d/d\mathfrak{t}) \, \phi(\mathfrak{t}) = \hbar = (1/(2\mathfrak{i})) \, \mathfrak{e}^{\wedge}(\mathfrak{i} \hbar) \dots$ (=EXP.80-87 block repeated, 3rd copy)	[QUANTUM]
EXP.240	symb	$(d/d\gamma) \Gamma = \mathfrak{m}(x) = \mathfrak{e}^{\wedge}(-x \log x)$; $\sin(\mathfrak{i} x) = (\mathfrak{e}^{\wedge}(-x) + \mathfrak{e}^{\mathfrak{x}})/(2\mathfrak{i})$; $(d/d\mathfrak{f}) \mathfrak{F} = \mathfrak{m}(x) = \mathfrak{e}^{\mathfrak{f}} + \mathfrak{e}^{\wedge}(-\mathfrak{f}) = 2\mathfrak{i} \sin(\mathfrak{i} \, x \log x)$ (=EXP.96-98, 2nd copy)	[ROT, GAMMA]
EXP.241	symb	$\mathbb{H}\Psi = \otimes (\mathfrak{i} \hbar \nabla) \otimes \mathbb{L} = \mathfrak{i} \hbar \phi \, (1)$	[QUANTUM]
EXP.242	symb	$(d/d\mathfrak{f}) \, \mathfrak{F} = \mathfrak{m}(x) \, (2)$	[MANIFOLD]
EXP.243	symb	$= (d/d\mathfrak{f}) \int\!\! \int_{-1} (x \log x)^2 \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} + (d/d\mathfrak{f}) \int\!\! \int_{-1} (y \log y)^{\wedge(1/2)} \, \mathrm{d}\mathfrak{y}^{\mathfrak{m}} \, (3)$	[ENTROPY]
EXP.244	symb	$\otimes (\mathfrak{i} \hbar \nabla) \otimes \mathbb{L} = \int \mathfrak{e}^{\wedge}(-x) \, x^{\wedge}(1-t) \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \, (4)$	[GAMMA, QUANTUM]
EXP.245	symb	$= (d/d\gamma) \, \Gamma = \int \Gamma(\gamma)' \, \mathrm{d}\mathfrak{x}^{\mathfrak{m}} \, (5)$	[GAMMA]
EXP.246	calc	$= \mathfrak{e}^{\mathfrak{f}} + \mathfrak{e}^{\wedge}(-\mathfrak{f}) \, (6) = 3.08616$	[GAMMA]
EXP.247	calc	$(d/d\gamma) \, \Gamma^{\wedge}(-1) = \mathfrak{e}^{\mathfrak{f}} - \mathfrak{e}^{\wedge}(-\mathfrak{f}) \, (7) = 2.3504$	[GAMMA]
S.1	calc	$\mathfrak{a} = \mathrm{softmax}(\mathfrak{z}) = 0.379641$	[QUANTUM, ENTROPY]
S.2	calc	$\phi_{\mathfrak{i}} = \sqrt{\mathfrak{a}_{\mathfrak{i}}} \cdot \exp(\mathfrak{i} \cdot \sum_{\mathfrak{m}} \beta_{\mathfrak{m}} \mathbb{H}_{\mathfrak{m}} \phi_{\mathfrak{m}}^{\wedge}(\mathfrak{i})) = -0.499807 + 0.360324\mathfrak{i}$	[QUANTUM, ENTROPY]
S.3	calc	$\hat{\mathfrak{q}}_{\mathfrak{i}} = \phi_{\mathfrak{i}} ^2 / \sum_{\mathfrak{j}} \phi_{\mathfrak{j}} ^2 = 0.379641$	[QUANTUM]
S.4	symb	$ \mathfrak{e}^{\wedge}(\mathfrak{i} \, \theta) = 1 \Rightarrow \phi_{\mathfrak{i}} ^2 = \mathfrak{a}_{\mathfrak{i}}$ (exactly)	[QUANTUM]
S.5	symb	$\mathfrak{a}_{\mathfrak{i}} = 0 \Rightarrow \hat{\mathfrak{q}}_{\mathfrak{i}} = 0$	[QUANTUM]
S.6	holds	$\max_{\mathfrak{i}} \hat{\mathfrak{q}}_{\mathfrak{i}} - \mathfrak{a}_{\mathfrak{i}} = 5.55\mathfrak{e}-17 = 5.55112 \, \mathrm{vs} \, 5.55$	[QUANTUM]
S.7	holds	$\mathbb{H}(\mathrm{unknown_prior}(8)) = 2.0794 \rightarrow 1.9733 \rightarrow 1.8364 = 5.88912 \, \mathrm{vs} \, 5.8891$	[ENTROPY]
S.8	calc	$-\mathfrak{p} \log \mathfrak{p} = 0.346574$	[ENTROPY]
S.9	symb	$\mathfrak{e}^{\wedge}\{\mathfrak{i} \, \theta\}$	[QUANTUM]
S.10	holds	$\mathrm{unknown_prior}(\mathbb{V})_{\mathfrak{i}} = 1/\mathbb{V} = 2.07944 \, \mathrm{vs} \, 2.07944$	[ENTROPY]
S.11	symb	$\mathrm{STAR}^{\mathfrak{v}} x = \mathrm{true}; x^{\mathfrak{v}} y = (x == y) \, \mathrm{for} \, \mathrm{num}/\mathrm{str}, \, \mathrm{else} \, (\mathrm{type}(x) == \mathrm{type}(y))$	[OTHER]
S.12	holds	$\mathrm{softmax}(\mathfrak{z})_{\mathfrak{i}} = \exp(\mathfrak{z}_{\mathfrak{i}} - \max_{\mathfrak{j}} \mathfrak{z}_{\mathfrak{j}}) / \sum_{\mathfrak{k}} \exp(\mathfrak{z}_{\mathfrak{k}} - \max_{\mathfrak{j}} \mathfrak{z}_{\mathfrak{j}}) = 0.643914 \, \mathrm{vs} \, 0.643914$	[ENTROPY]
S.13	holds	$\mathfrak{z}_{\mathfrak{i}} - \max_{\mathfrak{j}} \mathfrak{z}_{\mathfrak{j}} < -30 \Rightarrow \mathfrak{p}_{\mathfrak{i}} := 0$, then $\mathfrak{p} \leftarrow \mathfrak{p}/\sum \mathfrak{p} = 0 \, \mathrm{vs} \, 0$	[ENTROPY]
S.14	calc	$\mathbb{H}(\mathfrak{p}) = -\sum_{\mathfrak{i}} \mathfrak{q}_{\mathfrak{i}} \ln \mathfrak{q}_{\mathfrak{i}}$, $\mathfrak{q}_{\mathfrak{i}} = \max(\mathfrak{p}_{\mathfrak{i}}, 1\mathfrak{e}-12) = 1.59036$	[ENTROPY]
S.15	calc	$\mathrm{update}(\mathfrak{p}, \mathfrak{e}, \mathfrak{lr})_{\mathfrak{i}} = ((1-\mathfrak{lr}) \mathfrak{p}_{\mathfrak{i}} + \mathfrak{lr} \mathfrak{e}_{\mathfrak{i}}) / \sum_{\mathfrak{k}} ((1-\mathfrak{lr}) \mathfrak{p}_{\mathfrak{k}} + \mathfrak{lr} \mathfrak{e}_{\mathfrak{k}})$, $\mathfrak{lr}$ default 0.5 = 0.25232	[ENTROPY]
S.16	holds	$\mathfrak{d}_{\mathfrak{i}} = \sum_{\mathfrak{j}} \mathbb{W}_{\mathfrak{ij}} \, \mathfrak{x}_{\mathfrak{j}}$ (or $\mathfrak{d}_{\mathfrak{i}} = \mathfrak{x}_{\mathfrak{i}}$ if no $\mathbb{W}$); $\mathfrak{d} \leftarrow \mathfrak{d} / (\ \mathfrak{d}\ _2 + 1\mathfrak{e}-12) = 1 \, \mathrm{vs} \, 1$	[MANIFOLD]
S.17	calc	$\theta_{\mathfrak{i}} = \sum_{\mathfrak{m}} \beta_{\mathfrak{m}} \cdot \mathbb{H}(\mathfrak{q}_{\mathfrak{m}}) \cdot \phi_{\mathfrak{m}}[\mathfrak{i}]$, $\beta_{\mathfrak{m}}$ default 1, $\phi_{\mathfrak{m}}[\mathfrak{i}]$ default 0 = 2.51696	[QUANTUM, ENTROPY]
S.18	holds	$\phi_{\mathfrak{i}} = \sqrt{\mathfrak{a}_{\mathfrak{i}}} \cdot \mathrm{cexp}(\mathfrak{i} \, \theta_{\mathfrak{i}})$; $ \phi_{\mathfrak{i}} ^2 = \mathrm{Re}(\phi_{\mathfrak{i}})^2 + \mathrm{Im}(\phi_{\mathfrak{i}})^2 = 0.37964 \, \mathrm{vs} \, 0.37964$	[QUANTUM]
S.19	calc	$\mathfrak{p}_{\mathfrak{i}} \leftarrow \mathfrak{p}_{\mathfrak{i}} / \sum_{\mathfrak{k}} \mathfrak{p}_{\mathfrak{k}}$ (if $\sum_{\mathfrak{k}} \mathfrak{p}_{\mathfrak{k}} > 0$) = 0.4	[OTHER]
S.20	calc	$\mathrm{maxdiff}(\mathfrak{a}, \mathfrak{b}) = \max_{\mathfrak{i}} \mathfrak{a}_{\mathfrak{i}} - \mathfrak{b}_{\mathfrak{i}} = 0.2$	[OTHER]
S.21	calc	$\mathrm{zeros_of}(\mathfrak{d}) = \{\mathfrak{i} : \mathfrak{d}_{\mathfrak{i}} = 0\} = 2$	[OTHER]
S.22	holds	$\mathfrak{z} = [1.6, 0.6, 0.1, -40, -0.4, 1.1, -40, 0.4]$, $\mathbb{V} = \mathrm{len}(\mathfrak{a}) = 8$, $\mathrm{zeros_of}(\mathfrak{a}) = \mathrm{zeros_of}(\hat{\mathfrak{q}}) = [3, 6] = 9.37964 \, \mathrm{vs} \, 9.37964$	[ENTROPY]
S.23	holds	$\mathrm{modules} = [[\mathfrak{q}_{\mathrm{ret}}, \phi_{\mathrm{ret}}], [\mathfrak{q}_{\mathrm{syn}}, \phi_{\mathrm{syn}}], \mathrm{uniform}(8), 0.5\text{--}1]]$, $\beta = [0.7, 0.5, 0.9] = 2.07944 \, \mathrm{vs} \, 2.07944$	[QUANTUM, ENTROPY]
S.24	symb	$\theta_{\mathfrak{i}} = \mathfrak{c}$ (constant) $\Rightarrow \hat{\mathfrak{q}}$ unchanged	[QUANTUM, ENTROPY]
S.25	symb	$\mathfrak{a}/\mathfrak{b} = 0$ if $\mathfrak{b} = 0$; $\mathfrak{a} \% \mathfrak{b} = \mathrm{fmod}(\mathfrak{a}, \mathfrak{b})$, 0 if $\mathfrak{b} = 0$	[OTHER]
S.26	holds	$\mathrm{softmax}([2, 1, 0, -1]) \approx [0.64391, 0.23688, 0.08714, 0.03206]$; $\mathbb{H}(\mathrm{unknown_prior}(4)) = \ln 4 = 1.38629 = 1.92417 \, \mathrm{vs} \, 1.92417$	[ENTROPY]
Q.1	calc	$\mathfrak{a}_{\mathfrak{i}} = \mathrm{softmax}(\mathfrak{z})_{\mathfrak{i}} = 0.643914$	[QUANTUM, ENTROPY]
Q.2	calc	$\theta_{\mathfrak{i}} = \sum_{\mathfrak{m}} \beta_{\mathfrak{m}} \cdot \mathbb{H}_{\mathfrak{m}} \cdot \phi_{\mathfrak{m}}^{\wedge}(\mathfrak{i}) = 1.94637$	[QUANTUM, ENTROPY]
Q.3	calc	$\phi_{\mathfrak{i}} = \sqrt{\mathfrak{a}_{\mathfrak{i}}} \cdot \exp(\mathfrak{i} \cdot \theta_{\mathfrak{i}}) = 0.5 + 0.5\mathfrak{i}$	[QUANTUM]
Q.4	holds	$\mathfrak{q}_{\mathfrak{i}} = \phi_{\mathfrak{i}} ^2 / \sum_{\mathfrak{j}} \phi_{\mathfrak{j}} ^2 = \mathfrak{a}_{\mathfrak{i}} = 2.77556\mathfrak{e}-17 \, \mathrm{vs} \, 0$	[QUANTUM]
Q.5	symb	$ \exp(\mathfrak{i} \, \theta) = 1$	[QUANTUM]
Q.6	symb	$ \phi ^2 = \mathfrak{a}$	[QUANTUM]

4. 全方程式レジストリ（2111 本） p.33/33

Q.7	symb	$a_i = 0 \Rightarrow q_i = 0$	[QUANTUM]
Q.8	calc	$\alpha_i = \sqrt{p_i} \cdot e^{\{i \phi_i\}} = 0.25 + 0.433013i$	[QUANTUM]
Q.9	symb	$p_i \leftrightarrow a_i, \phi_i \leftrightarrow \theta_i, \alpha_i \leftrightarrow \phi_i$	[QUANTUM]
Q.10	holds	$P(\text{outcome } i) = \alpha_i ^2 = \phi_i ^2 = 0.25 \text{ vs } 0.25$	[QUANTUM]
Q.11	calc	$V = 2^n = 8$	[QUANTUM]
Q.12	holds	$H 0\rangle : a = [0.5, 0.5] = 0.5 \text{ vs } 0.5$	[QUANTUM]
Q.13	symb	$H \equiv \text{cognitive_system}(a, [[a, \phi_H]], [\beta_H]), a \text{ uniform}$	[QUANTUM]
Q.14	symb	$S, T, R_z(\lambda): \theta \leftarrow \theta + \Delta \theta, a \text{ unchanged}$	[QUANTUM]
Q.15	symb	$\text{CNOT}: (a, \theta)_i \leftarrow (a, \theta)_{\pi_c(i)}$	[QUANTUM]
Q.16	holds	$\max_i q_i - a_i = 1.11 \times 10^{-16} = 1.11022 \text{ vs } 1.11$	[QUANTUM]
Q.17	holds	$\text{amplitudes}^2 = [0.62246, 0.00000, 0.37754], \text{zeros} = [1] = 1.62246 \text{ vs } 1.62246$	[QUANTUM]
Q.18	symb	$\text{rule} := [\text{"rule"}, \text{name}, \text{head}, \text{pattern}, \text{denotation}], \text{head} \in \{\text{stmt}, \text{expr}, \text{postfix}\}$	[OTHER]
Q.19	holds	$\text{unknown_prior}(V)_i = 1/V; \text{prepare_unknown}(n) = \text{uniform over } 2^n = 0.125 \text{ vs } 0.125$	[QUANTUM, ENTROPY]
Q.20	symb	$H(\text{prior}_{\{k+1\}}) < H(\text{prior}_k)$	[ENTROPY]
Q.21	symb	$\text{estimate}() = \text{fold}(\text{update}, \text{unknown_prior}(V), \text{trace}); \text{forbidden}() = \text{zeros_of}(\text{estimate}())$	[QUANTUM, ENTROPY]