

Topology theorem from string theory and duality element of differential structure

Masaaki Yamaguchi

$$\begin{aligned}
\int d\tau &= f(x), \tau = (\alpha \circ \beta)^{-1} \cdot \alpha\beta \\
&\geq \frac{\alpha\beta}{\alpha^{-1}\beta^{-1}} \\
(\alpha \circ \beta)^{-1} &\leq \alpha^{-1} + \beta^{-1} \\
\beta(p, q) &= \frac{\Gamma(p)\Gamma(q)}{\Gamma(p+q)} \\
\frac{\int d\tau}{\int d\tau} &= \left| \frac{f}{f} \frac{d\tau}{d\tau} \right| d\psi = \left(\frac{\nabla_i \nabla_j f}{\Delta f} \right)^{-1} \\
(f(x)g(x))' &= f'(x)g(x) + f(x)g'(x) \\
\int (f(x)g(x))' &= \int f'(x)g(x) + \int f(x)g'(x) = \int f(x)g(x) = [f(x)g(x)] - \int f'(x)g(x) \\
[f(x), g(x)] &= f(x)g(x) - g(x)f(x) \\
(fg)' &= fg' + g'f, \left(\frac{f}{g} \right)' = \frac{f'g - g'f}{g^2} \\
\frac{\{f, g\}}{[f, g]} &= \frac{(fg)' \otimes dx_{fg}}{\left(\frac{f}{g} \right)' \otimes g^{-2} dx_{fg}} \\
\left(\frac{\{f, g\}}{[f, g]} \right) &= (i)'
\end{aligned}$$

これらの方程式は、フロベニウスの定理と実数条件からの大域的微分方程式の最小単位による積分でもある。これらは、演算子同士による計算、記号論に行き着く。非可換代数方程式、Lie 代数の部分群、フェルミオンの部分群、どこから始めたらいいか。神の人刺し、神の見えざる手、フェルミオンとボソンの商代数による積と商の微分部分群による準同型写像の計算についての幾通りの算術式。

$$\begin{aligned}
V(M) &= \frac{\partial}{\partial f} \left({}^N \int [f \setminus M]^{\oplus N} \right)^{\mu\nu} dx^\mu dx^\nu \\
\bigoplus_{k=0}^{\infty} [f \setminus g] &= \vee(M \wedge N) \\
[f \setminus g] &= (F - g) \otimes \frac{1}{M}
\end{aligned}$$

$$V-E+F=2, \log \frac{\sin \theta}{i \cos \theta} = \log(\sin \theta) - \log(i \cos \theta)$$

$$=-2R_{ij}, \frac{1}{i} \tan \theta = H \psi$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$\vee \int \frac{C^+ \nabla M_m}{\Delta (M_-^+ \nabla C_-^+)} i = \exists (M_-^+ \nabla R^+)$$

$$\vee (R + \nabla_i \nabla_j f)^n = \int \frac{\wedge (R + \nabla_i \nabla_j f)^2}{\exists (R + \Delta f)^n}$$

$$\wedge (R + \nabla_i \nabla_j f)^2 = \frac{d}{df} \int \int \frac{1}{(y \log y)^{\frac{1}{2}}} dy_m$$


$$\frac{d}{dt}g_{ij}(t)=-2R_{ij}$$

$$\vee \int \wedge (R + \nabla_i \nabla_j f)^x = \frac{\wedge (R + \nabla_i \nabla_j f)^n}{\exists (R + \nabla_i \nabla_j f \circ g)}$$

$$\bigoplus f_n^m = f_1 \oplus f_2 \oplus f_3 \oplus \cdots \oplus f_n^m$$

$$\cong \wedge[f \oplus g] \cong \wedge[f \bigoplus M_1^+]$$

$$\cong \wedge[f \bigoplus M_{-}^{+}], \sum_{k=0}^{\infty} \frac{1}{n^s}$$



$$\frac{\partial}{\partial f} \left(\int \int \int \left[\left(\frac{1}{f} + \left| \left(f + g \right) + \sigma + \kappa \sigma \right| e^{\frac{1}{f}} \right) \right] d\tau \right) = G^{\mu \nu} - \frac{1}{2} \wedge g_{ij} = T^{\mu \nu}$$

$$\partial_9(X) \cong \left(\quad \right) + \left(+ \cup + \wedge \right)_{ij}$$

$$\left(\begin{array}{c} \text{diagram of two overlapping circles with a square in the intersection} \end{array} \right) = (x, y, z) \cdot (u, v, w) / \sqrt{\quad}$$

$$\left(\begin{array}{c} \text{diagram of a square with a circle around it} \end{array} \right) \cong \left[\left| \left(\begin{array}{c} \text{diagram of a circle with a dot} \end{array} \right) + \left(\begin{array}{c} \text{diagram of a circle with a dot} \end{array} \right) \right| + \sqrt{\quad} + \left(\begin{array}{c} \text{diagram of a circle with a dot} \end{array} \right) \right] \Big|_{\frac{d\tau}{dt}}$$