

State-feedback stabilization of Markov jump linear systems with randomly observed Markov states

Masaki Ogura University of Pennsylvania



Ahmet Cetinkaya Tokyo Institute of Technology

Victor M. Preciado University of Pennsylvania

Markov jump linear systems

$$x(k+1) = A_{r(k)}x(k) + B_{r(k)}u(k)$$


- ▣ r : time-homogeneous Markov chain (**Markov state**)
- ▣ Example: Networked control system



- ▣ Feedback control
 - ▣ Need to measure both the state and the Markov state


State-feedback control

- If the controller knows the Markov state:



- Well-developed theory
 - ▣ Stabilization, H^2 control, H^∞ control, ... [Costa et al. '05]
 - ▣ Linear matrix inequalities
- Can we always measure the Markov state?

Observation through filters



□ Filter models

▣ Deterministic case

- Cluster observation [Val et al. '02], Periodic observation [Cetinkaya & Hayakawa '15]


▣ Stochastic case

- Unreliable observation [Costa et al. '15]

▣ Deterministic + Stochastic?


- Periodic observation with probabilistic failures

General filter model



□ Sampler

- has its own Markov chain
- samples *when and only when* the chain hits a subset



□ Examples





$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

Sampling with
period three


$$\begin{bmatrix} p & 1-p \\ p & 1-p \end{bmatrix}$$

Sampling with
success probability p

General filter model




State-feedback control (cont'd)



- σ is **not** a Markov chain
 - ▣ cannot use the theory of Markov jump linear systems
- But, the triple (r, s, σ) **is** a Markov chain
 - ▣ **Extended Markov chain**
 - ▣ Embed the closed-loop system into **another** Markov jump linear system with a big mode space

Equivalent system



Stabilization

- Theorem Assume that the matrices $R_{\alpha,\beta,\gamma}$, G_γ , and F_γ satisfy the linear matrix inequalities

$$\begin{bmatrix} R_{\alpha,\beta,\gamma} & A_\alpha G_\gamma + B_\alpha F_\gamma \\ G_\gamma^\top A_\alpha^\top + F_\gamma^\top B_\alpha^\top & G_\gamma + G_\gamma^\top - \mathcal{D}_{\alpha,\beta,\gamma}(R) \end{bmatrix} > 0$$

$$\mathcal{D}_{\alpha,\beta,\gamma}(R) = \sum_{\alpha',\beta',\gamma'} \bar{p}_{(\alpha',\beta',\gamma'),(\alpha,\beta,\gamma)} R_{\alpha',\beta',\gamma'}$$

Define the feedback gains by

$$K_\gamma = F_\gamma G_\gamma^{-1}$$

transition prob. of
extended Markov chain

Then, $E[\|x(k)\|^2]$ converges to zero exponentially fast.

- ▣ Proof: Use the idea in cluster observation [Val et al. '02]

Example

- Markov jump linear system with 3 modes

$$A_1 = \begin{bmatrix} -0.45 & -0.3 \\ 1.2 & 0.45 \end{bmatrix}, \quad A_2 = A_3 = \begin{bmatrix} -0.7 & 0.7 \\ 0.2 & 0.8 \end{bmatrix},$$

$$B_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad B_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad B_3 = \begin{bmatrix} -1 \\ 0 \end{bmatrix}.$$


- Transition probabilities

$$\begin{bmatrix} 0.6 & 0.2 & 0.2 \\ 0.2 & 0.6 & 0.2 \\ 0.2 & 0.2 & 0.6 \end{bmatrix}$$

- Observation: Periodic ($T=4$) observation with failure probability 0.5

Numerical simulation

- Sample paths



Conclusion

- State feedback stabilization of Markov jump linear systems
 - ▣ General model of Markov-state observation
 - Extends various frameworks in the literature
 - ▣ Feedback gains via solving linear matrix inequalities