

A
Top 30 of 138 catalog genes
(— : not tested in that tissue; bold: ≥ 0.5)
Ndufs5 3.8 3.3 5.1 10.2
Rnf26 2.7 2.8 1.9 6.0
Rps2 1.1 1.4 1.1 5.4
Atp5g2 1.6 1.6 1.6 4.2
Rps18 1.4 1.4 1.4 4.5
Rpl34 1.4 1.3 1.4 4.6
Rpl3 0.9 0.8 0.9 5.8
Acp1 2.0 1.6 1.7 2.4
Carnmt1 2.5 1.4 2.0 1.8
Rpl35a 1.2 1.6 1.2 3.5
Gstp2 2.6 1.5 1.2 2.2
Gm55594 2.2 1.2 1.4 2.1
Ndufa12 2.0 1.1 1.6 2.1
Hmgn2 1.8 1.5 1.5 1.9
Rpl30 0.7 0.8 0.7 4.4
Rsph3b 1.4 1.6 1.4 1.9
Gbp6 1.8 1.6 1.5 1.4
Atp6v0c 0.9 1.2 1.3 2.6
Rnps1 1.4 1.2 1.2 1.8
Ldhb 1.9 1.2 1.1 1.4
Gm14410 1.4 1.3 1.3 1.3
Psmb5 1.3 1.1 1.2 1.4
Ppih 1.0 1.0 1.1 1.8
5830417I10Rik 0.8 1.2 1.2 1.6
Hmgb1 1.1 0.8 0.9 2.0
Hdac1 1.0 0.7 0.8 2.1
Gnpda1 0.9 1.0 1.0 1.5
Mfap1b 1.2 1.0 1.1 0.9
Hmga1 1.1 0.9 0.8 1.4
Tdg 0.9 1.2 1.1 0.9
Adipose
Hypo-thalamus
Liver
Skin
Bias (log₂FC FVB ref − log₂FC naive)
4
2
0
−2
−4
B
Catalog overview (138 genes;
point size ∝ mean bias; labelled: top 8
four-tissue genes and mean bias ≥ 3)
Mean bias across tested tissues (log₂)
Tissues with bias ≥ 0.5 (of 4)
Ndufs5
Gm25939
Gm49708
Mup17
H2-Q5
Rnf26
H2ac4
Rps2
Atp5g2
Rps18
Rpl34
Rpl3
Acp1
Biotype
Protein-coding (n = 102)
lncRNA (n = 15)
Pseudogene (n = 18)
Other (n = 3)