

デジタルドキュメント(2)

高久 雅生

2014年4月24日(木)3・4時限

本日のお品書き

- 前回の復習及び質疑
- 本日の内容
 - デジタルドキュメントとは？（続）
 - 学術分野におけるデジタルドキュメント
 - Eジャーナル
 - 最近の動向
 - ドキュメントの構成

（前回の出席カードから）

- 多くの感想コメントと質問あり：
- いくつか主要な質問を取り上げて返答します
- 紹介を希望しないコメントの場合は、その旨を記載しておいてください

前回の出席カードから (1)

- 早めにレポートの回数や時期が分かると予定が立てやすくて助かります。

レポート回数は昨年度は4回でした。
今年度も4～5回を予定しています。

- 5限が三学で行われる授業「情報科指導法」なので、4限は定時に終わっていただけると助かります。

はい、できるだけ定時までには終わるようにします。

前回の出席カードから (2)

- 配布資料は両面印刷でもよいと思う。さらに、Webサイトで公開しているなら、各自で印刷してもよい。
- レジюмеは毎回印刷したものをいただけるのでしょうか。

検討してみましたが、次回以降、印刷版の配布をやめようと思います。

→ 次回以降は、授業資料サイトを確認の上、各自で印刷してくること。

※資料は前日昼までにアップロードします。

<http://masao.jpn.org/lecture/2014/digital-document/>

前回の出席カードから (3)

- 紙が好きと言っていましたが、研究等の雑記は紙に書きますか？ それともデータで保存していますか？ データの場合、バックアップはどうしていますか。

わたし個人はデータファイルの形でメモを取っています。ちなみに、メモには a) 即時的・リアルタイムによる速記メモ, b) 読書・論文概要メモ, c) 研究ノート の3種類があり、速記メモはテキストエディタでの速記とテキストファイルへの保存とし、後者2つはブログ形式のテキストファイル(日時, リンク情報付)で保存しています。データファイルの保存は、内容に応じて、Dropbox, Github, Google Driveといった複数のストレージサービス等を使い分けています(PC上のファイルはバックアップ保存しない)。

前回の出席カードから (4)

- ドキュメントとデータは異なるものなのか、それともドキュメントはある意味データに含まれるのか？ 加工していない情報と加工済みの情報というように階層が違うのかもしれない。ドキュメントとコンテンツの関係についてはドキュメント<コンテンツというイメージがある。コンテンツというと基本的にデジタルである。
- ドキュメントとデータの違いがよく分からなかった。ドキュメントとデジタルメディアが混合していたが、とてもよく分かった。

ドキュメントとデータは、その「目的」「文脈」が異なるという話を前回にしました。もう少し詳しく述べると、

データ (Data) ← ラテン語 [dare] = “given” 「所与」
cf. 「情報」, 「知識」, 「データ」, そして「ドキュメント」

これらは互いに同じモノを指すこともあるので紛らわしいが、それぞれ違う側面を述べている。

前回の復習

- デジタルドキュメントとは？
 - あまり定まったものは無いが、デジタルメディア上で展開されるドキュメント。この授業では、作成、流通、利用を扱う。
- 広義と狭義のデジタルドキュメント
- 類縁概念との関係
 - メディア、デジタル、ドキュメント、データ、コンテンツ
 - デジタルライブラリ、電子出版
- 文脈と用途, ジャンル
- 事例：
 - 私文書、公文書
 - 学術分野の専門書、論文誌（Eブック、Eジャーナル）
 - 電子書籍


Digital Documents in Scholarly Communication

学術分野のデジタルドキュメント

学術分野のデジタルドキュメント


- 科学研究：新しい発見、知見を伝える営み
- 刊行された文書（論文）メディアを通じた相互理解と情報流通
- 実際の例
 - 雑誌論文
 - もっとも普遍的な学術文献。
 - （査読プロセス）
 - 書籍

学術情報のライフサイクル



Science

26 August 2011 | \$10



HAYABUSA

Dust from Itokawa

AAAS

『Science』誌
2011年8月26日号

<http://www.sciencemag.org/content/333/6046.cover-expansion>

Itokawa Dust Particles: A Direct Link Between S-Type Asteroids and Ordinary Chondrites

Tomoki Nakamura,^{1,*} Takaaki Noguchi,² Masahiko Tanaka,³ Michael E. Zolensky,⁴ Makoto Kimura,² Akira Tsuchiyama,⁵ Aiko Nakato,¹ Toshihiro Ogami,¹ Hatsumi Ishida,¹ Masayuki Uesugi,⁶ Toru Yada,⁶ Kei Shirai,⁶ Akio Fujimura,⁶ Ryuji Okazaki,⁷ Scott A. Sandford,⁸ Yukihiro Ishibashi,⁶ Masanao Abe,⁶ Tatsuaki Okada,⁶ Munetaka Ueno,⁶ Toshifumi Mukai,⁶ Makoto Yoshikawa,⁶ Junichiro Kawaguchi⁹

The Hayabusa spacecraft successfully recovered dust particles from the surface of near-Earth asteroid 25143 Itokawa. Synchrotron-radiation x-ray diffraction and transmission and scanning electron microscope analyses indicate that the mineralogy and mineral chemistry of the Itokawa dust particles are identical to those of thermally metamorphosed LL chondrites, consistent with spectroscopic observations made from Earth and by the Hayabusa spacecraft. Our results directly demonstrate that ordinary chondrites, the most abundant meteorites found on Earth, come from S-type asteroids. Mineral chemistry indicates that the majority of regolith surface particles suffered long-term thermal annealing and subsequent impact shock, suggesting that Itokawa is an asteroid made of reassembled pieces of the interior portions of a once larger asteroid.

The Hayabusa spacecraft arrived at S(IV)-type asteroid 25143 Itokawa (formerly 1998 SF36) in September 2005 (1). Remote-sensing measurements from the spacecraft suggest that Itokawa consists of rocks similar to LL5 and LL6 ordinary chondrites (2, 3), confirming ground-based spectral characterization (4). On 20 and 26 November 2005, the spacecraft descended to touchdown and capture dust particles from MUSES-C Regio. This area consists of dust and gravel deposits dominated by grains up to 1 cm in diameter (5). Although the sampler did not operate as planned, an elastic sampling horn impacted onto the asteroid surface, directing dust particles into the spacecraft's sample catcher device (5). The Hayabusa sample capsule successfully landed in the Woomera Prohibited Area in South Australia on 13 June 2010. Dust particles collected at the second touchdown were recovered by two methods. In one method, we used a Teflon spatula to sweep particles from about 10% of the surface of a sample catcher. In the other method, we gently tapped on the exterior of the sample catcher, causing particles to drop onto a pure silica glass slide (6).

On the Teflon spatula, we identified 1534 rocky particles by means of a field-emission scanning electron microscope. The particles have diameters ranging from 3 to 40 μm but are mostly smaller than 10 μm (7). Most Itokawa particles are angular and are probably broken pieces of larger rocks. Among the 1534 harvested rocky particles, 1087 are monomineralic, including 580 olivine particles, 126 low-Ca pyroxenes, 56 high-Ca pyroxenes, 186 feldspars (172 plagioclase and 14 K-feldspar), 113 troilites, 13 chromites, 10 Ca phosphates, and 3 Fe-Ni metal

grains. The remaining 447 particles are poly-mineralic mixtures, mainly silicates. Several other particles are silica minerals and K-bearing halite, all of uncertain origin.

Of the 40 particles removed by tapping (diameters ranging from 30 to 180 μm) that were analyzed by x-ray computed microtomography (7) and x-ray diffraction, 38 were subjected to more detailed mineralogic analysis. Backscattered electron images of selected particles are shown in Fig. 1, A to D. RA-QD02-0030 (Fig. 1A) and RA-QD02-0024 (Fig. 1B) have a platy morphology, are polymineralic, and have many mineral grains 1 to 10 μm in diameter adhering to their surfaces. Their appearance is typical of most Itokawa particles. Two particles show different morphologies. RA-QD02-0013 (Fig. 1C) has a smoother soccer-ball shape, whereas RA-QD02-0027 (Fig. 1D) consists of a large troilite crystal and smaller silicates. Particles that contain troilite or taenite as major components like RA-QD02-0027 are rare.

Mineralogical analysis of individual "tapped" particles indicates that they consist mainly of coarse [typically 10 to 50 μm in diameter (7)] crystalline silicates, the most abundant being olivine. The next most abundant minerals are low- and high-Ca pyroxene and plagioclase (fig. S6A). Low-Ca pyroxene is exclusively composed of orthopyroxene, except for RA-QD02-0060, which is dominated by low-Ca clinopyroxene (monoclinic structure was confirmed by x-ray diffraction). The degree of crystallinity of silicates differs between and within particles, particularly for plagioclase. Some particles contain chromite, chlorapatite, merrillite, and troilite up to 25 μm in size. Small inclusions (up to 10 μm) of taenite, kamacite, troilite, and




Fig. 1. (A to D) Backscattered electron (BSE) images of RA-QD02-0030 (A), RA-QD02-0024 (B), RA-QD02-0013 (C), and RA-QD02-0027 (D).

論文の実例(1)

Tomoki Nakamura, Takaaki Noguchi, Masahiko Tanaka, et al.: "Itokawa Dust Particles: A Direct Link Between S-Type Asteroids and Ordinary Chondrites". *Science*, Vol.333, No.6046, 2011, p.1113-1116.

<http://dx.doi.org/10.1126/science.1207758>

¹Department of Earth and Planetary Material Sciences, Faculty of Science, Tohoku University, Aoba, Sendai, Miyagi 980-8578, Japan. ²College of Science, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan. ³Synchrotron X-ray Station at SPring-8, National Institute for Materials Science, Sayo, Hyogo 679-5198, Japan. ⁴ARES, NASA Johnson Space Center, Houston, TX 77058, USA. ⁵Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka 560-0043, Japan. ⁶JAXA-ISAS, 3-1-1 Yoshinodai, Sagamihara, Kanagawa 229-8510, Japan. ⁷Department of Earth and Planetary Science, Faculty of Science, Kyushu University, Hakozaki, Fukuoka 812-8581, Japan. ⁸NASA Ames Research Center, Moffett Field, CA 94035, USA.

*To whom correspondence should be addressed. E-mail: tomoki@rm.tohoku.ac.jp

Article Views

> Abstract

> Full Text

> Full Text (PDF)

> Figures Only

> Supporting Online Material

> Podcast Interview

Article Tools

> Save to My Folders

> Download Citation

> Alert Me When Article is Cited

> Post to CiteULike

> E-mail This Page

> Rights & Permissions

> Commercial Reprints and E-Prints

> View PubMed Citation

Related Content

In Science Magazine

> Science News & Analysis by Kerr

> Science This Week in Science

> More Information on Related Content

Similar Articles In:

> Science Magazine

> PubMed

Search Google Scholar for:

> Articles by Nakamura, T.

> Articles by Kawaguchi,

Science 26 August 2011:

Vol. 333 no. 6046 pp. 1113-1116

DOI: 10.1126/science.1207758

REPORT

Itokawa Dust Particles: A Direct Link Between S-Type Asteroids and Ordinary Chondrites

Tomoki Nakamura^{1,*}, Takaaki Noguchi², Masahiko Tanaka³, Michael E. Zolensky⁴, Makoto Kimura², Akira Tsuchiyama⁵, Aiko Nakato¹, Toshihiro Ogami¹, Hatsumi Ishida¹, Masayuki Uesugi⁶, Toru Yada⁶, Kei Shirai⁶, Akio Fujimura⁶, Ryuji Okazaki⁷, Scott A. Sandford⁸, Yukihiro Ishibashi⁶, Masanao Abe⁶, Tatsuaki Okada⁶, Munetaka Ueno⁶, Toshifumi Mukai⁶, Makoto Yoshikawa⁶, Junichiro Kawaguchi⁶

* Author Affiliations

* To whom correspondence should be addressed. E-mail: tomoki@m.tohoku.ac.jp

ABSTRACT

The Hayabusa spacecraft successfully recovered dust particles from the surface of near-Earth asteroid 25143 Itokawa. Synchrotron-radiation x-ray diffraction and transmission and scanning electron microscope analyses indicate that the mineralogy and mineral chemistry of the Itokawa dust particles are identical to those of thermally metamorphosed LL chondrites, consistent with spectroscopic observations made from Earth and by the Hayabusa spacecraft. Our results directly demonstrate that ordinary chondrites, the most abundant meteorites found on Earth, come from S-type asteroids. Mineral chemistry indicates that the majority of regolith surface particles suffered long-term thermal annealing and subsequent impact shock, suggesting that Itokawa is an asteroid made of reassembled pieces of the interior portions of a once larger asteroid.

The Hayabusa spacecraft arrived at S(IV)-type asteroid 25143 Itokawa (formerly 1998 SF36) in September 2005 (1). Remote-sensing measurements from the spacecraft suggest that Itokawa consists of rocks similar to LL5 and LL6 ordinary chondrites (2, 3), confirming ground-based spectral characterization (4). On 20 and 26 November 2005, the spacecraft descended to touchdown and capture dust particles from MUSE-C Regio. This area consists of dust and gravel deposits dominated by grains up to 1 cm in diameter (5). Although the sampler did not operate as planned, an elastic sampling horn impacted onto the asteroid surface, directing dust particles into the spacecraft's sample catcher device (5). The Hayabusa sample capsule successfully landed in the Woomera Prohibited Area in South Australia on 13 June 2010. Dust particles collected at the second touchdown were recovered by two methods. In one method, we used a Teflon spatula to sweep particles from about 10% of the surface of a sample catcher. In

< Prev | Table of Contents | Next >

ADVERTISEMENT

Science Webinar
SERIES

Register for upcoming webinars.

Access Now!

ADVERTISEMENT

Science MOBILE

Now Available for Android Phones



The latest news from ScienceNOW, career advice articles from Science Careers

Related Resources

In Science Magazine

NEWS & ANALYSIS
PLANETARY SCIENCE

Hayabusa Gets Bottom of Decade Asteroid Cloak

Richard A. Kerr

Science 26 August 2011: 1081.

THIS WEEK IN SCIENCE

Extraterrestrial Dust Collection

Science 26 August 2011: 1067.

<http://www.sciencemag.org/content/333/6046/1113.full>



Scan the barcode to download from the Android Market.

[To Advertise](#) [Find Products](#)

論文の実例(2)

江村優花, 関洋平: マイクロブログにおける感情・コミュニケーション・動作タイプの推定に基づく顔文字の推薦. 自然言語処理. Vol.19, No.5, 2012, p.401-418.

マイクロブログにおける感情・コミュニケーション・動作タイプの推定に基づく顔文字の推薦

江村 優花^{†,††}・関 洋平^{†††}

現在, 電子メール, チャット, マイクロブログなどのメディアで, 顔文字は日常的に使用されている。顔文字は, 言語コミュニケーションで表現できない, ユーザの感情やコミュニケーションの意図を表すのに便利であるが, 反面, その種類は膨大であり, 場面に合った顔文字を選ぶことは難しい。本研究では, ユーザの顔文字選択支援を目的として, ユーザが入力したテキストに現れる感情, コミュニケーション, 動作のタイプ推定を行い, 顔文字を推薦する方法を提案する。感情, コミュニケーション, 動作のタイプは, *Twitter* から収集したコーパスを用いてカテゴリを定義し, 推定システムは, k -NN に基づき実現した。また, システムが推薦する顔文字がユーザの意図にどの程度適合しているか, 5名の被験者により評価した結果, 91件のつぶやきに対して 66.6%の顔文字が適切に推定されており, 感情カテゴリのみを用いて推薦された結果と比べて, 提案手法の顔文字推薦の精度が有意に向上していることがわかった。

キーワード: 顔文字推薦, 感情カテゴリ, コミュニケーションタイプ, 動作タイプ, マイクロブログ

Facemark Recommendation based on Emotion, Communication, and Motion Type Estimation in Microblogs

YUKA EMURA^{†,††} and YOHEI SEKI^{†††}

Many users use facemarks everyday in recent computer mediated communication environments such as e-mail, chatting, and Microblogs. Although facemarks are useful to express the emotion or communication intentions beyond natural language communication, many users feel difficult to choose the right one from lots of candidates according to the situation. We propose a method to recommend facemarks based on the estimation of emotions, communication, or motion types in texts written by users. Emotion, communication, or motion types are defined with *Twitter* corpus, and estimation system is implemented with k -NN. Five assessors evaluated the relevance of recommended facemarks for their intention, and found that 66.6% of facemarks for

[†] 筑波大学情報学群知識情報・図書館学類, College of Knowledge and Library Sciences, School of Informatics, University of Tsukuba

^{††} 現在, フコク情報システム株式会社, Presently with Fukoku Information Systems

^{†††} 筑波大学図書館情報メディア系, Faculty of Library, Information and Media Science, University of Tsukuba

https://www.jstage.jst.go.jp/article/jnlp/19/5/19_401/article/-char/ja/

マイクロブログにおける感情・コミュニケーション・動作タイプの推定に基づく顔文字の推薦

江村 優花¹⁾²⁾, 関 洋平³⁾

1) 筑波大学情報学群知識情報・図書館学類 2) 現在, フコク情報システム株式会社 3) 筑波大学図書館情報メディア系

J-STAGE公開日 20130319

キーワード: 顔文字推薦, 感情カテゴリ, コミュニケーションタイプ, 動作タイプ, マイクロブログ

 本文PDF [400K]

現在, 電子メール, チャット, マイクロブログなどのメディアで, 顔文字が用いられている。顔文字は, 言語コミュニケーションで表現できない, ユーザーの意図を表すのに便利であるが, 反面, その種類は膨大であり, 適切な顔文字を選ぶことは難しい。本研究では, ユーザーの顔文字選択支援を目的として, マイクロブログから収集した感情, コミュニケーション, 動作のタイプ推定を行い, 顔文字を推薦する方法を提案する。感情, コミュニケーション, 動作のタイプは, Twitter から収集したコーパスを用いてカテゴリを定義し, 推定システムは, k-NN に基づき実現した。また, システムが推薦する顔文字が, ユーザーの意図にどの程度適合しているか, 5名の被験者により評価

記事ツール

★ お気に入り登録

🔔 被引用アラート

🔔 認証解除アラート

🔔 追加情報アラート

📄 URLコピー

✉ 著者にメール

📄 書誌事項をダウンロード

RIS

BibTeX

[ヘルプ]

🔍 問い合わせ

この記事を共有

✉ メールで知らせる

https://www.jstage.jst.go.jp/article/jnlp/19/5/19_401/_article/-char/ja/

論文の実例(3)

薬袋秀樹: 図書館協議会の可能性—草の根からの図書館振興. 社会教育. No.792, 2012, p.20-25.

<http://www.tulips.tsukuba.ac.jp/dspace/handle/2241/117267>

ARTICLE

図書館協議会の可能性

草の根からの図書館振興

筑波大学図書館情報メディア系教授 薬袋秀樹

はじめに

近年、公立図書館における図書館協議会の活動に関するニュースを聞くことが多く、以前と比べて、図書館協議会の活動は活発化していると言われるが、課題も多い。

図書館協議会の設置の現状については、三年ごとに、文部科学省の「社会教育調査」で県別、地方公共団体の種類別に設置図書館数の調査が行われている^(注1)。

全国の図書館協議会の詳しい実態については、一九八五年に日本図書館協会^(注2)、二〇一二年に平山陽菜・池内淳(筑波大学図書館情報メディア系)が調査を行っている^(注3)。公立図書館の特定事項に関する調査で取り上げられる場合もある。

図書館協議会の現状、可能性、課題について、関係文献と筆者の協議会委員の経験をもとに論じてみたい。

一、図書館協議会設置の趣旨

(一) 現行図書館法(二〇一一年改正)

図書館協議会については、図書館法第一四―一六条で定められている。その任務は「図書館の運営に関し館長の諮問に応ずるとともに、図書館の行う図書館奉仕につき、館長に対して意見を述べる」ことである(一四条)。

図書館法制定時の解説書では、「住民の具体的な図書館に対する要望なり意見なりを、図書館奉仕を実施する責任者とも言ふべき館長に対して反映せしめるために置かれる」と書かれている^(注4)。

図書館協議会の設置は任意で(一四条)、協議会の委員は、当該図書館を設置する地方公共団体の教育委員会が任命する(二五条)。協議会の設置、委員の定数、任期等必要な事項は地方公共団体

の条例で定めなければならない(二六条)。

(二〇一一年に、地域の自主性及び自立性を高めるための改革の推進を図るための関係法律の整備に関する法律(平成二十三年法律第一〇五号)によって図書館法が改正され、「委員の任命の基準については、文部科学省令で定める基準を参酌するものとする」が付け加えられた(一六条)。

併せて、図書館法施行規則(文部科学省令)が改正され、参酌すべき基準として、「学校教育及び社会教育の関係者、家庭教育の向上に資する活動を行う者並びに学識経験のある者の中から任命することとする」と定められた(二二条)。

(二) 図書館法の改正(二〇〇八年)
二〇〇八年の図書館法改正で、第三条が次のように改正された(傍線が改

検索

[詳細検索](#)

[→ ホーム](#)

ブラウズ

[→ コミュニティ
& コレクション](#)

[→ 発行日](#)

[→ 著者](#)

[→ タイトル](#)

[→ 主題](#)

[つくばリポジトリ \(Tulips-R\) >](#)
[0 コンテンツタイプ別 \(Content type\) >](#)
[01 雑誌発表論文等 \(Journal article, etc.\) >](#)
[社会教育 \(Social education\) >](#)

このアイテムの引用には次の識別子を使用してください: <http://hdl.handle.net/2241/117267>

タイトル: 図書館協議会の可能性: 草の根からの図書館振興

著者: 葉袋, 秀樹
[Minai, Hideki](#)
[ミナイ, ヒデキ](#)

発行日: 6月-2012

出版者: 全日本社会教育連合会

誌名: 社会教育

号: 792

開始ページ: 20

終了ページ: 25

URI: <http://hdl.handle.net/2241/117267>

テキストバージョン: publisher

出現コレクション: [葉袋 秀樹 \(Minai Hideki\)](#)
[社会教育 \(Social education\)](#)

このアイテムのファイル:

ファイル	記述	サイズ	フォーマット
社会教育 792.pdf		672.23 kB	Adobe PDF 見る/開く


[このリポジトリに保管されているアイテムはすべて著作権により保護されています。](#)

■ HOME

■ 調査・研究事業


平成20年度

子どもゆめ基金助成活動




平成21年度

子どもゆめ基金助成活動



平成22年度

子どもゆめ基金助成活動



平成23年度

子どもゆめ基金助成活動



■ 今月の社会教育



■ 出版書籍案内

■ 調査・研究報告書

■ 一般書籍



CONTENTS



購読案内

前年<< 前号<

2012年 Index

>>次号 >>次年

○○○○○ ◆ 2012年6月号(792号) 780円 ◆ ○○○○○

今月のことば 「老後の初心、忘るべからず」
佛教大学教育学部 教授 白石克己

特集

社会教育における委員制度は機能しているのか

[論文]

社会教育の振興に向けた行政からの委嘱委員への期待

東京家政大学・同大学院教授 山本和人

社会教育委員の今日的役割
—活動を活性化するために—

常葉学園大学教育学部教授 上條秀元

図書館協議会の可能性
—草の根からの図書館改革—

筑波大学大学院図書館情報メディア系教授 薬袋秀樹

[提言]

各種委員を活性化するための提案
公民館運営審議会

千葉県公民館連絡協議会前顧問 朱膳寺宏一

各種委員を活性化するための提案
社会教育委員

前全国社会教育委員連合常務理事 大西康之

図書館協議会は、公共図書館を評価し
支援する友好的な委員会組織
—東京都・杉並区立図書館協議会の活性化について—

杉並区立図書館協議会副会長 渥美恵子

博物館経営の有効運用のために
～博物館協議会委員の位置づけの明確化～

長崎歴史文化博物館館長 大堀 哲

スポーツ新時代とスポーツ推進委員への期待
公益社団法人全国スポーツ推進委員連合専務理事

相模横浜大学教授・フジ・山・健康政策学部長 関山和夫

<http://www.social-edu.com/magazine/y2012/y1206.html>


教育評論家・墨田区男女共同参画推進委員 武笠和夫

社会教育委員の役割

学術分野における電子ジャーナルの歴史

- 1980年代頃から、電子出版の試み
 - → 作成プロセスの電子化
 - 当初はCD-ROM等による、テキストの提供など
- 1990年代頃から、電子版提供の試み
- 1990年代半ば～終わり頃、ウェブを通じたオンラインジャーナルの提供開始
 - はじめは紙版の代替(簡易版)
 - PDF
 - 次第に本格版へ
 - E-onlyの登場(オンライン雑誌)
- 多様化・先端化・変容へ
 - XML / HTML+ α による提供
 - Future of article (2011) の登場

電子ジャーナルの規模・変遷



出典: Ulrich's Periodicals Directory.

倉田敬子:「学術情報流通とオープンアクセス」. 勁草書房, 2007, p.117. 図5.2より。

日本国内における電子ジャーナルの提供状況

	2012年2月			2008年1月		
資料種別	雑誌総数	電子ジャーナル数 (%)		雑誌総数	電子ジャーナル数 (%)	
学術誌・学会誌	2,478	1,480	60%	2,207	1,036	47%
研究報告・技術報告	3,138	1,836	59%	3,085	1,555	50%
会議論文集	662	167	25%	614	112	18%
実用誌・解説誌	985	373	38%	917	244	27%
会議要旨集	1,494	346	23%	1,306	198	15%
その他	902	490	54%	969	413	43%
合計	9,659	4,692	49%	9,098	3,558	39%

- 国内の電子ジャーナルは発行総数の約半


出典: 佐藤正樹ほか. JST国内収録誌の電子化状況調査報告 (2012). 情報管理. 2013, Vol.56, No.2, p.93-101. <http://dx.doi.org/10.1241/johokanri.56.93>

電子ジャーナルの受容

- 1990年代終わりに始まった電子ジャーナルは急速に受け入られた。
- 2011年に国内の大学・研究機関を対象とした学術図書館研究委員会によるアンケート調査(SCREAL調査)によれば
 - 薬学、化学、生物学、物理学、医学の分野では、研究者の半数以上が「ほぼ毎日」電子ジャーナルを使う
 - 人文社会科学系の分野でも、7割以上が「月に1回以上」利用している。
- 週に1回以上電子ジャーナルを利用する研究者の割合は、2001年調査：44%→2011年調査：76%まで増加(自然科学系)

電子ジャーナル利用の浸透度(Q15)

電子ジャーナルをどの程度利用していますか
(目次の閲覧を含む)



電子ジャーナル利用の浸透度（自然科学系）

■ 週に1回以上使う ■ 月に1回以上使う ■ 使わない

0% 20% 40% 60% 80% 100%

2001年国立大学

44.2

22.3

33.5

2003年国立大学

62.3

22.4

15.5

2005年公立私立大学

43.8

17.7

38.5

2007年国内大学・研究機関

82.4

12.5

5.2

2011年国内大学・研究機関

76.1

16.1

7.8

論文の要素と構造

論文の要素、構造

- (書誌情報)
 - タイトル
 - 著者
 - 抄録(要旨)
 - キーワード
- 本文
 - 章
 - 節
 - 段落
 - » 文
- 図
- 表
- 参考文献(リスト)

REPORTS

Itokawa Dust Particles: A Direct Link Between S-Type Asteroids and Ordinary Chondrites

Tomoki Nakamura,^{1,*} Takaaki Noguchi,² Masahiko Tanaka,³ Michael E. Zolensky,⁴ Makoto Kimura,² Akira Tsuchiyama,⁵ Aiko Nakato,¹ Toshihiro Ogami,¹ Hatsumi Ishida,¹ Masayuki Uesugi,⁶ Toru Yada,⁶ Kei Shirai,⁶ Akio Fujimura,⁶ Ryuji Okazaki,⁷ Scott A. Sandford,⁸ Yukihiko Ishibashi,⁶ Masanao Abe,⁶ Tatsuaki Okada,⁶ Munetaka Ueno,⁶ Toshifumi Mukai,⁶ Nakoto Yoshikawa,⁶ Junichiro Kawaguchi⁶

The Hayabusa spacecraft successfully recovered dust particles from the surface of near-Earth asteroid 25143 Itokawa. Synchrotron-radiation x-ray diffraction and transmission and scanning electron microscope analyses indicate that the mineralogy and mineral chemistry of the Itokawa dust particles are identical to those of the thermally metamorphosed LL chondrites, consistent with spectroscopic observations made from Earth and by the Hayabusa spacecraft. Our results directly demonstrate that ordinary chondrites, the most abundant meteorites found on Earth, come from S-type asteroids. Mineral chemistry indicates that the majority of regolith surface particles suffered long-term thermal annealing and subsequent impact shock, suggesting that Itokawa is an asteroid made of reassembled pieces of the interior portions of a once larger asteroid.

The Hayabusa spacecraft arrived at (SIV) type asteroid 25143 Itokawa (formerly 1998 SF36) in September 2005 (1). Remote-sensing measurements from the spacecraft suggest that Itokawa consists of rocks similar to LL5 and LL6 ordinary chondrites (2, 3), confirming ground-based spectral characterization (4). On 20 and 26 November 2005, the spacecraft descended to touchdown and capture dust particles from MUSES-C Regio. This area consists of dust and gravel deposits dominated by grains up to 1 cm in diameter (5). Although the sampler did not operate as planned, an elastic sampling horn impacted onto the asteroid surface, directing dust particles into the spacecraft's sample catcher device (5). The Hayabusa sample capsule successfully landed in the Woomera Prohibited Area in South Australia on 13 June 2010. Dust particles collected at the second touchdown were recovered by two methods. In one method, we used a Teflon spatula to sweep particles from about 10% of the surface of a sample catcher. In the other method, we gently tapped on the exterior of the sample catcher, causing particles to drop onto a pure silica glass slide (6).

On the Teflon spatula, we identified 1534 rocky particles by means of a field-emission scanning electron microscope. The particles have diameters ranging from 3 to 40 μm but are mostly smaller than 10 μm (7). Most Itokawa particles are angular and are probably broken pieces of larger rocks. Among the 1534 harvested rocky particles, 1087 are monomineralic, including 580 olivine particles, 126 low-Ca pyroxenes, 56 high-Ca pyroxenes, 186 feldspars (172 plagioclase and 14 K-feldspar), 113 troilites, 13 chromites, 10 Ca phosphates, and 3 Fe-Ni metal

grains. The remaining 447 particles are polyminerally mixed, mainly silicates. Several other particles are silica minerals and K-bearing halite, all of uncertain origin.

Of the 40 particles removed by tapping (diameters ranging from 30 to 180 μm) that were analyzed by x-ray computed microtomography (7) and x-ray diffraction, 38 were subjected to more detailed mineralogical analysis. Backscattered electron images of selected particles are shown in Fig. 1. A to D. RA-QD02-0030 (Fig. 1A) and RA-QD02-0024 (Fig. 1B) have a platy morphology, are polyminerally mixed, and have many mineral grains 1 to 10 μm in diameter adhering to their surfaces. Their appearance is typical of most Itokawa particles. Two particles show different morphologies. RA-QD02-0013 (Fig. 1C) has a smoother soccer-ball shape, whereas RA-QD02-0027 (Fig. 1D) consists of a large troilite crystal and smaller silicates. Particles that contain troilite or taenite as major components like RA-QD02-0027 are rare.

Mineralogical analysis of individual "tapped" particles indicates that they consist mainly of coarse [typically 10 to 50 μm in diameter (7)] crystalline silicates, the most abundant being olivine. The next most abundant minerals are low- and high-Ca pyroxene and plagioclase (fig. S6A). Low-Ca pyroxene is exclusively composed of orthopyroxene, except for RA-QD02-0060, which is dominated by low-Ca clinopyroxene (monoclinic structure was confirmed by x-ray diffraction). The degree of crystallinity of silicates differs between and within particles, particularly for plagioclase. Some particles contain chromite, chlorapatite, merillite, and troilite up to 25 μm in size. Small inclusions (up to 10 μm) of taenite, kamacite, troilite, and

Fig. 1. (A to D) Backscattered electron (BSE) images of RA-QD02-0030 (A), RA-QD02-0024 (B), RA-QD02-0013 (C), and RA-QD02-0027 (D).

¹Department of Earth and Planetary Material Sciences, Faculty of Science, Tohoku University, Araba, Sendai, Miyagi 980-8578, Japan. ²College of Science, Ibaraki University, 2-1-1 Bunkyo, Mito, Ibaraki 310-8512, Japan. ³Synchrotron X-ray Station at Spring-8, National Institute for Materials Science, Sayo, Hyogo 679-5198, Japan. ⁴ARES, NASA Johnson Space Center, Houston, TX 77058, USA. ⁵Department of Earth and Space Science, Graduate School of Science, Osaka University, Toyonaka 560-0043, Japan. ⁶JAXA-ISAS, 3-1-1 Yoshinodai, Sagamihara, Kanagawa 229-8510, Japan. ⁷Department of Earth and Planetary Science, Faculty of Science, Kyushu University, Hakozaki, Fukuoka 812-8581, Japan. ⁸NASA Ames Research Center, Moffett Field, CA 94035, USA.

*To whom correspondence should be addressed. E-mail: tomoki@m.tohoku.ac.jp

www.sciencemag.org SCIENCE VOL 333 26 AUGUST 2011

Downloaded from www.sciencemag.org on April 23, 2013

論文の要素、構造(図表)

Of the 40 particles removed by tapping (diameters ranging from 30 to 180 μm) that were analyzed by x-ray computed microtomography (7) and x-ray diffraction, 38 were subjected to more detailed mineralogic analysis. Backscattered electron images of selected particles are shown in Fig. 1, A to D. 0030 (Fig. 1A) and RA-QD02-00 have a platy morphology, are polyn have many mineral grains 1 to 10 eter adhering to their surfaces. This is typical of most Itokawa particles show different morphologies. 0013 (Fig. 1C) has a smoother soc whereas RA-QD02-0027 (Fig. 1D) large troilite crystal and smaller si cles that contain troilite or taenite a onents like RA-QD02-0027 are rare

- 図表⇔本文のリンク
- 図表番号
- キャプション




Fig. 1. (A to D) Backscattered electron (BSE) images of RA-QD02-0030 (A), RA-QD02-0024 (B), RA-QD02-0013 (C), and RA-QD02-0027 (D).

論文の要素、構造(文献参照)

The Hayabusa spacecraft arrived at S(IV)-type asteroid 25143 Itokawa (formerly 1998 SF36) in September 2005 (1). Remote-sensing measurements from the spacecraft suggest that Itokawa consists of rocks similar to LL5 and LL6 ordinary chondrites (2, 3), confirming ground-based spectral characterization (4). On 20 and 26 November 2005, the spacecraft descended to touchdown at Itokawa. Particles from MUSES-C Regio of dust and gravel deposits collected up to 1 cm in diameter (5).

参考文献
(References)との
リンク

- 表
- 参考文献(リスト)

REPORTS

greatly between particles (fig. S7), which is typical of moderately shocked astromaterial corresponding shock stages up to S4 (6, 26).

MUSES-C Regio probably formed by segregation and accumulation of fine gravel into areas close to the gravitational center of Itokawa due to global-scale electrostatic grain levitation, vibration-induced granular migration, and deposition of slow moving ejecta launched from surface impacts (27–29). Therefore, particles in MUSES-C Regio originally derived from diverse regions of Itokawa. Fortunately, despite the small mass of the recovered Itokawa samples, they record the critical steps in the history of this asteroid. Itokawa was classified as an S-type asteroid from terrestrial remote sensing, and it has been commonly suggested that S-type asteroids, the most abundant asteroids in the inner asteroid belt, are the parent bodies of ordinary chondrites. Our petrologic data from MUSES-C Regio confirm that Itokawa is indeed an ordinary chondrite (LL4 to LL6), thereby finally linking these asteroids and meteorites.

References and Notes

1. A. Fujimura *et al.*, *Science* **312**, 1330 (2006).
2. M. Abe *et al.*, *Science* **312**, 1334 (2006).

3. T. Okada *et al.*, *Science* **312**, 1338 (2006).
4. R. Binzel, A. S. Rivkin, S. J. Bus, J. M. Sunshine, T. H. Burbine, *Meteorit. Planet. Sci.* **36**, 1167 (2001).
5. H. Yano *et al.*, *Science* **312**, 1350 (2006).
6. See supporting material on Science Online.
7. A. Tsuchiyama *et al.*, *Science* **333**, 1125 (2011).
8. A. J. Brearley, R. H. Jones, *Rev. Mineral.* **36**, 3–1 (1998).
9. A. E. Rubin, *Geochim. Cosmochim. Acta* **54**, 1217 (1990).
10. J. B. Brady, D. J. Cherniak, In *Diffusion in Minerals and Melts*, Y. Zhang, D. J. Cherniak, Eds. (Mineralogical Society of America, Chantilly, VA, 2010), pp. 899–920.
11. W. R. Van Schmus, J. A. Wood, *Geochim. Cosmochim. Acta* **31**, 747 (1967).
12. E. A. Jobbins *et al.*, *Mineral. Mag.* **35**, 881 (1966).
13. T. J. McCoy, E. R. D. Scott, R. H. Jones, K. Keil, G. J. Taylor, *Geochim. Cosmochim. Acta* **55**, 601 (1991).
14. G. R. Huss, A. E. Rubin, J. N. Grossman, in *Meteorites and the Early Solar System II*, D. S. Lauretta, H. Y. McSween Jr., Eds. (University of Arizona Press, Tucson, AZ, 2006), pp. 567–586.
15. T. E. Bunch, K. Keil, K. G. Snetsinger, *Geochim. Cosmochim. Acta* **31**, 1569 (1967).
16. M. Kimura, H. Nakajima, H. Hiyaogon, M. K. Weisberg, *Geochim. Cosmochim. Acta* **70**, 5634 (2006).
17. J. V. Smith, *J. Geol.* **80**, 505 (1972).
18. Y. Nakamura, Y. Motomura, *Meteorit. Planet. Sci.* **34**, 763 (1999).
19. V. Slater-Reynolds, H. Y. McSween Jr., *Meteorit. Planet. Sci.* **40**, 745 (2005).
20. D. H. Lindsay, *Am. Mineral.* **68**, 477 (1983).
21. J. Fabriès, *Contrib. Mineral. Petrol.* **69**, 329 (1979).

22. K. Ozawa, *Geochim. Cosmochim. Acta* **48**, 2597 (1984).
23. M. Miyamoto, N. Fujii, H. Takeda, *Proc. Lunar Planet. Sci.* **12B**, 1145 (1981).
24. M. Trieloff *et al.*, *Nature* **422**, 502 (2003).
25. H. Y. McSween *et al.*, in *Asteroids III*, W. Bottke *et al.*, Eds. (Univ. of Arizona Press, Tucson, AZ, 2002), pp. 559–571.
26. D. Stöffler, K. Keil, E. R. D. Scott, *Geochim. Cosmochim. Acta* **55**, 3845 (1991).
27. H. Miyamoto *et al.*, *Science* **316**, 1011 (2007).
28. P. Lee, *Icarus* **124**, 181 (1996).
29. D. G. Korycansky, E. Asphaug, *Icarus* **171**, 110 (2004).

Acknowledgments: We thank the Hayabusa project team for sample return; KEK for synchrotron experiments; H. Nakano, Y. Yamazaki, K. Shimada, Y. Kakazu, T. Hashimoto, M. Konno, Y. Katsuya, and Y. Matsushita, for technical support; and J. Grossman, T. Ikeda, T. Hokada, K. Ozawa, Y. Nakamura, and S. Wakita for helpful discussions. Supported by NASA grant 769583.07.03 (M.E.Z. and S.A.S.).

Supporting Online Material

www.sciencemag.org/cgi/content/full/333/6046/1113/DC1
Figs. S1 to S8
Tables S1 to S5
References (30–40)

2 May 2011; accepted 2 August 2011
10.1126/science.1207758

論文の要素、構造

- (書誌情報)
 - タイトル
 - 著者
 - 抄録(要旨)
 - キーワード
- 本文
 - 章
 - 節
 - 段落
 - » 文
- 図
- 表
- 参考文献(リスト)
- 付録資料
(Supplementary material)
 - 図表
 - ビデオ
 - 音声
 - ...

デジタルドキュメントならではの要素が多く出てくる



International Journal for Parasitology

Volume 40, Issue 1, April 2010, Pages 543-554



Exposed proteins of the *Schistosoma japonicum* tegument

Jason Muijenia ^{a,*}, Luke Moorlat ^a, Malcolm K. Jones ^a, Supree Nisaratna ^a, Erica M. Loran ^a, Geoffrey N. Gobert ^a, Michelle Colgrave ^a, Alan Jones ^a, Alex Loukas ^a, Donald P. McMahon ^b

^a Corresponding author. Tel.: +61 7 3945 2726; fax: +61 7 3945 2007.

^a Immunology Laboratory, Division of Infectious Diseases, Queensland Institute of Medical Research, Qld 4030, Australia

^b Molecular Parasitology Laboratory, Division of Infectious Diseases, Queensland Institute of Medical Research, Qld 4030, Australia

^c Parasite Cell Biology Laboratory, Division of Infectious Diseases, Queensland Institute of Medical Research, Qld 4030, Australia

^d The University of Queensland, School of Veterinary Sciences, Qld 4072, Australia

^e CSIRO Livestock Industries, Brisbane, Qld 4037, Australia

^f The University of Queensland, Institute for Molecular Biosciences, Qld 4072, Australia


^g These authors contributed equally to the manuscript.

Research highlights

- Proteins exposed on the surface of parasitic worms are an important source of novel drug and vaccine targets.
- These proteins are the most accessible to the host and likely to possess functions important for the survival of the worm.
- Biotinylation (labelling of lysine side-chain residues with biotin) of whole worms is a useful technique for separating exposed proteins from other protein constituents of the tegument.
- Using this technique in combination with LC-MS/MS we identified 54 proteins as putatively host-exposed in *Schistosoma japonicum*.
- Using confocal and electron microscopy, the in vivo localization of these proteins was observed.



Protein Identification v. Subcellular location



Subcellular content

proteins

5 of 54 (9.1%)

NCBI



elongation factor 1- α [*Schistosoma japonicum*]

543 aa protein

Download protein structure (PDB)



Open in new window

Subcellular Location

Cytoplasm/Nucleus

Accession

AAK37918

Version

AAK37918.1 GI 14288828

Schistosoma japonicum [Eukaryota; Metazoa; Platyhelminthes; Trematoda; Digenea; Strigidae; Schistosomatidae; Schistosomatidae; Schistosomatidae]

Similar proteins (PDB.org)

<http://www.articleofthefuture.com/>

本日のまとめ

- デジタルドキュメントとは？（続き）
- 学術分野のデジタルドキュメント
 - オンラインジャーナル論文の事例、歴史
- 次回は、学術分野のデジタルドキュメントの続き：
 - 電子ジャーナルコンテンツの作成、利用の実際
 - 論文以外の学術分野のドキュメント
 - ドキュメントフォーマット、標準規格
- Readings（参考文献）
 - 倉田敬子: 学術情報流通とオープンアクセス. 勁草書房, 2007, 196p. ISBN:978-4-326-00032-6
 - 佐藤翔: 電子リソースの普及と研究活動への影響. カレントアウェアネス. 2010, No.304, CA1720, p.17-20. <http://current.ndl.go.jp/ca1720>