

Deligne–Lusztig theory and the local Langlands correspondence

Masao Oi (National Taiwan Univ.)
j. w. with Charlotte Chan (Univ. of Michigan)

January 15, 2025
2025 TMS Annual Meeting

What is the local Langlands correspondence?

- **Langlands correspondence** (conj.) predicts a natural connection between automorphic representations and Galois representations.
- **Local Langlands correspondence** (conj.) is the local version of the Langlands correspondence.
- F : a local field of characteristic 0 (i.e., a finite extension of \mathbb{Q}_p or \mathbb{R}).
- W_F : the Weil group of F .
- G : a reductive group over F .
- \hat{G} : the Langlands dual group of G over \mathbb{C} ($\curvearrowright W_F$).

G	GL_n	U_n	SO_{2n+1}	Sp_{2n}	SO_{2n}
\hat{G}	GL_n	GL_n	Sp_{2n}	SO_{2n+1}	SO_{2n}

- $\Pi(G)$: the set of irreducible admissible representations of $G(F)$.
- $\Phi(G)$: the set of L -parameters of G ($W_F \times \mathrm{SL}_2(\mathbb{C}) \rightarrow \hat{G} \rtimes W_F$).

Local Langlands correspondence (rough form)

Local Langlands correspondence (conjectural/rough form)

There exists a “natural” map from $\text{LLC}_G: \Pi(G) \rightarrow \Phi(G)$.

- The meaning of “natural” is not clear in general.
(\rightsquigarrow It is not known even how to characterize LLC_G !)
- Known for $F = \mathbb{R}, \mathbb{C}$ due to Langlands ('89).
(\rightsquigarrow We focus on a p -adic F in the following.)
- \exists various results for specific G :
 - $G = \text{GL}_n$: Harris–Taylor ('01), Henniart ('00).
 - $G = \text{Sp}_{2n}, \text{SO}_n$: Arthur ('13).
 - $G = \text{U}_n$: Mok ('15).
 - ...
- But \exists still many things to study even in these cases.

Want: an explicit description (construction) of LLC_G .

Regular supercuspidal representations

Another approach to LLC: Instead of specifying a group G , restrict ourselves to some specific class of representations.

- $\Pi_{\text{sc}}(G)$: the set of supercuspidal representations of $G(F)$.
- Kaletha ('19) introduced a subclass $\Pi_{\text{reg}}(G)$ of $\Pi_{\text{tame}}(G)$ (called “regular supercuspidal representations”) and parametrized $\Pi_{\text{reg}}(G)$ via “regular pairs” (S, θ) .
 - S : a tamely ramified elliptic maximal torus of G over F .
 - $\theta: S(F) \rightarrow \mathbb{C}^\times$: a character satisfying a certain “regularity” condition.

$$\begin{aligned}\Pi(G) \supset \Pi_{\text{sc}}(G) \supset \Pi_{\text{reg}}(G) &\xleftarrow{1:1} \{(S, \theta) \mid \text{regular pairs}\} \\ \pi_{(S, \theta)} &\longleftrightarrow (S, \theta)\end{aligned}$$

LLC for regular supercuspidal representations

- Then Kaletha constructed “ LLC_G ” on $\Pi_{\text{reg}}(G)$:
 - Apply the LLC for S (\equiv local class field theory) to θ to get the L -parameter $\phi_\theta: W_F \rightarrow \hat{S} \rtimes W_F$.
 - Apply the “Langlands–Shlestad construction” to get an embedding $\hat{S} \rtimes W_F \hookrightarrow \hat{G} \rtimes W_F$.
 - By composing them, get an L -parameter of G (say ϕ):
$$\phi: W_F \xrightarrow{\phi_\theta} \hat{S} \rtimes W_F \hookrightarrow \hat{G} \rtimes W_F$$
 - Define $\phi := \text{LLC}_G(\pi_{(S,\theta)})$.

Theorem (O.–Tokimoto, 2021)

Kaletha’s LLC = Harris–Taylor and Henniart’s LLC for $\Pi_{\text{reg}}(\text{GL}_n)$.

Theorem (O., 2023)

Kaletha’s LLC = Arthur’s LLC for “toral” regular supercuspidal representations of Sp_{2n} or SO_n when $p \gg 0$.

\rightsquigarrow provides one answer to the problem of explicit LLC!

Deligne–Lusztig theory

Questions

1. The parametrization $(S, \theta) \leftrightarrow \pi_{(S, \theta)}$ is constructive (based on Yu's theory ('01)). Is there a more conceptual way to understand it?
2. Is it possible to establish such a parametrization outside $\Pi_{\text{reg}}(G)$?

We may seek a hint in **Deligne–Lusztig theory**.

- Deligne–Lusztig ('76) constructed all irreducible representations of $\mathbb{G}(\mathbb{F}_q)$ for any reductive group \mathbb{G} over \mathbb{F}_q .
 - Take any \mathbb{F}_q -rational maximal torus \mathbb{S} of \mathbb{G} .
 - Introduce a variety $X_{\mathbb{S}}^{\mathbb{G}} \hookrightarrow \mathbb{G}(\mathbb{F}_q) \times \mathbb{S}(\mathbb{F}_q)$ (“Deligne–Lusztig variety”).
 - Take its ℓ -adic cohomology $H_c^i(X_{\mathbb{S}}^{\mathbb{G}}, \overline{\mathbb{Q}}_{\ell})$ (repn of $\mathbb{G}(\mathbb{F}_q) \times \mathbb{S}(\mathbb{F}_q)$).
 - Take any character $\theta: \mathbb{S}(\mathbb{F}_q) \rightarrow \mathbb{C}^{\times} (\cong \overline{\mathbb{Q}}_{\ell}^{\times})$.
 - Cut & take the alternating sum: $R_{\mathbb{S}}^{\mathbb{G}}(\theta) := \sum_i (-1)^i H_c^i(X_{\mathbb{S}}^{\mathbb{G}}, \overline{\mathbb{Q}}_{\ell})[\theta]$.
- Input/output look similar to $(S, \theta) \leftrightarrow \pi_{(S, \theta)}$!

p -adic version of Deligne–Lusztig theory

- Chan–Ivanov ('21) established a p -adic version of the story of Deligne–Lusztig theory.
- To any pair (S, θ) of
 - any **unramified** F -rational maximal torus S of G ,
 - any character $\theta: S(F) \rightarrow \mathbb{C}^\times$,they associated a virtual representation $\pi_{(S, \theta)}^{\text{CI}}$ of $G(F)$.

Both Kaletha–Yu & Chan–Ivanov associate a repn of $G(F)$ to (S, θ) . But,

- Kaletha–Yu: S must be tame and elliptic; θ must be regular.
- Chan–Ivanov: S must be unramified, but can be non-elliptic; θ can be any.

Q. Relation between those on the “intersection” of their domains?

Comparison result

Theorem (Chan–O., '21, '24)

Suppose that S is an unramified elliptic maximal torus of G and $\theta: S(F) \rightarrow \mathbb{C}^\times$ is a regular character. Then $\pi_{(S,\theta)} \cong \pi_{(S,\theta)}^{\text{CI}}$ under some assumptions on q (the order of the residue field of F).

Idea of proof.

- \exists explicit character formulas:
 - $\pi_{(S,\theta)}$: Adler–DeBacker ('09), DeBacker–Spice ('18), Spice ('18, '21).
 - $\pi_{(S,\theta)}^{\text{CI}}$: Chan–Ivanov ('21).
- Both are too complicated to be compared in general, but are drastically simplified on “very regular” semisimple elements.
- If $q \gg 0$, we have enough many very regular semisimple elements so that the coincidence of the characters only on such elements implies the coincidence of representations.



Q. Is there a unified parametrization $(S, \theta) \leftrightarrow \pi_{(S, \theta)}$ which works on the “union” of the domains of Kaletha–Yu and Chan–Ivanov?

