

Notes on the Derivation of the Chern-Simons Form

mastex

September 29, 2025

Abstract

A memorandum. My goal is to be able to derive the Chern-Simons 3-form $\omega_3 = \text{tr} \left(AdA + \frac{2}{3} A^3 \right)$ from scratch.

1 Covariant Derivative of a p -form

We write the exterior derivative as d , the connection as A , the covariant derivative as $D = d + A$, and the curvature as $F = D^2 = dA + A^2$.

For a p -form C and an appropriate differential form ϕ , we have:

$$d(C\phi) = (dC)\phi + (-1)^p C d\phi$$

Now, if we introduce the connection A and replace the exterior derivative with the covariant derivative, $d \rightarrow D$, we get:

$$D(C\phi) = (DC)\phi + (-1)^p CD\phi$$

Rearranging the terms, we get:

$$(DC)\phi = D(C\phi) - (-1)^p CD\phi$$

$$= (dC)\phi + (AC)\phi - (-1)^p CA\phi$$

$$= (dC + [A, C])\phi$$

where we have defined

$$[A, C] = AC - (-1)^p CA$$

Therefore, the covariant derivative D acting on a p -form C is

$$DC = dC + [A, C]$$

This is the result.

2 Introduction of a Parameterized Connection

Here, we introduce a parameter $s \in \mathbb{R}$ and set $A_s = sA$.

Correspondingly, we have:

$$\begin{cases} D_s &= d + sA \\ F_s = (D_s)^2 &= sdA + s^2A^2 \end{cases}$$

In this case, we also get

$$\begin{cases} \frac{dF_s}{ds} = dA + 2sA^2 = D_sA \\ D_sF_s = 0 \end{cases}$$

and so on.

3 Derivation of the Chern-Simons $2n - 1$ Form

The Chern-Simons $2n - 1$ form ω_{2n-1} is defined by

$$d\omega_{2n-1} = \text{tr}F^n$$

Here we use the fact that

$$\text{tr}F^n = \int_0^1 ds \frac{d}{ds} \text{tr}F_s^n$$

The integrand of the right-hand side is

$$\begin{aligned} \frac{d}{ds} \text{tr}F_s^n &= \text{tr} \frac{dF_s^n}{ds} = \text{tr} \frac{dF_s^n}{ds} n F_s^{n-1} \\ &= n \text{tr}(D_sA) F_s^{n-1} \\ &= n \text{tr}D_s(AF_s^{n-1}) \end{aligned}$$

Here we used

$$D_sF_s^{n-1} = 0$$

Furthermore, using

$$D_sC = dC + [A_s, C]$$

we get

$$n \text{tr}D_s(AF_s^{n-1}) = n \text{tr}D_s(AF_s^{n-1}) + [D_s, (AF_s^{n-1})]$$

The second term is zero, so we finally get

$$\frac{d}{ds} \text{tr}F_s^n = d \left(n \text{tr}(AF_s^{n-1}) \right)$$

Returning to the definition of the Chern-Simons form,

$$d\omega_{2n-1} = d \left(\int_0^1 ds n \text{tr}(AF_s^{n-1}) \right)$$

$$\omega_{2n-1} = \int_0^1 ds n \text{tr}(AF_s^{n-1}) + \text{exact form}$$

This can be written.

When $n = 2$,

$$\omega_3 = \int_0^1 ds 2 \text{tr}A(sdA + s^2A^2) = \text{tr} \left(AdA + \frac{2}{3}A^3 \right)$$