

A Unified View of Monoids, Groups, Topological Groups, and Lie Groups

Masaru Okada

October 1, 2025

Abstract

We'll review the definitions of monoids, groups, topological groups, and Lie groups, and see how writing them in terms of category-theoretic diagrams makes them magically unified and easy to understand.

1 Naive Definitions

First, let's define monoids, groups, topological groups, and Lie groups naively.

1.1 Definition of a Monoid

A monoid is a semigroup with an identity element, satisfying associativity and having an identity.

A set M with a binary operation $\cdot : M \times M \rightarrow M$

1. Associativity: For any elements of M , $(a \cdot b) \cdot c = a \cdot (b \cdot c)$.
2. Existence of an identity element: There exists an element e in M such that for any element a in M , $e \cdot a = a \cdot e = a$.

1.2 Definition of a Group

A group is a monoid that also has an inverse element.

A set G with a binary operation $\cdot : G \times G \rightarrow G$

1. Associativity: For any elements of G , $(a \cdot b) \cdot c = a \cdot (b \cdot c)$.
2. Existence of an identity element: There exists an element e in M such that for any element a in G , $e \cdot a = a \cdot e = a$.
3. Existence of an inverse element: For any element a in G , there exists an element a^{-1} that satisfies $a^{-1} \cdot a = a \cdot a^{-1} = e$.

1.3 Definition of a Topological Group

A set G is a topological group if it is a group and satisfies the axioms of a topological space, and there exist continuous maps for the product and inverse.

1. G is a group.
2. G is a topological space.
3. The maps $\mu : G \times G \rightarrow G$ and $\nu : G \rightarrow G$, defined as $\mu(x, y) = xy$ and $\nu(x) = x^{-1}$ respectively, are continuous.

1.4 Definition of a Lie Group

A set G is a Lie group if it is a differentiable manifold, and satisfies the axioms of a topological group, and the maps for the product and inverse are differentiable.

1. G is a differentiable manifold.
2. The maps $\mu : G \times G \rightarrow G$ and $\nu : G \rightarrow G$, defined as $\mu(x, y) = xy$ and $\nu(x) = x^{-1}$ respectively, are differentiable.

2 A Unified View

2.1 Monoid

Next, let's try to understand these concepts in a unified way using elementary category theory.

A one-point set $1 = \{0\}$ acts as an identity element for the product (Cartesian product), considering the bijections

$$1 \times X \xrightarrow{\lambda} X \xleftarrow{\rho} X \times 1$$

given by $\lambda < 0, x >= x$, $\rho < x, 0 >= x$.

A monoid M is a set M equipped with two functions

$$\mu : M \times M \rightarrow M, \quad \eta : 1 \rightarrow M$$

that make the following diagrams commute with respect to μ and η .

$$\begin{array}{ccccc}
 M \times M \times M & \xrightarrow{1 \times \mu} & M \times M & & \\
 \downarrow \mu \times 1 & & \downarrow \mu & & \\
 M \times M & \xrightarrow{\mu} & M & & \\
 \\
 1 \times M & \xrightarrow{\eta \times 1} & M \times M & \xleftarrow{1 \times \eta} & M \times 1 \\
 & \searrow \lambda & \downarrow \mu & \swarrow \rho & \\
 & & M & &
 \end{array}$$

2.2 Introducing a Map for the Inverse Element

A group has an inverse element. A group is a monoid that makes the following diagram commute using a map $\xi : M \rightarrow M$ where $\xi : x \mapsto x^{-1}$.

$$\begin{array}{ccccc}
M & \xrightarrow{\delta} & M \times M & \xrightarrow{1 \times \xi} & M \times M \\
\downarrow \eta \circ ! & & & & \downarrow \mu \\
1 & \xrightarrow{\eta} & & & M
\end{array}$$

where η is the arrow to the identity element and $!$ is the unique arrow from M to the one-point set 1.

$$\begin{array}{ccccc}
x & \mapsto & \langle x, x \rangle & \mapsto & \langle x, x^{-1} \rangle \\
\downarrow & & & & \downarrow \\
0 & \mapsto & & & xx^{-1}
\end{array}$$

Here, $\delta : M \rightarrow M \times M$ is the diagonal map given by $\delta : x \mapsto \langle x, x \rangle$.

A group is what you get when M is a set and each morphism is just a function that makes the above diagrams commute.

A topological group is what you get when M is a topological space and each morphism is a continuous map that makes the diagrams commute.

A Lie group is what you get when M is a differentiable manifold and each morphism is a smooth map of manifolds that makes the diagrams commute.