Performance of a 250L liquid Argon TPC for sub-Gev charged particle identification

J-PARC T32 collaboration

Abstract

We have constructed a Liquid Argon time projection chamber (LArTPC) detector with fiducial mass of 150 kg (250L Detector) as a part of the R&D program of the next generation neutrino and nucleon decay detector.

This paper describes a study of particle identification performance of the 250L Detector using well-defined charged particles (pions, kaons, and protons) with momentum of \sim 800 MeV/c obtained at J-PARC K1.1Br beamline.

Keywords:

1. Introduction

2. Experimental Apparatus

- 2.1. K1.1Br Beamline
- 2.2. 250L Detector
- 2.3. Beamline Equipment
- 2.4. Oct/2010 Beam Test

3. Data Sample

- 3.1. Collected Data
- 3.2. Noise Reduction
- 3.3. Hit Finding/Clustering

4. Detector Calibration

- 4.1. Channel-by-Channel Calibration
- 4.2. Liquid Argon Purity
- 4.3. ("Cross Talk")

5. Simulation Tuning

- 6. Pion Result
- 7. Proton Result
- 7.1. (Recombination)?
- 8. Kaon Result
- 9. Summary

References

- [1] O. Araoka et al., J. Phys. Conf. Ser. 308, 012008 (2011) [arXiv:1105.5818 [physics.ins-det]].
- [2] S. Mihara [MEG Collaboration], Nucl. Instrum. Meth. A 518, 45 (2004).
- [3] S. Amoruso et al. [ICARUS Collaboration], Nucl. Instrum. Meth. A 523, 275 (2004).
- [4] S. Amoruso, M. Antonello, P. Aprili, F. Arneodo, A. Badertscher, B. Baibusinov, M. Baldo-Ceolin and G. Battistoni et al., Nucl. Instrum. Meth. A 516, 68 (2004).
- [5] A. Bettini et al., Nucl. Instrum. Meth. A 305, 177 (1991).
- [6] P.V.C Hough 'Method and means for recognizing complex patterns', United States Patent Office 3069654(1962)

Preprint submitted to NIMA February 9, 2012