

ALTRES PUBLICACIONS
OTRAS PUBLICACIONES
OTHER PUBLICATIONS

- Valero J, España J, Parra-Damas A, Martín E, Rodríguez-Álvarez J, Saura CA.
Short-term Environmental enrichment rescues adult neurogenesis and memory deficits in APP(Sw,Ind) transgenic mice.
PLoS One. **2011**;6(2).
- Badiola N, Penas C, Miñano-Molina A, Barneda-Zahonero B, Fadó R, Sánchez-Opazo G, Comella JX, Sabriá J, Zhu C, Blomgren K, Casas C, Rodríguez-Álvarez J.
Induction of ER stress in response to oxygen-glucose deprivation of cortical cultures involves the activation of the PERK and IRE-1 pathways and of caspase-12.
Cell Death Dis. **2011**;2:e149
- España J, Giménez-Llort L, Valero J, Miñano A, Rábano A, Rodríguez-Álvarez J, LaFerla FM, Saura CA.
Intraneuronal beta-Amyloid Accumulation in the Amygdala Enhances Fear and Anxiety in Alzheimer's Disease Transgenic Mice.
Biol Psychiatry. **2010**;67(6):513-21.
- España J, Valero J, Miñano-Molina AJ, Masgrau R, Martín E, Guardia-Laguarta C, Lleó A, Giménez-Llort L, Rodríguez-Álvarez J, Saura CA.
Beta-Amyloid disrupts activity-dependent gene transcription required for memory through the CREB coactivator CRTC1.
J Neurosci. **2010**;30(28):9402-10.
- Badiola N, Malagelada C, Llecha N, Hidalgo J, Comella JX, Sabriá J, Rodríguez-Alvarez J.
Activation of caspase-8 by tumor necrosis factor receptor 1 is necessary for caspase-3 activation and apoptosis in oxygen-glucose deprived cultured cortical cells.
Neurobiol Dis. **2009**;35(3):438-47.
- Barneda-Zahonero B, Miñano-Molina A, Badiola N, Fadó R, Xifró X, Saura CA, Rodríguez-Álvarez J.
Bone morphogenetic protein-6 promotes cerebellar granule neurons survival by activation of the MEK/ERK/CREB pathway.
Mol Biol Cell. **2009**;20(24):5051-63.
- Pérez V, Sosti V, Rubio A, Barbanoj M, Gich I, Rodríguez-Álvarez J, Kulisevsky J.
Noradrenergic modulation of the motor response induced by long-term levodopa administration in Parkinsonian rats.

J Neural Transm. **2009**;116(7):867-74.

- Chaïb-Oukadour I, Gil C, Rodríguez-Alvarez J, Ortega A, Aguilera J.
Tetanus toxin H(C) fragment reduces neuronal MPP+ toxicity.
Mol Cell Neurosci. **2009**;41(3):297-303.

- Raffaella G., Sole,C., Llecha,N, Segura, M.F., Moubarak,R., Iglesias-Guimaraes,V., Perez-Garcia,M.J., Reix,S., Zhang,J., Badiola,N., Sanchis,D., Rodríguez-Álvarez,J.,Trullas,R., Yuste,V.J., and J.X. Comella.
BCL-XL regulates TNF- α -mediated cell death independently of NF- κ B, FLIP and IAPs.
Cell Res. **2008**;18(10):1020-36.

- Miñano A, Xifró X, Pérez V, Barneda-Zahonero B, Saura CA, Rodríguez-Alvarez J.
Estradiol facilitates neurite maintenance by a Src/Ras/ERK signalling pathway.
Mol Cell Neurosci. **2008**;39(2):143-51.

- Miñano A, Caballero-Benítez A, Lluch M, Morán J, Rodríguez-Alvarez J.
C2-ceramide mediates cerebellar granule cells apoptosis by activation of caspase-2,-9 and -3.
J Neurosci Res. **2008**;86(8):1734-47.

- Badiola N, Segura M F, Comella J X, Rodríguez-Álvarez J.
Receptores de muerte e isquemia cerebral.
En Fisiopatología de la isquemia cerebral pp 77-89 (**2008**)
Editor J. Montaner. Merge Medica Books

- Segura,M.F., Sole,C., Pascual,M., Moubarak,R., Perez-Garcia,J.M., Gozzelino,R., Iglesias,V., Badiola,N., Bayascas,J.R., Llecha,N., Rodríguez-Alvarez,J., Soriano,E., Yuste, V.J. y J.X. Comella
The long form of Fas apoptotic inhibitory molecule (FAIML) is specifically expressed in neurons and protects them against death-receptor-triggered apoptosis.
J. Neurosci. 42: 11228-11241-**2007**

- Miñano,A., Cerbón, M.A., Xifró,X., Malagelada,C., Aguilera,J. y J. Rodríguez-Alvarez
Estradiol does not protect cerebellar granule cells from neither excitotoxicity nor apoptosis.
J. Neurochem. 102: 354-364-**2007**

- Pérez,V., Sosti,V., Rubio,A., BarbanoJ,M., Rodríguez-Alvarez,J. y J. Kulisevsky
Modulation of the motor response to dopaminergic drugs in a parkinsonian model of combined dopaminergic and noradrenergic degeneration.
Eur. J. Pharmacol. 576:83-90- **2007**

- Xifró,X., Miñano,A y J. Rodríguez-Alvarez
Survival of cerebellar granule neurons. Role of protein kinases pathways.
In Neurochemistry of Neuronal Death pp 157-174 -**2007**
Eds. Massieu, Arias y Morán. Research Singpost
- Aubert N, Falluel-Morel A, Vaudry D, Xifro X, Rodriguez-Alvarez J, Fisch C,de Jouffrey S, Lebigot JF, Fournier A, Vaudry H, Gonzalez BJ.
PACAP and C2-ceramide generate different AP-1 complexes through a MAP-kinase-dependent pathway: involvement of c-Fos in PACAP-induced Bcl-2 expression.
J Neurochem. 99:1237-50-**2006**
- Xifro X, Falluel-Morel A, Minano A, Aubert N, Fado R, Malagelada C, Vaudry D, Vaudry H, Gonzalez B, Rodriguez-Alvarez J.
N-methyl-D-aspartate blocks activation of JNK and mitochondrial apoptotic pathway induced by potassium deprivation in cerebellar granule cells.
J Biol Chem. 281:6801-12-**2006**
- Malagelada C, Xifro X, Minano A, Sabria J, Rodriguez-Alvarez J.
Contribution of caspase-mediated apoptosis to the cell death caused by oxygen-glucose deprivation in cortical cell cultures.
Neurobiol Dis. 20:27-37-**2005**
- Xifro X, Malagelada C, Minano A, Rodriguez-Alvarez J.
Brief exposure to NMDA produces long-term protection of cerebellar granule cells from apoptosis.
Eur J Neurosci. 27:827-40 -**2005**
- Malagelada C, Xifro X, Badiola N, Sabria J, Rodriguez-Alvarez J.
Histamine H2-receptor antagonist ranitidine protects against neural death induced by oxygen-glucose deprivation.
Stroke. 35:2396-401-**2004**
- Dot J, Lluch M, Blanco I, Rodriguez-Alvarez J.
Polyamine uptake in cultured cerebellar granule neurons.
Neurochem Int. 44:549-56 -**2004**