

קורס שפת סי ענת פרטוש פתרון מטלה מס' 2 – אופרטורים

פתרון לשאלה 1

```
z1=0 , z2=-1 , z3=0 , c1=12.0 , c2=4.0 , c3=7.0 : נימוק : z1=x%y - a=7%4 - 2.5=3 - 2.5=0.5\Rightarrow 0 z2=1+y*b=1+4*(-0.5)=1+(-2.0)=-1.0\Rightarrow -1 z3=b / z3=-0.5 / z3=-0.1667\Rightarrow 0 z3=x+a*2=7+2.5*2=7+5.0=12.0 z3=x+3/y = z3=x/y = z3=x
```

<u>פתרון לשאלה 2</u>

```
z1=1 , z2=1 , z2=1 , z2=0.5 , z2=0.0 : נימוק : z1=x+y/a=1+2/3.0=1+0.667=1.667 \Rightarrow z1=1 z1=x+y/a=1+2/3.0=1+0.667=1.667 \Rightarrow z1=1 z1=x+y/a=(1+2)/3.0=3/3.0=1.0 \Rightarrow z2=1 z1=x*((a-b)/y+x)=1*((3.0-4.0)/2+1)=1*((-1.0)/2+1)=1*(-0.5+1)=0.5 z1=1 z1=1 z1=1 z1=1 z2=1 z1=1 z1
```

<u>פתרון לשאלה 3</u>

```
a=b=c=1\;,d=1\;;בסיום הקטע : a=b=3<5\;;\qquad \  \  \, | a=b=1\;; \\ c=!5\;||1\;; \qquad \  \  \, | c=!1\;||1=0\;||1=1\;; \\ d=a\;!=(b+1)\;; \qquad \  \  \, | d=1!=(1+1)=1\;
```

<u>פתרון לשאלה 4</u> (חובה)

```
Ş
```

<u>פתרון לשאלה 5</u> (חובה)

```
א. a<b && c<b = (5<2) && (לא משנה) = 0

ב. a<!b || !!a = (5<0) || !!(5) = false || true = 1

ג. a+b < !c+c = (5+2) < (0+3) = 7 < 3 = 0

T. a-x || b*c && b/a = (5-1.0) || ( (לא משנה) ) && (לא משנה) ) = 1

a- (x || b*c && b/a)=5 - (1.0 || ( (לא משנה) ) && (לא משנה) ) )=5-1=4
```

פתרון לשאלה <u>6</u> (חובה)

```
נימוק: x1 = !num1; // x1 = !1 = 0 x2 = !(num2 >= num3); // x2 = !(2>=3) = !(0) = 1 x3 = (num1+num2 == num3) || (num2 > num3); // x3 = (1+2==3)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(?) = (1)||(
```

x1 = 0, x2 = 1, x3 = 1, x4 = 1, x5 = 1: בסיום בקטע יהיו הערכים

<u>פתרון לשאלה 7 (חובה)</u>

num1=-1 num2=311 : למסך יודפס נימוק :

```
num1+=num2\%10 = 32 + 10\%10 = 32 + 0 = 32

num2*=num1++ - 1 = 10 * (32-1) = 310    num1=31

num1/=num2 + 3 = 31/(310+3) = 0

num2-=--num1 = 310 - (-1) = 311    num1=-1
```

8 פתרון לשאלה

a=0 , b=2 , x=1 , y=-2 : בסיום הקטע נימוק :

int
$$a=0$$
, $b=2$, x , y ;
 $x = (++a*a--)*(--b*b++)$; // $x=(1*1)*(1*1)=1$, $a=0$, $b=2$
 $y = (!(a || b)) ? (a+b) : (a-b)$; // $y=(!(0||2))?(a+b):(a-b) => y=a-b=0-2=-2$



פתרון לשאלה 9

: נימוק

x = -1, m = 1, n = 5: ערך המשתנים

```
int m=-5, n=2 , x ; 
 x = ++m \Rightarrow m=-4 , x=-4 
 x /= n+1 \Rightarrow x=x/(n+1) = -4/(2+1) = -4/3 = -1 
 n = (x-m) % n \Rightarrow n=(-1-(-4)) % 2 = (-1+4)%2 = 3%2 = 1 
 m += x+n && n++ \Rightarrow m=m+(x+n&&n++) = -4+((-1+1)&&(n++)) = -4+(0&&(n++)) = -4+0 = -4
```

פתרון לשאלה 10

```
x=10 , m=-4 , n=1 : נימוק : x=10 , x=
```

```
int m=-6, n=0 , x ; 
 x = --m \Rightarrow m=-7 , x=-7 
 x *= ++n-3 \Rightarrow n=1 , x = x * (++n-3) = -7 * (1-3) = -7 * (-2) = 14 
 n = (x+m) \% x \Rightarrow n = (14+(-7)) \% 10 = 7 \% 10 = 7 
 m = x-n==n \&\& -m==n \Rightarrow m = (14-7==7)\&\&(--7==7) \Rightarrow (1)\&\&(1) = 1
```