

STA 674

Regression Analysis And Design Of Experiments

Measuring Association between Two Variables – Lecture 4

STA 674, RADOE:

Measuring Association between Two Variables

- Interpretation of Correlation
 - Last time – sign
 - This time – magnitude

STA 674: Measuring Association between Two Variables

Correlation

- The correlation, $R_{X,Y} = \frac{\sum_{i=1}^n [(x_i - \bar{x})(y_i - \bar{y})]}{(n-1)s_X s_Y}$, always lies between -1 and 1, inclusive.
- Suggested language:

If $R_{X,Y}$ is between	then we may say* X and Y have ...
$0.5 \leq R_{X,Y} \leq 1$	a strong positive correlation
$0 < R_{X,Y} < 0.5$	a weak positive correlation
$-0.5 < R_{X,Y} < 0$	a weak negative correlation
$-1 \leq R_{X,Y} \leq -0.5$	a strong negative correlation


STA 674: Measuring Association between Two Variables

Correlation

- Example: Average January temperature in US cities

Previously, we found $R_{XY} = -0.94$

Average Jan. Temp. vs Latitude




City		Latitude	Average Jan Temp
Louisville, KY		39	27
Key West, FL		25	65
New Orleans, LA		30.8	45
Atlanta, GA		33.9	37
Charlotte, NC		35.9	34
Harrisburg, PA		40.9	24
Omaha, NE		41.9	13
Detroit, MI		43.1	21
Burlington, VT		45	7
Spokane, WA		48.1	19
Means		38.4	29.2
Standard deviations		7	16.9

STA 674: Measuring Association between Two Variables

Correlation

- Practice (?)


Graph	A	B	C	D
\bar{x}	50	50	50	50
\bar{y}	54	54	58	27
Graph	E	F	G	H
\bar{x}	50	50	31	50
\bar{y}	44	54	31	58




STA 674: Measuring Association between Two Variables

Correlation


- True values:




$$R_{X,Y} = .96$$




$$R_{X,Y} = .82$$




$$R_{X,Y} = .44$$




$$R_{X,Y} = -.05$$




$$R_{X,Y} = -.23$$



$$R_{X,Y} = .96$$



$$R_{X,Y} = .89$$



$$R_{X,Y} = -.45$$

STA 674: Measuring Association between Two Variables

Correlation

- Next time – *cautions* about interpretations of the correlation coefficient.