

Applying the CDC's Building Resilience Against Climate Effects (BRACE) Framework in Massachusetts

Tuesday, October 3, 2014

MIT Endicott House, Dedham, MA

Marc A. Nascarella, Ph.D.
Director, Environmental Toxicology Program




Outline



1. Climate Change Health Hazards
2. Introduction to the CDC BRACE Framework
3. Applying BRACE to Evaluate Public Health Risk
4. Available Tools
5. Questions

Climate Change Hazards


3



(Source: Nicholas Stern (2006), "Stern Review on the Economics of Climate Change")

Climate Change Effects

4



(Source: Nicholas Stern (2006), "Stern Review on the Economics of Climate Change")



BRACE Framework

6

PURPOSE: Couple climate projections with health data to more effectively **anticipate, prepare, and respond** to climate sensitive health impacts.

- STEP 1:** Forecast Climate Impact and Assess Vulnerabilities
- STEP 2:** Project Disease Burdens from Climate Change
- STEP 3:** Assess Public Health Interventions
- STEP 4:** Develop & Implement Climate Adaptation Plan
- STEP 5:** Evaluate Impact and Improve Framework



BRACE Framework

7

Int. J. Environ. Res. Public Health **2014**, *11*, 6433–6458; doi:10.3390/ijerph110606433

OPEN ACCESS

International Journal of
Environmental Research and
Public Health
ISSN 1660-4601
www.mdpi.com/journal/ijerph

Article

Building Resilience against Climate Effects—A Novel Framework to Facilitate Climate Readiness in Public Health Agencies

Gino D. Marinucci ¹, George Luber ¹, Christopher K. Uejio ^{1,2}, Shubhayu Saha ¹ and Jeremy J. Hess ^{1,3,4,*}

¹ Climate and Health Program, Division of Environmental Hazards and Health Effects, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, GA 30341, USA; E-Mails: ipx1@cdc.gov (G.D.M.); gcl4@cdc.gov (G.L.); cuejio@fssu.edu (C.K.U.); hsf5@cdc.gov (S.S.)

² Department of Geography, Florida State University, 113 Collegiate Loop, Tallahassee, FL 32306, USA


³ Department of Environmental Health, Rollins School of Public Health at Emory University, Atlanta, GA 30322, USA

⁴ Department of Emergency Medicine, School of Medicine, Emory University, Atlanta, GA 30322, USA

* Author to whom correspondence should be addressed; E-Mail: jhess@emory.edu; Tel.: +1-404-251-8851; Fax: +1-404-688-6351.


Received: 8 April 2014; in revised form: 3 June 2014 / Accepted: 6 June 2014 /

Published: 20 June 2014



Assessing the health effects of climate change...where to begin?

8



Basic Public Health Risk Assessment

- Public health **risks** are a function of the type of **hazard** and our **exposure** to them.
- e.g., health hazard such as an air contaminant:
 - ▣ STEP 1 - Determine inherent toxicity of the contaminant (e.g., “imminent **hazard** value”)
 - ▣ STEP 2 - Determine **exposure** (i.e., what level are people breathing).
 - ▣ Estimate **risk** based on:

$$\text{RISK} = \text{HAZARD} \times \text{EXPOSURE}$$



Climate Risk Assessment

- Public health **risks** of climate change are similar
- **Risks** are directly related to specific climate **hazards** and our **exposure** to them.
- Using the **CDC BRACE** framework we:
 - Predict climate-related **hazards**
 - Predict human vulnerability of **exposure**
 - Estimate **risk** based on:

$$\text{RISK} = \text{HAZARD} \times \text{EXPOSURE}$$

Risk of Climate Effects

11



HAZARD

Probability
Magnitude

EXPOSURE

Vulnerability
Susceptibility

RISK

Damage
Valuation

Hazard x Exposure = Risk

Example: Risk of Flooding



RISK = **HAZARD** x **VULNERABILITY**

Risk = Flooding in a Coastal Home

Hazard = Sea Level

Vulnerability = Elevation of Home

Predicting Hazards


13

1. Current Health Effects

Determine relationship between suspected hazard (i.e., extreme heat) and population experiencing adverse health outcomes.

2. Future Health Effects

Based on relationship between vulnerable population and number of hot days in the future (i.e., population and temperature in 2030).



Predicting Hazards: Extreme Heat



$$\text{RISK} = \text{HAZARD} \times \text{VULNERABILITY}$$

Risk = Acute Heat Injury Requiring Medical Services


Hazard = Days where Temp. $\geq 93^{\circ}\text{F}$

Vulnerability = Social Isolation + Age ≥ 65 years old*

*see Health Vulnerability Index; Reid et al. (2012). Environmental Health Perspectives, 120(5), 715-719.


Massachusetts Weather Data

Maximum Temperature Data May-September 2002




***DRAFT* ILLUSTRATION FOR PRESENTATION PURPOSES**

Maximum Temperature in MA



***DRAFT* ILLUSTRATION FOR PRESENTATION PURPOSES**

Combine Weather and Health Data



***DRAFT* ILLUSTRATION FOR PRESENTATION PURPOSES**

Massachusetts Health Data




Heat-related Health Burden in MA Communities

- Community level heat-related hospital admissions, per community, per day
 - Population Data
 - Percent age 65 or older
 - Percent > 65 and live alone
 - Percent of homes with no Air Conditioning
 - Health Data
 - Heat related illness
 - Electrolyte imbalance
 - Diabetes
 - Nephritis and nephritic syndrome
 - Respiratory Disease

Risk of Heat-Related Hospital Visits


- Estimated 1°- 3°F temperature increase by 2030
- 0 - 162 additional heat-related hospital visits/town/year



***DRAFT* ILLUSTRATION FOR PRESENTATION PURPOSES**

Planning for Climate Adaptation

20



Parameter	Current Conditions (1961–1990)	Predicted Range of Change by 2050	Predicted Range of Change by 2100
Annual temperature ¹ (°C/°F)	8/46	2 to 3 / 4 to 5	3 to 5/5 to 10**
Winter temperature ¹ (°C/°F)	-5/23	1 to 3 / 2 to 5	2 to 5 / 4 to 10**
Summer temperature ¹ (°C/°F)	20/68	2 to 3 / 4 to 5	2 to 6 / 4 to 10**
Over 90 °F (32.2 °C) temperature ² (days/yr)	5 to 20	—	30 to 60
Over 100 °F (37.7 °C) temperature ² (days/yr)	0 to 2	—	3 to 28
Ocean pH ^{3,4}	7 to 8	—	-0.1 to -0.3 [†]
Annual sea surface temperature (°C/°F)	12/53 ⁵	2/3 (in 2050) ⁵	4/8
Annual precipitation ¹	103 cm/41 in.	5% to 8%	7% to 14%**
Winter precipitation ¹	21 cm/8 in.	6% to 16%	12% to 30%**
Summer precipitation ¹	28 cm/11 in.	-1% to -3%	-1% to 0%**
Streamflow—timing of spring peak flow ¹ (number of calendar days following January 1)	85	-5 to -8	-11 to -13**
Droughts lasting 1–3 months ¹ (#/30 yrs)	13	5 to 7	3 to 10**
Snow days (number of days/month) ¹	5	-2	-2 to -4**
Length of growing season ¹ (days/year)	184	12 to 27	29 to 43

Table 1: Changes in Massachusetts' Climate


Sources: 1-Hayhoe et al., 2006; 2-Frumhoff et al., 2007; 3-IPCC, 2007; 4-MWRA, unpublished; 5-Nixon et al., 2004
 Note: All numbers have been rounded to the nearest whole number. Unless otherwise indicated, the predictions for the year listed as 2050 are for the period between 2035–2064. [†] Global data; **Predictions for period between 2070–2099



Capacity to Respond to Climate Adaptation

21

“...comprehensive survey aimed at assessing the capacity of local health departments in Massachusetts to respond to the public health impacts of climate change...”






<http://www.mass.gov/eohhs/docs/dph/environmental/exposure/climate-change-report-2014.pdf>

MA Environmental Public Health Tracking

<https://matracking.ehs.state.ma.us/#>

22



Research

Evaluation of a Heat Vulnerability Index on Abnormally Hot Days: An Environmental Public Health Tracking Study

Colleen E. Reid,¹ Jennifer K. Mann,¹ Ruth Alfasso,² Paul B. English,³ Galatea C. King,³ Rebecca A. Lincoln,⁴ Helene G. Margolis,⁵ Dan J. Rubado,⁶ Joseph E. Sabato,² Nancy L. West,⁷ Brian Woods,⁸ Kathleen M. Navarro,¹ and John R. Balmes^{1,9}

¹Environmental Health Sciences, School of Public Health, University of California-Berkeley, Berkeley, California, USA; ²Massachusetts Department of Public Health, Bureau of Environmental Health, Boston, Massachusetts, USA; ³Environmental Health Investigations Branch, California Department of Public Health, Richmond, California, USA; ⁴Department of Health and Human Services, Maine Center for Disease Control and Prevention, Augusta, Maine, USA; ⁵Department of Internal Medicine, School of Medicine, University of California-Davis, Davis, California, USA; ⁶Office of Environmental Public Health, Oregon Health Authority, Portland, Oregon, USA; ⁷Division of Environmental Health, Washington State Department of Health, Olympia, Washington, USA; ⁸Environmental Health Epidemiology Bureau, New Mexico Department of Health, Santa Fe, New Mexico, USA; ⁹Division of Occupational and Environmental Medicine, Department of Medicine, University of California, San Francisco, California, USA

BACKGROUND: Extreme hot weather conditions have been associated with increased morbidity and mortality, but risks are not evenly distributed throughout the population. Previously, a heat vulnerability index (HVI) was created to geographically locate populations with increased vulnerability to heat in metropolitan areas throughout the United States.

OBJECTIVES: We sought to determine whether areas with higher heat vulnerability, as characterized by the HVI, experienced higher rates of morbidity and mortality on abnormally hot days.

for specific metropolitan areas (Johnson et al. 2009; Lindley et al. 2006; Loughnan et al. 2009; Rinner et al. 2009; Sister et al. 2009; Vescovi et al. 2005), the HVI is the only vulnerability map that is national in scope. The HVI suggests substantial variability in heat

Thank you

Questions?

Marc A. Nascarella, Ph.D.
Director, Environmental Toxicology Program

617-624-5757 | marc.nascarella@state.ma.us

