

# Breast lesion Segmentation in Ultra-Sound images based on super-pixels and high-level descriptors

Joan Massich<sup>a</sup> and Guillaume Lemaître<sup>a,b</sup> and Joan Martí<sup>b</sup> and Fabrice Mériaudeau<sup>a</sup>

<sup>a</sup>LE2I-UMR CNRS 6306, Université de Bourgogne, 12 rue de la Fonderie, 71200 Le Creusot, France;

<sup>b</sup>ViCOROB, Universitat de Girona, Campus Montilivi, Edifici P4, 17071 Girona, Spain

## ABSTRACT

This document shows the desired format and appearance of a manuscript prepared for the Proceedings of the SPIE. It contains general formatting instructions and hints about how to use LaTeX. The LaTeX source file that produced this document, `article.tex` (Version 3.3), provides a template, used in conjunction with `spie.cls` (Version 3.3).

**Keywords:** Breast Ultra-Sound, Machine-Learning based Segmentation

## 1. INTRODUCTION

Breast cancer is the second most common cancer (1.4 million cases per year, 10.9% of diagnosed cancers) after lung cancer, followed by colorectal, stomach, prostate and liver cancers.<sup>1</sup> In terms of mortality, breast cancer is the fifth most common cause of cancer death. However, it place as the leading cause of cancer death among females both in western countries and in economically developing countries.<sup>2</sup>

Medical imaging plays an important role in breast cancer mortality reduction, contributing to its early detection through screening, diagnosis, image-guided biopsy, treatment follow-up and suchlike procedures.<sup>3</sup> Although Digital Mammography (DM) remains the reference imaging modality, US imaging has proven to be a successful adjunct image modality for breast cancer screening,<sup>3,4</sup> specially as a consequence of the discriminative capabilities that US offers for differentiating between solid lesions that are benign or malignant<sup>5</sup> so that the amount of unnecessary biopsies, which is estimated to be between 65 ~ 85% of the prescribed biopsies,<sup>6</sup> can be reduced<sup>7</sup> in replacing them by short-term US screening follow-up.<sup>8</sup>

Figure ... shows ... what doctors look for.

Analysing figure ... it can be observed that most of the markers depend on the lesion delineation. Therefore in order to develop releable Computer Aided Diagnose (CAD) systems accurate segmentations to properly delineate the lesions are needed. This article presents a segmentation technique based on classifying superpixels based on their appearance.

## 2. SEGMENTATION METHODOLOGY DESCRIPTION

Optimization methodologies offer a standardized manner to approach segmentation by minimizing an application-driven cost function.<sup>9</sup> The segmentation strategy here adopted is formulated as a *metric labelling* problem. The image is conceived as a discrete set of elements  $\mathcal{S}$  that need to be labelled using a label  $l$  from a labelling set  $\mathcal{L}$ . Let  $\mathcal{W}$  be all the possible labelling configurations of the set  $\mathcal{S}$  given  $\mathcal{L}$ , and let  $U(\cdot)$  be a cost function encoding how good is a labelling configuration  $\omega \in \mathcal{W}$  based on the appearance of the elements in  $\mathcal{S}$ , their relation and some designing constrains. Then the desired segmentation  $\hat{\omega}$ , corresponds to the labelling configuration that minimizes this cost function  $U(\cdot)$ , further described in Equation (1).

$$U(\omega) = \sum_{s \in \mathcal{S}} D_s(\omega_s) + \sum_s \sum_{r \in \mathcal{N}_s} V_{s,r}(\omega_s, \omega_r) \quad (1)$$

Figure 1 illustrates a generic version of the segmentation strategy here proposed. The image

---

Further author information: (Send correspondence to J.M.)

J.M.: E-mail: joan.massich@u-bourgogne.fr



Figure 1. Conceptual block representation of the segmentation methodology

### 3. FEATURE DESCRIPTION

Features description

### ACKNOWLEDGMENTS

This unnumbered section is used to identify those who have aided the authors in understanding or accomplishing the work presented and to acknowledge sources of funding.

### REFERENCES

1. J. Ferlay, H.-R. Shin, F. Bray, D. Forman, C. Mathers, and D. M. Parkin, "Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008," *International Journal of Cancer* **127**(12), pp. 2893–2917, 2010.
2. A. Jemal, F. Bray, M. M. Center, J. Ferlay, E. Ward, and D. Forman, "Global cancer statistics," *CA: A Cancer Journal for Clinicians* **61**(2), pp. 69–90, 2011.
3. R. A. Smith, D. Saslow, K. A. Sawyer, W. Burke, M. E. Costanza, W. Evans, R. S. Foster, E. Hendrick, H. J. Eyre, and S. Sener, "American cancer society guidelines for breast cancer screening: update 2003," *CA: a cancer journal for clinicians* **53**(3), pp. 141–169, 2003.
4. W. A. Berg, L. Gutierrez, M. S. NessAiver, W. B. Carter, M. Bhargavan, R. S. Lewis, and O. B. Ioffe, "Diagnostic accuracy of mammography, clinical examination, US, and MR imaging in preoperative assessment of breast cancer," *Radiology* **233**(3), pp. 830–849, 2004.
5. A. T. Stavros, D. Thickman, C. L. Rapp, M. A. Dennis, S. H. Parker, and G. A. Sisney, "Solid breast nodules: Use of sonography to distinguish between benign and malignant lesions," *Radiology* **196**(1), pp. 123–34, 1995.
6. Y. Yuan, M. L. Giger, H. Li, N. Bhooshan, and C. A. Sennett, "Multimodality computer-aided breast cancer diagnosis with ffdm and dce-mri.," *Academic radiology* **17**(9), p. 1158, 2010.
7. S. Ciatto, M. Rosselli del Turco, S. Catarzi, D. Morrone, *et al.*, "The contribution of ultrasonography to the differential diagnosis of breast cancer.," *Neoplasma* **41**(6), p. 341, 1994.
8. P. B. Gordon and S. L. Goldenberg, "Malignant breast masses detected only by ultrasound. A retrospective review," *Cancer* **76**(4), pp. 626–630, 1995.
9. D. Cremers, M. Rousson, and R. Deriche, "A review of statistical approaches to level set segmentation: integrating color, texture, motion and shape," *International journal of computer vision* **72**(2), pp. 195–215, 2007.