

Overlapping: An R package for Estimating Overlapping in Empirical Distributions

23 July 2018

Summary

Overlapping can be defined as the area intersected by two or more probability density functions. The idea of overlapping was introduced in a formal way by Gini and Livada (1943) and, more recently, it has been applied in several research problems involving, for instance, data fusion (Moravec 1988), information processing (Viola and Wells III 1997), applied statistics (Inman and Bradley Jr 1989), economics (Milanovic and Yitzhaki 2001), psychology, as a basis for Cohen's U index (Cohen 1988), McGraw and Wong's CL measure (McGraw and Wong 1992), and Huberty's I degree of non-overlap index (Huberty and Lowman 2000).

`overlapping` is a R package for estimating the overlapping area of two or more empirical distributions. The main idea of the package is to offer an easy way to quantify the similarity (or the difference) between two or more empirical distributions. In addition, the package allows to plot density distributions, highlighting the overlapped area by using the `ggplot2` R package (Wickham 2009).

The package is available from GitHub (<https://github.com/masspastore/overlapping>) and CRAN (<https://cran.r-project.org/package=overlapping>). A full reference manual can be found at <https://cran.r-project.org/web/packages/overlapping/overlapping.pdf>.

Examples

Suppose we have collected data in two groups of 100 subjects each with respect to a generic variable Y , expressed by scores ranging between 0 and 30, and to be interested in assessing whether the two groups can be considered samples from populations with the same average.

We can simulate the groups scores as follows:

```
set.seed( 1 )
n <- 100
G1 <- sample( 0:30, size = n, replace = TRUE )
G2 <- sample( 0:30, size = n, replace = TRUE, prob = dbinom( 0:30, 31, .55 ) )
```

For Group 1 (G1) we randomly sampled $n = 100$ values from a uniform distribution; for Group 2 (G2) we randomly sampled 100 values from a binomial distribution. In the first group, scores range between 0 and 30 with mean 15.55 and standard deviation 8.32. In the second group, scores range between 10 and 24 with mean 16.72 and standard deviation 2.74.

We can display the scores distribution as follows:

```
library( ggplot2 )
Data <- data.frame( y = c(G1,G2), group = rep(c("G1","G2"),each=n) )
ggplot( Data, aes( y ) ) + facet_wrap( ~group ) + geom_histogram()
```

obtaining the figure 1. In the left panel are depicted the scores of group 1, in the right panel the scores of group 2. From this figure it is evident the heterogeneity of the variances in the two groups. In such a case, the statistical comparison between means can be biased and not very informative; for example, with a t -test, corrected for heterogeneity, we obtain the following result: $t(120.24) = -1.34$, $p = 0.18$, from which we cannot draw any conclusion.




Figure 1: Score distributions of simulated groups of 100 subjects each.

So, let us assume a different perspective: Rather than assessing the similarity between the two groups on the basis of averages (and standard deviations) only, we use all the information available in the data. In practice we estimate the degree of similarity by exploring the overlapping between group scores. We expect 0% to indicate the absence of overlapping (i.e., maximum distance between groups), and 100% to indicate the perfect overlap between the two distributions (i.e., groups are identically distributed). We can use the `overlapping` package in the following way:

```
library( overlapping )
dataList <- list( G1 = G1, G2 = G2 )
overlap( dataList )$OV * 100
```

```
##      G1-G2
## 43.21998
```

With the command `library()` we load the **overlapping** package, next we create a `list` containing the two groups scores, and finally, by using the `overlap()` function, we compute the overlap index. The index value (43.22) is an estimate of the percentage of overlapping between density scores. We can obtain a graphical representation by adding the option `plot = TRUE` as follows:

```
overlap( dataList, plot = TRUE )
```

obtaining the figure 2. In the figure are represented the estimated densities of the two groups scores, with different colors, and the shaded part is the overlapping area of densities.

Examples of real-world analysis

overlapping package was already used in different publications for many purposes, such as: 1) evaluating group invariance in questionnaires, by using parameters bootstrap distributions (Lionetti, Mastrotheodoros, and Palladino 2018, @marci+al:2018); 2) for computing a distance index in antropological measures (Altoè, D'Amore, and Scalfari, n.d.); 3) for identifying cut-off scores in questionnaires, estimating the intersection points of density distributions (Pluess et al. 2018, @lionetti+al:2018dandelions).




Figure 2: Comparison between densities of two groups. The overlap (43%) is represented by the shaded area.

References

- Altoè, Gianmarco, Giuseppe D'Amore, and Francesco Scalfari. n.d. "Skulls and transvariation."
- Cohen, Jacob. 1988. *Statistical power analysis for the behavioral sciences*. Psychology Press.
- Gini, C, and G Livada. 1943. *Nuovi contributi alla teoria della transvariazione*. Atti della VI Riunione della Società Italiana di Statistica, Roma, gennaio 1943.
- Huberty, Carl J, and Lauren L Lowman. 2000. "Group overlap as a basis for effect size." *Educational and Psychological Measurement* 60 (4). Sage Publications Sage CA: Thousand Oaks, CA: 543–63.
- Inman, Henry F, and Edwin L Bradley Jr. 1989. "The overlapping coefficient as a measure of agreement between probability distributions and point estimation of the overlap of two normal densities." *Communications in Statistics-Theory and Methods* 18 (10). Taylor & Francis: 3851–74.
- Lionetti, Francesca, Arthur Aron, Elaine N Aron, G Leonard Burns, Jadzia Jagiellowicz, and Michael Pluess. 2018. "Dandelions, Tulips and Orchids: Evidence for the Existence of Low-Sensitive, Medium-Sensitive and High-Sensitive Individuals." *Translational Psychiatry* 8 (1). Nature Publishing Group: 24.
- Lionetti, Francesca, Stefanos Mastrotheodoros, and Benedetta Emanuela Palladino. 2018. "Experiences in Close Relationships Revised Child Version (Ecr-Rc): Psychometric Evidence in Support of a Security Factor." *European Journal of Developmental Psychology* 15 (4). Taylor & Francis: 452–63.
- Marci, Tatiana, Francesca Lionetti, Ughetta Moscardino, Massimiliano Pastore, Vincenzo Calvo, and Gianmarco Altoè. 2018. "Measuring attachment security via the Security Scale: Latent structure, invariance across mothers and fathers and convergent validity." *European Journal of Developmental Psychology* 15 (4). Taylor & Francis: 481–92.
- McGraw, Kenneth O, and SP Wong. 1992. "A common language effect size statistic." *Psychological Bulletin* 111 (2). American Psychological Association: 361.
- Milanovic, Branko, and Shlomo Yitzhaki. 2001. *Decomposing world income distribution: Does the world have a middle class?* The World Bank.
- Moravec, Hans P. 1988. "Sensor fusion in certainty grids for mobile robots." *AI Magazine* 9 (2): 61.
- Pluess, Michael, Elham Assary, Francesca Lionetti, Kathryn J Lester, Eva Krapohl, Elaine N Aron, and Arthur Aron. 2018. "Environmental Sensitivity in Children: Development of the Highly Sensitive Child

Scale and Identification of Sensitivity Groups.” *Developmental Psychology* 54 (1). American Psychological Association: 51.

Viola, Paul, and William M Wells III. 1997. “Alignment by maximization of mutual information.” *International Journal of Computer Vision* 24 (2). Springer: 137–54.

Wickham, Hadley. 2009. *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. <http://ggplot2.org>.