

Exploring Biomechanical and Clinical Factors in Diabetic Foot Ulcer Risk Assessment

Helya Amiri

Supervised by: Pierre-Yves Rohan, Jerome Haddad

University of Strasbourg, France

August 24, 2025


Outline

- 1 Introduction
- 2 Study Objectives
- 3 Methodology
- 4 Conclusion

Introduction

- **Diabetes:** A long-term disease where blood sugar stays high because the body makes too little insulin or does not use it well. It can harm nerves and blood vessels.
- **Neuropathy:** Nerve damage from diabetes that can cause foot pain, numbness, or weakness.

Diabetic Foot Ulcers



Risk Factors for Diabetic Foot Ulcers

- **Neuropathy:** Less feeling in the feet, so injuries happen more easily.
- **Poor Circulation:** Blood flow is weak, so wounds heal slowly.
- **Foot Deformities:** Feet have unusual shapes, causing more pressure on some spots.




Figure: Plantar Pressure Map

Why Study Diabetic Foot Ulcers?

- Over 4 million adults in France are living with diabetes (2024).¹
- Diabetic Foot Ulcers (DFUs) cause more than 10,000 amputations each year.
- **Goal:** Create a tool to study diabetic foot data and help doctors identify patients at risk before ulcers appear.

¹World Health Organization, 2016; International Diabetes Federation, 2024

Objectives

- **Identify key risk markers** (e.g., tissue stiffness, plantar pressure).
- **Explore left-right foot differences** as warning signs.
- **Apply unsupervised learning** (e.g., clustering)
- **Compare clusters with IWGDF grades**²

²IWGDF (International Working Group on the Diabetic Foot)

Data Collection

- Study done in a day-hospital with ethical approval.
- Participants signed consent forms.
- Included adults 18–65 with diabetes.
- Excluded pregnant, breastfeeding, or legally protected people.

The DIAFOOT Dataset

- **Participants:** 21 diabetic patients (11 F, 10 M).
- **Age:** Mean 56.8 years.
- **BMI:** Mean 25.83 kg/m²






Figure: IWGDF Grades

Data Parameters


- **Anthropometric:** BMI, height, weight
- **Vascular:** Ankle and big toe blood pressure
- **Neurological:** Michigan score, SUDOSCAN (hands/feet)
- **Biomechanical:** Foot pressure (SESA, hallux, TM5), tissue stiffness
- **Imaging:** Ultrasound tissue thickness
- **Thermal:** Foot temperature and normalized values

Three zones Example

Fifth Metatarsal



Bunions also known as hallux valgus



Analysis Pipeline

Streamlit Dashboard

- **Platform:** Built using Streamlit, A dynamic, Python-driven platform for real-time data exploration and visualization.
- **Data Source:** Excel file with 'DIAFOOT' sheet containing patient metrics.
- **Key Features:** Descriptive statistics, normality tests, clustering, correlations and comparative analyses.
- **Tools:** Pandas (data handling), NumPy (numerical), Scikit-learn (ML, clustering), Matplotlib, Seaborn (visuals), `scipy.stats` (hypothesis testing), Statsmodels (MANOVA, regression),

Review of Statistics: p-value

- **p-value:** Shows if a result is real or by chance.
- Small p-value $< 0.05 \rightarrow$ **significant** (result likely real)
- Large p-value $\geq 0.05 \rightarrow$ **not significant** (result may be by chance)
- Used to decide whether to reject the null hypothesis.

Explore the DIAFOOT Analysis Dashboard

<https://diafoot-analysis.streamlit.app/>

Upload data, analyze it, and see important results easily.

Try it now!

Asymmetry Insights

Significant Left-Right Differences

- **TM5(fifth metatarsal) Pressure:** Right (36.93 kPa) > Left (26.34 kPa), $p = 0.0054$.
- **Hallux Stiffness:** Right higher, $p = 0.0498$.
- **Temperature (Plantar Arch, Lateral Sole):** Right warmer, $p < 0.01$.




Figure: Pressure Asymmetry

Group Differences: Low vs. High Risk

Parameter	Grades 0-1	Grades 2-3	p-value
Michigan Score	11.69	4.13	0.00053
Tissue Thickness SESA R (mm)	9.55	4.17	0.0099
Heel Temperature R (°C)	25.85	28.55	0.0188

Table: Significant Group Differences

- Higher-risk patients have lower Michigan scores, thinner tissue, and warmer heels, showing higher ulcer risk.

Clustering Analysis

- Clustering done with and without IWGDF grades
- **PCA:** 2D visualization of patient data.
- **Clustering:** Agglomerative, KMeans, GMM; 2–6 clusters.
- **Metrics:** Silhouette, Calinski–Harabasz, Davies–Bouldin, ARI, NMI.
- **Feature importance:** LASSO, Random Forest, ANOVA.
- **Group differences:** MANOVA, p-values indicate significance.

Clustering Metrics

Algorithm	Silhouette	CH Score	DB Score	ARI	NMI
With IWGDF Grade					
Agglomerative (Ward)	0.553	43.959	0.485	0.794	0.778
Agglomerative (Average)	0.553	43.959	0.485	0.794	0.778
KMeans	0.553	43.959	0.485	0.794	0.778
Gaussian Mixture Model	0.344	33.367	0.939	0.490	0.654
Without IWGDF Grade					
Agglomerative (Ward)	0.539	40.721	0.532	0.794	0.778
Agglomerative (Average)	0.539	40.721	0.532	0.794	0.778
KMeans	0.539	40.721	0.532	0.794	0.778
Gaussian Mixture Model	0.345	31.163	0.957	0.490	0.654

Clustering Results

Key Insights

- Focused parameters (e.g., Michigan Score, tissue thickness) yield better clusters (Silhouette 0.55) than full set (0.22), likely due to noise from less-informative features.
- Top predictors: Michigan Score, tissue thickness (HALLUX, TM5), temperature.
- IWGDF grades improve clustering but miss some variability.

Feature Importance

- **Logistic Regression (LASSO):** Michigan Score, Michigan Score 2, Tissue Thickness (SESA_R).
- **Random Forest:** Tissue Thickness (TM5_L, SESA_R), Michigan Score.
- **ANOVA (Eta^2):** Tissue Thickness (HALLUX_R), Michigan Score.

Clustering Rankings:

- **With IWGDF Grade:** Michigan Score + IWGDF grade, then Tissue Thickness (TM5_L, SESA_R).
- **Without IWGDF Grade:** Michigan Score + Tissue Thickness (SESA_R, HALLUX_R, TM5_L).

Correlations




Figure: Correlogram ($|r| > 0.7$)

To summarize

- **Left-right asymmetries** are important early signals.
- **Clustering** using selected features reveals meaningful patient groups.
- **Important indicators** include Michigan Score, tissue thickness, and temperature.
- **Take-home message:** Combining clinical and biomechanical data improves risk screening.

Limitations and Future Directions

Challenges

- Small sample (21 patients) limits statistics.
- Uneven patient numbers in IWGDF grades.
- Some measurements (ultrasound, pressure) may be biased.


Next Steps

- Include more patients for stronger results.
- Use better ways to handle missing data and track patients over time.
- Try deep learning to improve prediction.

Personal Reflections

Internship Experience

- Improved Python and data analysis skills.
- Collaborated with supervisors to ensure medical accuracy.
- Learned to apply data science in healthcare.



Useful URLs

- [Streamlit Cloud](#)
- [Scikit-learn Clustering Documentation](#)

References I



World Health Organization, *Global Report on Diabetes*, World Health Organization, April 2016, <https://www.who.int/publications/i/item/9789241565257>.



International Diabetes Federation, Number of adults (20–79 years) with diabetes in France, 2024, https://diabetesatlas.org/data-by-location/country/france/?utm_source=chatgpt.com.



International Diabetes Federation, Diabetes Facts & Figures, 2025, <https://idf.org/about-diabetes/diabetes-facts-figures/>.



Wikipedia contributors, Chronic wound, 2025, https://en.wikipedia.org/wiki/Chronic_wound?utm_source=chatgpt.com.

References II



S. Sivanandam, M. Anburajan, B. Venkatraman, M. Menaka, D. Sharath, *Medical thermography: a diagnostic approach for type 2 diabetes based on non-contact infrared thermal imaging*, Endocrine, vol. 42, no. 2, pp. 343–351, 2012.
<http://dx.doi.org/10.1007/s12020-012-9645-8>



S. Bagavathiappan, J. Philip, T. Jayakumar, B. Raj, P. N. S. Rao, M. Varalakshmi, V. Mohan, *Correlation between Plantar Foot Temperature and Diabetic Neuropathy: A Case Study by Using an Infrared Thermal Imaging Technique*, Journal of Diabetes Science and Technology, vol. 4, no. 6, pp. 1386–1392, 2010.
<http://dx.doi.org/10.1177/193229681000400613>



M. Oe, R. R. Yotsu, H. Sanada, T. Nagase, T. Tamaki, *Screening for Osteomyelitis Using Thermography in Patients with Diabetic Foot Ulcers*, vol. 2013, pp. 1–6, 2013.
<http://dx.doi.org/10.1155/2013/284294>

References III



J. J. van Netten, J. G. van Baal, C. Liu, F. van der Heijden, S. A. Bus, *Infrared Thermal Imaging for Automated Detection of Diabetic Foot Complications*, Journal of Diabetes Science and Technology, vol. 7, no. 5, pp. 1122–1129, 2013.
<http://dx.doi.org/10.1177/193229681300700504>



L. Fraiwan, M. AlKhodari, J. Ninan, B. Mustafa, A. Saleh, M. Ghazal, *Diabetic foot ulcer mobile detection system using smart phone thermal camera: a feasibility study*, BioMedical Engineering OnLine, vol. 16, no. 1, 2017.
<http://dx.doi.org/10.1186/s12938-017-0408-x>

References IV



B. W. S. Soh, A. Corrias, L. Tucker-Kellogg, *Computational modeling of the thin muscle layer, panniculus carnosus, demonstrates principles of pressure injury and prophylactic dressings*, In: Innovations and Emerging Technologies in Wound Care, Elsevier, pp. 41–54, 2020. <http://dx.doi.org/10.1016/B978-0-12-815028-3.00003-1>



N. J. M. Nasir, A. Corrias, H. Heemskerk, E. T. Ang, J. H. Jenkins, S. J. Sebastin, L. Tucker-Kellogg, *The panniculus carnosus muscle: a missing link in the chronicity of heel pressure ulcers?*, Journal of The Royal Society Interface, vol. 19, no. 187, 2022. <http://dx.doi.org/10.1098/rsif.2021.0631>

References V



S. L. Schnider, R. R. Kohn, *Effects of age and diabetes mellitus on the solubility of collagen from human skin, tracheal cartilage and dura mater*, Experimental Gerontology, vol. 17, no. 3, pp. 185–194, 1982.

[http://dx.doi.org/10.1016/0531-5565\(82\)90024-9](http://dx.doi.org/10.1016/0531-5565(82)90024-9)



J. J. van Netten, M. Prijs, J. G. van Baal, C. Liu, F. van der Heijden, S. A. Bus, *Diagnostic Values for Skin Temperature Assessment to Detect Diabetes-Related Foot Complications*, Diabetes Technology Therapeutics, vol. 16, no. 11, pp. 714–721, 2014.

<http://dx.doi.org/10.1089/dia.2014.0052>

References VI



L. Delbridge, C. S. Ellis, K. Robertson, L. P. Lequesne, *Non-enzymatic glycosylation of keratin from the stratum corneum of the diabetic foot*, British Journal of Dermatology, vol. 112, no. 5, pp. 547–554, 2006. <http://dx.doi.org/10.1111/j.1365-2133.1985.tb15262.x>



World Health Organization, *Global Report on Diabetes*, World Health Organization, Geneva, 2016. <https://www.who.int/publications/i/item/9789241565257>



International Diabetes Federation, *IDF Diabetes Atlas, 11th Edition*, International Diabetes Federation, Brussels, 2024. <https://idf.org/aboutdiabetes/diabetesatlas/>

References VII



T. Brink, *Induration of the Diabetic Foot Pad: Another Risk Factor for Recurrent Neuropathic Plantar Ulcers - Verhärtung der diabetischen Fußsohle - eine Prädisposition für rezidivierende neuropathische Fußgeschwüre*, Biomedizinische Technik/Biomedical Engineering, vol. 40, no. 7–8, pp. 205–209, 1995.
<http://dx.doi.org/10.1515/bmte.1995.40.7-8.205>

Thank You!

Questions?

Thank you for your attention!