

Exploring Biomechanical and Clinical Factors in Diabetic Foot Ulcer Risk Assessment

Helya Amiri

Supervised by: Pierre-Yves Rohan, Jerome Haddad

University of Strasbourg, France

August 20, 2025


Outline

- 1 Introduction
- 2 Methodology
- 3 Conclusion

Introduction

- **Diabetes:** A long-term disease where blood sugar stays high because the body makes too little insulin or does not use it well. It can harm nerves and blood vessels.
- **Neuropathy:** Nerve damage from diabetes that can cause foot pain, numbness, or weakness.

Diabetic Foot Ulcers



Risk Factors for Diabetic Foot Ulcers

- **Neuropathy:** Less feeling in the feet, so injuries happen more easily.
- **Poor Circulation:** Blood flow is weak, so wounds heal slowly.
- **Foot Deformities:** Feet have unusual shapes, causing more pressure on some spots.




Figure: Plantar Pressure Map

Why Study Diabetic Foot Ulcers?

- Over 4 million adults in France are living with diabetes (2024).
- Diabetic Foot Ulcers (DFUs) cause more than 10,000 amputations each year.
- **Goal:** Create a tool to study diabetic foot data and help doctors make decisions.

Objectives

Internship Goals

- Find key risk markers (e.g., tissue stiffness, plantar pressure).
- Study how tissue properties relate to foot loading.
- Explore left-right foot differences as warning signs.
- Group patients and compare with IWGDF risk levels (grades that show how likely a diabetic foot is to get ulcers).

Data Collection

- Study done in a day-hospital with ethical approval.
- Participants signed consent forms.
- Included adults 18–65 with diabetes.
- Excluded pregnant, breastfeeding, or legally protected people.

The DIAFOOT Dataset

- **Participants:** 21 diabetic patients (11 females, 10 males).
- **Age:** Mean 56.8 years.
- **BMI:** Mean 25.83 kg/m²
- **IWGDF Grades:** 0 (n=11), 1 (n=2), 2 (n=1), 3 (n=7) (Risk levels for diabetic foot ulcers.)




Figure: Age Distribution

Data Parameters

- **Anthropometric:** BMI, height, weight
- **Vascular:** Ankle and big toe blood pressure
- **Neurological:** Michigan score, SUDOSCAN (hands/feet)
- **Biomechanical:** Foot pressure (SESA, hallux, TM5), tissue stiffness
- **Imaging:** Ultrasound tissue thickness
- **Thermal:** Foot temperature and normalized values

Analysis Pipeline

Streamlit Dashboard

- **Platform:** Built using Streamlit, A dynamic, Python-driven platform for real-time data exploration and visualization.
- **Data Source:** Excel file with 'DIAFOOT' sheet containing patient metrics.
- **Key Features:** Descriptive statistics, normality tests, clustering, correlations and comparative analyses.
- **Tools:** Pandas (data handling), NumPy (numerical), Scikit-learn (ML), Matplotlib Seaborn (visuals).

Explore the DIAFOOT Analysis Dashboard

<https://diafoot-analysis.streamlit.app/>

Upload data, analyze it, and see important results easily.

Try it now!

Review of Statistics: p-value

- **p-value:** Shows if a result is real or by chance.
- Small p-value $< 0.05 \rightarrow$ **significant** (result likely real)
- Large p-value $> 0.05 \rightarrow$ **not significant** (result may be by chance)
- Used to decide whether to reject the null hypothesis.

Asymmetry Insights

Significant Left-Right Differences

- **TM5 Pressure:** Right (36.93 kPa) > Left (26.34 kPa), $p = 0.0054$.
- **Hallux Stiffness:** Right higher, $p = 0.0498$.
- **Temperature (Plantar Arch, Lateral Sole):** Right warmer, $p < 0.01$.




Figure: Pressure Asymmetry

Group Differences: Low vs. High Risk

Parameter	Grades 0-1	Grades 2-3	p-value
Michigan Score	11.69	4.13	0.00053
Tissue Thickness SESA R (mm)	9.55	4.17	0.0099
Heel Temperature R (°C)	25.85	28.55	0.0188

Table: Significant Group Differences

- Higher-risk patients have lower Michigan scores, thinner tissue, and warmer heels, showing higher ulcer risk.

Clustering Analysis

- Clustering done with and without IWGDF grades
- Analyses run on:
 - Key parameters: IWGDF grade, MESI, Michigan, pressure, stiffness, thickness, amplitude, temperature.
 - Full dataset: 70+ clinical and biomechanical variables.
- **PCA:** 2D visualization of patient data.
- **Clustering:** Agglomerative, KMeans, GMM; 2–6 clusters.
- **Metrics:** Silhouette, Calinski–Harabasz, Davies–Bouldin, ARI, NMI.
- **Feature importance:** LASSO, Random Forest, ANOVA.
- **Group differences:** MANOVA, p-values indicate significance.

Clustering Results

Key Insights

- Focused parameters (e.g., Michigan Score, tissue thickness) yield better clusters (Silhouette 0.55) than full set (0.22).
- Top predictors: Michigan Score, tissue thickness (HALLUX, TM5), temperature.
- IWGDF grades improve clustering but miss some variability.

Correlations




Figure: Correlogram ($|r| > 0.7$)

Key Takeaways

- **Asymmetries** in pressure, stiffness, and temperature show which patients are at risk.
- **Clustering** using selected features reveals meaningful patient groups.
- **Important indicators** include Michigan Score, tissue thickness, and temperature.
- **Take-home message:** Combining clinical and biomechanical data improves risk screening.

Limitations and Future Directions

Challenges

- Small sample (21 patients) limits statistics.
- Uneven patient numbers in IWGDF grades.
- Some measurements (ultrasound, pressure) may be biased.


Next Steps

- Include more patients for stronger results.
- Use better ways to handle missing data and track patients over time.
- Try deep learning to improve prediction.

Personal Reflections

Internship Experience

- Improved Python and data analysis skills.
- Collaborated with supervisors to ensure medical accuracy.
- Learned to apply data science in healthcare.



Useful URLs

- [Streamlit Cloud](#)
- [Scikit-learn Clustering Documentation](#)

References I

-  World Health Organization, *Global Report on Diabetes*, World Health Organization, April 2016, <https://www.who.int/publications/i/item/9789241565257>.
-  International Diabetes Federation, Number of adults (20–79 years) with diabetes in France, 2024, https://diabetesatlas.org/data-by-location/country/france/?utm_source=chatgpt.com.
-  International Diabetes Federation, Diabetes Facts & Figures, 2025, <https://idf.org/about-diabetes/diabetes-facts-figures/>.
-  Wikipedia contributors, Chronic wound, 2025, https://en.wikipedia.org/wiki/Chronic_wound?utm_source=chatgpt.com.

References II

- 

S. Sivanandam, M. Anburajan, B. Venkatraman, M. Menaka, D. Sharath, *Medical thermography: a diagnostic approach for type 2 diabetes based on non-contact infrared thermal imaging*, Endocrine, vol. 42, no. 2, pp. 343–351, 2012.
<http://dx.doi.org/10.1007/s12020-012-9645-8>
- 

S. Bagavathiappan, J. Philip, T. Jayakumar, B. Raj, P. N. S. Rao, M. Varalakshmi, V. Mohan, *Correlation between Plantar Foot Temperature and Diabetic Neuropathy: A Case Study by Using an Infrared Thermal Imaging Technique*, Journal of Diabetes Science and Technology, vol. 4, no. 6, pp. 1386–1392, 2010.
<http://dx.doi.org/10.1177/193229681000400613>
- 

M. Oe, R. R. Yotsu, H. Sanada, T. Nagase, T. Tamaki, *Screening for Osteomyelitis Using Thermography in Patients with Diabetic Foot, Ulcers*, vol. 2013, pp. 1–6, 2013.
<http://dx.doi.org/10.1155/2013/284294>

References III



J. J. van Netten, J. G. van Baal, C. Liu, F. van der Heijden, S. A. Bus, *Infrared Thermal Imaging for Automated Detection of Diabetic Foot Complications*, Journal of Diabetes Science and Technology, vol. 7, no. 5, pp. 1122–1129, 2013.

<http://dx.doi.org/10.1177/193229681300700504>



L. Fraiwan, M. AlKhodari, J. Ninan, B. Mustafa, A. Saleh, M. Ghazal, *Diabetic foot ulcer mobile detection system using smart phone thermal camera: a feasibility study*, BioMedical Engineering OnLine, vol. 16, no. 1, 2017.

<http://dx.doi.org/10.1186/s12938-017-0408-x>

References IV



B. W. S. Soh, A. Corrias, L. Tucker-Kellogg, *Computational modeling of the thin muscle layer, panniculus carnosus, demonstrates principles of pressure injury and prophylactic dressings*, In: Innovations and Emerging Technologies in Wound Care, Elsevier, pp. 41–54, 2020. <http://dx.doi.org/10.1016/B978-0-12-815028-3.00003-1>



N. J. M. Nasir, A. Corrias, H. Heemskerk, E. T. Ang, J. H. Jenkins, S. J. Sebastin, L. Tucker-Kellogg, *The panniculus carnosus muscle: a missing link in the chronicity of heel pressure ulcers?*, Journal of The Royal Society Interface, vol. 19, no. 187, 2022. <http://dx.doi.org/10.1098/rsif.2021.0631>

References V

-  S. L. Schnider, R. R. Kohn, *Effects of age and diabetes mellitus on the solubility of collagen from human skin, tracheal cartilage and dura mater*, Experimental Gerontology, vol. 17, no. 3, pp. 185-194, 1982.
[http://dx.doi.org/10.1016/0531-5565\(82\)90024-9](http://dx.doi.org/10.1016/0531-5565(82)90024-9)
-  J. J. van Netten, M. Prijs, J. G. van Baal, C. Liu, F. van der Heijden, S. A. Bus, *Diagnostic Values for Skin Temperature Assessment to Detect Diabetes-Related Foot Complications*, Diabetes Technology Therapeutics, vol. 16, no. 11, pp. 714-721, 2014.
<http://dx.doi.org/10.1089/dia.2014.0052>

References VI



L. Delbridge, C. S. Ellis, K. Robertson, L. P. Lequesne, *Non-enzymatic glycosylation of keratin from the stratum corneum of the diabetic foot*, British Journal of Dermatology, vol. 112, no. 5, pp. 547–554, 2006. <http://dx.doi.org/10.1111/j.1365-2133.1985.tb15262.x>



T. Brink, *Induration of the Diabetic Foot Pad: Another Risk Factor for Recurrent Neuropathic Plantar Ulcers - Verhärtung der diabetischen Fußsohle - eine Prädisposition für rezidivierende neuropathische Fußgeschwüre*, Biomedizinische Technik/Biomedical Engineering, vol. 40, no. 7–8, pp. 205–209, 1995. <http://dx.doi.org/10.1515/bmte.1995.40.7-8.205>

Thank You!

Questions?

Thank you for your attention!