

# Advanced statistics and modelling

5. week

## Parameter inference in statistics Hypothesis testing, MLE, Bootstrap

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed

# The basic statistical inference problem

Hypothesis  
testing, MLE,  
Bootstrap

- Recall: The basic statistical inference problem was the following:
  - We have some observed data:  $X_1, X_2, \dots, X_n \sim F$ .
  - Based on the observations we would like to **infer** (or estimate or learn) some parameters (e.g.  $p$  in a Binomial).
- In data science: based on the observations validate an assumption (hypothesis) on some parameters.
- Statistical decision: calculate the probability, that the observations are consistent with the hypothesis.
- Calculate some parameters (3 methods):
  - assume normal distribution,
  - assume the observation is the most probable realization
  - assume all possible values to be observed



# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter
- We collect data  $X \in \mathcal{X}$ , and we want to make a decision on the hypothesis.

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter
- We collect data  $X \in \mathcal{X}$ , and we want to make a decision on the hypothesis.
- The **rejection region** is  $R \subset \mathcal{X}$  if  
 $X \in R \Rightarrow \text{reject } H_0$   
 $X \notin R \Rightarrow \text{retain(not reject)} H_0$

# Formal definition

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter
- We collect data  $X \in \mathcal{X}$ , and we want to make a decision on the hypothesis.
- The **rejection region** is  $R \subset \mathcal{X}$  if  
 $X \in R \Rightarrow \text{reject } H_0$   
 $X \notin R \Rightarrow \text{retain(not reject)} H_0$
- A function  $T : \mathcal{X} \rightarrow \mathbb{R}$  is a **test statistics**

# Formal definition

Hypothesis  
testing, MLE,  
Bootstrap

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter
- We collect data  $X \in \mathcal{X}$ , and we want to make a decision on the hypothesis.
- The **rejection region** is  $R \subset \mathcal{X}$  if  
 $X \in R \Rightarrow \text{reject } H_0$   
 $X \notin R \Rightarrow \text{retain(not reject)} H_0$
- A function  $T : \mathcal{X} \rightarrow \mathbb{R}$  is a **test statistics**
- The **critical value** is  $c$  if the rejection region can be expressed as  
 $R = \{x \in \mathcal{X} \mid T(x) > c\}$



# Formal definition

Hypothesis  
testing, MLE,  
Bootstrap

- The **parameter space**  $\Theta$  is partitioned:  
 $\Theta_0$  and  $\Theta_1$  ( $\Theta_0 \cap \Theta_1 = \emptyset$ ).
- The **null hypothesis** is  $H_0 : \theta \in \Theta_0$
- The **alternative hypothesis** is  $H_1 : \theta \in \Theta_1$
- Note:  $\theta$  is a fixed parameter
- We collect data  $X \in \mathcal{X}$ , and we want to make a decision on the hypothesis.
- The **rejection region** is  $R \subset \mathcal{X}$  if  
 $X \in R \Rightarrow \text{reject } H_0$   
 $X \notin R \Rightarrow \text{retain(not reject)} H_0$
- A function  $T : \mathcal{X} \rightarrow \mathbb{R}$  is a **test statistics**
- The **critical value** is  $c$  if the rejection region can be expressed as  
 $R = \{x \in \mathcal{X} \mid T(x) > c\}$
- The **hypothesis test** is to find  $T$  and  $c$ , which leads the least harmful decision.  
(e.g. Do I have cancer? or Do I got the most points for home work?)

# Formal definition

- Possible outcomes:

	Retain $H_0$	Reject $H_0$
$H_0$ true	OK	type I error
$H_1$ true	type II error	OK

# Formal definition

- Possible outcomes: 

	Retain $H_0$	Reject $H_0$
$H_0$ true	OK	type I error
$H_1$ true	type II error	OK

- The **power function of the test** is a  $\Theta \rightarrow \mathbb{R}$

$$\beta(\theta) = P_{\theta}(X \in R)$$

where  $R$  is the rejection region and  $X$  are some collected data.

# Formal definition

- Possible outcomes: 

	Retain $H_0$	Reject $H_0$
$H_0$ true	OK	type I error
$H_1$ true	type II error	OK

- The **power function of the test** is a  $\Theta \rightarrow \mathbb{R}$

$$\beta(\theta) = P_{\theta}(X \in R)$$

where  $R$  is the rejection region and  $X$  are some collected data.

- The test is **two sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta \neq \theta_0$

# Formal definition

- Possible outcomes: 

	Retain $H_0$	Reject $H_0$
$H_0$ true	OK	type I error
$H_1$ true	type II error	OK

- The **power function of the test** is a  $\Theta \rightarrow \mathbb{R}$

$$\beta(\theta) = P_{\theta}(X \in R)$$

where  $R$  is the rejection region and  $X$  are some collected data.

- The test is **two sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta \neq \theta_0$
- The test is **one sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta > \theta_0$   
or  $H_0 : \theta = \theta_0$  and  $H_1 : \theta < \theta_0$

# Formal definition

- Possible outcomes: 

	Retain $H_0$	Reject $H_0$
$H_0$ true	OK	type I error
$H_1$ true	type II error	OK

- The **power function of the test** is a  $\Theta \rightarrow \mathbb{R}$

$$\beta(\theta) = P_{\theta}(X \in R)$$

where  $R$  is the rejection region and  $X$  are some collected data.

- The test is **two sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta \neq \theta_0$
- The test is **one sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta > \theta_0$   
or  $H_0 : \theta = \theta_0$  and  $H_1 : \theta < \theta_0$

- The **level** of the test is  $\alpha = \sup_{\theta \in \Theta_0} \beta(\theta)$

the maximum probability for rejecting the observations, when the parameter is in the retained set.

# Formal definition

	Retain $H_0$	Reject $H_0$
Possible outcomes:	$H_0$ true OK	type I error
	$H_1$ true type II error	OK

- The **power function of the test** is a  $\Theta \rightarrow \mathbb{R}$   $\beta(\theta) = P_\theta(X \in R)$  where  $R$  is the rejection region and  $X$  are some collected data.
- The test is **two sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta \neq \theta_0$
- The test is **one sided** if  $H_0 : \theta = \theta_0$  and  $H_1 : \theta > \theta_0$   
or  $H_0 : \theta = \theta_0$  and  $H_1 : \theta < \theta_0$
- The **level** of the test is  $\alpha = \sup_{\theta \in \Theta_0} \beta(\theta)$   
the maximum probability for rejecting the observations, when the parameter is in the retained set.

Note some alternatives in the literature:

- $H_0 \leq \theta_0$  for one sided tests (later we see problems with this)
- $1 - \beta(\theta) = P_\theta(X \in R)$  (though the **power** is the same!)

Example:

We observe  $X_1, X_2, \dots, X_n$  standard normally distributed, independent variables, and we want to test:

Do these values tend to be negative or positive?

.



Example:

We observe  $X_1, X_2, \dots, X_n$  standard normally distributed, independent variables, and we want to test:

Do these values tend to be negative or positive?

.

Possible solution: the variables are from a  $N(\mu = 0, \sigma)$  distribution and we test for  $\mu \leq 0$ .

Example:

We observe  $X_1, X_2, \dots, X_n$  standard normally distributed, independent variables, and we want to test:

Do these values tend to be negative or positive?

.

Possible solution: the variables are from a  $N(\mu = 0, \sigma)$  distribution and we test for  $\mu \leq 0$ .

Parameter space:  $\Theta = (-\infty, \infty)$ ,

partition:  $\Theta_0 = (-\infty, 0]$ ,  $\Theta_1 = (0, \infty)$ ,

Example:

We observe  $X_1, X_2, \dots, X_n$  standard normally distributed, independent variables, and we want to test:

Do these values tend to be negative or positive?

.

Possible solution: the variables are from a  $N(\mu = 0, \sigma)$  distribution and we test for  $\mu \leq 0$ .

Parameter space:  $\Theta = (-\infty, \infty)$ ,

partition:  $\Theta_0 = (-\infty, 0]$ ,  $\Theta_1 = (0, \infty)$ ,

Null hypothesis:  $H_0 : \mu = 0$ ,  $H_1 : \mu > 0$

(Note the different relations  $H_0 \leftrightarrow \Theta_0$  and  $H_1 \leftrightarrow \Theta_1$ !)

Example:

We observe  $X_1, X_2, \dots, X_n$  standard normally distributed, independent variables, and we want to test:

Do these values tend to be negative or positive?

.

Possible solution: the variables are from a  $N(\mu = 0, \sigma)$  distribution and we test for  $\mu \leq 0$ .

Parameter space:  $\Theta = (-\infty, \infty)$ ,

partition:  $\Theta_0 = (-\infty, 0]$ ,  $\Theta_1 = (0, \infty)$ ,

Null hypothesis:  $H_0 : \mu = 0$ ,  $H_1 : \mu > 0$

(Note the different relations  $H_0 \leftrightarrow \Theta_0$  and  $H_1 \leftrightarrow \Theta_1$ !)

Let us define  $T = \bar{X}$  (average of observed values), and find  $c$  which rejects  $H_0$  if  $T > c$ .

Rejection region:  $R = \{x : T(x) > c\}$ .

# Example

Hypothesis  
testing, MLE,  
Bootstrap

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu} \left( \frac{\bar{X} - \mu}{\sigma} \sqrt{n} > \sqrt{n} \frac{c - \mu}{\sigma} \right).\end{aligned}$$

# Example

Hypothesis  
testing, MLE,  
Bootstrap

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu} \left( \frac{\bar{X} - \mu}{\sigma} \sqrt{n} > \sqrt{n} \frac{c - \mu}{\sigma} \right).\end{aligned}$$

Because  $\sqrt{n} \frac{\bar{X} - \mu}{\sigma}$  is the  $z$ -transformed (normalized) value of the average,

# Example

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu}\left(\frac{\bar{X} - \mu}{\sigma}\sqrt{n} > \sqrt{n}\frac{c - \mu}{\sigma}\right).\end{aligned}$$

Because  $\sqrt{n}\frac{\bar{X} - \mu}{\sigma}$  is the  $z$ -transformed (normalized) value of the average, it is  $Z \sim N(0, 1)$  distributed:

$$\begin{aligned}\beta(\theta) = \beta(\mu) &= P_{\mu}\left(Z > \sqrt{n}\frac{c - \mu}{\sigma}\right) \\ &= 1 - \Phi\left(\sqrt{n}\frac{c - \mu}{\sigma}\right)\end{aligned}$$

# Example

Hypothesis  
testing, MLE,  
Bootstrap

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu}\left(\frac{\bar{X} - \mu}{\sigma}\sqrt{n} > \sqrt{n}\frac{c - \mu}{\sigma}\right).\end{aligned}$$

Because  $\sqrt{n}\frac{\bar{X} - \mu}{\sigma}$  is the  $z$ -transformed (normalized) value of the average, it is  $Z \sim N(0, 1)$  distributed:

$$\begin{aligned}\beta(\theta) = \beta(\mu) &= P_{\mu}\left(Z > \sqrt{n}\frac{c - \mu}{\sigma}\right) \\ &= 1 - \Phi\left(\sqrt{n}\frac{c - \mu}{\sigma}\right)\end{aligned}$$

Note, that  $\beta(\mu)$  is increasing and  $\Theta_0 = (-\infty, 0]$ ,



# Example

Hypothesis  
testing, MLE,  
Bootstrap

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu}\left(\frac{\bar{X} - \mu}{\sigma}\sqrt{n} > \sqrt{n}\frac{c - \mu}{\sigma}\right).\end{aligned}$$

Because  $\sqrt{n}\frac{\bar{X} - \mu}{\sigma}$  is the  $z$ -transformed (normalized) value of the average, it is  $Z \sim N(0, 1)$  distributed:

$$\begin{aligned}\beta(\theta) = \beta(\mu) &= P_{\mu}\left(Z > \sqrt{n}\frac{c - \mu}{\sigma}\right) \\ &= 1 - \Phi\left(\sqrt{n}\frac{c - \mu}{\sigma}\right)\end{aligned}$$

Note, that  $\beta(\mu)$  is increasing and  $\Theta_0 = (-\infty, 0]$ , so  $\sup \beta(\mu)$  for  $\mu \in \Theta_0$  is at  $\mu = 0$ .

# Example

The power function:

$$\begin{aligned}\beta(\theta) &= P_{\mu}(\bar{X} > c) \\ &= P_{\mu} \left( \frac{\bar{X} - \mu}{\sigma} \sqrt{n} > \sqrt{n} \frac{c - \mu}{\sigma} \right).\end{aligned}$$

Because  $\sqrt{n} \frac{\bar{X} - \mu}{\sigma}$  is the  $z$ -transformed (normalized) value of the average, it is  $Z \sim N(0, 1)$  distributed:

$$\begin{aligned}\beta(\theta) = \beta(\mu) &= P_{\mu} \left( Z > \sqrt{n} \frac{c - \mu}{\sigma} \right) \\ &= 1 - \Phi \left( \sqrt{n} \frac{c - \mu}{\sigma} \right)\end{aligned}$$

Note, that  $\beta(\mu)$  is increasing and  $\Theta_0 = (-\infty, 0]$ , so  $\sup \beta(\mu)$  for  $\mu \in \Theta_0$  is at  $\mu = 0$ .

Find the level of the test:

$$\begin{aligned}\alpha &= \sup_{\mu \in [-\infty, 0]} \beta(\mu) = \sup_{\mu \in [-\infty, 0]} \left\{ 1 - \Phi \left( \sqrt{n} \frac{c - \mu}{\sigma} \right) \right\} \\ &= 1 - \Phi \left( \sqrt{n} \frac{c}{\sigma} \right)\end{aligned}$$

*or vica – verse :* at given level find the critical value

$$c = \Phi^{-1}(1 - \alpha) \frac{\sigma}{\sqrt{n}}$$

# Example

Hypothesis  
testing, MLE,  
Bootstrap

We reject  $H_0$

$$\text{if } T = \bar{X} > c = \Phi^{-1}(1 - \alpha) \frac{\sigma}{\sqrt{n}}$$

# Example

Hypothesis  
testing, MLE,  
Bootstrap

We reject  $H_0$

$$\text{if } T = \bar{X} > c = \Phi^{-1}(1 - \alpha) \frac{\sigma}{\sqrt{n}}$$

Or

$$\frac{\sqrt{n}(\bar{X} - 0)}{\sigma} > z_{\alpha}$$

where  $z_{\alpha}$  is the tabulated critical value of the standard normal distribution.

.

# Example

Hypothesis  
testing, MLE,  
Bootstrap

We reject  $H_0$

$$\text{if } T = \bar{X} > c = \Phi^{-1}(1 - \alpha) \frac{\sigma}{\sqrt{n}}$$

Or

$$\frac{\sqrt{n}(\bar{X} - 0)}{\sigma} > z_{\alpha}$$

where  $z_{\alpha}$  is the tabulated critical value of the standard normal distribution.

Note the substituted value of  $\mu = 0$ , which is the  $H_0$  null hypothesis expressed as an equation, **instead of an inequality**.

# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

Test statistics: reject  $H_0$  if

# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

Test statistics: reject  $H_0$  if

$$\left| \frac{(\hat{\theta} - \theta)}{\hat{se}} \right| > z_{\alpha/2}$$

in case of a two sided test



# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

Test statistics: reject  $H_0$  if

$$\frac{(\hat{\theta} - \theta)}{\hat{se}} > z_{\alpha}$$

in case of a one sided test

# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

Test statistics: reject  $H_0$  if

$$\frac{(\hat{\theta} - \theta)}{\hat{se}} < z_{\alpha}$$

in an other case of a one sided test

# Standard test I.: Wald test

This test is valid for asymptotically normally distributed variables.

- $\theta$  denotes the real value of the parameter,
- $\hat{\theta}$  an estimated value of the parameter,
- $\hat{se}$  estimated standard error of  $\hat{\theta}$

The formal test:  $H_0 : \theta = \hat{\theta}$  and  $H_1 : \theta \neq \hat{\theta}$

Test statistics: reject  $H_0$  if

$$\left| \frac{(\hat{\theta} - \theta)}{\hat{se}} \right| > z_{\alpha/2}$$

$$\frac{(\hat{\theta} - \theta)}{\hat{se}} > z_{\alpha} \quad or \quad \frac{(\hat{\theta} - \theta)}{\hat{se}} < -z_{\alpha}$$

Note the difference between the one sided and the two sided test:

one sided compares with:  $z_{\alpha}$  and no  $| \cdot |$

two sided compares with:  $z_{\alpha/2}$  and takes absolute value

# Some properties of Wald test

Hypothesis  
testing, MLE,  
Bootstrap

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

# Some properties of Wald test

Hypothesis  
testing, MLE,  
Bootstrap

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$P(X \in R) = 1 - P(X \notin R)$$

where

$$P(X \notin R) = P(\theta_{\star} \text{ close to } \hat{\theta}) = \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} + z_{\alpha/2}\right) - \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} - z_{\alpha/2}\right)$$

so

# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$P(X \in R) = 1 - P(X \notin R)$$

where

$$P(X \notin R) = P(\theta_{\star} \text{ close to } \hat{\theta}) = \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} + z_{\alpha/2}\right) - \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} - z_{\alpha/2}\right)$$

so

$$\beta = 1 - \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} + z_{\alpha/2}\right) + \Phi\left(\frac{\hat{\theta} - \theta_{\star}}{\hat{se}} - z_{\alpha/2}\right)$$

or with the true value

$$\beta = 1 - \Phi\left(\frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} + z_{\alpha/2}\right) + \Phi\left(\frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} - z_{\alpha/2}\right)$$

# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$\beta = 1 - \Phi \left( \frac{\hat{\theta} - \theta_{\star}}{\hat{se}} + z_{\alpha/2} \right) + \Phi \left( \frac{\hat{\theta} - \theta_{\star}}{\hat{se}} - z_{\alpha/2} \right)$$

or with the true value

$$\beta = 1 - \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} + z_{\alpha/2} \right) + \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} - z_{\alpha/2} \right)$$

which is the probability correctly rejecting a false  $H_0$  under assuming  $\theta_{\star}$  as the parameter instead  $\theta$ .



# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$\beta = 1 - \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} + z_{\alpha/2} \right) + \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} - z_{\alpha/2} \right)$$

which is the probability correctly rejecting a false  $H_0$  under assuming  $\theta_{\star}$  as the parameter instead  $\theta$ .

The power increases if:

- difference between  $\theta_{\star}$  and  $\theta$  increases (note: we change only  $\theta_{\star}$ )

# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$\beta = 1 - \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} + z_{\alpha/2} \right) + \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} - z_{\alpha/2} \right)$$

which is the probability correctly rejecting a false  $H_0$  under assuming  $\theta_{\star}$  as the parameter instead  $\theta$ .

The power increases if:

- difference between  $\theta_{\star}$  and  $\theta$  increases (note: we change only  $\theta_{\star}$ )
- sample variance decreases (  $\hat{se} \sim 1/\sqrt{n}$  decreases as  $n$  grows)

# Some properties of Wald test

When the true value would be different what we expect as real parameter:

$$\theta_{\star} \neq \theta,$$

the power of the test ( $\beta = P(X \in R)$ ) will be:

$$\beta = 1 - \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} + z_{\alpha/2} \right) + \Phi \left( \frac{(\hat{\theta} - \theta) - (\theta_{\star} - \theta)}{\hat{se}} - z_{\alpha/2} \right)$$

which is the probability correctly rejecting a false  $H_0$  under assuming  $\theta_{\star}$  as the parameter instead  $\theta$ .

The power increases if:

- difference between  $\theta_{\star}$  and  $\theta$  increases (note: we change only  $\theta_{\star}$ )
- sample variance decreases ( $\hat{se} \sim 1/\sqrt{n}$  decreases as  $n$  grows)

Wald test is equivalent with a confidence interval:

$$C = (\hat{\theta} - z_{\alpha/2}\hat{se}, \hat{\theta} + z_{\alpha/2}\hat{se},)$$

$H_0 : \hat{\theta} = \theta_{\star}$  rejected at level  $\alpha$  if and only if  $\theta_{\star} \notin C$

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$   
and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.  
and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1 - p_1)/n)$

( $p_2$  similar)

and



# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1 - p_1)/n)$   
( $p_2$  similar)

If the two samples are independent, variances are additive.  
and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$   
( $p_2$  similar)

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$   
( $p_2$  similar)

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

$$\beta = P(z_{\text{sample}} > z_{\text{crit}})$$

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$   
( $p_2$  similar)

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

$$\beta = P(z_{\text{sample}} > z_{\text{crit}})$$

Conclusion: power increases as  $\hat{p}_1$  is farther from  $\hat{p}_2$   
or sample sizes ( $n$  and  $m$ ) increase

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

$$\beta = P(z_{\text{sample}} > z_{\text{crit}})$$

Conclusion: power increases as  $\hat{p}_1$  is farther from  $\hat{p}_2$   
or sample sizes ( $n$  and  $m$ ) increase

Note: the minimum of  $\min(n, m)$  dominates!

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

$$\beta = P(z_{\text{sample}} > z_{\text{crit}})$$

Conclusion: power increases as  $\hat{p}_1$  is farther from  $\hat{p}_2$   
or sample sizes ( $n$  and  $m$ ) increase

Note: the minimum of  $\min(n, m)$  dominates!

Note2:  $X$  and  $Y$  independent, two different samples

# Standard test II: Comparing two predictions

Hypothesis  
testing, MLE,  
Bootstrap

$W \sim \text{Binomial}(p_1, n)$  and  $Y \sim \text{Binomial}(p_2, m)$  are the number of incorrect predictions of two methods in two different samples.

Question: is  $p_1 = p_2$ ?

Test:  $H_0 : \delta = p_1 - p_2 = 0$  and  $H_1 : \delta \neq 0$

Power of the test:  $\beta = P(X \in R)$

For large samples: Binomial  $\sim$  Gauss distribution.

$\hat{p}_1 = \frac{W}{n}$  is approx. Normal distr.:  $N(\mu, \sigma^2) = N(p_1, p_1(1-p_1)/n)$

If the two samples are independent, variances are additive.

Results to:

$$z_{\text{sample}} = \frac{\delta - 0}{\hat{se}} = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\frac{\hat{p}_1(1-\hat{p}_1)}{n} + \frac{\hat{p}_2(1-\hat{p}_2)}{m}}}$$

and

$$\beta = P(z_{\text{sample}} > z_{\text{crit}})$$

Conclusion: power increases as  $\hat{p}_1$  is farther from  $\hat{p}_2$   
or sample sizes ( $n$  and  $m$ ) increase

Note: the minimum of  $\min(n, m)$  dominates!

Note2:  $X$  and  $Y$  independent, two different samples

Note3: if  $X$  and  $Y$  measured together (**paired samples**), then  $\delta = \overline{D}$   
and  $\hat{se}^2 = \overline{(D - \overline{D})^2} / \sqrt{n}$ , where  $D_i = X_i - Y_i$  for each case  $i$

# Yet another definition: p-value

- If a test rejects at level  $\alpha$ , then it rejects for all  $\alpha' > \alpha$ .
- The rejection region  $R$  depends on the  $\alpha$  level (smaller  $\alpha$  results smaller  $R$ ).
- The **p-value** of a test is the smallest  $\alpha$  where the test rejects.

$$\text{p-value} = \inf \{ \alpha \mid \exists X \in R_\alpha \}$$

Recall:  $X$  is a random variable taking values from the observed values  $\mathcal{X}$   
Note: the p-value is a measure **against**  $H_0$ , and different from  $\mathbb{P}(H_0|data)$  (which is the probability of  $H_0$  being true with the condition of observed data.) The latter will be discussed under Bayesian inference later.



# Properties of p-value

Hypothesis  
testing, MLE,  
Bootstrap

- The smaller the p-value, the stronger evidence against  $H_0$

# Properties of p-value

Hypothesis  
testing, MLE,  
Bootstrap

- The smaller the p-value, the stronger evidence against  $H_0$
- Large p-value can result when:
  - $H_0$  is true

# Properties of p-value

- The smaller the p-value, the stronger evidence against  $H_0$
- Large p-value can result when:
  - $H_0$  is true
  - $H_1$  is true and the test has low power

# Properties of p-value

- The smaller the p-value, the stronger evidence against  $H_0$
- Large p-value can result when:
  - $H_0$  is true
  - $H_1$  is true and the test has low power

Let's calculate the p-value:

$$\text{reject } H_0 : T(X) > c_\alpha \Leftrightarrow \text{p-value} = \sup_{\theta \in \Theta_0, \xi \in \mathcal{X}} \mathbb{P}_\theta(T(\xi) \geq T(X))$$

which is the probability of observing a test statistics as extreme or more extreme that was actually observed.

# Properties of p-value

- Why prefer the null hypothesis  $H_0 : \theta = \theta_0$  against  $H_0 : \theta \leq \theta_0$ ?
  - In practice, the probability is calculated at a given value of the parameter  $\theta$ , which is set according to the null hypothesis.
  - If the distribution of  $T(X)$  is continuous, and  $H_0 : \theta = \theta_0$ , then the p-value has a uniform distribution on  $[0,1]$

The test is **statistically significant** if  $\alpha > \text{p-value}$ .

# Properties of p-value

- Why prefer the null hypothesis  $H_0 : \theta = \theta_0$  against  $H_0 : \theta \leq \theta_0$ ?
  - In practice, the probability is calculated at a given value of the parameter  $\theta$ , which is set according to the null hypothesis.
  - If the distribution of  $T(X)$  is continuous, and  $H_0 : \theta = \theta_0$ , then the p-value has a uniform distribution on  $[0,1]$

The test is **statistically significant** if  $\alpha > \text{p-value}$ .

Note: a test can be statistically significant, but practically not significant, if the confidence interval is very small.

# Calculating $c$ and $T$ in general

Hypothesis  
testing, MLE,  
Bootstrap

The Ward test assumes normal distributions:  $N(\mu, \sigma^2)$  needs the expected value and the variance of the quantity.

How to calculate  $\overline{(\cdot)}$ ,  $\hat{se}$  and how to derive  $T(x)$  functions in general?

Some methods:

- Bootstrap
- Method of Moments
- Maximum Likelihood Estimation

# Bootstrap

Hypothesis  
testing, MLE,  
Bootstrap

Theory behind this method:

$$\mathbb{E}(Y) = \lim_{B \rightarrow \infty} \frac{1}{B} \sum_{i=1}^B Y_i$$

For each function  $h$  with a finite mean

$$\mathbb{E}(h(Y)) = \lim_{B \rightarrow \infty} \frac{1}{B} \sum_{i=1}^B h(Y_i) = \langle h \rangle_B$$

Here  $B$  is the number of independent measurements (sampling with replacement) on the original dataset.

.



# Bootstrap

Hypothesis  
testing, MLE,  
Bootstrap

Theory behind this method:

$$\mathbb{E}(Y) = \lim_{B \rightarrow \infty} \frac{1}{B} \sum_{i=1}^B Y_i$$

For each function  $h$  with a finite mean

$$\mathbb{E}(h(Y)) = \lim_{B \rightarrow \infty} \frac{1}{B} \sum_{i=1}^B h(Y_i) = \langle h \rangle_B$$

Here  $B$  is the number of independent measurements (sampling with replacement) on the original dataset.

• In case the function  $h$  has  $k > 1$  arguments, we simply draw  $k$  data with replacement (due to independence) from the original dataset.

• This delivers the expectation value as  $\langle \cdot \rangle_B$

The standard error is just another function:  $se^2 = \langle (\cdot - \langle \cdot \rangle_B)^2 \rangle_B$

# Bootstrap

Hypothesis  
testing, MLE,  
Bootstrap

Formal definition:

Bootstrap method is used for approximating the expected value and the standard error of any function from a measured dataset. E.g. if the  $T(X)$  test statistics is approximated, then  $X$  is distributed according to a fixed, but unknown  $F$  distribution.

Steps for any  $T(X)$  function:

- Draw  $k$  points from the measured dataset: this follows the  $F$  distribution, since the measured dataset has values according to  $F$ .
- Compute  $T(X)$  where  $X$  is  $k$  dimensional vector.
- Repeat  $B$  times the above steps
- $E_{bootstrap} = \frac{1}{B} \sum_b^B T_b$
- $SE_{bootstrap} = \frac{1}{B} \sum_b^B (T_b - \frac{1}{B} \sum_r^B T_r)^2$

Note: Jackknife was a similar, replica based method.

Note2: if  $T(X)$  is the test function for the mean,

then  $T(X) = \sum_i^n X_i/n$  is the average, using all values of the sample.

# Bootstrap for confidence intervals

Hypothesis  
testing, MLE,  
Bootstrap

Using bootstrap for confidence intervals  $C(a, b)$ :

# Bootstrap for confidence intervals

Using bootstrap for confidence intervals  $C(a, b)$ :

- Normal method:  $\hat{T} \pm z_{\alpha/2} \hat{se}_{bootstrap}$ , so

$$a = \hat{T} - z_{\alpha/2} \hat{se}_{bootstrap}, b = \hat{T} + z_{\alpha/2} \hat{se}_{bootstrap}$$

# Bootstrap for confidence intervals

Using bootstrap for confidence intervals  $C(a, b)$ :

- Normal method:  $\hat{T} \pm z_{\alpha/2} \hat{se}_{bootstrap}$ , so

$$a = \hat{T} - z_{\alpha/2} \hat{se}_{bootstrap}, b = \hat{T} + z_{\alpha/2} \hat{se}_{bootstrap}$$

- Pivotal method:  $C = (a, b)$

$$a = \hat{T} - H^{-1}(1 - \alpha/2) \text{ and } b = \hat{T} + H^{-1}(\alpha/2)$$

$$\text{where } H(r) = \mathbb{P}_F(\hat{T} - T \leq r)$$

# Bootstrap for confidence intervals

Using bootstrap for confidence intervals  $C(a, b)$ :

- Normal method:  $\hat{T} \pm z_{\alpha/2} \hat{se}_{bootstrap}$ , so

$$a = \hat{T} - z_{\alpha/2} \hat{se}_{bootstrap}, b = \hat{T} + z_{\alpha/2} \hat{se}_{bootstrap}$$

- Pivotal method:  $C = (a, b)$

$$a = \hat{T} - H^{-1}(1 - \alpha/2) \text{ and } b = \hat{T} + H^{-1}(\alpha/2)$$

where  $H(r) = \mathbb{P}_F(\hat{T} - T \leq r)$

- good approximation:  $a = 2\hat{T} - T_{1-\alpha/2}^*$ ,  $b = 2\hat{T} - T_{\alpha/2}^*$ ,  
where  $T_{\beta}^*$  is  $\beta$ -sample-quantile of  $T_1, T_2, \dots, T_B$  replicas.

# Bootstrap for confidence intervals

Using bootstrap for confidence intervals  $C(a, b)$ :

- Normal method:  $\hat{T} \pm z_{\alpha/2} \hat{se}_{bootstrap}$ , so
$$a = \hat{T} - z_{\alpha/2} \hat{se}_{bootstrap}, b = \hat{T} + z_{\alpha/2} \hat{se}_{bootstrap}$$
- Pivotal method:  $C = (a, b)$ 
$$a = \hat{T} - H^{-1}(1 - \alpha/2) \text{ and } b = \hat{T} + H^{-1}(\alpha/2)$$
where  $H(r) = \mathbb{P}_F(\hat{T} - T \leq r)$ 
  - good approximation:  $a = 2\hat{T} - T_{1-\alpha/2}^*$ ,  $b = 2\hat{T} - T_{\alpha/2}^*$ ,  
where  $T_{\beta}^*$  is  $\beta$ -sample-quantile of  $T_1, T_2, \dots, T_B$  replicas.
- Percentile interval:  $(T_{\alpha/2}^*, T_{1-\alpha/2}^*)$

# Parameter inference

Hypothesis  
testing, MLE,  
Bootstrap

How can we find  $T = T(t_1, t_2, \dots)$  functions?



# Parameter inference

Hypothesis  
testing, MLE,  
Bootstrap

How can we find  $T = T(t_1, t_2, \dots)$  functions?

E.g. by estimating the parameters  $t_1, t_2, \dots$

Two methods:

- moments (MME)
- max. likelihood (MLE)

# Method of moments (MME)

Hypothesis  
testing, MLE,  
Bootstrap

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

# Method of moments (MME)

Hypothesis  
testing, MLE,  
Bootstrap

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

# Method of moments (MME)

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

Expressing  $t$  at the left and substituting the data values at the right, one gets an approximation for the parameter  $t$ .

# Method of moments (MME)

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

Expressing  $t$  at the left and substituting the data values at the right, one gets an approximation for the parameter  $t$ .  
In case of more parameters, use so many moments as many unknown parameters.

# Method of moments (MME)

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

Expressing  $t$  at the left and substituting the data values at the right, one gets an approximation for the parameter  $t$ .

In case of more parameters, use so many moments as many unknown parameters.

Some properties of the method:

- An estimate for  $\hat{t}$  exists with  $\mathbb{P} = 1$

# Method of moments (MME)

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

Expressing  $t$  at the left and substituting the data values at the right, one gets an approximation for the parameter  $t$ .

In case of more parameters, use so many moments as many unknown parameters.

Some properties of the method:

- An estimate for  $\hat{t}$  exists with  $\mathbb{P} = 1$
- The estimate is consistent:  $\hat{t} \rightarrow t$

# Method of moments (MME)

Given a measured dataset, any moments of the distribution can be approximated:

$$\alpha_k = \int x^k dF = \frac{1}{n} \sum x_i^k$$

If the distribution  $F$  depends on the parameter  $t$ , the left hand side provides a formula for  $t$  based on the  $m$ -th moment.

$$\alpha_m(t) = \frac{1}{n} \sum x_i^m$$

Expressing  $t$  at the left and substituting the data values at the right, one gets an approximation for the parameter  $t$ .

In case of more parameters, use so many moments as many unknown parameters.

Some properties of the method:

- An estimate for  $\hat{t}$  exists with  $\mathbb{P} = 1$
- The estimate is consistent:  $\hat{t} \rightarrow t$
- The estimate is asymptotically normal:  $\sqrt{n}(\hat{t} - t) \rightarrow N(0, \sigma^2)$   
(allows Ward test)



# Maximum Likelihood Estimate (MLE)

Hypothesis  
testing, MLE,  
Bootstrap

We measure  $n$  values:  $X_1, X_2, \dots, X_n$

These are IID random variables with PDF  $f(X_i, \theta)$

- The **likelihood function** is  $\mathcal{L}(\theta) = \prod_i f(X_i, \theta)$
- The **log-likelihood function** is  $\ell(\theta) = \sum_i \log f(X_i, \theta)$
- The **Maximum Likelihood Estimator** is  $\hat{\theta}$  that maximizes  $\mathcal{L}$

# Properties of MLE

Hypothesis  
testing, MLE,  
Bootstrap

- The  $\mathcal{L}(\theta) \in [0, \infty)$ : it is the joint density of data, but not a probability:  
 $\int d\theta \mathcal{L}(\theta) \neq 1$
- consistent:  $\hat{\theta} \rightarrow \theta$
- equivariant: if  $\hat{\theta}$  MLE of  $\theta$  then  $g(\hat{\theta})$  MLE of  $g(\theta)$
- asymptotically normal:  $\sqrt{n}(\hat{\theta} - \theta) \rightarrow N(0, \hat{se})$
- asymptotically optimal: MLE has smallest variance for large samples