


6.2 Volumes

3D立體: 體積篇


英語教室: solid ['sɒlɪd] 立體, volume ['vɒljəm] 體積,
cylinder ['sɪlɪdər] 柱, cone [kɒn] 錐, circular ['sɜːkjələ] 圓形的,
box [bɒks] 盒, rectangular [ˌrɛkˈtæŋɡjələ] 矩形的,
sphere [sfɪə] 球, spherical ['sfɛrɪkl] 球狀的,
perpendicular [ˌpɜːpənˈdɪkjələ] 垂直的, cross-section [krɒs-ˈseɪʃən] 橫切面,
revolution [ˌrɛvəˈluːʃən] 旋轉. disk [dɪsk] 圓盤, washer ['wɑːʃə] 墊圈。

Cylinder




$$V = Ah$$

Circular cylinder



$$V = \pi r^2 h$$

Rectangular box




$$V = lwh$$

Note: $V(\text{錐}) = \frac{1}{3}V(\text{柱})$ 。


Define: Let S be a **solid** 立體 that lies between $x = a$ and $x = b$. If the cross-sectional area of S in the plane P_x , through x and perpendicular to the x -axis, is $A(x)$, where A is a continuous function, then the **volume** 體積 of S is (體積是近似柱體積和的極限)

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n A(x_i^*) \Delta x = \int_a^b A(x) dx$$



Note: 如果選擇的橫切面垂直 y -軸，用 $V = \int_c^d A(y) dy$.

Define: The ***solid of revolution*** 旋轉體 is obtained by revolving a region about a line.




Note: 如果繞:

- horizontal line 水平線 (x -軸, $y = k$), 用 $V = \int_a^b A(x) dx$;
- vertical line 垂直線 (y -軸, $x = \ell$), 用 $V = \int_c^d A(y) dy$.

(Why? 切面積好算!)

If the cross-sectional area $A(x)$ or $A(y)$ is:

- a ***disk*** 圓盤, then $A = \pi R^2$, or
- a ***washer*** 墊圈, then $A = \pi R^2 - \pi r^2$



Skill: 如何列式:
$$V = \int_{\text{③}}^{\text{③}} \underbrace{A(\text{②})}_{\text{截面積}} \underbrace{d\text{①}}_{\text{厚度}}$$

- ① 厚度(thickness) 是往 x/y -軸方向就 dx/dy 。
- ② 截面積(cross-sectional area) 跟著變成 x/y 的函數。
- ③ 上下限(upper/lower limits) 找 x/y 的範圍。


Example 0.1 Show that the volume of a sphere of radius r is $V = \frac{4}{3}\pi r^3$.

(把球想像成半圓繞直徑的旋轉體。)

$x^2 + y^2 = r^2$, $A(x) = \pi y^2 = \pi(r^2 - x^2)$. (對 x 積分, 要換成 x 的函數。)

$$\begin{aligned} V &= \int_{-r}^r A(x) dx = \int_{-r}^r \pi(r^2 - x^2) dx \\ &= \left\langle \begin{aligned} &\left[\pi r^2 x - \pi \frac{x^3}{3} \right]_{-r}^r = \left[\pi r^3 - \pi \frac{r^3}{3} \right] - \left[\pi r^2(-r) - \pi \frac{(-r)^3}{3} \right] \quad (\text{直接算}) \\ &2\pi \int_0^r (r^2 - x^2) dx = 2\pi \left[r^2 x - \frac{x^3}{3} \right]_0^r = 2\pi \left(r^3 - \frac{r^3}{3} \right) \quad (\text{偶函數}) \end{aligned} \right\rangle \\ &= \frac{4}{3}\pi r^3. \end{aligned}$$

■




Example 0.2 (*x*-axis) Find the volume of the solid obtained by rotating about the *x*-axis the region under the curve $y = \sqrt{x}$ from 0 to 1. Illustrate the definition of volume by sketching a typical approximating cylinder.

$$A(x) = \pi(\sqrt{x})^2 = \pi x.$$

the volume of the approximating cylinder is $A(x)\Delta x = \pi x \Delta x$.

$$V = \int_0^1 \pi x \, dx = \left[\frac{\pi x^2}{2} \right]_0^1 = \frac{\pi}{2}.$$

■




Note: $y = f(x)$ 繞 x -軸 旋轉體: 面積是 πy^2 , 體積是 $\int_a^b \pi [f(x)]^2 \, dx$.
 不要背! 用畫圖找圓半徑。

Example 0.3 (*y*-axis) Find the volume of the solid obtained by rotating the region bounded by $y = x^3$, $y = 8$, and $x = 0$ about the y -axis.

$A(y) = \pi x^2 = \pi y^{2/3}$, (解反函數: $y = x^3 \rightarrow x = \sqrt[3]{y} = y^{1/3}$.)
the volume of the approximating cylinder is $A(x)\Delta x = \pi y^{2/3}\Delta y$.

$$V = \int_0^8 \pi y^{2/3} dy = \left[\frac{3}{5} \pi y^{5/3} \right]_0^8 = \frac{96\pi}{5}.$$

■



Note: $y = f(x)$ 繞 y -軸 旋轉體: 面積是 πx^2 , 體積是 $\int_a^b \pi [f^{-1}(y)]^2 dy$.
還是不要背! 用畫圖找圓半徑。

Example 0.4 (*washer*) The region R enclosed by the curves $y = x$ and $y = x^2$ is rotated about the x -axis. Find the volume of the resulting solid.


Solve $x = y = x^2$, $(x, y) = (0, 0), (1, 1)$. (上下界是從 0 到 1。)

$x \geq x^2$ on $[0, 1]$, 外圈半徑是 $y = x$, 內圈半徑是 $y = x^2$ 。

$A(x) = \pi x^2 - \pi (x^2)^2 = \pi(x^2 - x^4)$, (繞 x -軸, 對 x 積。)


$$V = \int_0^1 \pi(x^2 - x^4) dx = \left[\pi \left(\frac{x^3}{3} - \frac{x^5}{5} \right) \right]_0^1 = \frac{2\pi}{15}.$$

■



Question: 什麼時候會是圓盤/墊圈?

Answer: 用長方形去近似區域, 長方形繞出的體積近似區域繞出的體積。
當長方形 (區域) 貼著旋轉軸, 會繞出圓盤; 否則會繞出墊圈。




Example 0.5 (horizontal line) Find the volume of the solid obtained by rotating the region in Example 4 about the line $y = 2$.

$2 - x^2 \geq 2 - x$ on $[0, 1]$, 外圈半徑變成 $2 - x^2$, 內圈半徑變成 $2 - x$ 。

$A(x) = \pi(2 - x^2)^2 - \pi(2 - x)^2 = \pi(x^4 - 5x^2 + 4x)$, (繞 $y = 2$, 對 x 積。)

$$V = \int_0^1 \pi(x^4 - 5x^2 + 4x) dx = \left[\pi\left(\frac{x^5}{5} - \frac{5x^3}{3} + 2x^2\right) \right]_0^1 = \frac{8\pi}{15}. \quad \blacksquare$$




Example 0.6 (vertical line) Find the volume of the solid obtained by rotating the region in Example 4 about the line $x = -1$.

$x = \sqrt{y}$ and $x = y$. (解反函數)

$1 + \sqrt{y} \geq 1 + y$ on $[0, 1]$, 外圈半徑變成 $1 + \sqrt{y}$, 內圈半徑變成 $1 + y$ 。

$A(y) = \pi(1 + \sqrt{y})^2 - \pi(1 + y)^2 = \pi(2\sqrt{y} - y - y^2)$, (繞 $x = -1$, 對 y 積。)


$$V = \int_0^1 \pi(2\sqrt{y} - y - y^2) dy = \left[\pi\left(\frac{4}{3}y^{3/2} - \frac{1}{2}y^2 - \frac{1}{3}y^3\right) \right]_0^1 = \frac{\pi}{2}. \quad \blacksquare$$



Example 0.7 A solid with a circular base of radius 1. Parallel cross-sections perpendicular to the base are equilateral(等邊) triangles. Find the volume of the solid.

$$\begin{aligned}
 x^2 + y^2 &= 1, \quad A(x) = \frac{1}{2} \cdot 2y \cdot \sqrt{3}y = \sqrt{3}(1 - x^2), \\
 V &= \int_{-1}^1 A(x) \, dx = \int_{-1}^1 \sqrt{3}(1 - x^2) \, dx \\
 &= \left\langle \begin{aligned} &\left[\sqrt{3} \left(x - \frac{x^3}{3} \right) \right]_{-1}^1 = [\sqrt{3}(1 - \frac{1}{3})] - [\sqrt{3}(-1 + \frac{1}{3})] \quad (\text{直接算}) \\ &2 \int_0^1 \sqrt{3}(1 - x^2) \, dx = 2\sqrt{3} \left[x - \frac{x^3}{3} \right]_0^1 = 2\sqrt{3}(1 - \frac{1}{3}) \quad (\text{偶函數}) \end{aligned} \right\rangle \\
 &= \frac{4\sqrt{3}}{3}.
 \end{aligned}$$

■




Example 0.8 Find the volume of a pyramid whose base is a square with side L and whose height is h .



[Sol 1] 把頂點放在原點, x -軸是中心軸, 假設在 x 時的截方形邊長是 s .
 $\frac{x}{h} = \frac{s/2}{L/2}, A(x) = s^2 = \frac{L^2}{h^2}x^2$.

$$V = \int_0^h A(x) dx = \int_0^h \frac{L^2}{h^2}x^2 dx = \frac{L^2}{h^2} \left[\frac{x^3}{3} \right]_0^h = \frac{L^2h}{3}.$$


[Sol 2] 把底部中心放在原點, 假設 y 高時的截方形邊長是 t .
 $\frac{h-y}{h} = \frac{t/2}{L/2}, A(y) = t^2 = \frac{L^2}{h^2}(h-y)^2. V = \int_0^h A(y) dy = \dots = \frac{L^2h}{3}.$ ■



(可不可以倒過來算? 可以!)



(可不可以用其他方式切? 可以! 好算嗎?)



◆: Volume of cone = $\frac{1}{3}$ volume of cylinder

$$\text{錐體積} = \lim_{n \rightarrow \infty} \sum \text{四角錐體積} = \lim_{n \rightarrow \infty} \sum \frac{1}{3} \text{四角柱體積} = \frac{1}{3} \text{柱體積}。$$

