

Behavioural syndromes and social insects: personality at multiple levels

Jennifer M. Jandt^{1,*†}, Sarah Bengston^{2,†}, Noa Pinter-Wollman³, Jonathan N. Pruitt⁴, Nigel E. Raine⁵, Anna Dornhaus² and Andrew Sih⁶

¹*Department of Ecology, Evolutionary and Organismal Biology, Iowa State University, Ames, IA 50011, USA*

²*Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ 85721, USA*

³*BioCircuits Institute, University of California, San Diego, La Jolla, CA 92093-0328, USA*

⁴*Department of Biological Science, University of Pittsburgh, Pittsburgh, PA 15260, USA*

⁵*School of Biological Sciences, Royal Holloway University of London, Surrey, TW20 0EX, UK*

⁶*Department of Environmental Science and Policy, University of California, Davis, CA 95616, USA*

ABSTRACT

Animal personalities or behavioural syndromes are consistent and/or correlated behaviours across two or more situations within a population. Social insect biologists have measured consistent individual variation in behaviour within and across colonies for decades. The goal of this review is to illustrate the ways in which both the study of social insects and of behavioural syndromes has overlapped, and to highlight ways in which both fields can move forward through the synergy of knowledge from each. Here we, (i) review work to date on behavioural syndromes (though not always referred to as such) in social insects, and discuss mechanisms and fitness effects of maintaining individual behavioural variation within and between colonies; (ii) summarise approaches and principles from studies of behavioural syndromes, such as trade-offs, feedback, and statistical methods developed specifically to study behavioural consistencies and correlations, and discuss how they might be applied specifically to the study of social insects; (iii) discuss how the study of social insects can enhance our understanding of behavioural syndromes—research in behavioural syndromes is beginning to explore the role of sociality in maintaining or developing behavioural types, and work on social insects can provide new insights in this area; and (iv) suggest future directions for study, with an emphasis on examining behavioural types at multiple levels of organisation (genes, individuals, colonies, or groups of individuals).

Key words: behavioral syndromes, behavioural types, levels of organisation, behavioural carryover, behavioural consistency, temperament, repeatability.

CONTENTS

I. Introduction	2
(1) Defining ‘behavioural syndromes’	2
(2) Mechanisms affecting behavioural type	3
(3) Fitness effects of behavioural syndromes	3
II. Social insects and a hierarchy of behavioural types	4
(1) Variation within colonies	4
(a) Variation in morphological castes	5
(b) Variation in monomorphic species	5
(c) Mechanisms underlying intra-colony variation in behavioural type	5
(i) Genetics	5
(ii) Physiology	8
(iii) Environment	8

* Address for correspondence (Tel: +520-603-5850; Fax: +515-294-1337; E-mail: jjandt2@gmail.com)

† Authors contributed equally.

(d) Fitness consequences of intra-colony behavioural variation	8
(2) Variation among colonies	9
(a) Mechanisms underlying variation among colonies in behavioural type	9
(i) Colony genetics	9
(ii) Colony composition and emergent behaviour	9
(iii) Colony environment	10
(b) Fitness consequences of colony behavioural syndromes	11
III. Insights from behavioural syndromes for the study of social insects	11
(1) Trade-offs	11
(2) Feedback	11
(3) Statistical methods	12
IV. Insights from social insects for behavioural syndromes	12
(1) Group-living might affect behavioural type	12
(2) Mechanisms affecting behavioural type	13
(3) Development of behavioural types across levels of organisation	13
V. Future directions	13
(1) What are the adaptive reasons for colonies, or groups of individuals, to maintain mixtures of behavioural types?	13
(2) Do colonies differ in their degree of cooperation?	14
(3) How does consistency in behavioural type affect task performance when individuals switch task?	14
VI. Conclusions	15
VII. Acknowledgements	15
VIII. References	15

I. INTRODUCTION

In recent years, numerous studies have found that animals exhibit ‘personalities’ or behavioural syndromes—consistent and/or correlated behaviour across two or more situations (Gosling, 2001; Dall, Houston & McNamara, 2004; Sih, Bell & Johnson, 2004a; Sih *et al.*, 2004b; Réale *et al.*, 2007; Biro & Stamps, 2008; Sih & Bell, 2008; Dall *et al.*, 2012; Garamszegi & Herczeg, 2012). For example, although most individuals alter their level of aggression depending on the ecological situation (e.g. resource availability or predation risk), some are consistently more aggressive than others (Riechert & Hedrick, 1993; Wilson *et al.*, 1994; Sih, Kats & Maurer, 2003; Duckworth, 2006; Bell & Sih, 2007; Johnson & Sih, 2007). Furthermore, aggressive behaviour is often correlated with bold behaviour or high activity level (Huntingford, 1982; Riechert & Hedrick, 1993; Bell, 2005; Johnson & Sih, 2005). To date, much of the work on behavioural syndromes has focused on variation in boldness, aggressiveness, or activity level (Réale *et al.*, 2007), referred to here as ‘classical’ behavioural types.

Consistent individual differences in behaviour have long been recognized and studied in several taxa, including humans, primates, laboratory rodents, and a few domesticated mammals (Gosling, 2001). Intraspecific differences in behavioural type have also been described when individuals differ in morphology. This includes classic stereotypical ideas on differences in behavioural type between males and females, and between alternative mating morphs [e.g. larger, aggressive, territorial males *versus* smaller, sneaky, satellite males (Gross & Charnov, 1980)]. Recent studies, however, have emphasized that individual behavioural differences

also exist in animal taxa that lack phenotypic polymorphism (Sih *et al.*, 2004b).

Social insects can exhibit behavioural variation at multiple levels of organisation: between species, colonies, castes, individuals, and genetic lines (Keller, 1999). Social insect species have long been known to differ in behavioural type (e.g. more aggressive species outcompete and may even displace subordinate species; Davidson, 1998; Holway & Suarez, 1999). Social insect castes (i.e. reproductive or morphologically distinct individuals) also differ in behavioural tendencies often associated with their specialized task (e.g. soldiers are more aggressive than other workers). Thus, social insect researchers have always considered, and quantified, behavioural syndromes. However, until recently, few social insect studies have investigated differences in behavioral type among colonies (within a species) or among monomorphic individuals within a colony (but see Jeanne, 1988). In addition, few social insect studies have investigated whether variation in behavioural type carries over across tasks. For example, in species that exhibit temporal polyethism (workers transition between tasks over time), it would be interesting to determine whether young nurses (brood-care workers) with a highly aggressive behavioural type are also more aggressive as older foragers or more likely to guard the nest. Such consistency in behavioural type carried over across task transitions could have implications for colony fitness.

(1) Defining ‘behavioural syndromes’

Behavioural syndromes are defined as consistent individual differences, within a population, in behaviour within and across context (Table 1). If an individual differs consistently in a behaviour (e.g. is consistently more aggressive than others),

Table 1. A comparison of terms used in animal behavior and social insect biology that describe similar phenomena

Animal behaviour terms	Social insect biology terms
Behavioural syndrome Individuals within a population exhibit consistency/correlation in two or more functionally different behaviours	
Animal personality Individuals within a population exhibit consistency in a single behaviour	Morphological/reproductive caste Physiological constraints (genetics, morphology, hormones, etc.) predispose individuals to perform specific behaviours (e.g. queens and workers)
Behavioural type Individual exhibits consistent behaviour across contexts and/or over time. Repeatability or consistency of the magnitude of behaviour often observed. This can be a continuous or discrete variable.	Behavioural specialisation Individual exhibits consistent task performance (that is not necessarily determined by its morphology)
Episodic personality Response to one set of stimuli predicts the response to other sets of stimuli over a short time scale. Responses are inconsistent over longer time scales.	Temporal polyethism In a predictable order, individuals switch among tasks as they develop. Task switching Individuals switch among tasks throughout their lifetime.
Keystone individuals An individual that significantly alters the behaviour of other individuals within the group.	Elite workers/activators Individuals that stimulate others, perform multiple tasks, or perform a disproportionate amount of work in the colony.

then that individual has a behavioural type (Table 1), and the population has a behavioural syndrome (Sih *et al.*, 2004a; Dingemanse *et al.*, 2010). Recently, Garamszegi & Herczeg (2012) differentiated animal personalities ('consistency in single behaviours') and behavioural syndromes ('consistency in two or more functionally different behaviours') (Table 1). These distinctions are most important when developing experimental design and performing statistical analyses (see Section III.3). In either case, to test whether individuals differ in behaviour in a particular context, it is necessary to quantify the behaviour of a set of individuals in standardized conditions, e.g. while foraging or building a nest (Réale *et al.*, 2007). Although most of the recent interest has been on differences among individuals in behavioural type, one can also look for consistent differences among populations or species in behavioural type. Social insects allow for additional levels of analysis—caste (within colonies) and colony.

(2) Mechanisms affecting behavioural type

A growing literature examines the mechanisms associated with differences among individuals of a behavioural type. Numerous studies have quantified physiological or neuroendocrine correlates of behavioural types (Koolhaas *et al.*, 2007; Biro & Stamps, 2008; Careau *et al.*, 2008; van Oers & Mueller, 2010). Genetic studies show that behavioural types typically exhibit low to moderate heritability (i.e. heritability (H^2) = 0.2–0.5; van Oers *et al.*, 2005; Penke, Denissen & Miller, 2007; van Oers & Mueller, 2010), with recent studies identifying candidate genes, often associated with neuroendocrine pathways, that explain some of the variation among

behavioural types (e.g. Ben-Shahar *et al.*, 2002; van Oers & Mueller, 2010). Genetic effects on behavioural types are also typically moderated by individual experience. In particular, early experience can often have strong effects on the development of a later behavioural type and associated hormonal 'stress response' systems (Stamps & Groothuis, 2010; Del Giudice, Ellis & Shirtcliff, 2011; Sih, 2011). Many of these mechanisms have been studied in social insects; however, as highlighted below, much remains to be discovered.

(3) Fitness effects of behavioural syndromes

Recent behavioural syndrome research has focused on the ecological and evolutionary relevance of within-population variation in behavioural type. An individual's behavioural type can clearly affect its fitness, typically in context-dependent ways (Smith & Blumstein, 2008); e.g. in a safe environment, selection might favour bold individuals that enjoy high feeding or mating success, but in a dangerous environment, cautious individuals might be more likely to survive (Fig. 1A). Notably, if individuals of a particular behavioural type exhibit limited behavioural plasticity, their behaviour could be suboptimal in situations that do not fit their behavioural type well. For example, bolder individuals can be inappropriately bold when predators are present (Sih *et al.*, 2003), and aggressive individuals can be inappropriately aggressive towards mates (Johnson & Sih, 2005), or can exhibit poor parental care (Duckworth, 2006). Given the potentially strong effects of behavioural types on fitness, many studies have asked how variation in behavioural types is maintained, e.g. *via* environmental variation in selection




Fig. 1. Concepts that have been borne out in behavioural syndrome (A) and social insect (B) research that could benefit the other field. (A) Behavioural syndromes—trade-offs of behavioural type across ecological context. Trade-offs associated with behavioural syndromes help us understand why populations maintain correlated behavioural types. For example (i) a bold cricket will be a better procurer of food, but (ii) a shy cricket will be less likely to be seen and eaten by a predator. (B) Social insects—multiple levels of selection/organisation. (i) Development of an individual's behavioural type is affected by internal and external mechanisms. (ii) Intra-colony composition (variation within and among castes) and development may affect colony behaviour (performance). (iii) Colony performance may feedback and influence intra-colony composition. (iv) Inter- and intra-colony interactions can feed back and affect colony-level behaviour.

pressures (Dingemanse *et al.*, 2004), frequency dependence (Dall *et al.*, 2004; Pruitt & Riechert, 2009), or condition dependence (Luttbeg & Sih, 2010). Behavioural syndromes can also have important ecological implications, e.g. effects on species interactions, population dynamics, ecological invasions, etc. (Sih *et al.*, 2012).

II. SOCIAL INSECTS AND A HIERARCHY OF BEHAVIOURAL TYPES

Social insects are unique in that selection acts on multiple levels (Fig. 1B; Folse & Roughgarden, 2010). In a social insect colony, workers and queens are individual organisms, but the colony itself can be considered a 'superorganism' (Hölldobler & Wilson, 2009). In general, colonies produce more colonies: the queen as the primary reproductive resembles a 'germ cell', whereas non-reproductive workers resemble 'somatic cells' in that they care for their sisters and queen but do not directly contribute to the next generation. However, in many species,

workers might also compete with one another and the queen to lay unfertilized eggs which can develop into reproductive males. Consistent individual differences (i.e. behavioural types) are found at both levels of organisation: within a colony (among workers and queens), and across colonies. Here we review the current knowledge of individual variation within and among colonies. We discuss both the mechanisms and fitness consequences of variation at each of these two levels.

(1) Variation within colonies

Individuals in social insect colonies commonly differ consistently in behaviour. This is usually interpreted as an effect of their allocation to specialized roles or 'tasks', e.g. nurses, guards, foragers, etc. Such specialization produces a division of labour, and is thought to be adaptive at the colony level (Robinson, 1992; Beshers & Fewell, 2001). When there is variation in morphology (castes; Table 1), such as soldiers *versus* 'minor' workers (Wilson, 1976), behavioural variation might be correlated with caste differences, although variation could also be found within a caste. The majority

of social insect species, however, are monomorphic, i.e. there are no morphological differences between workers (Hölldobler & Wilson, 1990). Nevertheless, workers that look the same might vary greatly in their behaviour. In some cases, these behavioural differences might lead to task groups or behavioural specialisations (Table 1), and still variation can be observed within these behavioural groups. Thus, social insects provide ample opportunities for studying the effects of morphology and other factors on behavioural syndromes.

(a) Variation in morphological castes

Morphological castes are an important functional ‘grouping’; variation in morphology often predisposes individuals to behave in a certain manner or to perform particular tasks (Oster & Wilson, 1978; Hölldobler & Wilson, 1990). Even in non-insect social groups, individuals often differ in behavioural tendencies based on morphology (e.g. males and females; Del Giudice, 2011) or body size (e.g. courting *versus* coercive mating tactics in large and small mosquitofish males; Hughes, 1985). In social insects, castes are thought to be a colony-level adaptation (Oster & Wilson, 1978; Hölldobler & Wilson, 1990): morphologically specialized individuals are able to perform functions that would otherwise be inaccessible (e.g. defensive soldier castes in *Cephalotes* spp. ants; Powell, 2008).

Within a caste, a range of behavioural variation might also persist. For example, task preference or efficiency can vary within caste. Within the forager/worker caste of leaf-cutter ants, as mandibles wear away, the individual can no longer cut leaves efficiently and thus perform the function to which this particular morphology predisposes it (Schofield *et al.*, 2011). This wear not only creates variation within the caste in terms of cutting efficiency, but it also prompts a switch in task preference to carrying leaves instead of cutting (Schofield *et al.*, 2011).

To date there has been little work exploring the extent to which individuals within a caste vary in other behaviours (Table 2). For example, are some soldiers more aggressive than others, and are they also aggressive in other situations aside from nest defence? While studies have highlighted the relative importance of castes to colony function (Porter & Jorgensen, 1981; Powell & Franks, 2006; Powell, 2008), little has been done to explore the importance of behavioural variation observed within castes.

(b) Variation in monomorphic species

Variation in behavioural type has also been observed in species without clear morphological castes. In social insects, social behavioural types (e.g. foraging, defence, etc.; Table 2) are the most commonly studied. Still, individuals within a colony could vary in behavioural type along a variety of axes such as those studied in the classic behavioural syndrome literature and/or the cognitive ecology literature (Sih & Del Giudice, 2012; Table 2). For example, *Myrmica* spp. ant workers vary not only in the task they are most likely to perform, but also in general behavioural type: patrollers are

bolder, more aggressive, and more exploratory across various contexts than nurses or foragers (Chapman *et al.*, 2011).

In social insect research, a lot is known about how individual workers differ in behaviour in ways that affect their role in the colony, and how these behaviours correlate with others across contexts (i.e. behavioural syndromes). A well-studied case is variation among honey bee (*Apis mellifera*) foragers. Some individuals have lower sucrose response thresholds (i.e. are more likely to respond to less-concentrated nectar), and are more likely to collect pollen, whereas those with higher sucrose response thresholds are more likely to collect nectar while foraging (Page, Erber & Fondrk, 1998; Scheiner, Erber & Page, 1999; Scheiner, Page & Erber, 2004). Sucrose response thresholds also correlate with reproductive development in workers (Amdam & Page, 2010), forager decision-making (Cakmak *et al.*, 2009), the likelihood of collecting non-food rewards, such as water or resin (Simone-Finstrom, Gardner & Spivak, 2010) and learning ability (Scheiner, 2012). In this sense, the cluster of correlated behaviours associated with sucrose response threshold is a broad behavioural syndrome affecting a series of seemingly unrelated behaviours across a variety of contexts.

We also observe variation in behaviours not directly corresponding to particular tasks. For example, although more aggressive workers might be more likely to engage in colony defence (i.e. more likely to attack non-nestmates: Crosland, 1990; Modlmeier & Foitzik, 2011), they could also be inappropriately aggressive to nestmates (van Doorn, 1989; Powell & Tschinkel, 1999; Robson *et al.*, 2000; Foster *et al.*, 2004; Molina & O'Donnell, 2009).

(c) Mechanisms underlying intra-colony variation in behavioural type

(i) *Genetics.* Multiple mating by a queen leads to an increase in within-colony genetic variation (i.e. workers are less related to one another) that can result in intra-colony variation in behavioural types (Table 3). In harvester ants (*Pogonomyrmex* spp.), genotype could predispose an individual to develop into a queen or a worker (Volny & Gordon, 2002; Cahan & Keller, 2003; Schwander, Cahan & Keller, 2007; Schwander *et al.*, 2010). Genotypic differences in honey bees might affect individual learning differences (Chandra, Hosler & Smith, 2000) and foraging preferences (Page *et al.*, 1998). Genetic factors are also linked to how quickly a honey bee worker transitions between tasks (Page *et al.*, 1998; Amdam & Page, 2010). Page, Rueppell & Amdam (2012b) have summarized many of these genetic effects in terms of the ‘pollen-hoarding syndrome’, and have shown how overlapping genetic architecture could provide a mechanism for behavioural syndromes, both at the individual and colony level.

Behavioural variation also readily occurs within colonies where queens only mate with one male, making all workers full sisters and intra-colony genetic variation low [e.g. bumble bees (Jandt & Dornhaus, 2009), rock ants (Dornhaus *et al.*, 2008), social spiders (Pruitt *et al.*, 2011)]. In these cases, gene expression difference might underlie behavioural variation (Table 3). For example, differential expression of the *for*

Table 2. Traits in which individual (= Intra-colony) and colony (= Inter-colony) level behavioural variation, i.e. different behavioural types, have been measured in social insects

	Behavioral type	Intra-colony-level examples	Inter-colony-level examples
Classical	Aggression	Ants Chapman <i>et al.</i> (2011), Crosland (1990), Modlmeier & Foitzik (2011) and Powell & Tschinkel (1999)	Ants Buczkowski & Silverman (2006), Crosland (1990) and Suarez <i>et al.</i> (2002)
		Bees Pearce <i>et al.</i> (2001)	Bees Pearce <i>et al.</i> (2001)
		Wasps Cant <i>et al.</i> (2006), Reeve & Nonacs (1997)	Termites Adams <i>et al.</i> (2007)
		Termites Adams <i>et al.</i> (2007)	
		Ants Chapman <i>et al.</i> (2011) and Modlmeier & Foitzik (2011)	
Cognitive	Exploration	Ants Chapman <i>et al.</i> (2011) and Modlmeier & Foitzik (2011)	
	Boldness/shyness	Ants Chapman <i>et al.</i> (2011)	
	Proactive/reactive		Ants Gordon <i>et al.</i> (2011) and Pinter-Wollman <i>et al.</i> (2012a)
	Sensory bias/preference	Bees Page <i>et al.</i> (2012b), Spaethe <i>et al.</i> (2007) and Spaethe & Chittka (2003)	Bees Ings <i>et al.</i> (2009), Raine & Chittka (2005, 2007b) and Raine <i>et al.</i> (2006a)
	Learning	Bees Latshaw & Smith (2005), Lavery (1994), Lihoreau <i>et al.</i> (2012b), Molet <i>et al.</i> (2009), Muller <i>et al.</i> (2010), Page <i>et al.</i> (2012b), Raine & Chittka (2012), Scheiner <i>et al.</i> (2003) and Worden <i>et al.</i> (2005)	Bees Chittka <i>et al.</i> (2004), Ings <i>et al.</i> (2009), Raine & Chittka (2008, 2012) and Raine <i>et al.</i> (2006a,b)
Social	Speed and accuracy in decision-making	Bees Burns & Dyer (2008), Chittka <i>et al.</i> (2003), Dyer & Chittka (2004) and Kulahci <i>et al.</i> (2008)	Ants Pinter-Wollman <i>et al.</i> (2012a) Bees Wray & Seeley (2011)
	Cooperation	Ants Cahan & Fewell (2004)	Ants Cahan (2001)
		Social Spiders Pruitt & Riechert (2009)	Bees Robinson <i>et al.</i> (1990) Social Spiders Pruitt & Riechert (2011)
	Communication	Ants Pinter-Wollman <i>et al.</i> (2011)	Bees Mattila <i>et al.</i> (2008)
	Hygienic behaviour	Ants Diez <i>et al.</i> (2011)	Ants Scharf <i>et al.</i> (2012)
	House-hunting	Bees Arathi <i>et al.</i> (2000) and Visscher (1988)	Bees Paleolog (2009) and Wray <i>et al.</i> (2011)
		Ants Pinter-Wollman <i>et al.</i> (2012b)	Bees Wray & Seeley (2011)
	Foraging	Bees Liang <i>et al.</i> (2012)	
		Ants Beverly <i>et al.</i> (2009), Gordon <i>et al.</i> (2005) and Pinter-Wollman <i>et al.</i> (2012b)	Ants Bengston & Dornhaus (2013), Cole <i>et al.</i> (2010), Gordon <i>et al.</i> (2011), Gordon <i>et al.</i> (2013) and Pinter-Wollman <i>et al.</i> (2012a)
	Defense	Bees Fewell & Page (1993), Liang <i>et al.</i> (2012), Lihoreau <i>et al.</i> (2010, 2011, 2012a,b), Muller <i>et al.</i> (2010), Page <i>et al.</i> (2012b), Raine & Chittka (2007a), Robinson & Page (1989), Saleh & Chittka (2007) and Spaethe & Weidenmüller (2002)	Bees Gill <i>et al.</i> (2012), Ings <i>et al.</i> (2006), Page <i>et al.</i> (2012b), Raine & Chittka (2007b, 2008) and Wray <i>et al.</i> (2011)
		Bees Grüter <i>et al.</i> (2012)	Ants Pinter-Wollman <i>et al.</i> (2012a) Bees Wray <i>et al.</i> (2011)

We divide these into: ‘classical’—those behavioural axes most commonly measured in the behavioural syndrome literature, ‘cognitive’—those most commonly measured in the psychology/learning literature, and ‘social’—those most commonly measured in the social insect literature. We do not presume this list to be exhaustive, but instead wish to provide examples for researchers interested in studying behavioural syndromes in social insects.

Table 3. Mechanisms that affect behavioural variation presented in the behavioural syndrome and social insect literature

	Behavioral syndrome literature	Social insect literature
Genetic mechanisms	Aggression Kralj-Fiser <i>et al.</i> (2010) Boldness/shyness Fairbanks <i>et al.</i> (1999), Jones <i>et al.</i> (2011) and Kinnally <i>et al.</i> (2006)	Learning Chandra <i>et al.</i> (2000) Division of labour (general) Herb <i>et al.</i> (2012), Fewell & Page (1993), Page <i>et al.</i> (1992), Robinson & Page (1989), Robinson <i>et al.</i> (1994), Smith <i>et al.</i> (2008) and Stuart & Page (1991) Hygienic behaviour Robinson & Page (1988) Foraging Ben-Shahar <i>et al.</i> (2002), Fewell & Page (1993), Hunt <i>et al.</i> (2007), Ingram <i>et al.</i> (2005), Page <i>et al.</i> (1998, 2012b), Pankiw <i>et al.</i> (2002), Robinson & Page (1989), Scheiner <i>et al.</i> (2001) and Waddington <i>et al.</i> (1998) Defence Breed <i>et al.</i> (2004), Hunt <i>et al.</i> (2007), Moore <i>et al.</i> (1987) and Robinson & Page (1988) Reproductive Cahan & Keller (2003), Graff <i>et al.</i> (2007), Page <i>et al.</i> (2012b), Schwander <i>et al.</i> (2007, 2010), Smith <i>et al.</i> (2008), Volny & Gordon (2002) and Weiner & Toth (2012)
Physiological mechanisms	Aggression Brown <i>et al.</i> (2003), Bryant & Newton (1994), Cutts <i>et al.</i> (1998) and Koolhaas <i>et al.</i> (2007) Exploratory (activity) Careau <i>et al.</i> (2008) and Chappell <i>et al.</i> (2004) Boldness/shyness Huntingford <i>et al.</i> (2010) Proactive/reactive Careau <i>et al.</i> (2008) and Koolhaas <i>et al.</i> (2007) Foraging Mathot <i>et al.</i> (2009)	Aggressive Tibbetts & Huang (2010) Cooperative Lin <i>et al.</i> (1999) Division of labour (general) Blanchard <i>et al.</i> (2000), Dolezal <i>et al.</i> (2012), Robinson (2009), Robinson & Huang (1998) and Toth <i>et al.</i> (2009) Hygienic behaviour Spivak <i>et al.</i> (2003) Foraging Dolezal <i>et al.</i> (2012), Fewell & Page (1993), Giray <i>et al.</i> (2005), Robinson & Page (1989) and Seid & Traniello (2006) Reproductive Brent <i>et al.</i> (2006), Giray <i>et al.</i> (2005), Page <i>et al.</i> (2012b) and Smith <i>et al.</i> (2008) Exploratory (activity) Seeley <i>et al.</i> (1998) Cooperative Lin <i>et al.</i> (1999) Division of labour (general) Greene & Gordon (2003, 2007), Powell & Tschinkel (1999), Robinson & Huang (1998) and Wilson (1985) Foraging Anderson & Ratnieks (1999), Fewell & Page (1993), Pankiw <i>et al.</i> (2002) and Robinson & Page (1989) Reproductive Kamakura (2011) and Suryanarayanan <i>et al.</i> (2011a)
Experiential mechanisms	Aggression Del Giudice <i>et al.</i> (2011) and Niemelä <i>et al.</i> (2012) Exploration (activity) Del Giudice <i>et al.</i> (2011), Heifetz & Applebaum (1995) and Lihoreau <i>et al.</i> (2009) Boldness/shyness Del Giudice <i>et al.</i> (2011) and Niemelä <i>et al.</i> (2012) Foraging Lihoreau <i>et al.</i> (2009)	

Mechanisms are not mutually exclusive. Note that each literature has focused on defining mechanisms for very different sets of behavioural axes: behavioural syndrome research has focused on mechanisms for the classically studied behavioural types whereas social insect literature has focused on mechanisms for primarily social behavioural types.

(*foraging*) gene in honey bees and harvester ants affects whether an individual will be more likely to engage in brood care or foraging tasks, irrespective of age (Ben-Shahar *et al.*, 2002; Ingram, Oefner & Gordon, 2005). Differences in gene expression might be the result of experience, nutrition, age, or genotype; and often expression patterns are conserved across species (e.g. expression of *hexamerin* storage proteins often correlates with reproductive caste differentiation: Smith *et al.*, 2008). Epigenetics, i.e. the effects of DNA methylation, is a relatively new area of research, and its role on behavioural type is beginning to be investigated in social insects, both in terms of reproductive (Weiner & Toth, 2012) and non-reproductive (Herb *et al.*, 2012) division of labour.

(ii) *Physiology*. Social insect workers often change their role or specialisation over the course of their life, referred to as temporal polyethism (Table 1; Seeley, 1982). In the behavioural syndrome literature, switching between behavioural types is referred to as episodic personality (Table 1; Pronk, Wilson & Harcourt, 2010). In honey bees, transitions tend to occur in a particular order (Seeley, 1982), but this is not necessarily the case for all social insects. In some species, task repertoire (the number of tasks an individual performs) expands with age (Seid & Traniello, 2006), whereas in others, the order of task-switching is seemingly random (Gordon *et al.*, 2005; Jandt, Huang & Dornhaus, 2009).

Often, juvenile hormone (JH) influences the task an individual performs (Table 3; Robinson, 1987; Hartfelder, 2000; Giray, Giovanetti & West-Eberhard, 2005; Brent *et al.*, 2006; Dolezal *et al.*, 2012) (but see Cameron & Robinson, 1990), and in some cases, change in corpulence (fat content) also correlates with the performance of certain tasks (Table 3; Toth & Robinson, 2005; Robinson, 2009; Robinson *et al.*, 2009; Daugherty, Toth & Robinson, 2011), but this is not necessarily true for all social insects (Couvillon *et al.*, 2011). Research on *Polistes dominulus* wasps has shown that JH concentration changes during aggressive encounters (Tibbetts & Huang, 2010), and so it might be directly related to an individual's aggressive behavioural type.

It is indeed possible that temporal polyethism, caused by physiological change over an individual's lifetime, might also be a type of 'maturation' or 'ageing' syndrome—that is, several associated traits are ultimately caused by ageing, and the ageing process could differ among individuals (Dingemanse *et al.*, 2010). Therefore, it is important to consider maturation, as well as unavoidable senescence, as plausible hypotheses both to explain the evolution of temporal polyethism and of behavioural syndromes in general.

(iii) *Environment*. There are a variety of environmental factors that can influence behavioural type (Table 3): abiotic factors (e.g. temperature), ecological environment (e.g. food availability and/or competition), and social environment (e.g. interaction rates and communication signals). Early developmental influences are often particularly important. For example, early developmental influences, such as larval feeding or maternal effects can lead to size, reproductive, and in some cases distinct caste differences (Wheeler, 1986; Winston, 1987; Couvillon & Dornhaus,

2009; Kamakura, 2011; Suryanarayanan *et al.*, 2011a; Suryanarayanan, Hermanson & Jeanne, 2011b). Thus, certain individuals that take care of brood could determine the behavioural types within a colony either through direct communication (Suryanarayanan & Jeanne, 2008) or by manipulating the way in which larvae are raised (Pereboom, Velthuis & Duchateau, 2003; Couvillon & Dornhaus, 2009). Such manipulations during development will likely have a strong influence on the behavioural type of an individual which might then carry over across contexts or over time (Bergmüller, Schürch & Hamilton, 2010).

In addition to events that happen during development, the interactions of adult workers (or lack thereof) in a colony might also determine the behaviour of individuals. Studies in which individuals performing a certain task are removed from a colony show that other individuals change which tasks they perform (Gordon, 1989, 1996; Pinter-Wollman *et al.*, 2012b, and references therein). Thus, the social environment is an important factor in determining how individuals behave.

(d) *Fitness consequences of intra-colony behavioural variation*

Because individuals might not be infinitely flexible, and because switching between behavioural types (see Table 2 for examples) can be costly or inefficient, colonies can circumvent these constraints by maintaining mixtures of individuals with different behavioural types. For example, if the distribution of flowering plants changes within a season or even within a day (O'Neal & Waller, 1984), maintaining a mixture of foragers that vary along the speed-accuracy behavioural axis could allow colonies to respond more quickly to environmental fluctuation (Burns & Dyer, 2008). That is, slower, more accurate individuals can bring large quantities of food back to the colony when food abundance is constant, whereas faster, 'sloppier' individuals might be more efficient at exploiting resources in more frequently changing environments (Raine & Chittka, 2008, 2012; Chittka, Skorupski & Raine, 2009). A colony containing a mixture of these types might thus be able to handle different environments despite individual inflexibility or carryover.

In the ant *Temnothorax longispinosus* and the social spider *Anelosimus studiosus*, groups with a mixture of aggression types tend to have higher fitness than groups with only one type (Modlmeier & Foitzik, 2011; Pruitt & Riechert, 2011). In ant colonies, aggressive individuals defend the nest whereas docile individuals take care of brood. In the spiders, more aggressive individuals are faster to attack and subdue prey (leading to an increase in foraging performance), whereas less aggressive (more docile) individuals promote group cohesiveness by not escalating aggression during foraging and/or contest encounters (the absence of docile individuals can lead to excessive within-group fighting, cannibalism of colony mates, and group disbandment).

However, maintaining a mixture of relatively inflexible behavioural types could also incur costs to the colony. For example, aggressive *Rhytidoponera confusa* ants are not only more aggressive towards other ants, they are also more likely to lunge at or bite their own nestmates (Crosland, 1990).

This suggests a clear trade-off in aggressive behavioural differences, as well as a limited flexibility in the ability to 'shut-off' the aggressive phenotype across contexts.

Not all behaviours are inflexible, however. Pearce, Huang & Breed (2001) showed that although nurse and guard bees differ in their level of aggressiveness in the summer, these task groups could not be distinguished in the winter months, suggesting that aggressive behaviour can be somewhat plastic.

Numerous studies have found benefits to maintaining variation within colonies (Porter & Tschinkel, 1985; Cnaani & Hefetz, 1994; Page *et al.*, 1995a; Weidenmüller, 2004; Powell & Franks, 2005; Billick & Carter, 2007; Mattila & Seeley, 2007; Dornhaus, 2008; Powell, 2008; Modlmeier, Liebmann & Foitzik, 2012; J. M. Jandt & A. Dornhaus, in preparation), but it is not always clear why exactly variation (as opposed to all individuals being of one optimal type) is beneficial. There is a need to develop theory that helps us to understand under which circumstances we would expect variation to be a better solution than one stable optimal type—not only with regards to division of labour, but also in terms of variation in other behavioural types such as aggression or exploratory behaviour.

(2) Variation among colonies

The reproductive unit of social insects on which natural selection acts is the colony. Therefore, consistent differences among colonies could have important implications on the success of certain phenotypes over others, influencing population structure. Variation among colonies in a certain behaviour might in itself have important ecological consequences. For example, colonies of more aggressive species could out-compete other colonies through interference competition (Davidson, 1998; Rowles & O'Dowd, 2007). If colony behaviour is also consistent across contexts, we might find interesting relationships between the success of a colony in one situation and its success or failure in another. For example, if there is a relationship between aggression and exploration, colonies that are both exploratory and aggressive might be better at invading new habitat. However, when these two behaviours are not correlated, exploratory colonies may find new habitat, but be quickly displaced by aggressive colonies (Cote *et al.*, 2010; Fogarty, Cote & Sih, 2011; Sih *et al.*, 2012).

Aside from work to investigate the heritability of pollen-foraging behaviour (Page *et al.*, 1995a,b; Page, Fondrk & Rueppell, 2012a), colony-level behavioural syndromes have received comparatively little attention. However, the behavioural repertoires of colonies include traits measured in classical behavioural syndromes (e.g. aggressiveness), cognitive behavioural syndromes (e.g. learning ability), and social behavioural syndromes (e.g. communication: see Table 2). For example, Crosland (1990) showed that colonies of the ant *Rhytidoponera confusa* consistently exhibited high or low levels of aggression towards non-nestmates. This classical behavioural type of aggression has also been shown to vary among colonies of the invasive Argentine ants (Suarez *et al.*, 2002; Buczkowski & Silverman, 2006), a colony trait that

might help to explain the highly invasive success of certain colonies. Colonies also vary in cognitive behavioural types. For example, bumble bee colonies vary in their learning speed and foraging success (Raine *et al.*, 2006b; Raine & Chittka, 2008, 2012; Ings, Raine & Chittka, 2009; Gill, Ramos-Rodriguez & Raine, 2012). Colonies also differ in the degree to which they perform tasks such as foraging, defence, brood care, or hygienic behaviour. For example, colonies of harvester ants and honey bees vary in foraging activity levels (Gordon, 1991; Cole *et al.*, 2010; Gordon *et al.*, 2011; Wray, Mattila & Seeley, 2011) and in how they regulate foraging activity (Gordon *et al.*, 2011). Colonies of honey bees and black harvester ants (*Messor andreii*) also differ in defensive response (Wray *et al.*, 2011; Pinter-Wollman, Gordon & Holmes, 2012a). Furthermore, Africanized honey bees differ from European honey bees in defensive response (Collins *et al.*, 1982), as well as a number of other traits related to the pollen-hoarding syndrome (Pankiw, 2003).

Like workers, colonies also exhibit consistent behavioural variation across situations. For example, in both honey bees and black harvester ants, colonies vary consistently in their response to food and to disturbance, which are correlated with one another, producing a behavioural syndrome (Wray *et al.*, 2011; Pinter-Wollman *et al.*, 2012a). That is, colonies that send out many workers to forage also send out many workers to defend the nest when disturbed. These behavioural syndromes might have important fitness consequences (Wray *et al.*, 2011) and could result from constraints on colony behaviour. For example, workers might follow similar local rules in various situations (e.g. patrolling and foraging; Gordon *et al.*, 2011), and external features such as nest structure could influence how a colony behaves in various situations (Pinter-Wollman, 2012; Pinter-Wollman *et al.*, 2012a).

(a) Mechanisms underlying variation among colonies in behavioural type

(i) *Colony genetics.* The genetic make-up of a social insect colony is generally represented by the genotype of the queen and the number of unrelated males she mates with. The extent to which workers vary genetically might have an effect on the behavioural type observed at the colony level (see Section II.1c). Colonies with greater genetic variation might achieve better rates of thermoregulatory homeostasis (Jones *et al.*, 2004; Oldroyd & Fewell, 2007), foraging ability (Page *et al.*, 1995a; Mattila, Burke & Seeley, 2008), or productivity and fitness (Mattila & Seeley, 2007). Whether the behavioural type or genotype of a queen can be used to predict colony behavioural type has yet to be studied (but see Section II.2a.iii), although there is evidence that honey bee male genotype can be used to predict behavioural type in an offspring colony (Page *et al.*, 2012a).

(ii) *Colony composition and emergent behaviour.* The frequency of individual behavioural types in a colony might affect colony behavioural type and/or colony performance (Hillesheim, Koeniger & Moritz, 1989; Tsuji, 1994; Paleolog, 2009; Michelena *et al.*, 2010; J. M. Jandt & A. Dornhaus,

in preparation). Consistent individual differences among colonies might emerge both from variation in the average behavioural types of their workers and from the distribution of worker behavioural types (Pinter-Wollman, 2012). For example, if ant colonies vary in aggressive behaviour, this could be caused by the number of highly aggressive ants present in each colony (Crosland, 1990). That is, there might be many individuals of an aggressive caste (e.g. high numbers of soldiers in an aggressive colony), or alternatively there could be a high frequency of aggressive individuals within each caste.

Because the collective behaviour of colonies is an emergent property of individual workers that follow simple, local rules (Camazine *et al.*, 2001; O'Donnell & Bulova, 2007a; Sumpter, 2010), careful consideration is needed when measuring colony behaviour. For example, to determine colony aggression, it is not enough to sample a few workers, put each in an aggression assay against one other worker and sum their response to generate a colony aggression level. A more appropriate test would be to place two colonies in a situation where they are competing over a scarce resource (such as food or a nest site). Using entire colonies allows us to observe colony-level behaviours without the confounding factor of subsampling heterogeneous colony populations. That is, the sum of the individual behaviours might not necessarily represent the behaviour of the whole colony.

The relationship between colony composition and emergent colony behaviour might be non-linear, i.e. just a few individuals could significantly change the behaviour of a group or population as a whole (Sih & Watters, 2005; Sih, Hanser & McHugh, 2009). The queen of a social insect colony, as the primary reproductive, might be considered a conspicuous type of keystone (or key) individual (Table 1), but rarely does she take an active role in influencing the behaviour of other workers in the colony in non-reproductive contexts (Jeanson, 2012). Workers might also take on key roles in the colony, either as highly active or aggressive individuals that affect the overall performance of a colony (Paleolog, 2009), or as informed individuals that influence nest-site selection decision-making (Couzin *et al.*, 2005). For example, in honey bees, the presence of only a few aggressive individuals drastically changes the behaviour of the colony, because the individuals performing the task are a non-random (more aggressive) subset of the colony (Paleolog, 2009). So, although there are only a few aggressive individuals, those are the ones performing the task resulting in an aggressive colony (Paleolog, 2009). In *Anelosimus studiosus* spiders, the presence of only a few aggressive individuals drastically changes the efficiency with which colonies feed, and alters the nature of a number of species interactions (Pruitt & Ferrari, 2011). In these examples a few aggressive individuals can create a bias towards the colony becoming highly aggressive.

Key individuals (also referred to as catalysts, performers, and organisers; Robson & Traniello, 1999) can assume various roles. As colony behavioural outcomes might be described in a number of ways (foraging success, fecundity, activity, etc.), who is recognized as a keystone individual

might differ based on which output is being measured. Individuals performing keystone roles could increase performance by better facilitating information flow through the colony (Couzin *et al.*, 2005; Pinter-Wollman *et al.*, 2011), facilitate an increase in the rate of activity in other workers (Cao *et al.*, 2009), motivate nestmates to begin foraging (O'Donnell, 2006; Molet *et al.*, 2008), communicate more avidly with others in the colony (Cao *et al.*, 2007), or increase cohesion within the colony. For example, in honey bee colonies, certain individuals (i.e. 'activators') are more likely to roam the nest and activate idle nestmates by using vibration signals (Cao *et al.*, 2009). An interesting next step will be to determine whether the number or proportion of activators in the nest can predict overall activity level of the colony. In yet another type of keystone role, elite workers (or 'generalist elites') perform many tasks (unlike task specialists that tend to focus on only one task; Table 1), with little or no help from other workers (Dornhaus, 2008; Pinter-Wollman *et al.*, 2012b). For example, in *Leptothorax allardycei*, over 50% of the colony has been described as inactive—leaving the majority of tasks to be performed by the remaining workers (Cole, 1986). If this percentage were to vary, the behavioural type of the colony might change in the degree to which tasks are performed. Understanding keystone roles, the types of individuals that engage in those roles, and the rate of turnover within colonies is an important issue in any social group (including humans), and detailed studies can readily be conducted using social insects, e.g. *via* removal of diligent workers (Pinter-Wollman *et al.*, 2012b, and references therein).

(iii) *Colony environment.* Once a colony is established, the role of the queen rarely extends beyond functioning as a reproductive organ. Therefore, in addition to the difficulty of studying queen behaviour in the nest, research on queens is limited to what they do while founding a colony (Fewell & Page, 1999; Cahan & Fewell, 2004). Foundresses (pre-egg-laying reproductive females) vary in their method of dispersal (founding, swarming, budding, joining) (Augustin, Santos & Elliot, 2011). For example, a *Temnothorax longispinosus* ant foundress can either fly off and found her own colony (independent foundress) or join an already established nest (dependent foundress) (Howard, 2006). The level of conspecific tolerance seen during the initial nesting phase can also vary. For example, *Polistes dominulus* paper wasp foundresses can independently initiate or co-found colonies, and the number of foundresses can vary across colonies (Zanette & Field, 2011).

What remains unclear is how variation in queen behaviour translates into variation in colony behaviour. If queens vary inherently in traits that affect their habitat selection, this could have long-term effects on the colony, e.g. bold queens might found colonies in a densely populated, but high-quality, location and shy queens might establish further from other colonies but perhaps in locations with scarcer resources. This variation in microhabitat could influence how early resources for initial brood-rearing are acquired, in turn, impacting colony behavioural type. Once established, the on-going role of the queen in the colony can vary. In

eusocial naked mole rats, *Heterocephalus glaber*, for example, queens have been shown to activate lazy workers (Reeve, 1992). If queens vary in their thresholds to tolerate inactive workers, this might impact the overall activity of the colony.

(b) Fitness consequences of colony behavioural syndromes

A colony's behavioural type (e.g. in terms of aggressiveness, defensive response, foraging activity, or undertaking) can influence its fitness (in terms of colony productivity or winter survival; Wray *et al.*, 2011). As a behavioural syndrome inherently suggests limited behavioural plasticity, behavioural syndromes are often associated with some kind of trade-off. Beneficial responses to one stimulus might be coupled with sub-optimal behaviour in response to others. These costs might not be apparent until a colony is put in an alternative context. For example, more aggressive or exploratory colonies might out-compete other colonies in areas of limited resources. However in areas with plentiful resources, excessive or unnecessary contests with other colonies could result in losses of brood or workers, with no additional resources gained.

The fitness consequences of behavioural syndromes at the colony level are difficult to study because of the long generation time of social insect colonies and the paucity of studies on their reproductive success. However, indirect measures of colony success such as growth and foraging activity might provide some insights about the importance of behavioural syndromes at the colony level. For example, *Pogonomyrmex barbatus* ant colonies vary in how they regulate their foraging behaviour, which affects how much they forage on a given day (Gordon, 1991; Gordon *et al.*, 2011) and potentially how much food the colony can store. Honey bee colonies bred for better hygienic behaviour (efficient removal of dead larvae) are less susceptible to disease (Lapidge, Oldroyd & Spivak, 2002), possibly increasing overall health. Other research is beginning to show that connections between colony traits or behavioural types might have significant impact on colony performance and fitness (Suarez *et al.*, 2002; Buczkowski & Silverman, 2006; Raine & Chittka, 2008; Paleolog, 2009; Gordon *et al.*, 2011; Wray *et al.*, 2011; Modlmeier *et al.*, 2012; Pinter-Wollman *et al.*, 2012a).

III. INSIGHTS FROM BEHAVIOURAL SYNDROMES FOR THE STUDY OF SOCIAL INSECTS

(1) Trade-offs

A common theme in the behavioural syndrome literature is measuring the benefits or costs of an inflexible behavioural type across contexts. For instance, consistently bold and aggressive individuals often enjoy increased foraging success (Sih *et al.*, 2003; Short & Petren, 2008), growth rates (Biro & Stamps, 2008), and superior performance during contests (Pruitt & Riechert, 2009), but also suffer increased susceptibility to predation or parasites (Fig. 1A; Pontier *et al.*,

1998; Biro *et al.*, 2006). Other trade-offs can be less intuitive, particularly those involving social interactions. For example, male western blue birds (*Sialia mexicana*) exhibit a syndrome of aggressiveness between nest defence and male-male competition that ultimately results in decreased fitness. Highly aggressive males engage in excessive male-male contests during the breeding season, and consequently fail to provision nesting females with necessary food. This results in disrupted incubation patterns, as females must leave the nest to forage, reducing offspring survival (Duckworth, 2006).

Although recent literature has challenged the potential of behavioural syndromes to set evolutionary constraints (Bell, 2005; Dingemanse *et al.*, 2007), a sizeable (and growing) body of literature continues to document numerous trade-offs associated with behavioural consistency. It is yet unclear whether these trade-offs are the result of proximate constraints such as genetic linkage. Optimality theory suggests selection should act against limited behavioural plasticity; however there might be adaptive benefits (Sih *et al.*, 2004a). For example, limited plasticity prevents animals from making more costly errors in an unpredictable environment (Sih, 1992). Conversely, the social insect literature shows examples of trade-offs being the result of genetic linkage. For example, the reproductive ground-plan hypothesis proposes that different levels of pollen-hoarding in honey bee foragers is the result of evolutionary co-option of the female reproductive system (Amdam & Page, 2010), and genetic linkage between foraging behaviour, reproduction (Graham *et al.*, 2011) and at least two genes (*HR46* and *PDK1*) has been demonstrated (Wang *et al.*, 2009).

Because consistent traits across different contexts can produce performance trade-offs, determining the source and magnitude of these trade-offs is a continuing goal of behavioural syndromes research. Furthermore, until recently, colony-level behavioural variation has not received as much attention as within-colony variation (see Table 2). It could be useful to examine under what conditions within-population variation can evolve if trade-offs are present, and how those conditions match those where variation persists in populations of non-social animals.

(2) Feedback

Recent work has addressed how behavioural syndromes are maintained in a population. Several authors have suggested that a key mechanism for maintaining consistent differences in behavioural types is the feedback between the individual's behaviour and its state (Luttbeg & Sih, 2010; Wolf & Weissing, 2010), such as the feedback linking an individual's behavioural type and its energy reserves, condition, or social rank. For example, colonies that are quick to exploit a new resource could gain an early benefit, but they might also be easily displaced by a more dominant or aggressive colony and must then locate a new resource pool. These differences in strategy might be the result of differing behavioural types, and are consistently reinforced by the social rank of a colony. A central premise of this frequency-dependent selection has been that the mix of behavioural types in social

groups influences social dynamics and the fitness of each behavioural type. Because social dynamics are important in social insects, understanding how variation in behavioural types in the colony influences individual and group outcomes is an exciting area for future study (Pinter-Wollman, 2012).

(3) Statistical methods

Studies in social insects have illustrated variation in classical, cognitive, and social behavioural types both within and among colonies (Table 2). Quantifying various behavioural types, as well as understanding how variation across multiple behavioural axes might affect colony fitness are topics that are beginning to receive attention in the social insect literature (e.g. see Modlmeier *et al.*, 2012). Scientists studying social insects can draw on advances in statistical methods from the general behavioral syndrome literature for identifying, quantifying, and correlating behavioural types and syndromes across time and context (Sih *et al.*, 2004a; Réale *et al.*, 2007; Garamszegi & Herczeg, 2012). Recently, statistical approaches have been reviewed to quantify animal personalities, behavioural syndromes, and behavioural plasticity (Dingemanse *et al.*, 2010; Dingemanse, Dochtermann & Nakagawa, 2012; Garamszegi & Herczeg, 2012). We briefly highlight some points here, but refer the reader to these reviews for detailed analyses of how and when to use various statistical approaches.

Two key analytical tools for behavioural syndrome research are principal component analyses (PCAs) and pairwise correlations. When used together these analyses provide a convenient way to tackle large data sets where some combination of variables could predict behavioural correlations. Factor analyses, structured equation modeling, generalized linear mixed models, and other statistics have also been applied to behavioural syndromes research (Bell & Stamps, 2004; Dingemanse *et al.*, 2010; Garamszegi & Herczeg, 2012), and these approaches can add to the tool kit available to social insect biologists. While these statistical approaches require very large datasets, they help address the point that variation in behavioural flexibility could be as vital as differences in mean trait values. Moreover, when paired with familiar theory on the trade-offs associated with phenotypic plasticity, these approaches provide us with clear and testable predictions on when trait flexibility *versus* consistency should be favoured.

IV. INSIGHTS FROM SOCIAL INSECTS FOR BEHAVIOURAL SYNDROMES

(1) Group-living might affect behavioural type

Although many studies of behavioural syndromes have been conducted on social animals, and many of the commonly studied behavioural axes are inherently social behaviours (see Table 2), the social ecology of behavioural types has not yet received much experimental attention. For example,

how does a mix of behavioural types in a group influence individual behaviours [social plasticity (Kурvers *et al.*, 2010) or group outcomes (Crosland, 1990; Sih & Watters, 2005; Eldakar *et al.*, 2009; Paleolog, 2009; Pruitt & Riechert, 2011; Schürch, Rothenberger & Heg, 2011)]? Work on social insects has shown that indirect genetic effects (whether offspring are raised by the parent or siblings) can influence colony phenotype (Linksvayer, 2006). Colonial living can also result in greater variation in individual experiences, which can further alter individual behavioural tendencies (Bergmüller *et al.*, 2010; Pruitt *et al.*, 2012). Studies of social insect colonies have a long history of looking at how individuals and their behaviours collectively produce group outcomes (Oster & Wilson, 1978), and there are a variety of examples where feedback between the individual and the rest of the group might affect colony behaviour (Fig. 1B; Crosland, 1990; Beverly *et al.*, 2009; Paleolog, 2009).

Variation in behavioural type within a group could reduce competitive interactions among group members (reviewed in Ratnieks & Reeve, 1992; Bergmüller *et al.*, 2010). Furthermore, the spatial organisation of individuals in a colony might be analogous to niche partitioning measured in behavioural ecology (Sendova-Franks & Franks, 1995; Beverly *et al.*, 2009; Jandt & Dornhaus, 2011; Dall *et al.*, 2012). Recently, social network theory has been employed to study the development and maintenance of behavioural types within a social group (Krause, Lusseau & James, 2009; Sih *et al.*, 2009). There are a variety of examples of the use of network theory to understand how interactions affect group behaviour in social insects as well (Blonder & Dornhaus, 2011; Pinter-Wollman *et al.*, 2011). We propose that a link between models developed on social networks and on behavioural syndromes could benefit both the study of social insects and animal behaviour in general.

The study of behavioural syndromes can also benefit from exploring models that show how division of labour emerges from feedback mechanisms and interactions among workers, and models that predict how the interactions of those workers produce complex patterns and colony behaviour (Fig. 1B; Camazine, 1991; Theraulaz & Bonabeau, 1995; Bonabeau, Theraulaz & Deneubourg, 1998; Theraulaz, Bonabeau & Deneubourg, 1998; Camazine *et al.*, 2001; O'Donnell & Bulova, 2007b). Social feedbacks can facilitate the maintenance of multiple behavioural types within the nest (Gordon, 1989, 1996; Reeve & Nonacs, 1997; Bergmüller *et al.*, 2010). For example, small differences in response thresholds can lead to stable long-term differences in behaviour, even among non-social animals (Fewell & Page, 1999), and the social environment might prevent or promote transitions among tasks (Gordon, 1989; Amdam *et al.*, 2005; Gordon *et al.*, 2005). Researchers of behavioural syndromes are beginning to explore how positive feedback loops might maintain consistent behavioural types (Luttbeg & Sih, 2010). These models can guide further study of how complementary behavioural types in non-social insects can affect group dynamics, and how these in turn can determine the behaviour and fitness of individuals.

(2) Mechanisms affecting behavioural type

A crucial question in behavioural syndrome research is: what are the mechanisms that affect behaviour? Genetic, hormonal, and experiential mechanisms that could interact and predispose an individual to exhibit a particular social behavioural type have been well studied in social insects (see Section II.1c and Table 3). Changes in hormone levels (Robinson, 1987; Giray *et al.*, 2005; Dolezal *et al.*, 2012) and gene expression (Hunt *et al.*, 2010), often associated with a change in social environment (Amdam *et al.*, 2005) affect caste determination and division of labor (Robinson, 1987; Wheeler, Buck & Evans, 2006; Toth *et al.*, 2009; Amdam & Page, 2010; Hunt *et al.*, 2010), cooperative behaviour, in terms of worker egg production (Lin *et al.*, 1999), and aggressive behaviour (Pearce *et al.*, 2001; Tibbetts & Huang, 2010). For example, genetic linkage between foraging and reproductive behaviour (Wang *et al.*, 2009) might not only limit an individual's behavioural plasticity (or act as a 'constraint'), but could also have been an adaptive precursor that allowed solitary ancestors to evolve into social groups (Amdam *et al.*, 2004; Hunt & Amdam, 2005; Smith *et al.*, 2008; Toth *et al.*, 2009; Hunt *et al.*, 2010; Page *et al.*, 2012b).

Morphological differences have also been implicated as a mechanism underlying behavioural syndromes (Sih *et al.*, 2004b). However, it is often difficult to modify the morphology of an organism to determine which aspects of an individual's morphology have the greatest effects on behaviour. Insect colonies provide a unique system in which the demography and composition of a colony can be manipulated, i.e. whole task groups or castes can be removed or proportions adjusted (Porter & Tschinkel, 1985; Cnaani & Hefetz, 1994; Roux & Korb, 2004; Billick & Carter, 2007; J. M. Jandt & A. Dornhaus, in preparation). The nest structure can be considered an extended phenotype of the colony (Tschinkel, 2004), and changes to this structure, i.e. to nest morphology, could have an effect on colony-level behaviours (Pinter-Wollman *et al.*, 2012a). Both colony composition and nest architecture can be manipulated to determine the relative role each plays in the maintenance of a behavioural type.

(3) Development of behavioural types across levels of organisation

When examining the behavioural type of an individual, its behaviour is often examined at different stages or in response to different stimuli (Bell & Stamps, 2004; Briffa, Rundle & Fryer, 2008; Bell, Hankison & Laskowski, 2009; Biro, Beckmann & Stamps, 2010; Dingemanse *et al.*, 2010). It would be considerably more difficult to monitor the changes of individual cells in a vertebrate brain or body to understand how the parts affect the behavioural type of the individual, but we concede that these often play a significant role. However, using social insects we can monitor how variation in worker development [e.g. change in physiology and/or gene expression patterns (Kocher *et al.*, 2010; Herb *et al.*, 2012)], change in colony composition (i.e. the development of the superorganism), and feedback between levels of organisation (Kocher

et al., 2010) can affect behavioral type (Fig. 1B). The insight that can be gained from understanding the biological interactions within a social group and how that translates to colony-level behaviours can provide important insights to those who study behavioural syndromes in solitary organisms.

V. FUTURE DIRECTIONS

In the field of social insects, it has long been acknowledged that variation among workers might affect colony fitness by providing a mechanism for division of labour. Here we have reviewed examples of variation along multiple behavioural axes both at the colony and individual levels. While division of labour has long been speculated as critically important in enhancing colony fitness (Oster & Wilson, 1978), the behavioural syndromes approach provides an alternative method for determining the extent to which variation in behavioural types within colonies (including, but not limited to, those associated with caste and division of labour) and among colonies might affect fitness and ecology. While we acknowledge there are a variety of ways in which the behavioural-syndrome approach can be used to create insightful and important questions, we propose in particular the following three areas that would benefit from the attention of eager social-insect/behavioural syndrome enthusiasts.

(1) What are the adaptive reasons for colonies, or groups of individuals, to maintain mixtures of behavioural types?

As mentioned above (Sections II.1d, and II.2a), many studies have found benefits to maintaining variation within colonies (Porter & Tschinkel, 1985; Cnaani & Hefetz, 1994; Page *et al.*, 1995a; Weidenmüller, 2004; Powell & Franks, 2005; Billick & Carter, 2007; Mattila & Seeley, 2007; Dornhaus, 2008; Powell, 2008; Modlmeier *et al.*, 2012; J. M. Jandt & A. Dornhaus, in preparation). However, it is not always clear why exactly variation (as opposed to all individuals being of one optimal type) is beneficial. Furthermore, these studies can be expanded to include non-social-insect group-dwelling species (Sih & Watters, 2005; Eldakar *et al.*, 2009; Pruitt & Riechert, 2011).

Trait complexes could be the product of genetic or physiological linkage, rather than adaptive combinations for optimal division of labour. It is possible that mixtures of behavioural types within a colony (e.g. aggressive or exploratory types) shaped the way in which task groups (e.g. guards or patrollers) have evolved. We find correlations among multiple physiological factors that affect the behaviour of individual workers—both within and between castes—that are not obviously explained by the requirements of the tasks themselves.

Mixtures of behavioural types within a colony might affect the flexibility of the colony to react to changing environmental conditions. For example, colonies could become more active when environmental conditions are

favourable (Gordon, 1991; Cole *et al.*, 2010; Pinter-Wollman, 2012). The more variation among workers, the more types of individuals are available for each ecological condition, and indeed, high variation among workers increases colony productivity (Page *et al.*, 1995a; Jones *et al.*, 2004; Mattila & Seeley, 2007; Oldroyd & Fewell, 2007; Mattila *et al.*, 2008; Modlmeier & Foitzik, 2011). Differences in colony ability to respond to changing conditions could be critical in the modern world of human-induced rapid environmental change (HIREC; Sih, Ferrari & Harris, 2011), including changes in climate and habitat, and exposure to novel species or contaminants.

Finally, behavioural variation might persist in a group *via* social heterosis (Nonacs & Kapheim, 2007, 2008). Social heterosis is a mutually beneficial process that occurs when individuals benefit by associating with genetically dissimilar individuals. Under such circumstances, groups (or demes) of mixed genotype outperform monotypic groups and singleton individuals of any genotype. Potential mechanisms behind social heterosis include reduced competition among group constituents, complimentary skill pools, or instances where individuals are incapable of approximating the optimal genotype for a population. Importantly, social heterosis is sufficient in itself to maintain genetic diversity at one or more loci in the face of genetic drift (Nonacs & Kapheim, 2007, 2008). The predicted outcomes of social heterosis are high genetic diversity (and heritability) of functional traits (Stirling, Réale & Roff, 2002; Nonacs & Kapheim, 2007), and a positive association between within-group trait variation and fitness for group constituents. Both patterns have been observed in numerous test systems, from microbes (Vos & Velicer, 2006), to primates (Wooding *et al.*, 2006). In social insects, social heterosis might be an explanation as to why unrelated individuals will cooperate with one another during nest-founding in some species (Tschinkel, 1998; Queller *et al.*, 2000; Jeanson & Fewell, 2008; Leadbeater *et al.*, 2010; Zanette & Field, 2011).

(2) Do colonies differ in their degree of cooperation?

A growing field in behavioural syndrome research is to identify the degree to which individuals or groups vary in cooperativeness. A cooperative behavioural type could be measured in terms of group cohesion, lack of within-group conflict, short time spent on tasks devoted to self- *versus* group-maintenance, and low effort spent on reproduction (or competing to reproduce). Social insect colonies are generally assumed to be highly cooperative, yet extreme aggression, worker reproduction, and over-policing of queen-laid eggs are a few examples of within-colony non-cooperative (or cheating) behaviour (Bourke, Have & Franks, 1988; Schmid-Hempel, 1990; D'Ettorre, Heinze & Ratnieks, 2004; Wenseleers & Ratnieks, 2006a). Along with the examples listed above, we might measure colony-level cooperation based on the proportion of individuals that are inactive, or regularly refrain from working (Lindauer, 1952; Herbers, 1983; Cole, 1986; Dornhaus *et al.*, 2008; Jandt & Dornhaus, 2009). These inactive individuals might be a reserve set of

workers, or they might be selfishly hoarding their fat reserves and avoiding risky tasks to gain an opportunity to reproduce (Lindauer, 1952; Michener, 1964; Jandt & Dornhaus, 2011). While inactive individuals have been identified in many social insect species, it is unclear whether the proportion of 'lazy' individuals varies across colonies and how that might affect colony fitness.

An additional method for measuring cooperativeness is to examine worker policing behaviour—that is, some workers will regularly inspect the eggs laid in the nest and will eat those that were not laid by the queen (Ratnieks & Reeve, 1992; Wenseleers & Ratnieks, 2006b). Some individuals occasionally 'make a mistake' and eat queen eggs too, although often at a lower rate (D'Ettorre *et al.*, 2004). Degree of cooperation might be measured by examining the variation across colonies in the ratio of policing queen eggs *versus* worker eggs, and how that might affect colony fitness.

If colonies differ in how cooperative they are, what will this mean ecologically? Will more cooperative colonies have an ecological advantage in some environments, whereas less cooperative colonies have an advantage in others? Argentine ant colonies that have successfully invaded urban areas are less aggressive (and, hence, more cooperative) with conspecific colonies than those that remain in their native range (Buczkowski, 2010). That is, colonies interact and cooperate with one another, and ultimately end up forming 'supercolonies', a phenotype that allows them aggressively to expand and dominate the areas they colonize (Holway, 1999; Suarez *et al.*, 1999; Holway & Suarez, 2004; Buczkowski, 2010). In native areas, on the other hand, the same species will coexist with a variety of other ant species, and are less likely to form these supercolonies that can aggressively out-compete other species for territory. The theoretical basis for the evolution, maintenance and consequences of individual variation in cooperation has been developed in the behavioural syndrome literature (Bergmüller *et al.*, 2010).

(3) How does consistency in behavioural type affect task performance when individuals switch task?

The extent to which behavioural type persists over time within an individual, and how this could affect future task performance, has received little attention. In honey bees, for example, genetic factors are linked to how quickly an individual transitions between tasks (Page *et al.*, 1998; Rueppell *et al.*, 2004; Amdam & Page, 2010). Research on the honey bee pollen-hoarding syndrome has shown that this transition time is also linked to learning performance, and a variety of other behavioural types (Page *et al.*, 2012b). Incorporating a behavioural-syndromes approach also allows us to ask whether this variation in transition time correlates with classical, cognitive, or social behavioural axes (i.e. do more shy workers take longer to transition between tasks? Do they vary in the time required to master that new task or skill set?). Ultimately, this could reveal why individuals vary

in the time scale over which they become specialised on a given task.

Consistency of behavioural type over time could also affect division of labour. For example, are individuals that exhibit high levels of boldness or aggressiveness as a young nurse more likely to guard when they are older, whereas those that exhibit high levels of exploratory behaviour more likely to forage? What role does senescence, or changes in task based on decreased ability, play in behavioural syndromes? Do individuals change behavioural type as they age? Studies using the behavioural-syndromes approach could be useful when considering how workers maintain task organisation in a decentralized system.

VI. CONCLUSIONS

(1) There are clearly two fields of knowledge, the study of social insects and the study of behavioural syndromes, working in parallel on complementary ideas, yet until recently little effort has been invested in bridging these two fields. For example, researchers interested in studying behavioural syndromes of group-living organisms would benefit from the rich literature in social insects on how social interactions can affect individual decision-making. On the other hand, social insect researchers exploring evolutionary causes and consequences of maintaining behavioural types at multiple levels of social organisation would benefit from theory developed in behavioural syndrome research on trade-offs (Fig. 1A), as well as on statistical methods to analyse these phenomena.

(2) One of the advantages of studying behavioural syndromes in social insects is the ability to compare causes and consequences at multiple levels of social organisation. Because of the tight interplay between behavioural type of individuals and groups, it is clear that to understand behavioural syndromes of social animals it is important to understand the feedback among the levels of social organisation (Fig. 1B).

(3) Monitoring the development of an individual's behavioural type (i.e. changes of individual cells in a vertebrate brain or body to understand how the parts affect the behavioural type of the individual) may be more feasible using social insects. Here, we can monitor how variation in worker development (e.g. change in physiology and/or gene expression patterns), change in colony composition (i.e. the development of the superorganism), and feedback between levels of organisation can affect behavioral type. Furthermore, mechanisms that affect the development of these types, as well as the fitness effects of maintaining syndromes—have been measured both within and between colonies.

(4) We propose that both fields take advantage of the knowledge gleaned from the other to avoid reinventing the wheel too many times.

VII. ACKNOWLEDGEMENTS

All authors wish to acknowledge the International Union for the Study of Social Insects (IUSSI) international conference held in Copenhagen, Denmark, August 2010. The symposium on behavioural syndromes in social insects provided the fodder for many of the authors to begin the discussion of bridging the two fields of behavioural ecology and social insects in a more formal way. We also wish to thank the Dornhaus Lab (University of Arizona, U.S.A.) and Toth Lab (Iowa State University, U.S.A.) for helpful discussions on manuscript revisions, and are grateful for the thoughtful suggestions from Dr William Foster and two anonymous reviewers to improve the manuscript. Funding for travel was supported by NSF IOS-0841756.

VIII. REFERENCES

- ADAMS, E. S., ATKINSON, L. & BULMER, M. S. (2007). Relatedness, recognition errors, and colony fusion in the termite *Nasutitermes corniger*. *Behavioral Ecology and Sociobiology* **61**, 1195–1201.
- AMDAM, G. V., AASE, A. L. T. O., SEEHUS, S.-C., FONDRK, M. K., NORBERG, K. & HARTFELDER, K. (2005). Social reversal of immunosenescence in honey bee workers. *Experimental Gerontology* **40**, 939–947.
- AMDAM, G. V., NORBERG, K., FONDRK, M. K. & PAGE, R. E. (2004). Reproductive ground plan may mediate colony-level selection effects on individual foraging behavior in honey bees. *Proceedings of the National Academy of Sciences of the United States of America* **101**, 11350–11355.
- AMDAM, G. V. & PAGE, R. E. J. (2010). The developmental genetics and physiology of honeybee societies. *Animal Behaviour* **79**, 973–980.
- ANDERSON, C. & RATNIEKS, F. L. W. (1999). Worker allocation in insect societies: coordination of nectar foragers and nectar receivers in honey bee (*Apis mellifera*) colonies. *Behavioral Ecology and Sociobiology* **46**, 73–81.
- ARATHI, H. S., BURNS, I. & SPIVAK, M. (2000). Ethology of hygienic behaviour in the honey bee *Apis mellifera* L. (Hymenoptera : Apidae): behavioural repertoire of hygienic bees. *Ethology* **106**, 365–379.
- AUGUSTIN, J. O., SANTOS, J. F. L. & ELLIOT, S. L. (2011). A behavioral repertoire of *Atta sexdens* (Hymenoptera, Formicidae) queens during the claustral founding and ergonomic stages. *Insectes Sociaux* **58**, 197–206.
- BELL, A. M. (2005). Behavioural differences between individuals and two populations of stickleback (*Gasterosteus aculeatus*). *Journal of Evolutionary Biology* **18**, 464–473.
- BELL, A. M., HANKISON, S. J. & LASKOWSKI, K. L. (2009). The repeatability of behaviour: a meta-analysis. *Animal Behaviour* **77**, 771–783.
- BELL, A. M. & SIH, A. (2007). Exposure to predation generates personality in threespined sticklebacks (*Gasterosteus aculeatus*). *Ecology Letters* **10**, 828–834.
- BELL, A. M. & STAMPS, J. A. (2004). Development of behavioural differences between individuals and populations of sticklebacks, *Gasterosteus aculeatus*. *Animal Behaviour* **68**, 1339–1348.
- BEN-SHAHAR, Y., ROBICHON, A., SOKOLOWSKI, M. B. & ROBINSON, G. E. (2002). Influence of gene action across different time scales on behavior. *Science* **296**, 741–744.
- BENGSTON, S. E. & DORNHAUS, A. (2013). Colony size does not predict foraging distance in the ant *Temnothorax rugatulus*: a puzzle for standard scaling models. *Insectes Sociaux* **60**, 93–96.
- BERGMÜLLER, R., SCHÜRCH, R. & HAMILTON, I. M. (2010). Evolutionary causes and consequences of consistent individual variation in cooperative behaviour. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 2751–2764.
- BESHES, S. N. & FEWELL, J. H. (2001). Models of division of labor in social insects. *Annual Review of Entomology* **46**, 413–440.
- BEVERLY, B. D., MCLENDON, H., NACU, S., HOLMES, S. & GORDON, D. M. (2009). How site fidelity leads to individual differences in the foraging activity of harvester ants. *Behavioral Ecology* **20**, 633–638.
- BILLICK, I. & CARTER, C. (2007). Testing the importance of the distribution of worker sizes to colony performance in the ant species *Formica obscuripes* Forel. *Insectes Sociaux* **54**, 113–117.
- BIRO, P. A., ABRAHAM, M. V., POST, J. R. & PARKINSON, E. A. (2006). Behavioural trade-offs between growth and mortality explain evolution of submaximal growth rates. *Journal of Animal Ecology* **75**, 1165–1171.

- BIRO, P. A., BECKMANN, C. & STAMPS, J. A. (2010). Small within-day increases in temperature affects boldness and alters personality in coral reef fish. *Proceedings of the Royal Society B: Biological Sciences* **277**, 71–77.
- BIRO, P. A. & STAMPS, J. A. (2008). Are animal personality traits linked to life-history productivity? *Trends in Ecology & Evolution* **23**, 361–368.
- BLANCHARD, G. B., ORLEDGE, G. M., REYNOLDS, S. E. & FRANKS, N. R. (2000). Division of labour and seasonality in the ant *Leptothorax albipennis*: worker corpulence and its influence on behaviour. *Animal Behaviour* **59**, 723–738.
- BLONDER, B. & DORNHAUS, A. (2011). Time-ordered networks reveal limitations to information flow in ant colonies. *PLoS One* **6**, e20298.
- BONABEAU, E., THERAULAZ, G. & DENEUBOURG, J.-L. (1998). Fixed response thresholds and the regulation of division of labor in insect societies. *Bulletin of Mathematical Biology* **60**, 753–807.
- BOURKE, A. F. G., VAN DER HAVE, T. M. & FRANKS, N. R. (1988). Sex ratio determination and worker reproduction in the slave-making ant *Harpagoxenus sublaevis*. *Behavioral Ecology and Sociobiology* **23**, 233–245.
- BREED, M. D., GUZMAN-NOVOA, E. & HUNT, G. J. (2004). Defensive behavior of honey bees: organization, genetics, and comparisons with other bees. *Annual Review of Entomology* **49**, 271–298.
- BRENT, C., PEETERS, C., DIETMANN, V., CREWE, R. & VARGO, E. (2006). Hormonal correlates of reproductive status in the queenless ponerine ant, *Streblognathus peetersi*. *Journal of Comparative Physiology. A* **192**, 315–320.
- BRIFFA, M., RUNDLE, S. D. & FRYER, A. (2008). Comparing the strength of behavioural plasticity and consistency across situations: animal personalities in the hermit crab *Pagurus bernhardus*. *Proceedings of the Royal Society B: Biological Sciences* **275**, 1305–1311.
- BROWN, J. H., ROSS, B., MCCAULEY, S., DANCE, S., TAYLOR, A. C. & HUNTINGFORD, F. A. (2003). Resting metabolic rate and social status in juvenile giant freshwater prawns, *Macrobrachium rosenbergii*. *Marine and Freshwater Behaviour and Physiology* **36**, 31–40.
- BRYANT, D. M. & NEWTON, A. V. (1994). Metabolic costs of dominance in dippers, *Cinclus cinclus*. *Animal Behaviour* **48**, 447–455.
- BUCZKOWSKI, G. (2010). Extreme life history plasticity and the evolution of invasive characteristics in a native ant. *Biological Invasions* **12**, 3343–3349.
- BUCZKOWSKI, G. & SILVERMAN, J. (2006). Geographical variation in Argentine ant aggression behaviour mediated by environmentally derived nestmate recognition cues. *Animal Behaviour* **71**, 327–335.
- BURNS, J. G. & DYER, A. G. (2008). Diversity of speed-accuracy strategies benefits social insects. *Current Biology* **18**, R953–R954.
- CAHAN, S. H. (2001). Cooperation and conflict in ant foundress associations: insights from geographical variation. *Animal Behaviour* **61**, 819–825.
- CAHAN, S. H. & FEWELL, J. H. (2004). Division of labor and the evolution of task sharing in queen associations of the harvester ant *Pogonomyrmex californicus*. *Behavioral Ecology and Sociobiology* **56**, 9–17.
- CAHAN, S. H. & KELLER, L. (2003). Complex hybrid origin of genetic caste determination in harvester ants. *Nature* **424**, 306–309.
- CAKMAK, I., SANDERSON, C., BLOCKER, T. D., PHAM, L. L., CHECOTAH, S., NORMAN, A. A., HARADER-PATE, B. K., REIDENBAUGH, R. T., NENCHEV, P., BARTHELL, J. F. & WELLS, H. (2009). Different solutions by bees to a foraging problem. *Animal Behaviour* **77**, 1273–1280.
- CAMAZINE, S. (1991). Self-organizing pattern formation on the combs of honey bee colonies. *Behavioral Ecology and Sociobiology* **28**, 61–76.
- CAMAZINE, S., DENEUBOURG, J.-L., FRANKS, N. R., SNEYD, J., THERAULAZ, G. & BONABEAU, E. (2001). Self-Organization in Biological Systems. Princeton University Press, Princeton.
- CAMERON, S. A. & ROBINSON, G. E. (1990). Juvenile hormone does not affect division of labor in bumble bee colonies (Hymenoptera: Apidae). *Annals of the Entomological Society of America* **83**, 626–631.
- CANT, M. A., LLOP, J. B. & FIELD, J. (2006). Individual variation in social aggression and the probability of inheritance: theory and a field test. *American Naturalist* **167**, 837–852.
- CAO, T. T., HYLAND, K. M., MALECHUK, A., LEWIS, L. A. & SCHNEIDER, S. S. (2007). The influence of the vibration signal on worker interactions with the nest and nest mates in established and newly founded colonies of the honey bee, *Apis mellifera*. *Insectes Sociaux* **54**, 144–149.
- CAO, T. T., HYLAND, K. M., MALECHUK, A., LEWIS, L. A. & SCHNEIDER, S. S. (2009). The effect of repeated vibration signals on worker behavior in established and newly founded colonies of the honey bee, *Apis mellifera*. *Behavioral Ecology and Sociobiology* **63**, 521–529.
- CAREAU, V., THOMAS, D., HUMPHRIES, M. M. & REALE, D. (2008). Energy metabolism and animal personality. *Oikos* **117**, 641–653.
- CHANDRA, S. B. C., HOSLER, J. S. & SMITH, B. H. (2000). Heritable variation for latent inhibition and its correlation with reversal learning in honeybees (*Apis mellifera*). *Journal of Comparative Psychology* **114**, 86–97.
- CHAPMAN, B. B., THAIN, H., COUGHLIN, J. & HUGHES, W. O. H. (2011). Behavioural syndromes at multiple scales in *Myrmica* ants. *Animal Behaviour* **82**, 391–397.
- CHAPPELL, M. A., GARLAND, T., REZENDE, E. L. & GOMES, F. R. (2004). Voluntary running in deer mice: speed, distance, energy costs and temperature effects. *Journal of Experimental Biology* **207**, 3839–3854.
- CHITTKA, L., DYER, A. G., BOCK, F. & DORNHAUS, A. (2003). Bees trade off foraging speed for accuracy. *Nature* **424**, 388.
- CHITTKA, L., INGS, T. C. & RAINE, N. E. (2004). Chance and adaptation in the evolution of island bumblebee behaviour. *Population Ecology* **46**, 243–251.
- CHITTKA, L., SKORUPSKI, P. & RAINE, N. E. (2009). Speed-accuracy tradeoffs in animal decision making. *Trends in Ecology & Evolution* **24**, 400–407.
- CNAANI, J. & HEFETZ, A. (1994). The effect of workers size frequency distribution on colony development in *Bombus terrestris*. *Insectes Sociaux* **41**, 301–307.
- COLE, B. J. (1986). The social behavior of *Leptothorax allardycei* (Hymenoptera, Formicidae): time budgets and the evolution of worker reproduction. *Behavioral Ecology and Sociobiology* **18**, 165–173.
- COLE, B. J., SMITH, A. A., HUBER, Z. J. & WIERNASZ, D. C. (2010). The structure of foraging activity in colonies of the harvester ant, *Pogonomyrmex occidentalis*. *Behavioral Ecology* **21**, 337–342.
- COLLINS, A. M., RINDERER, T. E., HARBO, J. R. & BOLLEN, A. B. (1982). Colony defense by Africanized and European honey bees. *Science* **218**, 72–74.
- COTE, J., CLOBERT, J., BRODIN, T., FOGARTY, S. & SHI, A. (2010). Personality-dependent dispersal: characterization, ontogeny and consequences for spatially structured populations. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 4065–4076.
- COUVILLON, M. J. & DORNHAUS, A. (2009). Location, location, location: larvae position inside the nest is correlated with adult body size in worker bumble-bees (*Bombus impatiens*). *Proceedings of the Royal Society B: Biological Sciences* **276**, 2411–2418.
- COUVILLON, M. J., JANDT, J. M., BONDS, J., HELM, B. R. & DORNHAUS, A. (2011). Percent lipid is associated with body size but not task in the bumble bee *Bombus impatiens*. *Journal of Comparative Physiology. A, Neuroethology Sensory Neural and Behavioral Physiology* **197**, 1097–1104.
- COUZIN, I. D., KRAUSE, J., FRANKS, N. R. & LEVIN, S. A. (2005). Effective leadership and decision-making in animal groups on the move. *Nature* **433**, 513–516.
- CROSLAND, M. W. J. (1990). Variation in ant aggression and kin discrimination ability within and between colonies. *Journal of Insect Behavior* **3**, 359–379.
- CUTTS, C. J., METCALFE, N. B. & TAYLOR, A. C. (1998). Aggression and growth depression in juvenile Atlantic salmon: the consequences of individual variation in standard metabolic rate. *Journal of Fish Biology* **52**, 1026–1037.
- DALL, S. R. X., BELL, A. M., BOLNICK, D. I. & RATNIEKS, F. L. W. (2012). An evolutionary ecology of individual differences. *Ecology Letters* **15**, 1189–1198.
- DALL, S. R. X., HOUSTON, A. I. & MCNAMARA, J. M. (2004). The behavioural ecology of personality: consistent individual differences from an adaptive perspective. *Ecology Letters* **7**, 734–739.
- DAUGHERTY, T. H. F., TOTH, A. L. & ROBINSON, G. E. (2011). Nutrition and division of labor: effects on foraging and brain gene expression in the paper wasp *Polistes metricus*. *Molecular Ecology* **20**, 5337–5347.
- DAVIDSON, D. W. (1998). Resource discovery versus resource domination in ants: a functional mechanism for breaking the trade-off. *Ecological Entomology* **23**, 484–490.
- DEL GIUDICE, M. (2011). Sex differences in romantic attachment: a meta-analysis. *Personality and Social Psychology Bulletin* **37**, 193–214.
- DEL GIUDICE, M., ELLIS, B. J. & SHIRTCLIFF, E. A. (2011). The Adaptive Calibration Model of stress responsivity. *Neuroscience and Biobehavioral Reviews* **35**, 1562–1592.
- D'ETTORRE, P., HEINZE, J. & RATNIEKS, F. L. W. (2004). Worker policing by egg eating in the ponerine ant *Pachycondyla inversa*. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**, 1427–1434.
- DIEZ, L., DENEUBOURG, J.-L., HOEBEKE, L. & DETRAIN, C. (2011). Orientation in corpse-carrying ants: memory or chemical cues? *Animal Behaviour* **81**, 1171–1176.
- DINGEMANSE, N. J., BOTH, C., DRENT, P. J. & TINBERGEN, J. M. (2004). Fitness consequences of avian personalities in a fluctuating environment. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **271**, 847–852.
- DINGEMANSE, N. J., DOCHTERMAN, N. A. & NAKAGAWA, S. (2012). Defining behavioural syndromes and the role of 'syndrome deviation' in understanding their evolution. *Behavioral Ecology and Sociobiology* **66**, 1543–1548.
- DINGEMANSE, N. J., KAZEM, A. J. N., REALE, D. & WRIGHT, J. (2010). Behavioural reaction norms: animal personality meets individual plasticity. *Trends in Ecology & Evolution* **25**, 81–89.
- DINGEMANSE, N. J., WRIGHT, J., KAZEM, A. J. N., THOMAS, D. K., HICKLING, R. & DAWNAY, N. (2007). Behavioural syndromes differ predictably between 12 populations of three-spined stickleback. *Journal of Animal Ecology* **76**, 1128–1138.
- DOLEZAL, A. G., BRENT, C. S., HÖLLDOBLER, B. & ANDAM, G. V. (2012). Worker division of labor and endocrine physiology are associated in the harvester ant, *Pogonomyrmex californicus*. *The Journal of Experimental Biology* **215**, 454–460.
- VAN DOORN, A. (1989). Factors influencing dominance behaviour in queenless bumblebee workers (*Bombus terrestris*). *Physiological Entomology* **14**, 211–221.
- DORNHAUS, A. (2008). Specialization does not predict individual efficiency in an ant. *PLoS Biology* **6**, 2368–2375.
- DORNHAUS, A., HOLLEY, J. A., POOK, V. G., WORSWICK, G. & FRANKS, N. R. (2008). Why do not all workers work? Colony size and workload during emigrations in the ant *Temnothorax albipennis*. *Behavioral Ecology and Sociobiology* **63**, 43–51.
- DUCKWORTH, R. A. (2006). Behavioral correlations across breeding contexts provide a mechanism for a cost of aggression. *Behavioral Ecology* **17**, 1011–1019.

- DYER, A. G. & CHITTKA, L. (2004). Bumblebees (*Bombus terrestris*) sacrifice foraging speed to solve difficult colour discrimination tasks. *Journal of Comparative Physiology. A, Neuroethology Sensory Neural and Behavioral Physiology* **190**, 759–763.
- ELDAKAR, O. T., DLUGOS, M. J., WILCOX, R. S. & WILSON, D. S. (2009). Aggressive mating as a tragedy of the commons in the water strider *Aquarius remigis*. *Behavioral Ecology and Sociobiology* **64**, 25–33.
- FAIRBANKS, L. A., FONTENOT, M. B., PHILLIPS-CONROY, J. E., JOLLY, C. J., KAPLAN, J. R. & MANN, J. J. (1999). CSF monoamines, age and impulsivity in wild grivet monkeys (*Cercopithecus aethiops aethiops*). *Brain, Behavior and Evolution* **53**, 305–312.
- FEWELL, J. H. & PAGE, R. E. (1993). Genotypic variation in foraging responses to environmental stimuli by honey bees, *Apis mellifera*. *Experientia* **49**, 1106–1112.
- FEWELL, J. H. & PAGE, R. E. (1999). The emergence of division of labour in forced associations of normally solitary ant queens. *Evolutionary Ecology Research* **1**, 537–548.
- FOGARTY, S., COTE, J. & SIH, A. (2011). Social personality polymorphism and the spread of invasive species: a model. *American Naturalist* **177**, 273–287.
- FOLSE, H. J. III & ROUGHGARDEN, J. (2010). What is an individual organism? A multilevel selection perspective. *The Quarterly Review of Biology* **85**, 447–472.
- FOSTER, R. L., BRUNSKILL, A., VERDIRAME, D. & O'DONNELL, S. (2004). Reproductive physiology, dominance interactions, and division of labour among bumble bee workers. *Physiological Entomology* **29**, 327–334.
- GARAMSZEGI, L. Z. & HERCZEG, G. (2012). Behavioural syndromes, syndrome deviation and the within- and between-individual components of phenotypic correlations: when reality does not meet statistics. *Behavioral Ecology and Sociobiology* **66**, 1651–1658.
- GILL, R. J., RAMOS-RODRIGUEZ, O. & RAINE, N. E. (2012). Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature* **491**, 105–108.
- GIRAY, T., GIOVANETTI, M. & WEST-EBERHARD, M. J. (2005). Juvenile hormone, reproduction, and worker behavior in the neotropical social wasp *Polistes canadensis*. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 3330–3335.
- GORDON, D. M. (1989). Dynamics of task switching in harvester ants. *Animal Behaviour* **38**, 194–204.
- GORDON, D. M. (1991). Behavioral flexibility and the foraging ecology of seed-eating ants. *American Naturalist* **138**, 379–411.
- GORDON, D. M. (1996). The organization of work in social insect colonies. *Nature* **380**, 121–124.
- GORDON, D. M., CHU, J., LILLIE, A., TISSOT, M. & PINTER, N. (2005). Variation in the transition from inside to outside work in the red harvester ant *Pogonomyrmex barbatus*. *Insectes Sociaux* **52**, 212–217.
- GORDON, D. M., DEKAR, K. N. & PINTER-WOLLMAN, N. (2013). Harvester ant colony variation in foraging activity and response to humidity. *PLoS One* in press.
- GORDON, D. M., GUETZ, A., GREENE, M. J. & HOLMES, S. (2011). Colony variation in the collective regulation of foraging by harvester ants. *Behavioral Ecology* **22**, 429–435.
- GOSLING, S. D. (2001). From mice to men: what can we learn about personality from animal research? *Psychological Bulletin* **127**, 45–86.
- GRAFF, J., JEMIELITY, S., PARKER, J. D., PARKER, K. M. & KELLER, L. (2007). Differential gene expression between adult queens and workers in the ant *Lasius niger*. *Molecular Ecology* **16**, 675–683.
- GRAHAM, A., MUNDAY, M., KAFTANOGLU, O., PAGE, R., AMDAM, G. & RUEPPELL, O. (2011). Support for the reproductive ground plan hypothesis of social evolution and major QTL for ovary traits of Africanized worker honey bees (*Apis mellifera* L.). *BMC Evolutionary Biology* **11**, 95.
- GREENE, M. J. & GORDON, D. M. (2003). Social insects—Cuticular hydrocarbons inform task decisions. *Nature* **423**, 32.
- GREENE, M. J. & GORDON, D. M. (2007). Interaction rate informs harvester ant task decisions. *Behavioral Ecology* **18**, 451–455.
- GROSS, M. R. & CHARNOV, E. L. (1980). Alternative male life histories in bluegill sunfish. *Proceedings of the National Academy of Sciences of the United States of America: Biological Sciences* **77**, 6937–6940.
- GRÜTER, C., MENEZES, C., IMPERATRIZ-FONSECA, V. L. & RATNIEKS, F. L. W. (2012). A morphologically specialized soldier caste improves colony defense in a neotropical eusocial bee. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 1182–1186.
- HARTFELDER, K. (2000). Insect juvenile hormone: from “status quo” to high society. *Brazilian Journal of Medical and Biological Research* **33**, 157–177.
- HEIFETZ, Y. & APPLEBAUM, S. W. (1995). Density-dependent physiological phase in a non-migratory grasshopper *Aiolopus thalassinus*. *Entomologia Experimentalis et Applicata* **77**, 251–262.
- HERB, B. R., WOLSCHIN, F., HANSEN, K. D., ARYEE, M. J., LANGMEAD, B., IRIZARRY, R., AMDAM, G. V. & FEINBERG, A. P. (2012). Reversible switching between epigenetic states in honeybee behavioral subcastes. *Nature* **15**, 1371–1373.
- HERBERS, J. M. (1983). Social organization in *Leptothorax* ants: within and between species patterns. *Psyche* **90**, 361–386.
- HILLESHEIM, E., KOENIGER, N. & MORITZ, R. F. A. (1989). Colony performance in honeybees (*Apis mellifera capensis* Esch.) depends on the proportion of subordinate and dominant workers. *Behavioral Ecology and Sociobiology* **24**, 291–296.
- HÖLDOBLER, B. & WILSON, E. O. (1990). *The Ants*. Harvard University Press, Cambridge.
- HÖLDOBLER, B. & WILSON, E. O. (2009). *The Superorganism: The Beauty, Elegance, and Strangeness of Insect Societies*. W. W. Norton & Company, New York.
- HOLWAY, D. A. (1999). Competitive mechanisms underlying the displacement of native ants by the invasive Argentine ant. *Ecology* **80**, 238–251.
- HOLWAY, D. A. & SUAREZ, A. V. (1999). Animal behavior: an essential component of invasion biology. *Trends in Ecology & Evolution* **14**, 328–330.
- HOLWAY, D. A. & SUAREZ, A. V. (2004). Colony-structure variation and interspecific competitive ability in the invasive Argentine ant. *Oecologia* **138**, 216–222.
- HOWARD, K. J. (2006). Three queen morphs with alternative nest-founding behaviors in the ant, *Tennothorax longispinosus*. *Insectes Sociaux* **53**, 480–488.
- HUGHES, A. L. (1985). Male size, mating success, and mating strategy in the mosquitofish *Gambusia affinis* (Poeciliidae). *Behavioral Ecology and Sociobiology* **17**, 271–278.
- HUNT, J. H. & AMDAM, G. V. (2005). Bivoltinism as an antecedent to eusociality in the paper wasp genus *Polistes*. *Science* **308**, 264–267.
- HUNT, G. J., AMDAM, G. V., SCHLIPALUS, D., EMORE, C., SARDESAI, N., WILLIAMS, C. E., RUEPPELL, O., GUZMAN-NOVOA, E., ARECHAVELETA-VELASCO, M., CHANDRA, S., FONDRK, M. K., BEYE, M. & PAGE, R. E. (2007). Behavioral genomics of honeybee foraging and nest defense. *Naturwissenschaften* **94**, 247–267.
- HUNT, J. H., WOLSCHIN, F., HENSHAW, M. T., NEWMAN, T. C., TOTH, A. L. & AMDAM, G. V. (2010). Differential gene expression and protein abundance evince ontogenetic bias toward castes in a primitively eusocial wasp. *PLoS One* **5**, e10674.
- HUNTINGFORD, F. A. (1982). Do interspecific and intraspecific aggression vary in relation to predation pressure in sticklebacks. *Animal Behaviour* **30**, 909–916.
- HUNTINGFORD, F. A., ANDREW, G., MACKENZIE, S., MORERA, D., COYLE, S. M., PILARCZYK, M. & KADRI, S. (2010). Coping strategies in a strongly schooling fish, the common carp *Cyprinus carpio*. *Journal of Fish Biology* **76**, 1576–1591.
- INGRAM, K. K., OEFNER, P. & GORDON, D. M. (2005). Task-specific expression of the foraging gene in harvester ants. *Molecular Ecology* **14**, 813–818.
- INGS, T. C., RAINE, N. E. & CHITTKA, L. (2009). A population comparison of the strength and persistence of innate colour preference and learning speed in the bumblebee *Bombus terrestris*. *Behavioral Ecology and Sociobiology* **63**, 1207–1218.
- INGS, T. C., WARD, N. L. & CHITTKA, L. (2006). Can commercially imported bumble bees out-compete their native conspecifics? *Journal of Applied Ecology* **43**, 940–948.
- JANDT, J. M. & DORNHAUS, A. (2009). Spatial organization and division of labour in the bumblebee *Bombus impatiens*. *Animal Behaviour* **77**, 641–651.
- JANDT, J. M. & DORNHAUS, A. (2011). Competition and cooperation: bumblebee spatial organization and division of labor may affect worker reproduction late in life. *Behavioral Ecology and Sociobiology* **65**, 2341–2349.
- JANDT, J. M., HUANG, E. & DORNHAUS, A. (2009). Weak specialization of workers inside a bumble bee (*Bombus impatiens*) nest. *Behavioral Ecology and Sociobiology* **63**, 1829–1836.
- JEANNE, R. L. (1988). *Interindividual Behavioral Variability in Social Insects*. Westview Press, Boulder.
- JEANSON, R. (2012). Long-term dynamics in proximity networks in ants. *Animal Behaviour* **83**, 915–923.
- JEANSON, R. L. & FEWELL, J. H. (2008). Influence of the social context on division of labor in ant foundress associations. *Behavioral Ecology* **19**, 567–574.
- JOHNSON, J. C. & SIH, A. (2005). Precooperative sexual cannibalism in fishing spiders (*Dolomedes triton*): a role for behavioral syndromes. *Behavioral Ecology and Sociobiology* **58**, 390–396.
- JOHNSON, J. C. & SIH, A. (2007). Fear, food, sex and parental care: a syndrome of boldness in the fishing spider, *Dolomedes triton*. *Animal Behaviour* **74**, 1131–1138.
- JONES, T. C., AKOURY, T. S., HAUSER, C. K., NEBLETT, M. F., LINVILLE, B. J., EDGE, A. A. & WEBER, N. O. (2011). Octopamine and serotonin have opposite effects on antipredator behavior in the orb-weaving spider, *Larinioides cornutus*. *Journal of Comparative Physiology. A, Neuroethology Sensory Neural and Behavioral Physiology* **197**, 819–825.
- JONES, J. C., MYERSCOUGH, M. R., GRAHAM, S. & OLDROYD, B. P. (2004). Honey bee nest thermoregulation: diversity promotes stability. *Science* **305**, 402–404.
- KAMAKURA, M. (2011). Royalactin induces queen differentiation in honeybees. *Nature* **473**, 478–483.
- KELLER, L. (1999). *Levels of Selection in Evolution*. Princeton University Press, Princeton.
- KINNALLY, E. L., JENSEN, H. A., EWING, J. H. & FRENCH, J. A. (2006). Serotonin function is associated with behavioral response to a novel conspecific in marmosets. *American Journal of Primatology* **68**, 812–824.
- KOCHER, S. D., AYROLES, J. F., STONE, E. A. & GROZINGER, C. M. (2010). Individual variation in pheromone response correlates with reproductive traits and brain gene expression in worker honey bees. *PLoS One* **5**, e9116.
- KOOLHAAS, J. M., DE BOER, S. F., BUWALDA, B. & VAN REENEN, K. (2007). Individual variation in coping with stress: a multidimensional approach of ultimate and proximate mechanisms. *Brain, Behavior and Evolution* **70**, 218–226.
- KRALJ-FISER, S., WEISS, B. M. & KOTRSCHAL, K. (2010). Behavioural and physiological correlates of personality in greylag geese (*Anser anser*). *Journal of Ethology* **28**, 363–370.
- KRAUSE, J., LUSSEAU, D. & JAMES, R. (2009). Animal social networks: an introduction. *Behavioral Ecology and Sociobiology* **63**, 967–973.

- KULAHCI, I. G., DORNHAUS, A. & PAPAJ, D. R. (2008). Multimodal signals enhance decision making in foraging bumble bees. *Proceedings of the Royal Society B: Biological Sciences* **275**, 797–802.
- KURVERS, R., PRINS, H. H. T., VAN WIEREN, S. E., VAN OERS, K., NOLET, B. A. & YDENBERG, R. C. (2010). The effect of personality on social foraging: shy barnacle geese scrounge more. *Proceedings of the Royal Society B: Biological Sciences* **277**, 601–608.
- LAPIDGE, K. L., OLDROYD, B. P. & SPIVAK, M. (2002). Seven suggestive quantitative trait loci influence hygienic behavior of honey bees. *Naturwissenschaften* **89**, 565–568.
- LATSHAW, J. S. & SMITH, B. H. (2005). Heritable variation in learning performance affects foraging preferences in the honey bee (*Apis mellifera*). *Behavioral Ecology and Sociobiology* **58**, 200–207.
- LAVERY, T. M. (1994). Bumble bee learning and flower morphology. *Animal Behaviour* **47**, 531–545.
- LEADBEATER, E., CARRUTHERS, J. M., GREEN, J. P., VAN HEUSDEN, J. & FIELD, J. (2010). Unrelated helpers in a primitively eusocial wasp: is helping tailored towards direct fitness? *PLoS One* **5**, e11997.
- LIANG, Z. S., NGUYEN, T., MATTILA, H. R., RODRIGUEZ-ZAS, S. L., SEELEY, T. D. & ROBINSON, G. E. (2012). Molecular determinants of scouting behavior in honey bees. *Science* **335**, 1225–1228.
- LIHOREAU, M., BREPSON, L. & RIVAU, C. (2009). The weight of the clan: even in insects, social isolation can induce a behavioural syndrome. *Behavioural Processes* **82**, 81–84.
- LIHOREAU, M., CHITTKA, L., LE COMBER, S. C. & RAINE, N. E. (2012a). Bees do not use nearest-neighbour rules for optimization of multi-location routes. *Biology Letters* **8**, 13–16.
- LIHOREAU, M., RAINE, N. E., REYNOLDS, A. M., STELZER, R. J., LIM, K. S., SMITH, A. D., OSBORNE, J. L. & CHITTKA, L. (2012b). Radar tracking and motion-sensitive cameras on flowers reveal the development of pollinator multi-destination routes over large spatial scales. *PLoS Biology* **10**, e1001392.
- LIHOREAU, M., CHITTKA, L. & RAINE, N. E. (2010). Travel optimization by foraging bumblebees through readjustments of traplines after discovery of new feeding locations. *American Naturalist* **176**, 744–757.
- LIHOREAU, M., CHITTKA, L. & RAINE, N. E. (2011). Trade-off between travel distance and prioritization of high-reward sites in traplining bumblebees. *Functional Ecology* **25**, 1284–1292.
- LIN, H., WINSTON, M. L., HAUNDERLAND, N. H. & SLESSOR, K. N. (1999). Influence of age and population size on ovarian development, and of trophallaxis on ovarian development and vitellogenin titres of queenless worker honey bee (Hymenoptera: Apidae). *The Canadian Entomologist* **131**, 695–706.
- LINDAUER, M. (1952). Ein Beitrag zur Frage der arbeitsteilung im bienenstaat. *Zeitschrift für Vergleichende Physiologie* **34**, 299–345.
- LINKSVAYER, T. A. (2006). Direct, maternal, and sibsocial genetic effects on individual and colony traits in an ant. *Evolution* **60**, 2552–2561.
- LUTTBEG, B. & SIH, A. (2010). Risk, resources and state-dependent adaptive behavioural syndromes. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3977–3990.
- MATHOT, K. J., GODDE, S., CAREAU, V., THOMAS, D. W. & GIRALDEAU, L. A. (2009). Testing dynamic variance-sensitive foraging using individual differences in basal metabolic rates of zebra finches. *Oikos* **118**, 545–552.
- MATTILA, H. R., BURKE, K. M. & SEELEY, T. D. (2008). Genetic diversity within honeybee colonies increases signal production by waggle-dancing foragers. *Proceedings of the Royal Society B: Biological Sciences* **275**, 809–816.
- MATTILA, H. R. & SEELEY, T. D. (2007). Genetic diversity in honey bee colonies enhances productivity and fitness. *Science* **317**, 362–364.
- MICHELENA, P., JEANSON, R., DENEUBOURG, J. L. & SIBBALD, A. M. (2010). Personality and collective decision-making in foraging herbivores. *Proceedings of the Royal Society B: Biological Sciences* **277**, 1093–1099.
- MICHENER, C. (1964). Reproductive efficiency in relation to colony size in Hymenopterous societies. *Insectes Sociaux* **11**, 317–341.
- MODLMEIER, A. P. & FOITZIK, S. (2011). Productivity increases with variation in aggression among group members in *Temnothorax* ants. *Behavioral Ecology* **22**, 1026–1032.
- MODLMEIER, A. P., LIEBMANN, J. E. & FOITZIK, S. (2012). Diverse societies are more productive: a lesson from ants. *Proceedings of the Royal Society B: Biological Sciences* **279**, 2142–2150.
- MOLET, M., CHITTKA, L. & RAINE, N. E. (2009). How floral odours are learned inside the bumblebee (*Bombus terrestris*) nest. *Naturwissenschaften* **96**, 213–219.
- MOLET, M., CHITTKA, L., STELZER, R. J., STREIT, S. & RAINE, N. E. (2008). Colony nutritional status modulates worker responses to foraging recruitment pheromone in the bumblebee *Bombus terrestris*. *Behavioral Ecology and Sociobiology* **62**, 1919–1926.
- MOLINA, Y. & O'DONNELL, S. (2009). Worker reproductive competition affects division of labor in a primitively social paperwasp (*Polistes instabilis*). *Insectes Sociaux* **56**, 14–20.
- MOORE, A. J., BREED, M. D. & MOOR, M. J. (1987). The guard honey bee: ontogeny and behavioural variability of workers performing a specialized task. *Animal Behaviour* **35**, 1159–1167.
- MULLER, H., GROSSMANN, H. & CHITTKA, L. (2010). 'Personality' in bumblebees: individual consistency in responses to novel colours? *Animal Behaviour* **80**, 1065–1074.
- NIEMELÄ, P., VAINIKKA, A., LAHDENPERÄ, S. & KORTET, R. (2012). Nymphal density, behavioral development, and life history in a field cricket. *Behavioral Ecology and Sociobiology* **66**, 645–652.
- NONACS, P. & KAPHEIM, K. M. (2007). Social heterosis and the maintenance of genetic diversity. *Journal of Evolutionary Biology* **20**, 2253–2265.
- NONACS, P. & KAPHEIM, K. M. (2008). Social heterosis and the maintenance of genetic diversity at the genome level. *Journal of Evolutionary Biology* **21**, 631–635.
- O'DONNELL, S. (2006). *Polybia* wasp biting interactions recruit foragers following experimental worker removals. *Animal Behaviour* **71**, 709–715.
- O'DONNELL, S. & BULOVA, S. J. (2007a). Worker connectivity: a review of the design of worker communication systems and their effects on task performance in insect societies. *Insectes Sociaux* **54**, 203–210.
- O'DONNELL, S. & BULOVA, S. J. (2007b). Worker connectivity: a simulation model of variation in worker communication and its effects on task performance. *Insectes Sociaux* **54**, 211–218.
- VAN OERS, K., DE JONG, G., VAN NOORDWIJK, A. J., KEMPENAEERS, B. & DRENT, P. J. (2005). Contribution of genetics to the study of animal personalities: a review of case studies. *Behaviour* **142**, 1185–1206.
- VAN OERS, K. & MUELLER, J. C. (2010). Evolutionary genomics of animal personality. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3991–4000.
- OLDROYD, B. P. & FEWELL, J. H. (2007). Genetic diversity promotes homeostasis in insect colonies. *Trends in Ecology & Evolution* **22**, 408–413.
- O'NEAL, R. J. & WALLER, G. D. (1984). On the pollen harvest by the honey bee (*Apis mellifera* L.) near Tucson, Arizona (1976–1981). *Desert Plants* **6**, 81–109.
- OSTER, G. F. & WILSON, E. O. (1978). *Caste and Ecology in the Social Insects*. Princeton University Press, Princeton.
- PAGE, R. E. J., ERBER, J. & FONDRK, M. K. (1998). The effect of genotype on response thresholds to sucrose and foraging behavior of honey bees (*Apis mellifera* L.). *Journal of Comparative Physiology* **A182**, 489–500.
- PAGE, R. E., FONDRK, M. K. & RUEPPELL, O. (2012a). Complex pleiotropy characterizes the pollen hoarding syndrome in honey bees (*Apis mellifera* L.). *Behavioral Ecology and Sociobiology* **66**, 1459–1466.
- PAGE, R. E., RUEPPELL, O. & AMDAM, G. V. (2012b). Genetics of reproduction and regulation of honeybee (*Apis mellifera* L.) social behavior. *Annual Review of Genetics* **46**, 97–119.
- PAGE, R. E., ROBINSON, G. E., BRITTON, D. S. & FONDRK, M. K. (1992). Genotypic variability for rates of behavioral development in worker honeybees (*Apis mellifera* L.). *Behavioral Ecology* **3**, 173–180.
- PAGE, R. E., ROBINSON, G. E., FONDRK, M. K. & NASR, M. E. (1995a). Effects of worker genotypic diversity on honey bee colony development and behavior (*Apis mellifera* L.). *Behavioral Ecology and Sociobiology* **36**, 387–396.
- PAGE, R. E., WADDINGTON, K. D., HUNT, G. J. & FONDRK, M. K. (1995b). Genetic determinants of honey bee foraging behaviour. *Animal Behaviour* **50**, 1617–1625.
- PALEOLOG, J. (2009). Behavioural characteristics of honey bee (*Apis mellifera*) colonies containing mix of workers of divergent behavioural traits. *Animal Science Papers and Reports* **27**, 237–248.
- PANKIWI, T. (2003). Directional change in a suite of foraging behaviors in tropical and temperate evolved honey bees (*Apis mellifera* L.). *Behavioral Ecology and Sociobiology* **54**, 458–464.
- PANKIWI, T., TARPY, D. R. & PAGE, R. E. (2002). Genotype and rearing environment affect honeybee perception and foraging behaviour. *Animal Behaviour* **64**, 663–672.
- PEARCE, A. N., HUANG, Z. Y. & BREED, M. D. (2001). Juvenile hormone and aggression in honey bees. *Journal of Insect Physiology* **47**, 1243–1247.
- PENKE, L., DENNIS, J. J. A. & MILLER, G. F. (2007). The evolutionary genetics of personality. *European Journal of Personality* **21**, 549–587.
- PEREBOOM, J. J. M., VELTHUIS, H. H. W. & DUCHATEAU, M. J. (2003). The organisation of larval feeding in bumblebees (Hymenoptera, Apidae) and its significance to caste differentiation. *Insectes Sociaux* **50**, 127–133.
- PINTER-WOLLMAN, N. (2012). Personality in social insects: how does worker personality determine colony personality? *Current Zoology* **58**, 580–588.
- PINTER-WOLLMAN, N., GORDON, D. M. & HOLMES, S. (2012a). Nest site and weather affect the personality of harvester ant colonies. *Behavioral Ecology* **23**, 1022–1029.
- PINTER-WOLLMAN, N., HUBLER, J., HOLLEY, J.-A., FRANKS, N. & DORNHAUS, A. (2012b). How is activity distributed among and within tasks in *Temnothorax* ants? *Behavioral Ecology and Sociobiology* **66**, 1407–1420.
- PINTER-WOLLMAN, N., WOLLMAN, R., GUETZ, A., HOLMES, S. & GORDON, D. M. (2011). The effect of individual variation on the structure and function of interaction networks in harvester ants. *Journal of the Royal Society, Interface* **8**, 1562–1573.
- PONTIER, D., FROMONT, E., COURCHAMP, F., ARTOIS, M. & YOCOZ, N. G. (1998). Retroviruses and sexual size dimorphism in domestic cats (*Felis catus* L.). *Proceedings of the Royal Society of London. Series B: Biological Sciences* **265**, 167–173.
- PORTER, S. D. & JORGENSEN, C. D. (1981). Foragers of the harvester ant, *Pogonomyrmex ovyei*: a disposable caste? *Behavioral Ecology and Sociobiology* **9**, 247–256.
- PORTER, S. D. & TSCHINKEL, W. R. (1985). Fire ant polymorphism: the ergonomics of brood production. *Behavioral Ecology and Sociobiology* **16**, 323–336.
- POWELL, S. (2008). Ecological specialization and the evolution of a specialized caste in *Cephalotes* ants. *Functional Ecology* **22**, 902–911.

- POWELL, S. & FRANKS, N. R. (2005). Caste evolution and ecology: a special worker for novel prey. *Proceedings of the Royal Society B: Biological Sciences* **272**, 2173–2180.
- POWELL, S. & FRANKS, N. R. (2006). Ecology and the evolution of worker morphological diversity: a comparative analysis with *Eciton* army ants. *Functional Ecology* **20**, 1105–1114.
- POWELL, S. & TSCHINKEL, W. R. (1999). Ritualized conflict in *Odontomachus brunneus* and the generation of interaction-based task allocation: a new organizational mechanism in ants. *Animal Behaviour* **58**, 965–972.
- PRONK, R., WILSON, D. R. & HARCOURT, R. (2010). Video playback demonstrates episodic personality in the gloomy octopus. *Journal of Experimental Biology* **213**, 1035–1041.
- PRUITT, J. N. & FERRARI, M. C. O. (2011). Intraspecific trait variants determine the nature of interspecific interactions in a habitat-forming species. *Ecology* **92**, 1902–1908.
- PRUITT, J. N., ITURRALDE, G., AVILES, L. & RIECHERT, S. E. (2011). Amazonian social spiders share similar within-colony behavioural variation and behavioural syndromes. *Animal Behaviour* **82**, 1449–1455.
- PRUITT, J. N., OUFIERO, C. E., AVILES, L. & RIECHERT, S. E. (2012). Iterative evolution of increased behavioral variation characterizes the transition to sociality in spiders and proves advantageous. *The American Naturalist* **180**, 496–510.
- PRUITT, J. N. & RIECHERT, S. E. (2009). Frequency-dependent success of cheaters during foraging bouts might limit their spread within colonies of a socially polymorphic spider. *Evolution* **63**, 2966–2973.
- PRUITT, J. N. & RIECHERT, S. E. (2011). How within-group behavioural variation and task efficiency enhance fitness in a social group. *Proceedings of the Royal Society B: Biological Sciences* **278**, 1209–1215.
- QUELLER, D. C., ZACCHI, F., CERVO, R., TURILLAZZI, S., HENSHAW, M. T., SANTORELLI, L. A. & STRASSMANN, J. E. (2000). Unrelated helpers in a social insect. *Nature* **405**, 784–787.
- RAINE, N. E. & CHITTKA, L. (2005). Colour preference in relation to the foraging performance and fitness of the bumblebee *Bombus terrestris*. *Uludag Bee Journal* **5**, 145–150.
- RAINE, N. E. & CHITTKA, L. (2007a). Pollen foraging: learning a complex motor skill by bumblebees (*Bombus terrestris*). *Naturwissenschaften* **94**, 459–464.
- RAINE, N. E. & CHITTKA, L. (2007b). The adaptive significance of sensory bias in a foraging context: floral colour preferences in the bumblebee *Bombus terrestris*. *PLoS One* **2**, e556.
- RAINE, N. E. & CHITTKA, L. (2008). The correlation of learning speed and natural foraging success in bumble-bees. *Proceedings of the Royal Society B: Biological Sciences* **275**, 803–808.
- RAINE, N. E. & CHITTKA, L. (2012). No trade-off between learning speed and associative flexibility in bumblebees: a reversal learning test with multiple colonies. *PLoS One* **7**, e45096.
- RAINE, N. E., INGS, T. C., DORNHAUS, A., SALEH, N. & CHITTKA, L. (2006a). Adaptation, genetic drift, pleiotropy, and history in the evolution of bee foraging behavior. *Advances in the Study of Behavior* **36**, 305–354.
- RAINE, N. E., INGS, T. C., RAMOS-RODRIGUEZ, O. & CHITTKA, L. (2006b). Intercolony variation in learning performance of a wild British bumblebee population (Hymenoptera: Apidae: *Bombus terrestris* audax). *Entomologia Generalis* **28**, 241–256.
- RATNIEKS, F. L. W. & REEVE, H. K. (1992). Conflict in single-queen Hymenopteran societies: the structure of conflict and processes that reduce conflict in advanced eusocial species. *Journal of Theoretical Biology* **158**, 33–65.
- RÉALE, D., READER, S. M., SOL, D., McDOUGALL, P. T. & DINGEMANSE, N. J. (2007). Integrating animal temperament within ecology and evolution. *Biological Reviews* **82**, 291–318.
- REEVE, H. K. (1992). Queen activation of lazy workers in colonies of the eusocial naked mole-rat. *Nature* **358**, 147–149.
- REEVE, H. K. & NONACS, P. (1997). Within-group aggression and the value of group members: theory and a field test with social wasps. *Behavioral Ecology* **8**, 75–82.
- RIECHERT, S. E. & HEDRICK, A. V. (1993). A test for correlations among fitness-linked behavioral traits in the spider *Agelenopsis aperta* (Araneae, Agelenidae). *Animal Behaviour* **46**, 669–675.
- ROBINSON, G. E. (1987). Regulation of honey bee age polyethism by juvenile hormone. *Behavioral Ecology and Sociobiology* **20**, 329–338.
- ROBINSON, G. E. (1992). Regulation of division of labor in insect societies. *Annual Review of Entomology* **37**, 637–665.
- ROBINSON, E. J. H. (2009). Physiology as a caste-defining feature. *Insectes Sociaux* **56**, 1–6.
- ROBINSON, G. E. & HUANG, Z. Y. (1998). Colony integration in honey bees: genetic, endocrine and social control of division of labor. *Apidologie* **29**, 159–170.
- ROBINSON, G. E. & PAGE, R. E. (1988). Genetic determination of guarding and undertaking in honeybee colonies. *Nature* **333**, 356–358.
- ROBINSON, G. E. & PAGE, R. E. (1989). Genetic determination of nectar foraging, pollen foraging, and nest-site scouting in honey bee colonies. *Behavioral Ecology and Sociobiology* **24**, 317–323.
- ROBINSON, G. E., PAGE, R. E. & ARENSEN, N. (1994). Genotypic differences in brood rearing in honey bee colonies: context-specific? *Behavioral Ecology and Sociobiology* **34**, 125–137.
- ROBINSON, G. E., PAGE, R. E. J. & FONDRE, M. K. (1990). Intracolony behavioral variation in worker oviposition, oophagy, and larval care in queenless honey bee colonies. *Behavioral Ecology and Sociobiology* **26**, 315–323.
- ROBINSON, E., RICHARDSON, T., SENDOVA-FRANKS, A., FEINERMAN, O. & FRANKS, N. (2009). Radio tagging reveals the roles of corpulence, experience and social information in ant decision making. *Behavioral Ecology and Sociobiology* **63**, 627–636.
- ROBSON, S. K. A., BEAN, K., HANSEN, J., NORLING, K., ROWE, R. J. & WHITE, D. (2000). Social and spatial organisation in colonies of a primitively eusocial wasp, *Ropalidia revolutionalis* (de Saussure) (Hymenoptera: Vespidae). *Australian Journal of Entomology* **39**, 20–24.
- ROBSON, S. K. & TRANIELLO, J. F. A. (1999). Key individuals and the organisation of labor in ants. In *Information Processing in Social Insects* (eds C. DETRAIN and J. L. DENEUBOURG), pp. 239–260. Birkhäuser Verlag, Basel.
- ROUX, E. A. & KORB, J. (2004). Evolution of eusociality and the soldier caste in termites: a validation of the intrinsic benefit hypothesis. *Journal of Evolutionary Biology* **17**, 869–875.
- ROWLES, A. D. & O'DOWD, D. J. (2007). Interference competition by Argentine ants displaces native ants: implications for biotic resistance to invasion. *Biological Invasions* **9**, 73–85.
- RUEPPELL, O., PANKIW, T., NIELSEN, D. I., FONDRE, M. K., BEYE, M. & PAGE, R. E. (2004). The genetic architecture of the behavioral ontogeny of foraging in honeybee workers. *Genetics* **167**, 1767–1779.
- SALEH, N. & CHITTKA, L. (2007). Traplining in bumblebees (*Bombus impatiens*): a foraging strategy's ontogeny and the importance of spatial reference memory in short-range foraging. *Oecologia* **151**, 719–730.
- SCHARF, I., MODLMEIER, A. P., FRIES, S., TIRARD, C. & FOITZIK, S. (2012). Characterizing the collective personality of ant societies: aggressive colonies do not abandon their home. *PLoS One* **7**, e33314.
- SCHNEIDER, R. (2012). Birth weight and sucrose responsiveness predict cognitive skills of honeybee foragers. *Animal Behaviour* **84**, 305–308.
- SCHNEIDER, R., BARNERT, M. & ERBER, J. (2003). Variation in water and sucrose responsiveness during the foraging season affects proboscis extension learning in honey bees. *Apidologie* **34**, 67–72.
- SCHNEIDER, R., ERBER, J. & PAGE, R. E. (1999). Tactile learning and the individual evaluation of the reward in honey bees (*Apis mellifera* L.). *Journal of Comparative Physiology. A, Sensory Neural and Behavioral Physiology* **185**, 1–10.
- SCHNEIDER, R., PAGE, R. E. & ERBER, J. (2001). The effects of genotype, foraging role, and sucrose responsiveness on the tactile learning performance of honey bees (*Apis mellifera* L.). *Neurobiology of Learning and Memory* **76**, 138–150.
- SCHNEIDER, R., PAGE, R. E. & ERBER, J. (2004). Sucrose responsiveness and behavioral plasticity in honey bees (*Apis mellifera*). *Apidologie* **35**, 133–142.
- SCHMID-HEMPPEL, P. (1990). Reproductive competition and the evolution of work load in social insects. *American Naturalist* **135**, 501–526.
- SCHOFIELD, R., EMMETT, K., NIEDEBALA, J. & NESSON, M. (2011). Leaf-cutter ants with worn mandibles cut half as fast, spend twice the energy, and tend to carry instead of cut. *Behavioral Ecology and Sociobiology* **65**, 969–982.
- SCHÜRCH, R., ROTHENBERGER, S. & HEG, D. (2011). The building-up of social relationships: behavioural types, social networks and cooperative breeding in a cichlid. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 4089–4098.
- SCHWANDER, T., CAHAN, S. H. & KELLER, L. (2007). Characterization and distribution of *Pogonomyrmex* harvester ant lineages with genetic caste determination. *Molecular Ecology* **16**, 367–387.
- SCHWANDER, T., LO, N., BEEKMAN, M., OLDROYD, B. P. & KELLER, L. (2010). Nature versus nurture in social insect caste differentiation. *Trends in Ecology & Evolution* **25**, 275–282.
- SEELEY, T. D. (1982). Adaptive significance of the age polyethism schedule in honeybee colonies. *Behavioral Ecology and Sociobiology* **11**, 287–293.
- SEELEY, T. D., WEIDENMÜLLER, A. & KÜHNHOLZ, S. (1998). The shaking signal of the honey bee informs workers to prepare for greater activity. *Ethology* **104**, 10–26.
- SEID, M. A. & TRANIELLO, J. F. A. (2006). Age-related repertoire expansion and division of labor in *Pheidole dentata* (Hymenoptera: Formicidae): a new perspective on temporal polyethism and behavioral plasticity in ants. *Behavioral Ecology and Sociobiology* **60**, 631–644.
- SENDOVA-FRANKS, A. B. & FRANKS, N. R. (1995). Spatial relationships within nests of the ant *Leptothorax unifasciatus* (Latr.) and their implications for the division of labor. *Animal Behaviour* **50**, 121–136.
- SHORT, K. H. & PETREN, K. (2008). Boldness underlies foraging success of invasive *Lepidodactylus lugubris* geckos in the human landscape. *Animal Behaviour* **76**, 429–437.
- SIH, A. (1992). Prey uncertainty and the balancing of antipredator and feeding needs. *American Naturalist* **139**, 1052–1069.
- SIH, A. (2011). Effects of early stress on behavioral syndromes: an integrated adaptive perspective. *Neuroscience and Biobehavioral Reviews* **35**, 1452–1465.
- SIH, A. & BELL, A. M. (2008). Insights for behavioral ecology from behavioral syndromes. *Advances in the Study of Behavior* (Volume **38**), pp. 227–281. Elsevier Academic Press Inc., San Diego.
- SIH, A., BELL, A. & JOHNSON, J. C. (2004a). Behavioral syndromes: an ecological and evolutionary overview. *Trends in Ecology & Evolution* **19**, 372–378.

- SIH, A., BELL, A., JOHNSON, J. C. & ZIEMBA, R. E. (2004b). Behavioral syndromes: an integrative overview. *The Quarterly Review of Biology* **79**, 241–277.
- SIH, A., COTE, J., EVANS, M., FOGARTY, S. & PRUITT, J. (2012). Ecological implications of behavioural syndromes. *Ecology Letters* **15**, 278–289.
- SIH, A. & DEL GUDICE, M. (2012). Linking behavioural syndromes and cognition: a behavioural ecology perspective. *Philosophical Transactions of the Royal Society B: Biological Sciences* **367**, 2762–2772.
- SIH, A., FERRARI, M. C. O. & HARRIS, D. J. (2011). Evolution and behavioural responses to human-induced rapid environmental change. *Evolutionary Applications* **4**, 367–387.
- SIH, A., HANSER, S. F. & MCHUGH, K. A. (2009). Social network theory: new insights and issues for behavioral ecologists. *Behavioral Ecology and Sociobiology* **63**, 975–988.
- SIH, A., KATS, L. B. & MAURER, E. F. (2003). Behavioural correlations across situations and the evolution of antipredator behaviour in a sunfish-salamander system. *Animal Behaviour* **65**, 29–44.
- SIH, A. & WATTERS, J. V. (2005). The mix matters: behavioural types and group dynamics in water striders. *Behaviour* **142**, 1423–1437.
- SIMONE-FINSTROM, M., GARDNER, J. & SPIVAK, M. (2010). Tactile learning in resin foraging honeybees. *Behavioral Ecology and Sociobiology* **64**, 1609–1617.
- SMITH, B. R. & BLUMSTEIN, D. T. (2008). Fitness consequences of personality: a meta-analysis. *Behavioral Ecology* **19**, 448–455.
- SMITH, C. R., TOTH, A. L., SUAREZ, A. V. & ROBINSON, G. E. (2008). Genetic and genomic analyses of the division of labour in insect societies. *Nature Reviews Genetics* **9**, 735–748.
- SPAETHE, J., BROCKMANN, A., HALBIG, C. & TAUTZ, J. (2007). Size determines antennal sensitivity and behavioral threshold to odors in bumblebee workers. *Naturwissenschaften* **94**, 733–739.
- SPAETHE, J. & CHITTKA, L. (2003). Interindividual variation of eye optics and single object resolution in bumblebees. *Journal of Experimental Biology* **206**, 3447–3453.
- SPAETHE, J. & WEIDENMÜLLER, A. (2002). Size variation and foraging rate in bumblebees (*Bombus terrestris*). *Insectes Sociaux* **49**, 142–146.
- SPIVAK, M., MASTERMAN, R., ROSS, R. & MESCE, K. A. (2003). Hygienic behavior in the honey bee (*Apis mellifera* L.) and the modulatory role of octopamine. *Journal of Neurobiology* **55**, 341–354.
- STAMPS, J. A. & GROOTHUIS, T. G. G. (2010). Developmental perspectives on personality: implications for ecological and evolutionary studies of individual differences. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 4029–4041.
- STIRLING, D. G., RÉALE, D. & ROFF, D. A. (2002). Selection, structure and the heritability of behaviour. *Journal of Evolutionary Biology* **15**, 277–289.
- STUART, R. J. & PAGE, R. E. (1991). Genetic component to division of labor among workers of a Leptothoracine ant. *Naturwissenschaften* **78**, 375–377.
- SUAREZ, A. V., HOLWAY, D. A., LIANG, D. S., TSUTSUI, N. D. & CASE, T. J. (2002). Spatiotemporal patterns of intraspecific aggression in the invasive Argentine ant. *Animal Behaviour* **64**, 697–708.
- SUAREZ, A. V., TSUTSUI, N. D., HOLWAY, D. A. & CASE, T. J. (1999). Behavioral and genetic differentiation between native and introduced populations of the Argentine ant. *Biological Invasions* **1**, 43–53.
- SUMPTER, D. J. T. (2010). *Collective Animal Behavior*. Princeton University Press, Princeton.
- SURYANARAYANAN, S., HANTSCH, A. E., TORRES, C. G. & JEANNE, R. L. (2011a). Changes in the temporal pattern of antennal drumming behavior across the *Polistes fuscatus* colony cycle (Hymenoptera, Vespidae). *Insectes Sociaux* **58**, 97–106.
- SURYANARAYANAN, S., HERMANSON, J. C. & JEANNE, R. L. (2011b). A mechanical signal biases caste development in a social wasp. *Current Biology* **21**, 231–235.
- SURYANARAYANAN, S. & JEANNE, R. L. (2008). Antennal drumming, trophallaxis, and colony development in the social wasp *Polistes fuscatus* (Hymenoptera: Vespidae). *Ethology* **114**, 1201–1209.
- THERAULAZ, G. & BONABEAU, E. (1995). Modelling the collective building of complex architectures in social insects with lattice swarms. *Journal of Theoretical Biology* **177**, 381–400.
- THERAULAZ, G., BONABEAU, E. & DENEUBOURG, J.-L. (1998). Response threshold reinforcement and division of labour in insect societies. *Proceedings of the Royal Society B* **265**, 327–332.
- TIBBETTS, E. A. & HUANG, Z. Y. (2010). The challenge hypothesis in an insect: juvenile hormone increases during reproductive conflict following queen loss in *Polistes* wasps. *American Naturalist* **176**, 123–130.
- TOTH, A. L., BILOF, K. B. J., HENSHAW, M. T., HUNT, J. H. & ROBINSON, G. E. (2009). Lipid stores, ovary development, and brain gene expression in *Polistes metricus* females. *Insectes Sociaux* **56**, 77–84.
- TOTH, A. L. & ROBINSON, G. E. (2005). Worker nutrition and division of labour in honeybees. *Animal Behaviour* **69**, 427–435.
- TSCHINKEL, W. (1998). The reproductive biology of fire ant societies. *BioScience* **48**, 593–605.
- TSCHINKEL, W. R. (2004). The nest architecture of the Florida harvester ant, *Pogonomyrmex badius*. *Journal of Insect Science* **4**, 1–19.
- TSUJI, K. (1994). Inter-colonial selection for the maintenance of cooperative breeding in the ant *Pristomyrmex pungens*: a laboratory experiment. *Behavioral Ecology and Sociobiology* **35**, 109–113.
- VISSCHER, P. K. (1988). Undertaker specialists in honey bee colonies. In *Interindividual Behavioral Variability in Social Insects* (ed. R. L. JEANNE), pp. 359–383. Westview Press, Boulder.
- VOLNY, V. P. & GORDON, D. M. (2002). Genetic basis for queen-worker dimorphism in a social insect. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 6108–6111.
- VOS, M. & VELICER, G. J. (2006). Genetic population structure of the soil bacterium *Myxococcus xanthus* at the centimeter scale. *Applied and Environmental Microbiology* **72**, 3615–3625.
- WADDINGTON, K. D., NELSON, C. M. & PAGE, R. E. (1998). Effects of pollen quality and genotype on the dance of foraging honey bees. *Animal Behaviour* **56**, 35–39.
- WANG, Y., AMDAM, G. V., RUEPPEL, O., WALLRICH, M. A., FONDRIK, M. K., KAFTANOGLU, O. & PAGE, R. E. Jr. (2009). *PDK1* and *HR46* gene homologs tie social behavior to ovary signals. *PLoS One* **4**, e4899.
- WEIDENMÜLLER, A. (2004). The control of nest climate in bumblebee (*Bombus terrestris*) colonies: interindividual variability and self reinforcement in fanning response. *Behavioral Ecology* **15**, 120–128.
- WEINER, S. A. & TOTH, A. L. (2012). Epigenetics in social insects: a new direction for understanding the evolution of castes. *Genetics Research International* **2012**, 1–11.
- WENSELEERS, T. & RATNIEKS, F. L. W. (2006a). Comparative analysis of worker reproduction and policing in eusocial hymenoptera supports relatedness theory. *American Naturalist* **168**, E163–E179.
- WENSELEERS, T. & RATNIEKS, F. L. W. (2006b). Enforced altruism in insect societies. *Nature* **444**, 50.
- WHEELER, D. E. (1986). Developmental and physiological determinants of caste in social Hymenoptera: evolutionary implications. *American Naturalist* **128**, 13–34.
- WHEELER, D. E., BUCK, N. & EVANS, J. D. (2006). Expression of insulin pathway genes during the period of caste determination in the honey bee, *Apis mellifera*. *Insect Molecular Biology* **15**, 597–602.
- WILSON, E. O. (1976). Behavioral discretization and number of castes in an ant species. *Behavioral Ecology and Sociobiology* **1**, 141–154.
- WILSON, E. O. (1985). Between-caste aversion as a basis for division of labor in the ant *Pheidole pubiventris* (Hymenoptera: Formicidae). *Behavioral Ecology and Sociobiology* **17**, 35–37.
- WILSON, D. S., CLARK, A. B., COLEMAN, K. & DEARSTYNE, T. (1994). Shyness and boldness in humans and other animals. *Trends in Ecology & Evolution* **9**, 442–446.
- WINSTON, M. (1987). *The Biology of the Honey Bee*. Harvard University Press, Cambridge.
- WOLF, M. & WEISSING, F. J. (2010). An explanatory framework for adaptive personality differences. *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 3959–3968.
- WOODING, S., BUFE, B., GRASSI, C., HOWARD, M. T., STONE, A. C., VAZQUEZ, M., DUNN, D. M., MEYERHOF, W., WEISS, R. B. & BAMSHAD, M. J. (2006). Independent evolution of bitter-taste sensitivity in humans and chimpanzees. *Nature* **440**, 930–934.
- WORDEN, B. D., SKEMP, A. K. & PAPA, D. R. (2005). Learning in two contexts: the effects of interference and body size in bumblebees. *Journal of Experimental Biology* **208**, 2045–2053.
- WRAY, M. K., MATTILA, H. R. & SEELEY, T. D. (2011). Collective personalities in honeybee colonies are linked to colony fitness. *Animal Behaviour* **81**, 559–568.
- WRAY, M. K. & SEELEY, T. D. (2011). Consistent personality differences in house-hunting behavior but not decision speed in swarms of honey bees (*Apis mellifera*). *Behavioral Ecology and Sociobiology* **65**, 2061–2070.
- ZANETTE, L. R. S. & FIELD, J. (2011). Founders versus joiners: group formation in the paper wasp *Polistes dominulus*. *Animal Behaviour* **82**, 699–705.

(Received 28 April 2012; revised 9 April 2013; accepted 17 April 2013)