

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/269107469>

Assessing and influencing personality for improvement of animal welfare: A review of equine studies

Article in CAB Reviews Perspectives in Agriculture Veterinary Science Nutrition and Natural Resources · February 2013

DOI: 10.1079/PAVSNNR20138006

CITATIONS

18

READS

1,370

1 author:



Uta König von Borstel

G.A. Universität Göttingen

71 PUBLICATIONS **734** CITATIONS

SEE PROFILE

Review

Assessing and influencing personality for improvement of animal welfare: a review of equine studies

Uta König v. Borstel*

Address: Department of Animal Science, Georg-August Universität Göttingen, Albrecht-Thaer-Weg 3, 37075, Göttingen, Germany.

***Correspondence:** Email: koenigvb@gwdg.de

Received: 18 September 2012

Accepted: 20 November 2012

doi: 10.1079/PAVSNNR20138006

The electronic version of this article is the definitive one. It is located here: <http://www.cabi.org/cabreviews>

© CAB International 2013 (Online ISSN 1749-8848)

Abstract

Personality and its various sub-traits such as temperament can be defined as behavioural attitudes that remain relatively stable across time and situations. However, various personality models exist. Contrasting their strengths and weaknesses leads to the conclusion that there is presently no model that is truly superior to others in depicting animal personality. Furthermore, the present review highlights aspects in which animal, and in particular equine personality potentially relate to animal welfare. Three approaches are examined which may be taken to improve welfare: (1) genetically selecting for personality traits that make the horses better adapted to their designated work and/or husbandry conditions, (2) assessing personality in individual horses to optimize matching with owners and type of work and (3) influencing the ontogeny of personality traits, e.g. by specific training and husbandry regimes. Each of these strategies has its merit, but as a prerequisite, valid personality assessment methods are required. Although publication bias is likely present, most testing procedures yield acceptable inter-observer reliabilities and repeatabilities across time, even if repeatability across situations tends to be comparably low. Heritability estimates ranging mostly between $h^2=0.15$ and $h^2=0.40$ for traits assessed in personality tests are likewise promising. Research has paid less attention to intra-observer reliability, construct validity or discriminant ability, but based on pleiotropy it is argued that the latter is not essential for a trait to be valid. Therefore, some valid methods are available for use in assessment and subsequent selection of horses or personality-influencing strategies, ultimately resulting in reduction of stress in horses' everyday life.

Keywords: Personality, Welfare, Genetics of behaviour, Horse, Temperament, Human–animal interaction, Character

Review Methodology: The search machines and databases CAB Abstracts, Medline/PubMed, Agricola and Google Scholar were used for the literature search, using the following keyword search terms in different combinations: horse, equine, personality, temperament, character, behaviour, genetic, heritability, fear, reactivity, test, welfare, human–animal interaction, match, ontogeny. Furthermore, I used my personal literature database, which includes, in addition to English articles, articles written in several European languages. Once suitable articles had been identified, their references were used as well as the possibility to search within publishers' databases for related articles and further articles relating to the above keywords. Finally, I spoke to colleagues working in the area of equine personality to check for missed as well as upcoming studies.

Animal Personality

Methods to categorize human personality into distinct personality types has been documented as early as by Hippocrates (ca. 460–370 BC), although it was not until 2000 years later that research into animal personalities

has gained some acceptance. However, individual differences in personality are thought to have adaptive value in animals [1–3], and today there is little doubt that distinct animal personalities exist. While the first research in this area focused on species acknowledged to be capable of higher cognitive processes, such as primates [4], dogs

[5, 6] and later other mammals [7], present research indicates the existence of personality-like, individual-specific behavioural attitudes in other taxa and phyla such as birds [8], amphibians [9], fish [10, 11] and even invertebrates such as arthropods [12], molluscs, [13] and insects such as firebugs [14] and honey bees [15]. Gosling [16] undertook a comprehensive review of personality studies across various species. Given the relevance to human safety, economics, management and animal welfare, much interest has been devoted to the study of personality in livestock [17–19] and companion animals [6, 20–22]. The horse falls into both categories [23] and thus has received particular attention.

Definitions of Personality and Classification Schemes

For the purpose of the present review, personality is defined as the animal's basic stance towards environmental stimuli that is expressed by behaviour patterns and remains relatively stable across time and situations ([24, 25] based on definitions of temperament).

However, as frequently pointed out [16, 24, 26], a multitude of definitions of personality exists, and the use of terminology is rather inconsistent. Indeed, regarding animals the term personality itself is somewhat contradictory. Perhaps for fear of anthropomorphism, earlier researchers seemed to be reluctant to use the term personality in relation to animals, leading to suggestions for alternative descriptions, such as 'horsonality' [27]. More commonly, terms such as 'temperament' [24–26, 28–31] or 'character' [32] are used as an equivalent of personality, but a variety of other terms such as 'timidity' [33, 34], 'nervousness' [35], 'nervous balance', 'activity' or 'irritability' [36–38], 'emotionality' [39–41] or 'mental traits' [42, 43] has also been used. However, the use of the latter expressions is not very widespread. Furthermore with regard to horses, the terms 'temperament' and 'character' may be confusing as both are used in equine breeding [44, 45] to label specific sub-categories of personality. Therefore, it is suggested that also in animals 'personality' may be the most suitable term to embrace all behaviour-related traits.

While there is limited agreement regarding the labelling and definition of personality in animals, there is yet less agreement regarding the dimensions or sub-traits that comprise personality and the models used to categorize specific types of personality, as outlined in Table 1. The concepts have usually been adapted from personality research in human psychology or research in non-livestock species, but no consensus on terminology and model choice exists there either [74]. This is not surprising as researchers do not even agree on what constitutes behaviour [75], and for example behaviour ratings by scientists [76] or judges [77] may differ from ratings given by practitioners. These disagreements point to the

complexity of issues related to behaviour and personality. Some of the most commonly used classification schemes include the theory of active versus passive coping styles [46], the temperament-character inventory based on a model suggested by Cloninger [52], and the five-factor model [48]. Approaches based on multivariate analyses, such as factor analysis or principal component analysis used to define study-specific sets of traits that may or may not be fitted to an existing model, are also common. However, various other personality models exist in the literature on both human and animal personality. In view of this variety of approaches, many researchers call for a better standardization of classification schemes [16, 78]. Although comparability of results from different studies could be increased, based on a subjective evaluation of advantages and disadvantages (Table 1), it appears that there is no particular classification that is superior to all other models under all circumstances. For example, the classification based on active and passive coping strategies [46] is simple, but does not seem to fit well to most livestock personality traits as they tend to exhibit a continuous rather than binary or bimodal distribution [79–81]. Also, a direct comparison of the five factor model and temperament-character inventory did not reveal major advantages for either of the models regarding construct validity in humans [82]. Neither model was superior in predicting a real-life outcome, i.e. the development of personality disorders, from the presence of particular personality characteristics as assessed by the respective model. However, the temperament-character inventory [52, 53] appears to have particular merit in the context of animal breeding. The personality dimensions are each based on a distinct genetic background linked to neurotransmitter pathways: the trait novelty seeking is linked to basal dopaminergic activity, harm avoidance is linked to serotonergic activity, and reward dependence is linked to noradrenergic activity [52, 53]. These relationships could be confirmed in humans [83–85], and the model has also been successfully applied in a study aiming at detection of QTL affecting feather-pecking in chickens [86]. Similarly, genetic polymorphisms in the dopamine D4 receptor gene have been linked to differences in traits such as vigilance and curiosity in horses [55] and behaviour during handling in cattle [87]. Nevertheless, it remains open to debate as to which individual, phenotypically expressed behaviour traits are valid measurements of the underlying, genetically determined personality dimension, and some traits important to equine practice seem to be missing in that model. Of particular interest to breeders are also models that attempt to distinguish between the inherited component of personality (i.e. the form of behaviour: how quickly and for how long, etc. the animal conducts a behaviour pattern) and the component shaped by the present and previous environment during ontogeny of the trait (i.e. the content of the behaviour: what the animal decides to do), and this differentiation is planned to be adopted by Polish horse breeding authorities (Aleksandra

Table 1 Overview of different models used to describe animal personality including a brief, subjective evaluation of strengths and weaknesses of different models

Name/definition of personality	Traits/categories	Categories mutually exclusive?	Limitations	Advantages	References	
					Key ¹	Sample ²
Coping strategies/styles	Active Passive	Yes	Usually continuous rather than binary/bimodal distribution of behaviour profiles	Simple classification	[46]	[47]
Five factor model	Openness to experience Conscientiousness, Extraversion Agreeableness, Neuroticism	No	Some traits anthropomorphic and/or may be difficult to judge when assessing animals – e.g. conscientiousness	Widespread use in studies on humans	[48, 49]	[50, 51]
Temperament-character inventory	Harm avoidance Novelty seeking Reward dependence, Persistence Self-directedness Cooperativeness Self-transcendence	No	Traits important to practice seem to be missing	Widespread use; genetic basis partly in several species confirmed	[52, 53]	[54–56]
Four common aspects of different temperament definitions	Activity Reactivity Emotionality Sociability	No	Limited set of traits, other relevant traits missing	Straightforward definitions; traits highly relevant to practice	[26]	[57–59]
Using multivariate statistics to define traits	E.g. Gregariousness Reactivity	Partly	Limited consistency and agreement across studies	Optimal model for individual situation		[29, 47, 51, 60–69]
Behaviour traits evaluated in horse breeding programmes	E.g. Temperament, Character Willingness to work Ability to work Trainability Ride-/driveability Intelligence(see also Table 4b)	No	Poor definitions of traits, overlap, relevant traits missing, limited agreement between judges and breeding associations	Most traits highly relevant to practice	[70, 71]	see Table 4b
Form versus content; inherited versus acquired	Temperament (form of reaction, inherited) character (content of reaction, environmentally shaped)	No	Difficulties to distinguish between acquired and inherited components	Distinction highly relevant to animal breeding	*	
Four ‘humours’	Sanguinic Phlegmatic Melancholic Choleric	Yes	Unclear which specific behaviour patterns are indicative of which ‘humour’	Parallels to human behaviour may make concepts easily understood	[72]	[73]

*Personal communication: Aleksandra, Gorecka-Bruzda, 2012.

1=key references; 2=sample studies.

Górecka-Bruzda, 2012, personal communication). Similar trait distinctions are made by a few practitioners in other countries, too [88]. Therefore, based on the different strengths and weaknesses of different models, it is argued that it may actually be beneficial to maintain a variety of approaches until a universally valid model has been constructed. However, the parallel use of one common model and the subsequent development of a universally applicable model would be highly desirable.

Personality Traits in Livestock and Companion Animals

Of the different personality dimensions, temperament traits play a key role as functional traits in livestock. With particularly fearful and highly reactive animals, handling procedures may be slowed down [89], safety for both handler and animal decreases [90], and meat quality may be negatively affected e.g. in pigs [91], cattle [92] and poultry [93]. Aggressiveness and behaviour patterns leading to cannibalism are likewise traits of major importance to animal welfare and the livestock industry as they lead to stress, severe injuries and production losses and/or necessitate the routine introduction of painful procedures such as dehorning [94], castration [95, 96], tail-docking [97] and beak-trimming [98]. Maternal abilities [99] are of particular importance in animals managed under extensive husbandry conditions such as commonly the case with sheep and beef cattle. Depending on the intended use e.g. as a pet, guide dog, herd dog, hunting dog or a search dog, various personality traits beyond those related to temperament are of importance in dogs. They may include traits related to cognitive abilities [100], obedience [101] or protectiveness [102]. In horses, there are likewise a number of additional personality traits that are highly relevant to the animal's welfare and its suitability for different riding and work purposes. Unlike with other livestock species, behaviour traits can be considered to be performance traits, rather than functional traits, because of their direct relationship with the horses' suitability for different types of tasks [103]. Thus, it is for good reason that a large number of sport horse breeding associations (e.g. 11 out of 19 surveyed European breeding associations [70] or four out of five surveyed breeding associations from Europe and the USA [104] plus many additional breeding associations [105–108] not included in these surveys) evaluate personality traits along with the typical performance traits, i.e. the basic gaits and jumping ability. Evaluated personality traits include traits labelled character, temperament, willingness to work, ability to work, attitude to work, trainability, rideability/drivability and handleability [71, 80, 107, 109, 110]. Table 1 gives an overview of the different behaviour traits included in the breeding goals and/or evaluated by breeding associations. Notably the trait temperament is almost universally included in the evaluation. Some associations such as the

Swedish Warmblood breeding association even distinguish between temperament during dressage and show-jumping [111]. Both of the above facts point to the importance of this trait, which is, in spite of great differences between breeding associations, usually defined based on aspects of fear reactivity.

Validation of Personality Assessment Methods

Two major purposes may be distinguished for personality, or in particular temperament assessment: (a) the assessment of animals' suitability for a specific environment, i.e. how well a horse's personality suits to the specific task and (b) the assessment of animals' personality traits in breeding programmes, i.e. the use of individual animals' phenotypes to infer genotypes of their relatives. In any case, valid assessment methods are required. One problem in the validation of personality traits is that behaviour is, per definition, plastic. Therefore, traits defined on the basis of behavioural attitudes can be expected to have incomplete repeatabilities, thwarting attempts to validate assessment methods. For example, horses' reaction to novel stimuli is subject to habituation during repeated exposure [112–117], while generalization is limited to objects that are very similar in characteristics such as colour [118]. These facts imply that horses' reaction to one novel stimulus is not highly predictable of that horses' reaction to another novel stimulus. Nevertheless, three major prerequisites need to be fulfilled in order to ascertain validity of a particular method: reliability, repeatability and validity. Table 2 gives an overview of different measures undertaken to validate various methods for assessment of equine personality. Personality assessment methods most commonly include behaviour tests with or without measurements of physiological parameters, non-context specific ratings by owners or caretakers, and behaviour ratings by external observers during everyday situations. Just two studies focused on intra-observer reliability of equine behaviour traits [121, 137] although research in cattle suggests that high intra-observer reliability cannot be taken for granted, even if apparently objective measurements such as behaviour counts are taken [142]. In the majority of validation studies inter-observer reliabilities or repeatabilities across time are assessed. These figures are usually, though not always [47] high (Table 2), indicating that the horses' reactions can be assessed by independent observers with reasonable reliability and that they are consistent over time. However, publication bias may be present, such that studies yielding poor values may not have reported these values. Repeatability across situations tends to be considerably lower than across time [47, 57, 80]. Similarly, construct validity, such as e.g. correlation between different behavioural and physiological parameters used to measure the same trait is generally also rather weak (Table 2). Also correlations to real-life outcomes are

rarely assessed, except where owner ratings are considered real-life outcomes and subsequently used as a basis for comparison with results from behaviour tests [64]. Such mixed findings are also found in studies validating personality assessment methods in other livestock and companion species [18, 79, 143–150] as well as primates [4]. In addition to the above-mentioned validation steps, many authors argue that discriminant ability should also be fulfilled in order to consider a trait to be validated. However, the genetic architecture of most complex traits [151] is of a quantitative nature, implying that several thousand genes influence the trait in question as well as other traits simultaneously via genetic linkage or pleiotropy. For this reason, it appears unreasonable to insist on the absence of correlations of one trait with other traits. It is thus also not surprising that many personality studies [4] fail to show complete independence of different personality traits.

Assessment of Individual Horses' Personality to Infer Suitability for Specific Tasks

Personality tests are commonly suggested for use in matching animals with suitable owners [152] or for assessment of animals' suitability for specific tasks [145, 146, 153, 154]. In practice, for example specific 'calmness-tests' consisting of novel and startling stimuli are used to assess a horse's suitability as a leisure horse [155, 156], based on its level of fear reactivity. Although assessment is typically not based on such standardized tests, riding school [157], vaulting [158], mounted police [157] and therapy horses [125] are likewise usually assessed by the potential buyers for their fear reactivity. However, besides temperament, other personality traits play important roles in these tasks, too. For example, it has been suggested that one of the most important characteristics for a horse to be considered suitable for therapeutic riding would be the ability to form a positive relationship with humans along with an appropriate level of reactivity to environmental stimuli [159]. In vaulting horses, a high degree of sensitivity to tactile stimuli is undesirable as these horses have to cope with various, changing pressures exerted by the vaulters as they perform their exercises. Individual differences in tactile sensitivity exist [135], and more sensitive horses are likely to experience more pain or discomfort in this type of work. More recently, increasing attention has been diverted to tests allowing for optimization of the horse-rider match [160–162]. It is well known that the quality of human–animal interaction impacts both productivity and welfare of livestock [163], and there is some evidence that similar relationships exist with regard to human–horse interactions [164–166], as reviewed by Hausberger *et al.* [167]. Rider personality has been suggested to impact welfare and performance of the ridden horse, although this relationship does not exist universally but may only

apply to more emotionally reactive horses [162]. There is also some indication that with horse-rider dyads that are rated as a good match, horses perceive less stress during novel object tests [168], although in this case horses' level of reactivity and familiarity with the rider seem to play a role, too [157]. Links between pets' and their owners' personalities have been suggested for dogs [169] and horses [160] although the observed correlations between personality traits were rather weak. However, analyses of co-variation in personality traits of human couples likewise revealed only weak associations between partners' personalities, including those couples that declared to be happy with their relationship [170]. These findings may indicate that a complete match of personality traits in a given dyad is not necessary for satisfaction with the relationship and perhaps even counteracts contentment. Potentially, an optimum level of agreement rather than complete agreement yields highest satisfaction, allowing for some room to complement each other with different personality traits.

Assessment and Selection for Personality Traits in Equine Breeding

Breeding strategies may be used in many different ways to impact animal welfare [171]. Personality traits are commonly evaluated in horse breeding programmes. However, just a few breeding associations (e.g. some draft-horse breeds [172]) use the obtained information to derive breeding values for personality traits that could be used in sophisticated selection decisions. Also, a review of studies dealing with health disorders leading to death or euthanization in horses (Table 3) revealed that only in three out of nine studied populations horses were indeed retired or culled because of temperamental disorders. These three populations included a population of Icelandic horses [174], coldblood draft horses [179] and thoroughbred race horses [178]. Interestingly the former two breeds are known for their calm temperament. Therefore, results seem to reflect the rigorous selection strategies that in part may have resulted in the calm temperaments of these breeds, rather than the high prevalence of temperamental problems in these breeds. From an ethics point of view, such selection decisions have to be seen critically, as appropriate training and management methods are likely to prevent or alleviate existing problems. In contrast from a breeding point of view, the culling of horses with undesirable behaviour that in part may be based on a 'difficult' personality is an appropriate step towards a sustained progress in personality traits. However, the reluctance to base selection decisions solely on personality scores assigned during performance tests seems justified. The subjectivity of the evaluation systems and the lack of variance in personality scores have long been criticized [180, 181]. Our recent work revealed that these problems persist [71, 80, 182, 183] in spite of

Table 2 Overview of results from studies undertaken to validate different aspects of equine personality assessment

References ¹	Parameter	Test situation	Method ²	<i>n</i>	Repeated time	R. sit ³	Intra-obs. reliability ⁴	Inter-obs. reliability ⁵	Repeatability	Construct/content/ criterion validity	Discriminant validity
[119]	Rideability	PT ⁶ rider	VC	435	6.5, 8, 11 m				0.58–0.70		
[120]	Stereotypic behaviour	Rearing, training, competition	VC	278	Pre-/post weaning, 12, 18, 24 m	x			0.41		
[39]	Reactive behaviour	NOT ⁷	<i>r</i>	32						Tranquilizer lowered HR ⁸ during NOT	
[121]	Sexual behaviour	Video versus life recordings	<i>r_p</i> ; % agree.	12	1 w		0.47–1.0 60–100%	0.67–1.0			
[122]	Behaviour in barn	PT	VC	206	25, 50, 80, 100 days				0.71–0.89		
[45]	Temperament	PT	ICC		< 100 days				0.73–0.80		
	Character			247					0.61–0.83		
	Rideability								0.52–0.63		
[41]	Fearfulness	NOT	<i>r_s</i>	42		x				Arena-handling <i>r</i> =n.s.	
	Fearfulness	Handling								Handling-NOT <i>r</i> =0.48	
	Gregariousness	Arena test									
[31]	Fearfulness	Arena test, NOT, handling, instrumental task, memorization, detour, rating by rider	<i>r_s</i>	72		x				<i>r_s</i> =0.24–0.44 between tests and rider rating	
	Gregariousness										
[123]	Response time	Optical/acoustic stimuli	VC	155	1, 1.5, 2, 2.5 y				0.3–0.61	Fillies higher <i>r</i> than colts	
	behaviour score								0.32–0.68		
[124]	Affectionate	Subjective ratings by familiar observer	<i>r_s</i>	20				0.25			
	Alert							0.37			
	Bold							0.00			
	Confident							0.29			
	Enthusiastic							0.40			
	Fizzy							0.42			
	Flighty							0.70			
	Honest							0.22			
	Intelligent							0.30			
	Laid Back							0.40			
	Lazy							–0.11			
	Moddy							0.45			
	Sensitive							0.12			
	Sharp							0.66			
[125]		Survey, NOT, cortisol (nor-)epi-nephrene	<i>r</i> , <i>R</i> ²	103				0.01–0.52 in 37% of horses		<i>R</i> ² =0.07–0.11 for ability to predict personality score by hormone concentrations	
										Increase after exposure	
										Increase after exposure	
										Increase after exposure	
[126]	HR	Novel environment and NOT	ANOVA	8						No increase	
	(Nor-)adrenaline									No increase	
	Cortisol										
	Vasopressin										
	Oxytocin										

[61]	Temperament	NOT handling	r_s , PCA	41	9, 10, 21, 22 m		0.36–0.81	Only within years; limited r across years	
[127]	HR, HR variability, activity	NOT, trained or untrained horses		41	9, 10, 21, 22 m		HR consistent between years	Exposure to NOT leads to decrease in HR variability, effect stronger in untrained horses	
[50]	Big five, confidence in rating	Familiar (Ø 2.8 years) rater	PCA	210				Neuroticism + extraversion rated with most confidence, openness with least	
[51]	Neuroticism Extraversion Openness Agreeableness Conscientiousness	Rater familiar (Ø 2.8 years) with horse	r_s	10		0.62 0.45 0.26 0.49 0.36		$r=0.0-0.66$ for individual traits; not all judges and all traits equally good	$r=n.s.$ between five traits
[47]	Temperament, active/passive coping	Arena test, object test, human test, startle test questionnaire	r_s , PCA	33	3× after 9 days		only for arena significant r over time	Categorization in active/passive coping not possible; $r=n.s.$ between test + questionnaire	
[62]	Personality	NOT, handling, avoidance/reward learning	PCA	41				Personality traits predict show-jumping performance	No correlations between tests
[63]	Temperament	NOT, handling, unfamiliar rider assessment	r , PCA	18				HR in test correlates with rating by riders	Handling, but not NOT correlate with rider rating
[128]	Learning ability	Avoidance learning, reward learning	r_s , PCA	39	2× at 14–15 m, 2× 16–20 m later		r significant short term + for avoidance also long-term		No r between avoidance and reward learning
[28]	Temperament	Comparing survey and NOT + HR	r_s , r_p , PCA	86				Good agreement, e.g. horses rated more anxious had higher HR in test	Three independent traits: anxiety, novelty-seeking understanding
[129]	Temperament	Comparing a survey of caretakers in two groups of horses	FA Cronbach's α	69 and 70				Internal consistency ($\alpha=0.69-0.94$) for anxiety, trainability and affability	Four factors common to both groups: anxiety, trainability, affability, gate entrance
[130]	E.g. locomotion, vigilance, time eating, HR	Novel visual, auditory, olfactory objects	ANOVA	24				Time spent eating correlated with all other variables, HR from auditory test correlated with other tests	Behavioural response linked to type of stimulus
[131]	Reactivity	Social isolation (Passing) novel object Emotionality rating	Control with tranquilizer	40				No difference in HR between tranquilized/untranquilized horses $r>0.33$ with HR	
[64]	Behaviourally defined adjectives	Questionnaire + behaviour observations	r , PCA	61		Agreement in 72% of the horses		Significant correlation between behaviour test and rating	Six components identified, e.g. dominance

Table 2 (Continued)

References ¹	Parameter	Test situation	Method ²	<i>n</i>	Repeated time	R. sit ³	Intra-obs. reliability ⁴	Inter-obs. reliability ⁵	Repeatability	Construct/content/ criterion validity	Discriminant validity
[66]	Personality descriptive adjectives	Owner and trainer supplied information; Agreeableness Extraversion Neuroticism	PCA ICC3,1	100				0.72 0.52 0.45			Agreement with five-factor model
[132]	Behavioural reactivity	Veterinary examination	VC	4452					0.97–0.98	rg=0.9 between behaviour traits at different veterinary procedures	
[133]	Reactivity to humans	Familiar passive, familiar active, unfamiliar active human	<i>r_s</i>	110	8, 18, 30 m				For example time to touch: 0.40–0.67		
[134]	Reactivity to separation, isolation, attraction to, passing of/from conspecifics	Frequency of neighing, defecation, locomotion, etc.)	<i>r_p</i>	110	8, 18, 30 m	x			Time: 0.41 < <i>r</i> < 0.61	Medium <i>r</i> between situations	
[135]	Behaviour during visual, auditory, olfactory, gustatory, tactile stimulation	Novel stimuli challenging different senses	<i>r_s</i>	26	5 m	x			Generally within one sense: 0.27–0.59		No correlation between reactions to different senses
[136]	Fearfulness	NOT passing novel object startling test	<i>r</i>	66	0.6, 1.5, 2.5 years	x			Many parameters with medium correlation across times fewer correlations over time, but HR was correlated		<i>r</i> =n.s. to NOT
[67]	Responsiveness	Open field, 12 observers	GPA, PCA	20					Semantically consistent word charts		2 axes: tractable, attentive
[137]	Temperament	NOT, handling	Cohen's Kappa	56	Video recording		0.14–0.92	0.09–0.85		Significant correlations between tests	
[138]	Reactivity to humans, Fearfulness	NOT, handling, human, startling, questionnaire familiar rater	<i>r_s</i> , FA	53						Medium correlation between questionnaire, and both tests	<i>r</i> =0.36 human test – fear measures
[80]	Temperament, Reactivity	Ridden NOT	VC	224	2–3 or 18 w			0.89–0.95	0.69–0.75	Medium correlation between reactions to different stimuli	n.s. correlation to performance traits
[57]	Activity, reactivity, emotionality	NOT ridden, led, free-running	VC	65	3× in <10 min	x			0.10–0.45	Medium correlation between different measurements. No correlations to conventional personality traits from PT	
[139]	HR, HR variability Behaviour counts during riding	Dressage training	<i>r_p</i> , <i>R</i> ² , VC	36	4× in 4 w				0.57 0.23–0.25	<i>R</i> ² =0.26–0.46 for behaviour to explain PT personality scores	

[140]	Behaviour counts + evaluations, HR variability	PT riding training dressage, cross-country, NOT	VC, R^2	16	0.05–0.28	$R^2=51.5$ for behaviour (refusal jump, jumping manner, intensity of rider's aids) to explain 'willingness to work' score
[141]	Social rank, fearfulness, learning, fecal cortisol, HR	NOT, negative/positive reinforcement, (un)familiar environment	r_s	25	0.75	$r=-0.66$ between HR in novel environment and performance in negative reinforcement task; $r=-0.43$ between fecal cortisol from familiar environment and social rank; $r=n.s.$ between performance in negative and positive reinforcement

1=References; 2: r_p =Pearson correlation, r_s =Spearman correlation; agree.=agreement, VC=calculations based on variance components from analysis of variance, ANOVA=analysis of variance, ICC=intraclass correlation coefficient, R^2 =coefficient of determination from regression analysis, FA=factor analysis, PCA=principal component analysis; 3=repeated over altered test situations (x=yes; situations listed under methods); 4=intra-observer reliability; 5=inter-observer reliability; 6=performance test on station; 7=novel object test; 8=heart rate.

several attempts by breeding organizations to improve the situation [172]. In fact, the observed grade inflation [183] that can in part be attributed to economic competition between testing stations [71] suggests that problems continue to worsen. The lack of appropriate guidelines, the lack of reliance on existing guidelines and severe problems with the trait definitions in the existing guidelines [71, 182] allow for misuse of scores, e.g. to promote specific horses or to increase attractiveness of a testing station to breeders [71, 172]. However, judges and breeding authorities are aware of the problems [71, 182] and many attempts have been undertaken to develop assessment procedures that allow for a more objective assessment of personality traits [57, 58, 80, 137–139, 184–188]. Such a better consideration of personality traits in breeding programmes seems imperative, given the strong link between animal behaviour and rider and animal safety [189]. The importance of personality is further underlined by the value breeders [106, 190] and potential buyers both from the leisure riding sector [190–195] as well as from the professional or competition riding sector [190, 196, 197] assign to personality traits. At present a mismatch between purchasers' expectations and vendors breeding and sales strategies seems to exist [195, 198, 199], indicating that a greater emphasis of personality characteristics by vendors may be needed to align their and the customers' wishes. Although some differences in the relative importance of different personality characteristics exist between riders of different disciplines [200], groups or proficiency levels [190], remarkably, almost all of the above studies [106, 190–194, 196] invariably and independently from each other detected that personality traits were rated by the horse enthusiasts as more important than any of the conventional performance traits such as the gaits or jumping ability.

Genetics of Personality Traits

Domestication is known to be based largely on selection for tameness as impressively shown by selection experiments in foxes [201, 202], and there is thus little doubt that behaviour traits possess some genetic background. Breed [65, 80, 107, 203–205] and genetic-line [206, 207] or sire-line [41, 203, 208–211] differences in personality traits have frequently been reported in equine studies. Also, the level of inbreeding may [38] or may not [212] influence personality traits, depending on the breed and the traits in focus during (in)breeding. Yet, relatively few studies were specifically designed to estimate genetic parameters for equine behaviour traits (Table 4a). In comparison, various studies report genetic parameters for personality traits evaluated in horse breeding programmes (Table 4b). Heritability estimates for these personality scores from performance tests range from values not significantly different from 0 to 0.40. However, all values suggesting a low heritability are estimated based

Table 3 Prevalence of temperamental disorders leading to termination of functional and/or biological life in relation to other groups of health disorders in different riding, breeding or racing horse populations

Reference	[173]	[174]	[175]	[176]	[177]	[177]	[178]	[179]	[179]
Data:									
Type of horses	WB ¹	IC ²	WB	MI ³	WB	WB	TB ⁴	CB ⁵	WB
Gender of horses ⁶	♂	all	♀g (♂)	♀g	all	All	all	♀g	♀g
Use of horses ⁷	Breed	Ride	Ride	Ride	Ride Breed	Ride School	Race Breed	Work	Ride
Functional longevity? ¹³	✓	✓	✓	✓	—	✓	✓	✓	✓
Biological longevity?	✓	—	—	—	✓	—	—	—	—
Data source	Studbook	Survey	Insurance	Insurance	Insurance	Insurance	Survey	Survey	Survey
Time frame	1975 2010	ca. 1997 2002	1977 1987	1997 2002	1984 1994	1984 1994	2002 2003	1970 1975	1968 1982
Group of health disorder/ affected system	Frequencies of disorders (%)								
				8		10	11		12
Respiratory	10.3	0	16.3	4.1	7.8	10	<31	2.1	8.9
Reproductive/urinary	0.8	0	1.2	—	7.8	<5	<31	—	0.4
Cardiovascular	6.4	0	4.6	—	13.9	<5	<31	0	2.6
Infectious/nervous	1.6	1.0	2.8	—	2.3	5	<31	3.7	5.6 ¹²
Musculoskeletal	4.8	54.1	61.2	76.0	32.7	83	<31	13.4	55.5
Other/unknown	35.0	2.0	—	10.2	5.2	<5	16.8	15.5	4.2
Trauma/accident	4.0	12.2	—	17.7	⁹	⁹	<31	3.1	9.1
Gastrointestinal	7.9	3.1	9.8	10.1	35.5	<5	<31	6.2	5.6
Performance	29.4	11.2	—	—	—	—	36.5	—	—
Temperament	0.0	16.3	—	—	—	—	6.4	20.0	<5.6 ¹²

¹=Warmblood; ²=Icelandic horse; ³=mixed breeds; ⁴=thoroughbred; ⁵=coldblood; ⁶: ♂=stallions, ♀=mares, g=geldings; ⁷breed=breeding, ride=riding, school=riding school horses, work=draft and/or cavalry work; ⁸categories not mutually exclusive; ⁹respective category included with 'other'; ¹⁰approximate values; ¹¹detailed health disorders not further specified; illness and injury accounts together for a total of 31% of losses; ¹²temperament included with disorders of the nervous system; ¹³✓=yes, —=not recorded.

on data collected in the field. Unfortunately, individual variance components are not always given, but based on the differences between field and station data it could be speculated that the low values in the studies based on field records are mainly based on large phenotypic variances, which in turn are based on strong environmental influences, rather than a complete lack of additive genetic variance. However, as indicated above, and also highlighted earlier [172, 180] the present evaluation system for personality traits in horse breeding is flawed, so statistical properties of the scores are suboptimal and the corresponding genetic parameters have to be seen with great caution.

Based on specifically designed personality tests, horses' reactions to novel objects ($h^2=0.17-0.90$) during handling situations ($h^2=0.23-0.28$) or in learning tests ($h^2=0.15-0.89$) seem to be moderately to highly heritable (Table 4a). Similar to the estimates based on conventional personality scores, some restrictions apply such as low sample sizes and violation of parametric assumptions. However, the majority of estimates, including those from studies based on large sample sizes [132], are in a moderate range ($h^2=0.15-0.40$). These values support the view that selection schemes focusing on these traits could be successfully implemented into breeding programmes,

as already the case with a number of breeding organizations [187, 237, 238]. Furthermore, genetic correlations to other relevant performance traits appear to be generally low, and mostly of a synergistic nature. For example, according to our recent work [59], scores for reactivity in a novel object test show a genetic correlation of $r_g=0.41 \pm 0.33$ to the scores for the quality of the gait trot and of $r_g=0.69 \pm 0.26$ to scores for rideability, indicating that genetic selection for calm reactions in a novel object test will entail genetic improvement of the trot and rideability. Nevertheless, some consideration should be given to both ethical and economical aspects before setting up a rigorous selection scheme focusing on personality traits. For example, physical problems should be ruled out before attributing unwanted behaviour such as head-tossing during riding to poor personality characteristics of the horse [139].

Possibilities of Selection for Personality Traits other than Temperament-Related Traits

Compared with aspects of temperament, and in particular fear reactivity, little research has focused on other personality traits. However, some of these other personality

traits seem to be just as important to the riders [106, 190, 239], and at least indirectly also to equine welfare. Tables 1 and 4b list different personality traits currently considered in various breeding programmes, implying that they are considered to be highly relevant traits by practitioners. Surveys among riders [190, 194] and breeders [106, 190] revealed that traits such as rideability, willingness to work and sensitivity are just as important as temperament-related traits. Furthermore, a tight link between personality and performance seems to exist, at least based on the scores from conventional performance tests [80, 240]. Recent attempts of moving towards a more objective assessment of these additional personality traits include the assessment of rein-tension as an objective indicator of the intensity and quality of rein contact as an important component of rideability [241]. Considerable attention has also been paid to direct behavioural observations during different situations important to practitioners such as behaviour during ridden dressage training [139, 242], cross-country training [140], or during handling and leading [243] and veterinary inspections [30, 132]. Some of these studies detected links between specific behaviour patterns such as head-tossing [139, 242] and refusal to jump with the traits rideability and willingness to work, respectively, as assessed during conventional performance tests. Based on these links it is suggested that some of these detailed behaviour patterns should be specifically considered in personality evaluation as they would make the assessment more objective. At present, the situation is further complicated by traits bearing the same label, but which may be defined quite differently in different breeding organizations. For example, in Dutch warmblood breeding, the trait character is defined as a 'score for behaviour during the training of all traits. Character is clarified by concepts such as pleasant companionship, intelligence, courage and willingness to perform', while behaviour in the stable is defined as a separate trait [122]. In contrast, in German warmblood breeding, the trait character focuses on behaviour during handling and in the barn only, and is defined as the 'handleability and behaviour towards humans, the time taken to accustom to the testing station, the overall body posture and behaviour, and in particular behaviour during grooming and tacking' [244].

One of the major areas in which selective breeding on personality traits could help to improve equine welfare is the selection of animals better adapted to confined housing, as suggested also for zoo animals [245], and in particular to individual housing. Rebound effects in exercise after box housing [246] increases risks for injuries, and social separation is stressful to horses even if they are familiar with the surroundings [247]. Breed differences in responsiveness to arena tests involving social separation are present [203, 215], demonstrating that horses of some breeds may be better able to cope with the situation and thus experience less stress when isolated from conspecifics. Similarly, low levels of agonistic interactions

in specific groups of horses have been attributed in part to the temperament of that breed [248]. Although little is known about the within-breed additive genetic variance for these traits, it therefore seems likely that such aspects of social behaviour could be changed by selective breeding, and consequently could be used to positively influence animal welfare. In pigs, selection for social behaviour and in particular low aggressiveness has long been in the focus of research [148, 249]. Heritability estimates for aggressiveness and related indicator traits [249, 250], suggest that a selection on these traits may, in theory, be possible in pigs. However, as pointed out by Muir [251], traits related to establishing of dominance hierarchies may be difficult to change when focusing on selection based on individual's performance. Therefore strategies based on group selection, considering the performance of the entire group rather than just individuals within groups [251] may be a more successful strategy for changing traits related to social behaviour. In fact, other personality traits unrelated to social behaviour may also develop differently depending on the personalities of surrounding individuals. Personalities may best function, e.g. based on conflict avoidance [1], if there are large individual differences within an existing group, and thus personalities may form in dependence of personalities of group members to fill in free niches. These theories are supported by recent findings in pigs that genetic clones differed insignificantly less from each other in personality traits compared with a group of distantly related pigs reared and kept under the same husbandry conditions [252]. Likely, the same principles apply to horses, too, making selection for traits related to social behaviour difficult. Moreover, the standardized collection of data related to social behaviour is time-consuming, and costs in relation to profits may be prohibitive for inclusion in horse breeding goals. Even in a fairly long-term perspective it appears to be more economical to fit the environment to the animal rather than taking the opposite approach.

Personality traits related to intelligence and learning ability are likewise expected to impact equine welfare as slow learners will experience repeated negative reinforcement or punishment, while exceptionally intelligent or fast learning horses might likewise experience undue amounts of punishment as they might make undesired associations between unintentional cues and events. A large number of studies focused on various aspects of equine learning behaviour, although most commonly, studies on equine learning are based on positive reinforcement [253–259] rather than negative reinforcement [211, 260, 261], which is the most commonly used method in practical horse training [262, 263]. Since learning based on rewards via food or pleasurable tactile stimuli likely relies on fundamentally different underlying motivations compared to learning based on negative reinforcement via tactile stimuli, it is uncertain in how far results from the former studies are transferable to most practical settings. However, connections may nevertheless exist between

different types of learning as the same anatomical structure seems to be involved in processing of reward and unpleasant touch as occurs in negative reinforcement [264]. Either way, traits related to learning would deserve in future more attention in both research and practice. Some recent attempts to consider learning in breeding programmes include the assessment of speed of habituation in repeated novel object tests [80].

Potential Constraints in Selection for Personality Traits – Relationship between Personality, Performance and Welfare

A review of primate literature also suggests that individual differences in personality may impact welfare or call for individual adaptation of management and training strategies to secure a certain level of welfare [265]. Riders commonly believe that breeding for calm, less reactive horses will result in dull, less sensitive and low-performing horses. However, personality traits assessed during conventional performance tests almost invariably show desirable phenotypic and genetic correlations with performance traits (Table 4a, b [80, 266]). Equally, a number of studies using behaviour tests or ratings during working situations [58, 80, 267–269] found few antagonistic correlations and in many instances the presence of desirable correlations between aspects of temperament and performance in basic gaits, jumping or racing. In contrast, highly reactive horses may be more prone to injury: Lam *et al.* [270] suggested that higher tendon injury rates in stallions might be a result of a more difficult temperament caused by higher levels of male hormones. However, no such relationship was observed between temperament and susceptibility to colic [271], although stress is thought to be a predisposing factor for colic [272]. Therefore, overall a more rigorous selection for behaviour traits seems to be plausible. However, before specific traits are implemented into selection schemes of breeding programmes, these traits need to be validated, and genetic parameters estimated to assess the potential for a successful selection on these traits. Furthermore, economic weights would be necessary for reasonable weighing of the different breeding goal traits, but are notoriously difficult to obtain in horse breeding and for behaviour-related traits [106, 196, 273]. Although for a number of traits and testing regimens these prerequisites are fulfilled (see above and below chapters), and costs for assessment can be kept below the level, owners would be willing to pay additionally for an objective assessment [186], some breeding organizations remain reluctant to introduce new traits into their selection schemes as it would entail relaxing selection on the performance traits. Furthermore, breeders fear that deliberate manipulation may be an issue, in particular, with specific behaviour tests [71, 80]. Although specific training of behaviour traits is possible, the same applies to performance traits [274], and

may be used to indirectly assess horses' trainability. In the distant future, genomic selection might help to overcome trait manipulation. Provided that conventional breeding values with sufficient accuracies can be obtained, considerable gain in breeding value accuracies can be expected especially for lowly heritable traits [275].

Influencing Personality by Interventions during Ontogeny of Personality Traits

The influence of early experiences on cognitive development and on later expression of personality traits has long been known from laboratory animals [276], and some results from livestock species are also available. For example, in pigs, marginal differences in housing during rearing resulted in differences in pigs' subsequent reactions to novelty and handling [277]. Perhaps because of the requirement of long-term studies and their limited feasibility in equine research, comparably little research has been conducted regarding possibilities to take influence on the ontogeny of equine personality traits. Temporary effects on horses' fear reactions or learning performances regarding the influence of e.g. diet [278], specific calmatives [279, 280], predator odour [281], restriction of the visual field [282], the presence of other horses [283], or the possibility to observe another horse in a given situation [284], the presence or actions of humans [57, 115, 285–289] and the rider's riding style [290, 291] or training method [292] are well documented, but comparably little is known about possibilities to permanently influence equine behaviour traits. Nevertheless, some interesting results are available on how handling or housing regimens [293, 294] earlier in a horse's life can influence later behavioural attitudes or stress responses ([295–297], reviewed in [298]). Considerable research efforts have been undertaken to assess the impact of various foal or weanling handling regimens (including e.g. habituation to human contact and handling procedures such as haltering or picking up feet) on their handleability and trainability later in life [68, 73, 299–306] (Table 5). Somewhat disappointingly, most studies failed to reveal positive, long-lasting effects of these handling regimens [308], although in a few studies positive effects such as improved handleability and trainability were evident after periods of as much as 1.5 years [304] (Table 5). Positive effects were also transferred to the foals' reactions to unfamiliar humans [68]. On the other hand, negative effects of neonatal handling on social behaviour are also described. In the absence of additional experiences with humans that might overshadow past experiences, positive [309] or even one single negative experience with humans, such as a veterinary treatment [310], appears to have lasting effects on horses' reactions to humans. Similarly, positive handling of the mare rather than the foal appeared to have lasting (1 year) effects on a foal's behaviour towards humans [311]. Differences in juvenile

Table 4a Genetic parameters for equine behaviour assessed via specific behaviour observations taken during tests or everyday situations

Reference	Trait	Assessment/situation	Scale	Breed	n^{10}	h^2	Correlated to trait	r_g
[213]	Trainability	Owner rated	Score 1–5	SB ¹	159 (31 sires)	0.10 ± 0.32	n.c. ¹¹	n.c.
[120]	Stereotypic behaviour	Rearing + training	?	TB ²	278 (6 sires)	$0.13–0.79$	n.c.	n.c.
[214]	Timidity	Novel objects	Scores	MA ³	593, 424, 127	$0.39–0.90$	n.c.	
	Memorizing ability	Labyrinth	?	Wl ⁴		$0.15–0.89$		n.c.
				Sl ⁵				
[215]	Emotionality	Arena test	Index-scores	SF/16 dif. ⁶	246/703 (5583)	$0.39 \pm 0.26/0.42 \pm 0.11$	n.c.	n.c.
	Reactivity	Novel object				$0.40 \pm 0.24/0.29 \pm 0.12$		
	Reactivity	Bridge	Sec.			$0.81 \pm 0.17/0.24 \pm 0.10$		
	Intelligence	Chest opening	Sec.			$0.48 \pm 0.20/0.36 \pm 0.13$		
	Memorizing ability	Memorization	Sec.			$0.28 \pm 0.21/0.17 \pm 0.14$		
		chest opening						
[132]	Reaction to:	Routine veterinary examination	Score 0–3	TB ²	4452 (10 590)	$0.25–0.26$	Auscultation	$0.81–0.91$
	Inspection of conjunctiva					$0.23–0.26$		
	Auscultation					$0.27–0.28$	Blood sampling	$0.85–0.88$
	Blood sampling					0.26	Conjunctiva	$0.88–0.99$
[187]	Emotionality	Bridge – driving	Score/linear	FM ⁷	520	0.17	n.c.	n.c.
	Emotionality	Bridge – riding	1–9			0.23		
	Emotionality	Novel object – riding				0.27 ± 0.08		
[58, 59]	Reactivity	Novel object test	Score 1–10	WB ⁸ , PO ⁹	575 (10 970)	0.40 ± 0.22	Reactivity – basic gaits	$0.28–0.72$
	Activity		–5–5			0.26 ± 0.21		
	Emotionality		–5–5			0.32 ± 0.21		
	Time to calm down		Sec.			0.17 ± 0.29	Reactivity – rideability	0.67 ± 0.69
	Intensity of rider's aids		1–10			0.21 ± 0.13	Reactivity – free jumping	
	Attention to environment		%			0.18 ± 0.11		-0.04 ± 0.15
	Attention to rider		%			0.00 ± 0.01		
	Attention to stimulus		%					

1=Standardbred; 2=Thoroughbred; 3=Malopolski; 4=Wielkopolski; 5=Silesian; 6=Selle Français (first values for heritability) and 16 different breeds, including the 246 Selle Français (second values); 7=Franches Montagnes; 8=German Warmbloods; 9=mixed breeds of ponies; 10=values represent number of tested animals (and number of animals/sires in the pedigree); 11=not calculated.

Table 4b Genetic parameters for equine personality traits assessed during performance tests

Reference	Trait	Assessment/ type of PT	Scale	Breed	n ¹³	<i>h</i> ²	Correlated to trait	<i>r</i> _g
[216]	Temperament Character	Field	Score 5–10	IC ¹	1171	0.14 0.20	Performance, personality	Low but positive
[180]	Temperament Character Obedience	?		HB ²	284, 295 (31 sires)	0.03–0.11 0.07–0.38 0.12–0.09		
[217]	Temperament Character	Field	Score 5–10	IC ¹	3975	0.13 < <i>h</i> ² < 0.53		
[218]	Temperament Character	Station	Score 1–10	HA ³ , WB ⁴	730	ca. 0.10		
[219]	Temperament	Station	Score 1–10	HO ⁵	286	0.76	Influence of training status prior PT	
[220]	Character Rideability	Field	Score 1–10	KW ⁶	2023	0.06 ± 0.04 0.03	Gaits dressage in competition	0.42–0.87 0.83
[122]	Character Behaviour in stable	Station, 3 judges	Score 1–10	KW ⁶	206	0.24–0.52 0.31–0.65		
[111]	Riding temperament Jumping temperament		Score 1–10	SW ⁷ WB ⁴	3601	0.27 0.14	Jumping ability external conformation	0.42 0.74
[221]	Character Temperament	Station	Score 1–10	HA ³	994	0.27 0.30		
[222]	Willingness to work Character Temperament Willingness to work Constitution	Station	Score 1–10	WB ⁴	2207	0.24 0.18 0.26 0.14 0.10		
[223]	Temperament Docility	9 judges	Linear 0–10	HF ⁸	3902	0.06 0.02	Docility	– 0.84
[224]	Character Temperament Willingness to work Constitution	Station	Score 1–10	WB ⁴	2815	0.21 0.26 0.29 0.35		
[225]	Temperament		Score 1–10	PR ⁹	1273	0.08 ± 0.04	Performance traits	Not converged
[226]	Temperament	Field/station		HF ⁸	660	0.2 < <i>h</i> ² < 0.7	No antagonistic correlations	
[227]	Temperament Free-jumping Temperament Jumping	Station	Score 1–10	SW ⁷	359 (8918)	0.23 ± 0.14 0.33 ± 0.23	Jumping Jumping; n.s. to gaits	0.79 ± 0.19 0.90 ± 0.06
[228]	Handleability Willingness to work			HF ⁸		0.09 0.10		

[229]	Nerve strength Willingness to work Concentration Handleability	Field	Score 1–10	SG ¹⁰	0.14 0.14 0.17 0.22
[230]	Riding temperament	Field	Score 1–10	SW ⁷	0.38±0.11 0.51±0.12
[231]	Jumping temperament	Field	Score 1–10	SW ⁷	0.29 0.21
[232]	Jumping temperament	21 judges	Linear	IH ¹¹	0.26±0.03
[233]	Spirit	Field	Score 5–10	IC ¹	0.29±0.05–0.63±0.02
[234]	Character	Studbook inspection	Score 4–10	FI ¹²	0.12±0.02
[235, 236]	Cow sense/cutting ability	Competition	Score	SB ¹³ QH ¹⁴	0.17±0.06 0.04

1=Icelandic horses; 2=Halfbred; 3=Hanoverian; 4=Warmbloods; 5=Holstein; 6=Dutch Warmblood; 7=Swedish Warmblood; 8=Hafinger; 9=Spanish Pure Bred; 10=Southern German Draught Horse; 11=Italian Heavy Draught; 12=Finnhorse trotter; 13=Standardbred trotter; 14=values represent number of tested animals (and number of animals/sires in the pedigree).

or adult horses' behaviour during and following different training regimens generally revealed advantages of more sympathetic or positive reinforcement approaches [312–316], but again, studies on long-term effects are missing. Nonetheless, since horses seem to be capable of learning to learn [317], such training strategies might result in a permanent improvement of learning abilities.

Personality traits of dams may also impact welfare of their offspring. For example, less fearful mares showed higher levels of agitation during separation from their foals [318], perhaps indicating that these mares had better maternal abilities and better offspring survival, as shown e.g. in sheep [319]. More importantly, the weaning time and method [320–324] is another crucial management procedure that may have lasting effects on the animal [325–327]. Complete maternal deprivation impacts personality [328], though appropriate management may result in most behaviour traits not being affected [329]. However, orphan foals appeared to learn slower [330] and show more stress during social separation [331]. Weaning is considered to be one of the most stressful events in a horses' life as it generally entails the simultaneous break of the nutritional and social bond between foal and mare [323]. A large proportion of stereotypies in horses develop just after weaning. Although the true function of stereotypies is still unknown it has been suggested that they may act as a coping mechanism during stressful situations [332] including in particular the frustration of behavioural needs [333, 334]. As the predisposition to develop stereotypies seems to be heritable [120, 335, 336] and not all horses develop stereotypies under the same management conditions, and because there are links between the expression of stereotypic behaviour and personality characteristics such as learning [337] and perseverance, the expression of stereotypic behaviour might provide important information on equine personality. Interestingly, the prevalence of stereotypies in formerly feral horses was surprisingly low [338], possibly indicating that a stimulus-rich environment during rearing enables neural circuits to form that later help the individual to cope with the various stressful situations during confined housing and managed feeding. Links between type of work and personality [339] or stereotypies [340] have also been described, although the causal relationship is not always clear. Potentially, horses with certain personality characteristics are more likely to be selected for certain types of work rather than that factors such as work-specific handling or 'learning to learn' have a large effect. The type of housing is likewise suggested to influence handleability and trainability, although less is known about long-term effects. For example, group-housed horses seem to experience less stress during weaning [341], show improved social behaviour [342], are easier to handle [343], and are more efficiently trained [307, 344]. The general importance of the animal's social group to expression of individual personality has been reviewed for different species [345]. Obviously,

Table 5 Duration of effects of different handling/management regimens on personality-related parameters

Reference	Handling/management regimen ¹	Age ²	Parameter ³	No effect ⁴ (time)	Effect ⁵ (max. time)
[68]	1 m 45 min/day handling	ca. 16 m	Behaviour before/after handling	1 y	End
[73]	14 d daily handling	foals	Proportion of time resistant		6 m
[294]	Forest or stable reared; handled or non-handled	2 w–24 m	Behaviour score for various handling procedures		End
[297]	3 m single versus paired housing	2 y	Stereotypies, HPA-function ⁶		End
[302]	14 d (un-)forced contact	Weaning	Reaction to novel object	3 m	–
[303]	1 h separation from dam	0 d	Reaction to human, haltering	>4 m	4 m
[304]	0–1 y handling	2 y	Mare-foal-bonding, social behaviour		1 y
[305]	0–120 d, different intensities	2 y	Learning ability		End
[306]	0 versus 5 d daily handling	0 d	Learning ability, manageability	16 d	0
[307]	6–24 m single versus group housing	0 d	Calmness, friendly, handling		4 m
[308]	(Un-)forced contact	6 m	Handling, aggression towards humans		End
[309]	pos./neg. reinforcement	0 d	Human approach		2 w
[310]	12 d daily handling	1 y	Human-animal relationship		8 m
[311]	5 d human-mare contact	Weaning	Fitting halter, picking up feet, being led	18 m	18 m
		0 d	Human approach, handling		1 y

¹m=month; d=days; y=years; pos.=positive; neg=negative.²Age at which handling/management regimen started; w=weeks.³Parameters assessed during later tests.⁴Period of time after which no effect of handling/management regimen could be detected.⁵Maximum period of time after which an effect in at least one of the measured variables could be detected; end=there was an effect immediately at the end of the treatment regimen, but no tests were conducted at a later time.⁶HPA=hypothalamus-pituitary-axis.

behavioural expression of personality traits is also influenced by health and physical conditions [346]. For example, if a horse's vision is impaired it will have greater difficulties in learning tasks based on visual stimuli, and depending on the type of impairment, reactions to novel visual stimuli may be enhanced or lowered. Links between visual ability and general personality have also been suggested in humans [347], highlighting the importance of prior experiences on personality.

Conclusions

Some reliable and valid personality assessment methods are available, and an accurate assessment of animal personality could contribute to improvement of animal welfare, and equine welfare in particular, by better matching horses to certain tasks and owners, as well as by use as a selection tool in animal breeding. By these means, equine welfare can either on an individual level or in the long term on a population level be enhanced, as horses will be better suited to their training and husbandry regimens and thus experience less stress. Possibilities to influence personality via early handling or management interference also seem to exist, although little is known about long-term effects. A variety of different personality classification models exist, and there is no consensus on which model is best suited to describe equine personality. The majority of models entail specific advantages that may

be missing in other models and it is therefore argued that the frequent call for more standardized methods may be too early. However, it is suggested that the term personality is the most suitable descriptor of all traits relating to animal behaviour. To avoid confusion with existing personality traits in equine breeding, the term temperament should be reserved for a sub-trait of personality including but not limited to aspects of fear-reactivity.

Acknowledgements

Many thanks to María Díez León, Anne Ricard, Eva Søndergaard, Kathalijne Visser and Janne Winther Christensen for provision of their data, literature and/or help with the literature search. Very special thanks to Aleksandra Górecka-Bruzda for her valuable input and in-depth discussions on the topic.

References

1. Bergmüller R, Taborsky M. Animal personality due to social niche specialisation. *Trends in Ecology and Evolution* 2010;25:504–11.
2. Wolf M, van Doorn GS, Leimar O, Weissing FJ. Life-history trade-offs favour the evolution of animal personalities. *Nature* 2007;447:581–4.

3. Wolf M, Weissing FJ. Animal personalities: consequences for ecology and evolution. *Trends in Ecology and Evolution* 2012;27:452–61.
4. Freeman HD, Gosling SD. Personality in nonhuman primates: a review and evaluation of past research. *American Journal of Primatology* 2010;72:653–71.
5. Svartberg K, Forkman B. Personality traits in the domestic dog (*Canis familiaris*). *Applied Animal Behaviour Science* 2002;79:133–55.
6. Svartberg K, Tapper I, Temrin H, Radesater T, Thorman S. Consistency of personality traits in dogs. *Animal Behaviour* 2005;69:283–91.
7. Gosling SD, Olson Z. Animal personality. *Japanese Journal of Personality* 2008;17:111–9.
8. Houdelier C, Lumineau S, Bertin A, Guibert F, De Margerie E, Augery M, *et al.* Development of fearfulness in birds: genetic factors modulate non-genetic maternal influences. *PLoS ONE* 2011;6:e14604.
9. Koprivnikar J, Gibson CH, Redfern JC. Infectious personalities: behavioural syndromes and disease risk in larval amphibians. *Proceedings of the Royal Society B: Biological Sciences* 2012;279:1544–1550.
10. Raoult V, Brown C, Zuberi A, Williamson J. Blood cortisol concentrations predict boldness in juvenile mullet (*Argyrosomus japonicus*). *Journal of Ethology* 2012;30:225–32.
11. Chervet N, Zöttl M, Schürch R, Taborsky M, Heg D. Repeatability and heritability of behavioural types in a social cichlid. *International Journal of Evolutionary Biology* 2011;2011:15, 10.4061/2011/321729.
12. Briffa M, Rundle SD, Fryer A. Comparing the strength of behavioural plasticity and consistency across situations: animal personalities in the hermit crab *Pagurus bernhardus*. *Proceedings of the Royal Society B: Biological Sciences* 2008;275:1305–11.
13. Mather JA, Anderson RC. Personalities of octopuses (*Octopus rubescens*). *Journal of Comparative Psychology* 1993;107:336–40.
14. Gyuris E, Feró O, Barta Z. Personality traits across ontogeny in firebugs, *Pyrhocoris apterus*. *Animal Behaviour* 2012;84:103–109.
15. Liang ZS, Nguyen T, Mattila HR, Rodriguez-Zas SL, Seeley TD, Robinson GE. Molecular determinants of scouting behavior in honey bees. *Science* 2012;335:1225–8.
16. Gosling SD. From Mice to Men: what can we learn about personality from animal research? *Psychological Bulletin* 2001;127:45–86.
17. Müller R, Schrader L. Behavioural consistency during social separation and personality in dairy cows. *Behaviour* 2005;142:1289–306.
18. Brown JA, Dewey C, Delange CFM, Mandell IB, Purslow PP, Robinson JA, *et al.* Reliability of temperament tests on finishing pigs in group-housing and comparison to social tests. *Applied Animal Behavior Sciences* 2009;118:28–35.
19. Bagnator A, Rossoni A, Nicoletti C, Jakobsen J, Santus E. Milkability and temperament MACE correlation and pilot study in dairy cattle populations. In *Proceedings of the Interbull Meeting*, 23–26 August 2007 Dublin, Ireland; 2007. p. 95–7.
20. King T, Marston LC, Bennett PC. Breeding dogs for beauty and behaviour: why scientists need to do more to develop valid and reliable behaviour assessments for dogs kept as companions. *Applied Animal Behavior Science* 2012;137:1–12.
21. Saetre P, Strandberg E, Sundgren PE, Pettersson U, Jazin E, Bergström TF. The genetic contribution to canine personality. *Genes, Brain and Behavior* 2006;5:240–8.
22. Ley J, Bennett P. Measuring personality in dogs. *Journal of Veterinary Behavior* 2008;3:182.
23. Loving NS. *The Horse as a Companion Animal*. UFAW Animal Welfare Series. Wiley-Blackwell, Oxford; 2011. p. 255–74.
24. Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. Integrating animal temperament within ecology and evolution. *Biological Reviews* 2007;82:291–318.
25. Bates JE. Concepts and measures of temperament. In: Kohnstamm GA, Bates JE, Rothbart MK, editors. *Temperament in Childhood*. Wiley, Chichester, England; 1989. p. 4.
26. Goldsmith HH, Buss AH, Plomin R, Rothbart MK, Thomas A, Chess S, *et al.* Roundtable: what is temperament? four approaches. *Child Development* 1987;58:505–29.
27. Visser EK. *Horsonality: a Study on the Personality of the Horse*. PhD-thesis; Utrecht University, Utrecht, The Netherlands; 2002.
28. Momozawa Y, Ono T, Sato F, Kikusui T, Takeuchi Y, Mori Y, *et al.* Assessment of equine temperament by a questionnaire survey to caretakers and evaluation of its reliability by simultaneous behavior test. *Applied Animal Behavior Science* 2003;84:127–38.
29. Nagy K, Bodó G, Bárdos G, Bánszky N, Kabai P. Differences in temperament traits between crib-biting and control horses. *Applied Animal Behaviour Science* 2009;122:41–7.
30. Peeters M, Verwilghen D, Serteyn D, Vandenheede M. Relationships between young stallions' temperament and their behavioral reactions during standardized veterinary examinations. *Journal of Veterinary Behavior* 2012;7:311–321.
31. Le Scolan N, Hausberger M, Wolff A. Stability over situations in temperamental traits of horses as revealed by experimental and scoring approaches. *Behavioral Process* 1997;41:257–66.
32. Ettl R. *Cutting – Mit Rindern Arbeiten*. Müller Rüschlikon, Cham, Switzerland; 1998. p. 181.
33. Budzyński M. Timidity test to assess the degree of nervousness of horses. *Medycyna Weterynaryjna* 1984;40:156–8.
34. Sapula M, Kamieniak J, Budzyńska M, Gancarz J, Stefaniuk A. Evaluation of working abilities and nervous irritability in Małopolski stallions considering influence of sire breed. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 2002;20:249–55.
35. Budzyński M. Application of a sound test to evaluate the degree of nervousness of horses. *Medycyna Weterynaryjna* 1983;39:361–2.
36. Kamieniak J, Sapula M, Budzyński M, Budzyńska M, Sołtys L. The effect of nervous balance on the quality of stallions' behavioural reactions in colour perception test. *Annales*

- Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica 2004;22:207–14.
37. Baiguina EA, Gorbukov MA. Typological Particularities of Higher Nervous Activity of Russian Punch Stallions. National Academy of Science, Minsk; 2002. p. 125–8.
 38. Kamieniak J. Estimation of nervous irritability indexes conditioned by inbreeding level of purebred Arabian horses. I. The influence of inbreeding on the features of nervous irritability and race performance. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 1999;17:233–42.
 39. McCann JS, Heird JC, Bell RW, Lutherer LO. Normal and more highly reactive horses. I. Heart rate, respiration rate and behavioral observations. *Applied Animal Behavior Science* 1988;19:201–14.
 40. McCann JS, Heird JC, Bell RW, Lutherer LO. Normal and more highly reactive horses. II. The effect of handling and reserpine on the cardiac response to stimuli. *Applied Animal Behavior Science* 1988;19:215–26.
 41. Wolff A, Hausberger M, Le Sclan N. Experimental tests to assess emotionality in horses. *Behavioral Processes* 1997;40:209–21.
 42. Jönsson L, Viklund Å, Höög Y, Rundgren M, Philipsson J. Evaluation of mental traits at young horse performance test. In 61st Annual Meeting of the European Association for Animal Production, Crete, Greece; 2010. p. 150.
 43. Kamieniak J, Budzyński M, Sapuła M, Budzyńska M, Krupa W, Sołtys L. Level of mental traits of Arabian horses' progeny. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 2006;24:191–8.
 44. Sommer H, Barz A, Lindner A. Testing horses for character and temperament. *Tierärztliche Umschau* 1996;51:641–3.
 45. Schwark HJ, Petzold P, Karwath M. Studies on the objectivity and breeding efficiency of station performance tests on stallions of riding horse breeds. *Archiv für Tierzucht* 1992;35:97–106.
 46. Koolhaas JM, Korte SM, De Boer SF, Van Der Veegt BJ, Van Reenen CG, Hopster H, *et al.* Coping styles in animals: current status in behavior and stress-physiology. *Neuroscience and Biobehavioral Reviews* 1999;23:925–35.
 47. Seaman SC, Davidson HPB, Waran NK. How reliable is temperament assessment in the domestic horse (*Equus caballus*)? *Applied Animal Behavior Science* 2002;78:175–91.
 48. Goldberg LR. An alternative description of personality: the big-five factor structure. *Journal of Personality and Social Psychology* 1990;59:1216–29.
 49. Digman JM. Personality structure: emergence of the five-factor model. *Annual Review of Psychology* 1990;41:417–40.
 50. Morris PH, Gale A, Duffy K. Can judges agree on the personality of horses? *Personality and Individual Differences* 2002;33:67–81.
 51. Morris PH, Gale A, Howe S. The factor structure of horse personality. *Anthrozoos: A Multidisciplinary Journal of The Interactions of People and Animals* 2002;15:300–22.
 52. Cloninger CR, Svrakic DM, Przybeck TR. A psychobiological model of temperament and character. *Archives of General Psychiatry* 1993;50:975–90.
 53. Cloninger R. A unified biosocial theory of personality and its role in the development of anxiety states. *Psychiatric Developments* 1986;3:167–226.
 54. Momozawa Y, Kusunose R, Kikusui T, Takeuchi Y, Mori Y. A Search for Temperament-Associated Genetic Polymorphisms in Horses. *Purdue University Press, Indiana*; 2005. p. 92–7.
 55. Momozawa Y, Takeuchi Y, Kusunose R, Kikusui T, Mori Y. Association between equine temperament and polymorphisms in dopamine D4 receptor gene. *Mammalian Genome* 2005;16:538–44.
 56. Momozawa Y, Takeuchi Y, Tozaki T, Kikusui T, Hasegawa T, Raudsepp T, *et al.* SNP detection and radiation hybrid mapping in horses of nine candidate genes for temperament. *Animal Genetics* 2007;38:81–3.
 57. König von Borstel U, Euent S, Graf P, König S, Gauly M. Equine behaviour and heart rate in temperament tests with or without rider or handler. *Physiology and Behavior* 2011;104:454–63.
 58. Graf P, König von Borstel U, Gauly M. Interieurmerkmale für die Zuchtwertschätzung. In *Proceedings of the Göttinger Pferdetage '11*, 2011. p. 87–8.
 59. Graf P, Sitzenstock F, König V, Borstel U, Gauly M. "Genetische Parameter von Merkmalen eines Temperamenttests bei Pferden". *Proceedings of the Göttinger Pferdetage '13*; Göttingen, Germany. 2013; in press.
 60. Hausberger M, Sclan NI, Bruderer C, Pierre JS. Temperament in the horse: factors in play and practical implications. In French. *Proceedings of the 24th equine research day. Institut de Cheval, Paris, France*; 1998;159–69.
 61. Visser EK, van Reenen CG, Hopster H, Schilder MBH, Knaap JH, Barneveld A, *et al.* Quantifying aspects of young horses' temperament: consistency of behavioural variables. *Applied Animal Behavior Science* 2001;74:241–58.
 62. Visser EK, Van Reenen CG, Engel B, Schilder MBH, Barneveld A, Blokhuis HJ. The association between performance in show-jumping and personality traits earlier in life. *Applied Animal Behavior Science* 2003;82:279–95.
 63. Visser EK, Van Reenen CG, Rundgren M, Zetterqvist M, Morgan K, Blokhuis HJ. Responses of horses in behavioural tests correlate with temperament assessed by riders. *Equine Veterinary Journal* 2003;35:176–83.
 64. Lloyd AS, Martin JE, Bornett-Gauci HLI, Wilkinson RG. Evaluation of a novel method of horse personality assessment: rater-agreement and links to behaviour. *Applied Animal Behavior Science* 2007;105:205–22.
 65. Lloyd AS, Martin JE, Bornett-Gauci HLI, Wilkinson RG. Horse personality: variation between breeds. *Applied Animal Behavior Science* 2008;112:369–83.
 66. McGrogan C, Hutchison MD, King JE. Dimensions of horse personality based on owner and trainer supplied personality traits. *Applied Animal Behavior Science* 2008;113:206–14.
 67. Napolitano F, De Rosa G, Braghieri A, Grasso F, Bordini A, Wemelsfelder F. The qualitative assessment of responsiveness to environmental challenge in horses and ponies. *Applied Animal Behavior Science* 2008;109:342–54.
 68. Minero M, Tosi MV, Canali E, Wemelsfelder F. Quantitative and qualitative assessment of the response of foals to the presence of an unfamiliar human. *Applied Animal Behavior Science* 2009;116:74–81.

69. Creighton E. Matching horses for courses: development of robust tests of equine temperament in riding horses and ponies. *Animal Welfare* 2003;13(s1):S239.
70. Koenen EPC, Aldridge LI, Philipsson J. An overview of breeding objectives for warmblood sport horses. *Livestock Production Science* 2004;88:77–84.
71. Pasing S. Analyse des Status Quo und Ansätze zur Verbesserung der Interieurbeurteilung deutscher Pferdezuchten. MSc-thesis, University of Göttingen, Germany; 2010. p. 139.
72. Ruch W. Pavlov's types of nervous system, Eysenck's typology and the Hippocrates-Galen temperaments: an empirical examination of the asserted correspondence of three temperament typologies. *Personality and Individual Differences* 1992;13:1259–71.
73. Lansade L, Bertrand M, Bouissou M-F. Effects of neonatal handling on subsequent manageability, reactivity and learning ability of foals. *Applied Animal Behavior Science* 2005;92:143–58.
74. Uher J. Three methodological core issues of comparative personality research. *European Journal of Personality* 2008;22:475–96.
75. Levitis DA, Lidicker WZ, Freund G. Behavioural biologists don't agree on what constitutes behaviour. *Animal Behavior* 2009;78:103–10.
76. Visser K, Munsters C, Roost L, McGreevy P, Heleski C. Trainers and scientists differ when scoring stress-related behaviours in ridden horses. In 8th International Society for Equitation Science, Edinburgh, UK; 2012. p. 121.
77. Diverio S, Tami G, Marchei P, Tortiello C, Catalani MC, Barone A. Assessing horse performance: a comparison between trainers' and judges' behavior evaluations. *Journal of Veterinary Behavior* 2010;5:53–4.
78. Forkman B, Boissy A, Meunier-Salaün MC, Canali E, Jones RB. A critical review of fear tests used on cattle, pigs, sheep, poultry and horses. *Physiology and Behavior* 2007;92:340–74.
79. Janczak AM, Pedersen LJ, Bakken M. Aggression, fearfulness and coping styles in female pigs. *Applied Animal Behavior Science* 2003;81:13–28.
80. König v. Borstel U, Pirsich W, Gauly M, Bruns E. Repeatability and reliability of scores from ridden temperament tests conducted during performance tests. *Applied Animal Behavior Science* 2012;139:251–63.
81. Jensen P, Rushen J, Forkman B. Behavioural strategies or just individual variation in behaviour? – A lack of evidence for active and passive piglets. *Applied Animal Behavior Science* 1995;43:135–9.
82. De Fruyt F, De Clercq BJ, van de Wiele L, Van Heeringen K. The validity of Cloninger's psychobiological model versus the five-factor model to predict DSM-IV personality disorders in a heterogeneous psychiatric sample: domain facet and residualized facet descriptions. *Journal of Personality* 2006;74:479–510.
83. Ekelund J, Lichtermann D, Johnson LP, Wieckert DA. Association between novelty seeking and the type 4 dopamine receptor gene in a large Finnish cohort sample. *American Journal of Psychiatry* 1999;156:1453–5.
84. Manuck S, Flory J, Ferrell R, Dent K, Mann J, Muldoon M. Aggression and anger-related traits associated with a polymorphism of the tryptophan hydroxylase gene. *Biological Psychology* 1999;45:603–14.
85. Kumakiri C, Kodama K, Shimizu E, Yamanouchi N, Okada S, Noda S, *et al.* Study of the association between serotonin transporter gene regulatory region polymorphism and personality traits in a Japanese population. *Neuroscience Letters* 1999;263:205–7.
86. Flisikowski K, Schwarzenbacher H, Wysocki M, Weigend S, Preisinger R, Kjyer JB, *et al.* Variation in neighbouring genes of the dopaminergic and serotonergic systems affects feather pecking behaviour of laying hens. *Animal Genetics* 2008;40:192–9.
87. Schmutz SM, Stookey JM, Winkelman-Sim DC, Waltz CS, Plante Y, Buchanan FC. A QTL study of cattle behavioral traits in embryo transfer families. *Journal of Heredity* 2001;92:290–2.
88. Kreinberg P. Verhalten u. Interieur- Bedeutung für die praktische Pferdeausbildung. Göttinger Pferdetage, Göttingen, Germany; FN-Verlag, Warendorf 2009. p. 48–53.
89. Boissy A, Bouissou MF. Assessment of individual differences in behavioural reactions of heifers exposed to various fear-eliciting situations. *Applied Animal Behavior Science* 1995;46:17–31.
90. Boyle D, Gerberich SG, Gibson RW, Maldonado G, Robinson RA, Martin F, *et al.* Injury from dairy cattle activities. *Epidemiology* 1997;8:37–41.
91. D'Eath RB, Turner SP, Kurt E, Evans G, Thölking L, Looft H, *et al.* Pigs' aggressive temperament affects pre-slaughter mixing aggression, stress and meat quality. *Animal* 2010;4:604–16.
92. Voisinnet BD, Grandin T, O'Connor SF, Tatum JD, Deesing MJ. *Bos indicus*-cross feedlot cattle with excitable temperaments have tougher meat and a higher incidence of borderline dark cutters. *Meat Science* 1997;46:367–77.
93. Debut M, Berri C, Baeza E, Sellier N, Arnould C, Guemene D, *et al.* Variation of chicken technological meat quality in relation to genotype and preslaughter stress conditions. *Poultry Science* 2003;82:1829–38.
94. Duffield TF, Heinrich A, Millman ST, DeHaan A, James S, Lissemore K. Reduction in pain response by combined use of local lidocaine anesthesia and systemic ketoprofen in dairy calves dehorned by heat cauterization. *Canadian Veterinary Journal (La Revue Veterinaire Canadienne)* 2010;51:283–8.
95. Price J, Eager RA, Welsh EM, Waran NK. Current practice relating to equine castration in the UK. *Research in Veterinary Science* 2005;78:277–80.
96. Coetzee JF, Nutsch AL, Barbur LA, Bradburn RM. A survey of castration methods and associated livestock management practices performed by bovine veterinarians in the United States. *BMC Veterinary Research* 2010;6:12.
97. Taylor NR, Main DC, Mendl M, Edwards SA. Tail-biting: a new perspective. *Veterinary Journal* 2010;186:137–47.
98. Jendral MJ, Robinson FE. Beak trimming in chickens: historical, economical, physiological and welfare implications, and alternatives for preventing feather pecking and cannibalistic activity. *Avian and Poultry Biology Reviews* 2004;15:9–23.
99. Dwyer CM. Individual variation in the expression of maternal behaviour: a review of the neuroendocrine mechanisms in the sheep. *Journal of Neuroendocrinology* 2008;20:526–34.

100. Hare B, Brown M, Williamson C, Tomasello M. The domestication of social cognition in dogs. *Science* 2002;298:1634–6.
101. Haverbeke A, Laporte B, Depiereux D, Giffroy JM, Claire D, Diederich C. Training methods of military dog handlers and their effects on the team's performances. *Applied Animal Behavior Science* 2008;113:110–122.
102. Meyer F, Schawalder P, Gaillard C, Dolf G. Estimation of genetic parameters for behavior based on results of German Shepherd dogs in Switzerland. *Applied Animal Behavior Science* 2012;140:53–61.
103. von Borstel U. Verhaltenstests beim Pferd. Göttinger Pferdetage '09 Zucht und Haltung von Sportpferden, Göttingen, Germany; FN-Verlag, Warendorf. 2009. p. 54–66.
104. Könnel J. Vergleich der Durchführung und Ergebnisermittlung der Hengstleistungsprüfung in Europa sowie den USA und ihre Anerkennung in Deutschland. MSc-thesis, University of Göttingen, Göttingen, Germany; 2009. p. 101.
105. Catalano AL, Martuzzi F, Filippini S, Simonini FV. Performance test of Bardigiano breed stallions and mares for saddle and harness service. *Annali della Facoltà di Medicina Veterinaria, Università di Parma* 2006;26:119–26.
106. Teegen R, Edel C, Thaller G. Evaluation of the breeding objectives for the Trakehner Horse breed by a contingent valuation method (CV). In German; English abstract. *Züchtungskunde*. 2008;80:99–113.
107. Górecka-Bruzda A, Jezierski T. Breed differences in behaviour-related characteristics of stallions evaluated in performance tests. *Animal Science Papers and Reports* 2010;28:27–36.
108. Bene S, Giczi A, Szabó F. Performance test results of stallions of different breeds between 1998–2010 in Hungary. 1st paper. The warm blood breeds in cart. *Állattenyésztés és Takarmányozás* 2012;61:1–16.
109. Viklund Å, Thorén Hellsten E, Näsholm A, Strandberg E, Philipsson J. Genetic parameters for traits evaluated at field tests of 3- and 4-year-old Swedish Warmblood horses. *Animal* 2008;2:1832–41.
110. Huizinga HA, van der Werf JHJ, Korver S, van der Meij GJW. Stationary performance testing of stallions from the Dutch Warmblood riding horse population. 1. Estimated genetic parameters of scored traits and the genetic relation with dressage and jumping competition from offspring of breeding stallions. *Livestock Production Science* 1991;27:231–44.
111. Ohlsson L, Arnason T, Philipsson J. Quality evaluation of 4-year-old riding horses valuable performance indicator. *Fakta Husdjur* 1994;13:4–8.
112. Górecka A, Bakuniak M, Jezierski T. Behaviour of horses during habituation to a novel object. In *Proceedings of the 2nd International Equitation Science Symposium*. Fondazione Iniziative Zoorofilattiche e Zootechniche, Milano, Italy; 2006. p. 33.
113. Voith VL. Principles of learning. *Veterinary Clinics of North America Equine Practice* 1986;2:485–506.
114. Christensen JW, Rundgren M, Olsson K. Training methods for horses: habituation to a frightening stimulus. *Equine Veterinary Journal* 2006;38:439–43.
115. Górecka A, Bakuniak M, Chruszczewski MH, Jezierski TA. A note on the habituation to novelty in horses: handler effect. *Animal Science Papers and Reports* 2007;25:143–52.
116. König von Borstel U, Duncan IJH, Claesson-Lundin M, Keeling LJ. Fear reactions in trained and untrained horses from dressage and show-jumping breeding lines. *Applied Animal Behavior Science* 2010;125:124–31.
117. Leiner L, Fendt M. Behavioural fear and heart rate responses of horses after exposure to novel objects: effects of habituation. *Applied Animal Behavior Science* 2011;131:104–9.
118. Christensen JW, Zharkikh T, Ladewig J. Do horses generalise between objects during habituation? *Applied Animal Behavior Science* 2008;114:509–20.
119. Bruns E, Bade B, Haring H. Results on a more objective measurement of performance traits of stallions in performance testing at station. *Livestock Production Science* 1980;7:607–14.
120. Budzynski M. Repeatability and heritability of the assesment of nervous type in horses. *Annales Universitatis Mariae Curie-Skłodowska, EE (Zootechnica)* 1983;1:229–32.
121. McDonnell SM, Diehl NK. Computer-assisted recording of live and videotaped horse behavior: reliability studies. *Applied Animal Behavior Science* 1990;27:1–7.
122. Huizinga HA, Korver S, van der Meij GJW. Stationary performance testing of stallions from the Dutch Warmblood riding horse population. 2. Estimated heritabilities of and correlations between successive judgements of performance traits. *Livestock Production Science* 1991;27:245–54.
123. Budzynski M, Słomka Z, Sołtys L, Budzynska M, Pałyszka J. Estimation of the breeding value of purebred Arab stallions for excitability of their offspring during behavioural tests. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 1998;16:165–79.
124. Mills DS. Personality and individual differences in the horse, their significance, use and measurement. *Equine Veterinary Journal Supplement* 1998;30:10–3.
125. Anderson MK, Friend TH, Evans JW, Bushong DM. Behavioral assessment of horses in therapeutic riding programs. *Applied Animal Behavior Science* 1999;63:11–24.
126. Hada T, Onaka T, Kusunose R, Yagi K. Effects of novel environmental stimuli on neuroendocrine activity in Thoroughbred horses. *Journal of Equine Science* 2001;12:33–8.
127. Visser EK, van Reenen CG, van der Werf JTN, Schilder MBH, Knaap JH, Barneveld A, *et al.* Heart rate and heart rate variability during a novel object test and a handling test in young horses. *Physiology and Behavior* 2002;76:289–96.
128. Visser EK, van Reenen CG, Schilder MBH, Barneveld A, Blokhuis HJ. Learning performances in young horses using two different learning tests. *Applied Animal Behavior Science* 2003;80:311–26.
129. Momozawa Y, Kusunose R, Kikusui T, Takeuchi Y, Mori Y. Assessment of equine temperament questionnaire by comparing factor structure between two separate surveys. *Applied Animal Behavior Science* 2005;92:77–84.
130. Christensen JW, Keeling LJ, Nielsen BL. Responses of horses to novel visual, olfactory and auditory stimuli. *Applied Animal Behavior Science* 2005;93:53–65.

131. McCall CA, Hall S, McElhenney WH, Cummins KA. Evaluation and comparison of four methods of ranking horses based on reactivity. *Applied Animal Behavior Science* 2006;96:115–27.
132. Oki H, Kusunose R, Nakaoka H, Nishiura A, Miyake T, Sasaki Y. Estimation of heritability and genetic correlation for behavioural responses by Gibbs sampling in the Thoroughbred racehorse. *Journal of Animal Breeding and Genetics* 2007;124:185–91.
133. Lansade L, Bouissou M-F. Reactivity to humans: a temperament trait of horses which is stable across time and situations. *Applied Animal Behavior Science* 2008;114:492–508.
134. Lansade L, Bouissou M-F, Erhard HW. Reactivity to isolation and association with conspecifics: a temperament trait stable across time and situations. *Applied Animal Behavior Science* 2008;109:355–73.
135. Lansade L, Pichard G, Leconte M. Sensory sensitivities: components of a horse's temperament dimension. *Applied Animal Behavior Science* 2008;114:534–53.
136. Lansade L, Bouissou M-F, Erhard HW. Fearfulness in horses: a temperament trait stable across time and situations. *Applied Animal Behavior Science* 2008;115:182–200.
137. Visser K, Karlas K, Van Deurzen I, Workel I, Van Reenen K. Experts' assessment of temperament in sport horses. *Journal of Veterinary Behavior* 2010;5:214–5.
138. Górecka-Bruzda A, Jastrzebska E, Sosnowska Z, Jaworski Z, Jezierski T, Chruszczewski MH. Reactivity to humans and fearfulness tests: field validation in Polish Cold Blood Horses. *Applied Animal Behavior Science* 2011;133:207–15.
139. König von Borstel U, Pasing S, Gauly M. Towards a more objective assessment of equine personality using behavioural and physiological observations from performance test training. *Applied Animal Behavior Science* 2011;135:277–85.
140. König v. Borstel U, Peinemann V, Glißmann C, Euent S. Willing to work? Suitability of different riding situations in evaluation of equine personality. In 8th International Equitation Science Conference; 18th–20th July 2012, Edinburgh, UK; 2012. p. 129.
141. Christensen JW, Ahrendt LP, Lintrup R, Gaillard C, Palme R, Malmkvist J. Does learning performance in horses relate to fearfulness, baseline stress hormone, and social rank? *Applied Animal Behavior Science* 2012;140:44–52.
142. Theis S, Adita E, Hille K, König von Borstel U, Gauly M. Intra-observer reliability of different methods for recording temperament in beef and dairy calves. In 62nd Annual Meeting of the European Association for Animal Production, Bratislava, Slovakia; 2012. p. 175.
143. Kilgour R. The open-field test as an assessment of the temperament of dairy cows. *Animal Behaviour* 1975;23(Part 3):615–24.
144. Barnard S, Siracusa C, Reisner I, Valsecchi P, Serpell JA. Validity of model devices used to assess canine temperament in behavioral tests. *Applied Animal Behavior Science* 2012;138:79–87.
145. Duffy DL, Serpell JA. Predictive validity of a method for evaluating temperament in young guide and service dogs. *Applied Animal Behavior Science* 2012;138:99–109.
146. Serpell JA, Hsu Y. Development and validation of a novel method for evaluating behavior and temperament in guide dogs. *Applied Animal Behavior Science* 2001;72:347–64.
147. Sinn DL, Gosling SD, Hilliard S. Personality and performance in military working dogs: reliability and predictive validity of behavioral tests. *Applied Animal Behavior Science* 2010;127:51–65.
148. D'Eath RB. Consistency of aggressive temperament in domestic pigs: the effects of social experience and social disruption. *Aggressive Behavior* 2004;30:435–48.
149. Gibbons JM, Lawrence AB, Haskell MJ. Consistency of flight speed and response to restraint in a crush in dairy cattle. *Applied Animal Behavior Science* 2011;131:15–20.
150. Åkerberg H, Wilsen E, Sallander M, Hedhammar Å, Lagerstedt A-S, Larhammar D, *et al.* Test for personality characteristics in dogs used in research. *Journal of Veterinary Behavior* 2012;7:327–38.
151. Rauw WM, Kanis E, Noordhuizen-Stassen EN, Grommers FJ. Undesirable side effects of selection for high production efficiency in farm animals: a review. *Livestock Production Science* 1998;56:15–33.
152. French JM. Assessment of donkey temperament and the influence of home environment. *Applied Animal Behavior Science* 1993;36:249–57.
153. Flentje R, Creighton E. Can standardized behavior tests predict suitability for use in horses? *Journal of Veterinary Behavior* 2010;5:58–9.
154. Grajfoner DD, Austin EJ, Wemelsfelder F. Horse personality profiles and performance. *Journal of Veterinary Behavior* 2010;5:59.
155. Christmann L. Hannoveraner Erleben – Pferde für die Freizeit. *Der Hannoveraner* 2005;9:23–5.
156. Cavallo, FN (German Equestrian Federation). editors. GHP – Gelassenheitsprüfung für Sport- und Freizeitpferde. FN-Verlag/Redaktion Cavallo, Warendorf/Stuttgart; 2007.
157. König v. Borstel U, Krienert N. Influence of familiarity with the rider and type of work on horses' fear reactions. In 8th International Society for Equitation Science; Edinburgh, UK; 2012. p. 153.
158. Kaiser D. Mit eingebautem Uhrwerk – Die Anforderungen an Voltigierpferde sind in den vergangenen Jahren stetig gestiegen. *Der Hannoveraner* 2012;9:28–9.
159. Nicosia DMSID, Bacci ML. Study of therapeutic riding horses: a review. *Ippologia* 2009;20:9–14.
160. Wolframm IA, Meulenbroek RGJ. Co-variations between perceived personality traits and quality of the interaction between female riders and horses. *Applied Animal Behavior Science* 2012;139:96–104.
161. Zetterqvist M, Van Reenen CG, Visser EK, Hassmén P, Morgan K, Rundgren M, *et al.* Assessment of Co-Operation between the Rider and the Horse: the Relationship between Ratings Assigned by Riders and those Assigned by an External Judge. In Proceedings of the 37th International Congress of the International Society for Applied Ethology, Abano Terme, Italy; 2003. p. 56.
162. Visser EK, Reenen CGv, Blokhuis MZ, Morgan EKM, Hassmén P, Rundgren TMM, *et al.* Does horse temperament influence horse-rider cooperation? *Journal of Applied Animal Welfare Science* 2008;11:267–84.

163. Hemsworth PH, Coleman GJ. Human-livestock Interactions: The Stockperson and the Productivity and Welfare of Intensively Farmed Animals. CABI, Wallingford; 2011. ix + 187 pp.
164. Wolframm IA, Micklewright D. Rider anxiety, perception of equine temperament and ridden performance: do they relate? *Journal of Veterinary Behavior* 2010;5:210.
165. von Borstel UU. Fear in Horses and Riders: Two Hearts Living in just One Mind – The Influence of Rider, Training and Genetics on horses' fear. Verlag Dr Müller, Saarbrücken, Germany; 2008.
166. Bridgeman DJ. The working relationship between horse and rider during training and competition for equestrian sports. PhD-thesis, University of Southern Queensland, Australia; 2009. p. 278.
167. Hausberger M, Roche H, Henry S, Visser EK. A review of the human-horse relationship. *Applied Animal Behavior Science* 2008;109:1–24.
168. Munsters CCBM, Visser KEK, van den Broek J, Sloet van Oldruitenborgh-Oosterbaan MM. The influence of challenging objects and horse-rider matching on heart rate, heart rate variability and behavioural score in riding horses. *Veterinary Journal* 2012;192:75–80.
169. Turcsán B, Range F, Virányi Z, Miklósi Á, Kubinyi E. Birds of a feather flock together? Perceived personality matching in owner–dog dyads. *Applied Animal Behavior Science* 2012;140:144–60.
170. Watson D, Klohnen EC, Casillas A, Nus Simms E, Haig J, Berry DS. Match makers and deal breakers: analyses of assortative mating in newlywed couples. *Journal of Personality* 2004;72:1029–68.
171. Rodenburg TB, Turner SP. The role of breeding and genetics in the welfare of farm animals. *Animal Frontiers* 2012;2:16–21.
172. Kaiser H, Geuder U. 4.7. Pferdezucht. In: Götz K-U, Buitkamp J, editors. *Jahresbericht 2011: Bayerische Landesanstalt für Landwirtschaft*. Freising-Weihenstephan, Germany; 2012. p. 58–72.
173. König von Borstel U, Bernhard V. Impact of health disorders on functional and biological longevity in warmblood breeding stallions. *The Veterinary Journal* 2012;In press.
174. Björnsdóttir S, Arnason T, Lord P. Culling rate of Icelandic horses due to bone spavin. *Acta Veterinaria Scandinavica* 2003;44:161–9.
175. Clausen M, Preisinger R, Kalm E. Analyse von Krankheitsdaten in der deutschen Warmblutzucht. *Züchtungskunde* 1990;62:167–78.
176. Egenvall A, Lonnell C, Roepstorff L. Analysis of morbidity and mortality data in riding school horses, with special regard to locomotor problems. *Preventive Veterinary Medicine* 2009;88:193–204.
177. Hommerich G. Ausfallursachen und -frequenzen laut Schadensstatistik der Vereinigte Tierversicherung Gesellschaft a.G. Göttinger Pferdetage, Göttingen, Germany; 1995. p. 85–93.
178. Hayek AR, Jones B, Evans DL, Thomson PC, McGreevy P. Epidemiology of horses leaving the Thoroughbred and Standardbred racing industries. In 1st International Equitation Science Symposium 2005, Melbourne, Australia; 2005. p. 84–8.
179. Wallin L, Strandberg E, Philipsson J, Dalin G. Estimates of longevity and causes of culling and death in Swedish warmblood and coldblood horses. *Livestock Production Science* 2000;63:275–89.
180. Dusek J. Estimating heritability of some performance characters and body conformation of horses in small populations. *Zivocisna Vyroba* 1980;25:71–80.
181. Brockmann A, Bruns E. Estimation of genetic parameters for performance traits of riding horses. *Züchtungskunde* 2000;72:4–16.
182. Pasing S, Christmann L, Gauly M, König von Borstel U. Analysis of the status quo of personality evaluation in German horse breeding. In 62nd Annual Meeting of the European Association for Animal Production, Stavanger, Norway; 2011. p. 142.
183. Pasing S, König v. Borstel U. Evidence of grade inflation in personality trait scores from stallion performance tests. In 8th International Equitation Science Conference, 18–20 July 2012, Edinburgh, UK; 2012. p. 125.
184. Graf P, König von Borstel U, Gauly M. Implementation of equine temperament tests in performance test on station and field. In 61st Annual Meeting of the European Association for Animal Production 2010, 23–27 August, Crete, Greece; 2010. p. 151.
185. König von Borstel U, Euent S, Graf P, Gauly M. Should applied temperament tests in horses be conducted with or without a human handler or rider? In 61st Annual Meeting of the European Association for Animal Production, 23–27 August 2010, Crete, Greece; 2010. p. 150.
186. Graf P, Schneider T, König v. Borstel U, Gauly M. Economic evaluation of an objective temperament assessment in horses. *Züchtungskunde*. 2012; accepted (In German, English abstract) in press.
187. Baumgartner M, Burger D, Jallon L, Gerber V, Hagger C, Rieder S. Héritabilité de l'émotivité chez le cheval des Franches-Montagnes. *Schweiz Arch Tierheilk* 2010;152:194.
188. Lansade L, Dubois C, Gillot A, Delfosse A, Bon MI, Roche H, et al. An example of application of research in ethology from the horse's temperament tests and record of learning and behavior (LAC). In Proceedings of the 36th Equine Research Day, Le Pin au Haras, France; 2010;177–85.
189. Keeling LJ, Blomberg A, Ladewig J. Horse-riding accidents: When the human-animal relationship goes wrong! In Proceedings of the 33rd International Congress of the International Society for Applied Ethology, Lillehammer, Norway, 17–21 August 1999; 1999. p. 86.
190. Graf P. Akzeptanz und Eignung von Temperamenttests zur Integration in Pferdezuchtprogramme unter Berücksichtigung genetischer Parameter. PhD-thesis, University of Göttingen, Germany. 2012 in press.
191. Federici M, Gerber V, Haase B, Klopfenstein M, Poncet P-A, Burger D. Effekt von Grundfarbe und weißen Abzeichen auf die Gesundheit und die Vermarktung von Freibergpferden. *Schweiz Arch Tierheilk* 2010;152:197.
192. Gille C, Spiller A. Customer oriented horse breeding in Germany. Target group segmentation: an empirical analysis. *Züchtungskunde* 2010;82:229–40.
193. Gille C, Kayser M, Spiller A. Target group segmentation in the horse buyers' market against the background of equestrian experience. *Journal of Equine Science* 2010;21:67–72.

194. Górecka-Bruzda A, Chruszczewski MH, Jaworski Z, Golonka M, Jezierski T. Looking for an ideal horse: rider preferences. *Anthrozoös* 2011;24:379–92.
195. Hennessy KD, Quinn KM, Murphy J. Producer or purchaser: different expectations may lead to equine wastage and welfare concerns. *Journal of Applied Animal Welfare Science* 2008;11:232–5.
196. Bruns E, Bierbaum M, Frese D, Haring HJF. The development of selection criteria for the breeding of riding horses. IV. Estimation of relative economic values using sale prices. *Züchtungskunde* 1978;50:93–100.
197. Olsson K. A Review of Methods Used to Measure Temperamental Characteristics in Horses. BSc-thesis, Swedish University of Agricultural Science, Skara, Sweden; 2010.
198. Hawson LA, Oddie C, McLean AN, McGreevy PD. Is safety valued in the Australian pony market? *Journal of Veterinary Behavior* 2011;6:254–60.
199. Couzy C, Godet J. Mismatch between breeding elite sport horses and rider amateurs expectations. In 61st Annual Meeting of the European Association for Animal Production, 23–27 August 2010, Heraklion Greece; 2010. p. 341.
200. Axel-Nilsson M, Nyman S, Visser K, Blokhuis HJ. Do dressage and show jumping riders in Sweden differ in perception of optimal horse temperament? In 7th International Symposium on Equitation Science; Hooge Mierde, The Netherlands; 2011. p. 22.
201. Belyaev DK, Plyusnina IZ, Trut LN. Domestication in the silver fox (*Vulpes fulvus* Desm): changes in physiological boundaries of the sensitive period of primary socialization. *Applied Animal Behavior Science* 1985;13:359–70.
202. Trut LN, Plyusnina IZ, Oskina IN. An experiment on fox domestication and debatable issues of evolution of the dog. *Russian Journal of Genetics* 2004;40:644–55.
203. Hausberger M, Bruderer C, Scolan NL, Pierre J-S. Interplay between environmental and genetic factors in temperament/personality traits in horses (*Equus caballus*). *Journal of Comparative Psychology* 2004;118:434–46.
204. Lesimple C, Fureix C, LeScolan N, Richard-Yris M-A, Hausberger M. Housing conditions and breed are associated with emotionality and cognitive abilities in riding school horses. *Applied Animal Behavior Science* 2011;129:92–9.
205. Mader DR, Price EO. Discrimination learning in horses: effects of breed, age and social dominance. *Journal of Animal Science* 1980;50:962–5.
206. König von Borstel U, Duncan IJH, Lundin MC, Keeling LJ. Fear reactions in trained and untrained horses from dressage and show-jumping breeding lines. *Applied Animal Behavior Science* 2010;125:124–31.
207. Kamieniak J, Budzyński M, Sapuła M, Budzyńska M, Krupa W, Soltys L. Ethological variety of stud Małopolski horses considering genealogical lines. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 2005;23: 169–76.
208. Wolff A, Hausberger M. Behaviour of foals before weaning may have some genetic basis. *Ethology* 1994;96:1–10.
209. Wolff A, Hausberger M. Learning and memorisation of two different tasks in horses: the effects of age, sex and sire. *Applied Animal Behavior Science* 1995;46:137–43.
210. Wolff A, Hausberger M. Learning and memorisation of two different tasks in horses: the effects of age, sex and sire. *Applied Animal Behavior Science* 1996;46:137–43.
211. Warren-Smith AK, McLean AN, Nicol HI, McGreevy PD. Variations in the timing of reinforcement as a training technique for foals (*Equus caballus*). *Anthrozoös* 2005;18:255–72.
212. Dušek J. The effect of inbreeding on some physiological traits of black Kladruby horses. *Veterinární Medicina* 1980;25:349–58.
213. Cape L, van Vleck LD. Heritability of training scores for standardbred yearlings. *J Hered* 1981;72:437.
214. Budzynski M, Soltys L, Słomka Z, Rudzinski K, Kamieniak J, Chmiel K. Heritability of nervous balance in horses. *Annales Universitatis Mariae Curie-Skłodowska Sectio EE Zootechnica* 1994;12:121–7.
215. Ricard A. The Temperament of the Horse: Variation in Genetic Factors. Les Haras Nationaux Direction du Développement, Paris; 2004. p. 141–53.
216. Arnason TH. Studies on traits in the Icelandic toelter horses. I. Estimation of some environmental effects and genetic parameters. *Journal of Agricultural Research in Iceland* 1979;11:81–93.
217. Arnason T. Genetic studies on conformation and performance of Icelandic toelter horses. I. Estimation of non-genetic effects and genetic parameters. *Acta Agriculturae Scandinavica* 1984;34:409–27.
218. Bruns E, Bade B. Estimation of genetic parameters for traits from performance testing of stallions at station. 1984. p. 2
219. Nissen T, Kalm E. Analysis of station tests of Holstein mares. *Züchtungskunde* 1986;58:449–64.
220. Huizinga HA, Boukamp M, Smolders G. Estimated parameters of field performance testing of mares from the Dutch Warmblood riding horse population. *Livestock Production Science* 1990;26:291–9.
221. Christmann L. Zuchtwertschätzung für Merkmale der Stutbuchaufnahme und der Stutenleistungsprüfung im Zuchtgebiet Hannover. PhD-thesis, Georg-August-Universität Göttingen, Göttingen; 1996.
222. Schade W. Entwicklung eines Besamungszuchtprogrammes für die hannoversche Warmblutzuht. PhD-thesis, University of Göttingen, Göttingen, Germany; 1996.
223. Samoré AB, Pagnacco G, Miglior F. Genetic parameters and breeding values for linear type traits in the Haflinger horse. *Livestock Production Science* 1997;52:105–11.
224. Brockmann A. Entwicklung einer Eigenleistungsprüfung im Feld für Hengste unter Berücksichtigung der Turniersportprüfung. PhD-thesis, University of Göttingen 1998.
225. Molina A, Valera M, Dos Santos R, Rodero A. Genetic parameters of morphofunctional traits in Andalusian horse. *Livestock Production Science* 1999;60:295–303.
226. Zeiler M, Distl O. Estimation of genetic parameters for breeding of the Haflinger. *Züchtungskunde* 2000;72:241–57.
227. Gerber Olsson E, Arnason T, Nasholm A, Philipsson J. Genetic parameters for traits at performance test of stallions and correlations with traits at progeny tests in Swedish warmblood horses. *Livestock Production Science* 2000;65:81–9.

228. Sprengel D, Geuder U, Götz K-U. BLUP-Tiermodellzuchtwertschätzung beim Haflinger in Bayern. Institut für Tierzucht, Bayerische Landesanstalt für Landwirtschaft (LfL), Munich, Germany; 2003.
229. Geuder U, Götz K-U. Optimierte Zuchtwertschätzung beim Süddeutschen Kaltblut. Institut für Tierzucht, Bayerische Landesanstalt für Landwirtschaft, Munich, Germany; 2007.
230. Olsson E, Näsholm A, Strandberg E, Philipsson J. Use of field records and competition results in genetic evaluation of station performance tested Swedish warmblood stallions. *Livestock Science* 2008;117:287–97.
231. Viklund Å, Näsholm A, Strandberg E, Philipsson J. Effects of long-time series of data on genetic evaluations for performance of Swedish Warmblood riding horses. *Animal* 2010;4:1823–31.
232. Mantovani R, Sartori C, Pigozzi G. Genetics of temperament and productive traits in the Italian Heavy Draught horse. In *World Congress of Genetics Applied to Livestock Production*; Leipzig, Germany; 2010.
233. Albertsdóttir E, Eriksson S, Sigurdsson Á, Árnason T. Genetic analysis of 'breeding field test status' in Icelandic horses. *Journal of Animal Breeding and Genetics* 2011;128:124–32.
234. Suontama M, van der Werf JHJ, Juga J, Ojala M. The use of foal and studbook traits in the breeding programmes of Finnhorse and Standardbred trotters. *Journal of Animal Breeding and Genetics* 2011;128:114–23.
235. Kieffer NM. Heritability of Cutting in Horses. *Horse Short Course*. Texas A&M University, College Station, USA; 1968. p. 46.
236. Hintz RL. Genetics of performance in the horse. *Journal of Animal Science* 1980;51:582–94.
237. Beuing R, Matthe A, Pracht P, Erhardt G. Validation and application of a test for self-confidence in warmblood horses. In *49th Annual Meeting of the EAAP 1998*, Warsaw, Poland; 1998. p. 293.
238. Lansade L, Leconte M, Pichart G. Développement d'un outil de prediction du temperament et des aptitudes mentales du cheval aux différentes disciplines équestres. *Les Haras Nationaux*, 34ème journée d'étude, 28 February 2008; 2008. p. 1–12.
239. Rumpf C. Einflussfaktoren beim Pferdekauf und deren Bedeutung für Turnier- und Freizeitreiter. Eine Empirische Untersuchung auf Basis einer Discrete-Choice-Analyse. *Göttinger Pferdetage '11 – Zucht, Haltung und Ernährung von Sportpferden*. FN-Verlag, Warendorf, Göttingen, Germany; 2011. p. 24–5.
240. Górecka-Bruzda A, Chruszczewski MH, Jezierski T, Murphy J. Behaviour-related traits of Polish sport horse stallions participating in 100-day performance tests. *Livestock Science* 2011;136:207–15.
241. Glißmann C, König von Borstel U. Alternatives to conventional evaluation of rideability: rein tension and behaviour from performance test dressage training. In *8th International Equitation Science Conference*, 18–20 July 2012, Edinburgh, UK; 2012. p. 20.
242. Janssens S, Schaerlaeckens S, Vincentelli C, Laevens H, Dirk L, Vervaecke H. Can observed horse behaviours predict rideability scores in Belgian Warmblood Stallions? In *44th International Congress of the International Society for Applied Ethology*, Uppsala, Sweden; 2010. p. 207.
243. Rothmann J, Sondergaard E, Christensen OF, Ladewig J. Practical assessment of reactivity and associations to rideability and performance traits. In *61st Annual Meeting of the European Association for Animal Production*, 23–27 August 2010, Heraklion, Greece; 2010. p. 151.
244. BMVEL (Bundesministerium für Verbraucherschutz EuL. Leitlinien für die Veranlagungsprüfung von Hengsten der deutschen Reitpferdezuchten 2003; 2010(31.12.)). Available from: URL: <http://www.pferd-aktuell.de/Anlage16447/LeitlinienVeranlagungspruefung.pdf>
245. Creighton E, Pankhurst S. Animal personality and animal welfare. In *8th Annual Symposium on Zoo Research*, 24–25 July 2006, Colchester Zoo, Stanway, UK; 2006. p. 101.
246. Freire R, Buckley P, Cooper JJ. Effects of different forms of exercise on post inhibitory rebound and unwanted behaviour in stabled horses. *Equine Veterinary Journal* 2009;41:487–92.
247. Hartmann E, Christensen JW, Keeling LJ. Training young horses to social separation: effect of a companion horse on training efficiency. *Equine Veterinary Journal* 2011;43:580–4.
248. Tadich T, Pulido R. Preliminary behavioural study of Caballo Fino Chilote stallions with restricted access to space and water during summer. *Archivos de Medicina Veterinaria* 2010;42:195–201.
249. Turner SP, White IMS, Brotherstone S, Farnworth MJ, Knap PW, Penny P, *et al.* Heritability of post-mixing aggressiveness in grower-stage pigs and its relationship with production traits. *Animal Science* 2006;82:615–20.
250. Tönepöhl B. Untersuchung zur Erfassung und Genetik von Verhaltensmerkmalen beim Schwein unter Praxisbedingungen. PhD-thesis, University of Göttingen, Germany; 2012. p. 112.
251. Muir WM, Craig JV. Improving animal well-being through genetic selection. *Poultry Science* 1998;77:1781–8.
252. Søndergaard LV, Herskin MS, Ladewig J, Holm IE, Dagnæs-Hansen F. Effect of genetic homogeneity on behavioural variability in an object recognition test in cloned Göttingen minipigs. *Applied Animal Behavior Science* 2012;141:20–4.
253. Haag EL, Rudman R, Houpt KA. Avoidance, maze learning and social dominance in ponies. *Journal of Animal Science* 1980;50:329–35.
254. Mal ME, McCall CA, Newland C, Cummins KA. Evaluation of a one-trial learning apparatus to test learning ability in weanling horses. *Applied Animal Behavior Science* 1993;35:305–11.
255. Sappington BF, Goldman L. Discrimination learning and concept formation in the Arabian horse. *Journal of Animal Science* 1994;72:3080–7.
256. Flannery B. Relational discrimination learning in horses. *Applied Animal Behavior Science* 1997;54:267–80.
257. McCall CA, Burgin SE. Equine utilization of secondary reinforcement during response extinction and acquisition. *Applied Animal Behavior Science* 2002;78:253–62.
258. Williams JL, Friend TH, Nevill CH, Archer G. The efficacy of a secondary reinforcer (clicker) during acquisition and extinction of an operant task in horses. *Applied Animal Behavior Science* 2004;88:331–41.
259. Murphy J. Assessing equine prospective memory in a Y-maze apparatus. *Veterinary Journal* 2009;181:24–8.

260. Egenvall A, Eisersjö M, Roepstorff L. Pilot study of behavior responses in young riding horses using 2 methods of making transitions from trot to walk. *Journal of Veterinary Behavior* 2012;7:157–68.
261. McLean AN. The positive aspects of correct negative reinforcement. *Anthrozoos: A Multidisciplinary Journal of The Interactions of People and Animals* 2005;18:245–54.
262. McGreevy PD. The advent of equitation science. *Veterinary Journal* 2007;174:492–500.
263. Murphy J, Arkins S. Equine learning behaviour. *Behavioral Processes* 2007;76:1–13.
264. Rolls ET. The orbitofrontal cortex and reward. *Cerebral Cortex* 2000;10:284–94.
265. Coleman K. Individual differences in temperament and behavioral management practices for nonhuman primates. *Applied Animal Behavior Science* 2012;137:106–13.
266. Hellsten ET, Viklund Å, Koenen EPC, Ricard A, Bruns E, Philipsson J. Review of genetic parameters estimated at stallion and young horse performance tests and their correlations with later results in dressage and show-jumping competition. *Livestock Science* 2006;103:1–12.
267. Kršková L, Mlynek J, Halo M. Relationship between behavioural traits and performance test scores in sport horses. *Acta Veterinaria Brno* 2003;72:429–35.
268. Geringer H, Bek-Kaczowska I, Banasiewicz E. Behavioural tests of half-bred horses on racetrack. *Roczniki Naukowe Zootechniki*. 2001;14:27–34.
269. Kamińska K, Oedenberg HGd, Dobrowolski M. Assessment of behaviour in half-bred and purebred Arabian horses tested on racecourse. *Acta Scientiarum Polonorum – Zootechnica* 2007;6:19–24.
270. Lam KH, Parkin TDH, Riggs CM, Morgan KL. Descriptive analysis of retirement of Thoroughbred racehorses due to tendon injuries at the Hong Kong Jockey Club (1992–2004). *Equine Veterinary Journal* 2007;39:143–8.
271. Malamed R, Berger J, Bain MJ, Kass P, Spier SJ. Retrospective evaluation of crib-biting and windsucking behaviours and owner-perceived behavioural traits as risk factors for colic in horses. *Equine Veterinary Journal* 2010;42:686–92.
272. Leal BB, Alves GES, Douglas RH, Bringel B, Young RJ, Haddad JPA, *et al.* Cortisol circadian rhythm ratio: a simple method to detect stressed horses at higher risk of colic? *Journal of Equine Veterinary Science* 2011;31:188–90.
273. Langlois B, Minkema D, Bruns E. Genetic problems of horse breeding. 1981. p. 22
274. Santamaria S, Bobbert MF, Back W, Barneveld A, van Weeren PR. Can early training of show jumpers bias outcome of selection events? *Livestock Science* 2006;102:163–70.
275. Haberland AM, König von Borstel U, Simianer H, König S. Integration of genomic information into sport horse breeding programs for optimization of accuracy of selection. *Animal* 2012;6:1369–76.
276. Lindzey G, Winston HD, Manosevitz M. Early experience, genotype, and temperament in *Mus musculus*. *Journal of Comparative and Physiological Psychology* 1963;56:622–9.
277. Tönepöhl B, Appel AK, Welp S, Voß B, König von Borstel U, Gauly M. Effect of marginal environmental and social enrichment during rearing on pigs' reactions to novelty, conspecifics and handling. *Applied Animal Behavior Science* 2012;140:137–45.
278. Redondo AJ, Carranza J, Trigo P. Fat diet reduces stress and intensity of startle reaction in horses. *Applied Animal Behavior Science* 2009;118:69–75.
279. Malmkvist J, Christensen JW. A note on the effects of a commercial tryptophan product on horse reactivity. *Applied Animal Behavior Science* 2007;107:361–6.
280. Grimmett A, Sillence MN. Calmatives for the excitable horse: a review of L-tryptophan. *Veterinary Journal* 2005;170:24–32.
281. Christensen JW, Rundgren M. Predator odour *per se* does not frighten domestic horses. *Applied Animal Behavior Science* 2008;112:136–45.
282. Dziezyc J, Taylor L, Boggess MM, Scott HM. The effect of ocular blinkers on the horses' reactions to four different visual and audible stimuli: results of a crossover trial. *Veterinary ophthalmology* 2011;14:327–32.
283. Christensen JW, Malmkvist J, Nielsen BL, Keeling LJ. Effects of a calm companion on fear reactions in naive test horses. *Equine Veterinary Journal* 2008;40:46–50.
284. Ahrendt LP, Christensen JW, Ladewig J. The ability of horses to learn an instrumental task through social observation. *Applied Animal Behavior Science* 2012;139:105–13.
285. Jezierski T, Górecka A. Changes in the horses heart rate during different levels of social isolation. *Animal Science Papers and Reports – Polish Academy of Sciences, Institute of Genetics and Animal Breeding Jastrzebiec* 2000;18:33–41.
286. Heleski C, Wickens CL, Minero M, DallaCosta E, Czeszak E, König v. Borstel U. Do horses recognize the difference between harsh tones and soothing tones when using voice as a reinforcer for learning a frightening task? In 8th International Equitation Science Conference, 18–20 July 2012, Edinburgh, UK; 2012. p. 140.
287. Birke L, Hockenhull J, Creighton E, Pinno L, Mee J, Mills D. Horses' responses to variation in human approach. *Applied Animal Behavior Science* 2011;134:56–63.
288. Fureix C, Pagès M, Bon R, Lassalle J-M, Kuntz P, Gonzalez G. A preliminary study of the effects of handling type on horses' emotional reactivity and the human-horse relationship. *Behavioral Processes* 2009;82:202–10.
289. von Borstel UU, Keeling LJ, Duncan IJH. Transfer of nervousness from the rider to the horse. In 39th international congress of the International Society of Applied Ethology; Sagamihara, Japan; 2005. p. 84.
290. von Borstel UU, Duncan IJH, Shoveller AK, Merkies K, Keeling LJ, Millman ST. Impact of riding in a coercively obtained Rollkur posture on welfare and fear of performance horses. *Applied Animal Behavior Science* 2009;116:228–36.
291. König von Borstel U, Kassebaum L, Gauly M. Effect of discipline (Western versus Dressage) and skill level on learning and discomfort behaviour of riding horses. In 7th International Equitation Science Conference, 26–29 October 2011, Hooge Mierde, The Netherlands; 2011. p. 75.
292. König v. Borstel U, Euent S. Influence of type and intensity of riders' reinforcement on horses' fear reactions. In 8th International Equitation Science Conference, Edinburgh, UK; 2012. p. 100.
293. Parker M, Goodwin D, Redhead ES. Survey of breeders' management of horses in Europe, North America and

- Australia: Comparison of factors associated with the development of abnormal behaviour. *Applied Animal Behavior Science* 2008;114:206–15.
294. Jezierski T, Jaworski Z, Górecka A. Effects of handling on behaviour and heart rate in Konik horses: comparison of stable and forest reared youngstock. *Applied Animal Behavior Science* 1999;62:1–11.
 295. Hausberger M, Henry S, Richard-Yris MA. Early Experience and Behavioural Development in Foals. EAAP Publication No. 122. Wageningen Academic Publishers, Wageningen; 2007. p. 37–45.
 296. Lansade L, Boivin X, Bouissou MF. Effects of Period, Type and Duration of Handling on Manageability, Reactivity and Learning Ability of Horses. EAAP Publication No. 122. Wageningen Academic Publishers, Wageningen; 2007. p. 47–55.
 297. Visser EK, Ellis AD, Van Reenen CG. The effect of two different housing conditions on the welfare of young horses stabled for the first time. *Applied Animal Behavior Science* 2008;114:521–33.
 298. Ladewig J, Søndergaard E, Christensen JW. Ontogeny: Preparing the Young Horse for its Adult Life. Cambridge University Press, Cambridge; 2005. p. 139–49.
 299. González del Pino F, Escalante F, Wilde OR. Imprinting technique, and intensive socialization early in the foal. *REDVET* 2009;10:030924.
 300. Ligout Sv, Bouissou M-F, Boivin X. Comparison of the effects of two different handling methods on the subsequent behaviour of Anglo-Arabian foals toward humans and handling. *Applied Animal Behavior Science* 2008;113:175–88.
 301. Vincent. Imprint training behaviour: myth or reality? *Le Nouveau Praticien Vétérinaire – Équine Horse-Série*. 2007;3:121–4.
 302. Ligout S, Bouissou M-F, Boivin X. Comparison of the effects of two different handling methods on the subsequent behaviour of Anglo-Arabian foals toward humans and handling. *Applied Animal Behavior Science* 2008;113:175–88.
 303. Henry S, Richard-Yris MA, Tordjman S, Hausberger M. Neonatal handling affects durably bonding and social development. *PLoS ONE* 2009;4:e5216.
 304. Whitaker, Dean D. Effects of varying amounts of early handling at weaning and later ages in the subsequent learning ability of two-year-old horses [Dissertation/thesis]. Texas Tech University, United States – Texas; 1982.
 305. Mal ME, McCall CA, Cummins KA, Newland MC. Influence of preweaning handling methods on post-weaning learning ability and manageability of foals. *Applied Animal Behavior Science* 1994;40:187–95.
 306. Simpson BS. Neonatal foal handling. *Applied Animal Behavior Science* 2002;78:303–17.
 307. Søndergaard E, Ladewig J. Group housing exerts a positive effect on the behaviour of young horses during training. *Applied Animal Behavior Science* 2004;87:105–18.
 308. Henry S, Richard-Yris MA, Hausberger M. Influence of various early human-foal interferences on subsequent human-foal relationship. *Dev Psychobiol* 2006;48:712–8.
 309. Sankey C, Richard-Yris M-A, Leroy Hln, Henry Sv, Hausberger M. Positive interactions lead to lasting positive memories in horses, *Equus caballus*. *Animal Behaviour* 2010;79:869–75.
 310. Lansade L, Bertrand M, Boivin X, Bouissou MF. Effects of handling at weaning on manageability and reactivity of foals. *Applied Animal Behavior Science* 2004;87:131–49.
 311. Henry S, Hemery D, Richard MA, Hausberger M. Human-mare relationships and behaviour of foals toward humans. *Applied Animal Behavior Science* 2005;93:341–62.
 312. Polito R, Minero M, Canali E, Verga M. A pilot study on yearlings' reactions to handling in relation to the training method. *Anthrozoos: A Multidisciplinary Journal of The Interactions of People and Animals* 2007;20:295–303.
 313. Warren-Smith AK, McGreevy PD. The use of blended positive and negative reinforcement in shaping the halt response of horses (*Equus caballus*). *Animal Welfare* 2007;16:481–8.
 314. Innes L, McBride S. Negative versus positive reinforcement: an evaluation of training strategies for rehabilitated horses. *Applied Animal Behavior Science* 2008;112:357–68.
 315. Slater C, Dymond S. Using differential reinforcement to improve equine welfare: shaping appropriate truck loading and feet handling. *Behavioral Processes* 2011;88:329–39.
 316. Visser EK, VanDierendonck M, Ellis AD, Rijksen C, Van Reenen CG. A comparison of sympathetic and conventional training methods on responses to initial horse training. *Veterinary Journal* 2009;181:48–52.
 317. Martin TI, Zentall TR, Lawrence L. Simple discrimination reversals in the domestic horse (*Equus caballus*): effect of discriminative stimulus modality on learning to learn. *Applied Animal Behavior Science* 2006;101:328–38.
 318. Budzyńska M, Krupa W. Relation between fearfulness level and maternal behaviour in Arab mares. *Animal Science Papers and Reports* 2011;29:119–29.
 319. König von Borstel U, Moors E, Schichowski C, Gaulty M. Breed differences in maternal behaviour in relation to lamb (*Ovis orientalis aries*) productivity. *Livestock Science* 2011;137:42–8.
 320. Henry S, Zanella AJ, Sankey C, Richard-Yris M-A, Marko A, Hausberger M. Adults may be used to alleviate weaning stress in domestic foals (*Equus caballus*). *Physiology and Behavior* 2012;106:428–38.
 321. Houpt KA, Hintz HF, Butler WR. A preliminary study of two methods of weaning foals. *Applied Animal Behavior Science* 1984;12:177–81.
 322. Hoffman RM, Kronfeld DS, Holland JL, Greiwe-Crandell KM. Prewaning diet and stall weaning method influences on stress response in foals. *Journal of Animal Science* 1995;73:2922.
 323. Apter RC, Householder DD. Weaning and weaning management of foals: a review and some recommendations. *Journal of Equine Veterinary Science* 1996;16:428–35.
 324. Waters AJ, Nicol CJ, French NP. Factors influencing the development of stereotypic and redirected behaviours in young horses: findings of a four year prospective epidemiological study. *Equine Veterinary Journal* 2002;34:572–9.
 325. Moons CPH, Laughlin K, Zanella AJ. Effects of short-term maternal separations on weaning stress in foals. *Applied Animal Behavior Science* 2005;91:321–35.

326. Nicol CJ, Badnell-Waters AJ, Bice R, Kelland A, Wilson AD, Harris PA. The effects of diet and weaning method on the behaviour of young horses. *Applied Animal Behavior Science* 2005;95:205–21.
327. Waran NK, Clarke N, Farnworth M. The effects of weaning on the domestic horse (*Equus caballus*). *Applied Animal Behavior Science* 2008;110:42–57.
328. Grzimek B. Ein Fohlen, das kein Pferd kannte. *Zeitschrift für Tierpsychologie* 1949;6:391–405.
329. Glendinning SA. A system of rearing foals on an automatic calf feeding machine. *Equine Veterinary Journal* 1974;6:12–6.
330. Houpt KA, Parsons MS, Hintz HF. Learning ability of orphan foals, of normal foals and their mothers. *Journal of Animal Science* 1982;55:1027–31.
331. Houpt KA, Hintz HF. Some effects of maternal deprivation on maintenance behavior, spatial relationships and responses to environmental novelty in foals. *Applied Animal Ethology* 1983;9:221–30.
332. Cooper JJ, Albentosa MJ. Behavioural adaptation in the domestic horse: potential role of apparently abnormal responses including stereotypic behaviour. *Livestock Production Science* 2005;92:177–82.
333. Mason GJ, Turner MA, Bateson PPG, Klopfer PH, Thompson NS. Mechanisms involved in the development and control of stereotypies. In: *Behaviour and Evolution Perspectives in Ethology*, Vol. 10. Plenum Press, New York; 1993. p. 53–86.
334. Rushen J, Mason G. A decade-or-more's progress in understanding stereotypies. In: Mason G, Rushen J, editors. *Stereotypic Animal Behaviour – Fundamentals and Applications to Welfare*. CABI Publishing, Wallingford, UK; 2006. p. 1–18.
335. Vecchiotti GG, Galanti R. Evidence of heredity of cribbing, weaving and stall-walking in Thoroughbred horses. *Livestock Production Science* 1986;14:91–5.
336. Hosoda T. On the heritability of susceptibility to windsucking in horses. *Japanese Journal of Zootechnical Science* 1950;21:25–8.
337. Hausberger M, Gautier E, Muller C, Jégo P. Lower learning abilities in stereotypic horses. *Applied Animal Behavior Science* 2007;107:299–306.
338. Dodman NH, Normile JA, Cottam N, Guzman M, Shuster L. Prevalence of compulsive behaviors in formerly feral horses. *International Journal of Applied Research in Veterinary Medicine* 2005;3:20–4.
339. Hausberger M, Muller C, Lunel C. Does work affect personality? a study in horses. *PLoS ONE* 2011;6:e14659.
340. Normando S, Meers L, Samuels WE, Faustini M, Ödberg FO. Variables affecting the prevalence of behavioural problems in horses. Can riding style and other management factors be significant? *Applied Animal Behavior Science* 2011;133:186–98.
341. Heleski CR, Shelle AC, Nielsen BD, Zanella AJ. Influence of housing on weanling horse behavior and subsequent welfare. *Applied Animal Behavior Science* 2002;78:291–302.
342. Christensen JW, Ladewig J, Søndergaard E, Malmkvist J. Effects of individual versus group stabling on social behaviour in domestic stallions. *Applied Animal Behavior Science* 2002;75:233–48.
343. Søndergaard E, Halekoh U. Young horses' reactions to humans in relation to handling and social environment. *Applied Animal Behavior Science* 2003;84:265–80.
344. Rivera E, Benjamin S, Nielsen B, Shelle J, Zanella AJ. Behavioral and physiological responses of horses to initial training: the comparison between pastured versus stalled horses. *Applied Animal Behavior Science* 2002;78:235–52.
345. Webster MM, Ward AJW. Personality and social context. *Biological Reviews* 2010;86:759–773.
346. Hall C. The impact of visual perception on equine learning. *Equine Learning Behaviour* 2007;76:29–33.
347. Trevor-Roper P. *The World Through Blunted Sight*. Thames and Hudson, London, UK; 1970.