

Master Thesis, On the Absolute Anabelian Geometry of Hyperbolic Polycurves over Local Fields (Title, 仮)

Kazuki Masugi

May 6, 2026

Contents

1 Preliminaries on Logarithmic Geometry of Hyperbolic Polycurves	5
2 Cofinality of l -admissible finite étale coverings	23
3 Dehn-twist Pointed Stable Curves type Outer Representations	24
4 Reconstruction of Log-full Subgroups	25
5 Reconstruction of Dimensions via Corank 1 Intersections	26
6 Reconstruction of the Graph-theoretic Structure of Proper Hyperbolic Polycurves of Dimension 2	27

Introduction

なにか intro を書く必要がある。

Notations and Conventions

Numbers: The notation $\mathfrak{Primes}$ will be used to denote the set of prime numbers. The notation $\mathbb{N}$ will be used to denote the set of nonnegative integers. The notation $\mathbb{Z}$ will be used to denote the additive group or ring of integers. The notation $\mathbb{Q}$ will be used to denote the field of rational numbers. The notation $\mathbb{Q}_{\geq 0}$ will be used to denote the additive monoid of nonnegative rational numbers. The notation $\mathbb{R}$ will be used to denote the field of real numbers. For each $x \in \mathbb{R}$, the notation $\lfloor x \rfloor$ will be used to denote the greatest integer $\leq x$. If p is a prime number, then the notation $\mathbb{Q}_p$ will be used to denote the field of p -adic numbers; the notation $\mathbb{Z}_p$ will be used to denote the additive group or ring of p -adic integers; the notation $\overline{\mathbb{Q}_p}$ will be used to denote an algebraic closure of $\mathbb{Q}_p$; the notation $\mathbb{C}_p$ will be used to denote the p -adic completion of $\mathbb{Q}_p$. We shall refer to a finite extension field of $\mathbb{Q}_p$ as a **p -adic local field**.

Monoids: Let M be a commutative monoid. In this subsection, we regard the set of positive integers $\mathbb{N}_{\geq 1}$ as a directed set via its **multiplicative** structure, i.e., $i \leq j \Leftrightarrow i|j$. For $i \in \mathbb{N}_{\geq 1}$, write $M_i := M$. For $i, j \in \mathbb{N}_{\geq 1}$ such that $i \leq j$, write $\phi_{i,j} : M_i \rightarrow M_j$ for the homomorphism of monoids obtained by multiplication by $\frac{j}{i}$. Then we shall refer to the inductive limit of the inductive system $\{M_i, \phi_{i,j}\}$ on the directed set $\mathbb{N}_{\geq 1}$ as the **perfection** of M .

Rings and fields: Let R be a ring. Then we shall write $R^\times$ for the multiplicative group of units of the ring.

Let F be a perfect field, p a prime number. Then the notation $\overline{F}$ will be used to denote an algebraic closure of F ; $G_F := \text{Gal}(\overline{F}/F)$.

Suppose that F is a **valuation field**, i.e., a field equipped with a valuation map. Then we shall write $\mathcal{O}_F$ for the ring of integers of F ; $\mathfrak{m}_F$ for the maximal ideal of $\mathcal{O}_F$.

Topological groups: Let G be a topological group; $H \subseteq G$ a closed subgroup of G . Then we shall write G^{ab} for the abelianization of G ; $Z_G(H)$ for the **centralizer** of H in G , i.e., $Z_G(H) := \{g \in G \mid ghg^{-1} = h \text{ for all } h \in H\}$; $N_G(H)$ for the **normalizer** of H in G , i.e., $N_G(H) := \{g \in G \mid gHg^{-1} = H\}$; and $C_G(H)$ for the **commensurator** of $H \subseteq G$, i.e.,

$$C_G(H) := \{g \in G \mid H \cap gHg^{-1} \text{ is of finite index in } H \text{ and } gHg^{-1}\}.$$

We shall say that the closed subgroup H is **commensurably terminal** in G if $H = C_G(H)$. Let $\Sigma \subseteq \mathfrak{Primes}$ be a nonempty subset. Then we shall write G^Σ for the pro- Σ completion of G .

We shall write $\text{Aut}(G)$ for the group of continuous automorphisms of the topological group G , $\text{Inn}(G) \subseteq \text{Aut}(G)$ for the subgroup of inner automorphisms of G , and $\text{Out}(G) := \text{Aut}(G)/\text{Inn}(G)$. Suppose that G is **center-free**. Then we have a **natural exact sequence** of groups

$$1 \rightarrow \text{Inn}(G) \rightarrow \text{Aut}(G) \rightarrow \text{Out}(G) \rightarrow 1$$

where G is identified with $\text{Inn}(G)$ via the natural isomorphism $G \cong \text{Inn}(G)$ sending $g \in G$ to the inner automorphism $x \mapsto gxg^{-1}$ of G .

If J is a group, and $\rho : J \rightarrow \text{Out}(G)$ is a homomorphism, then we shall denote by $G \rtimes^{\text{out}} J$ the group obtained by pulling back the above exact sequence of groups via ρ . Thus, we have a natural exact sequence of groups

$$1 \rightarrow G \rightarrow G \rtimes^{\text{out}} J \rightarrow J \rightarrow 1.$$

Suppose further that the topology of G admits a **countable basis** consisting of **characteristic open subgroups** of G . Then one verifies immediately that the topology of G induces a **natural topology** on the group $\text{Aut}(G)$, hence on the group $\text{Out}(G)$. In particular, one verifies easily that if J is a **topological group**, and $\rho : J \rightarrow \text{Out}(G)$ is **continuous**, then $G \rtimes^{\text{out}} J$ admits a natural **topological group** structure.

Let G_1, G_2 be topological groups. Then we shall write $\text{Isom}(G_1, G_2)$ for the set of **isomorphisms** from G_1 to G_2 ;

$$\text{OutIsom}(G_1, G_2)$$

for the set of **outer isomorphisms** from G_1 to G_2 , i.e., the set of $\text{Inn}(G_2)$ -orbits of $\text{Isom}(G_1, G_2)$ under the natural action of $\text{Inn}(G_2)$ on $\text{Isom}(G_1, G_2)$ by post-composition.

Schemes: Let K be a field; $K \subseteq L$ a field extension; X an algebraic variety [i.e., a separated, geometrically integral scheme of finite type] over K . Then we shall write $X_L := X \times_K L$. Suppose that K is a valuation field. Let X be an $\mathcal{O}_K$ -scheme. Then we shall write X_s for the special fiber of X [i.e., the fiber of X over the closed point of $\text{Spec } \mathcal{O}_K$].

Let S_1, S_2 be schemes. Then we shall write $\text{Isom}(S_1, S_2)$ for the set of isomorphisms of schemes between S_1 and S_2 .

Log schemes: If $X^{\log}$ is a fine log scheme, then we shall write

- X for the underlying scheme of $X^{\log}$;
- $\mathcal{M}_{X^{\log}}$ for the étale sheaf of monoids on X that defines the log structure of $X^{\log}$;
- $\mathcal{C}_{X^{\log}} := \mathcal{M}_{X^{\log}} / \mathcal{O}_X^\times$, which we shall refer to as the characteristic of $X^{\log}$ [cf. [MT], Definition 5.1, (i)].

Let K be a complete discrete valuation field; Y a hyperbolic curve over K ; Y a compactified semistable model of Y over $\mathcal{O}_K$. Write $S := \text{Spec } \mathcal{O}_K$; $S^{\log}$ for the log scheme determined by the log structure associated $\text{Spec } \mathcal{O}_K$; $S^{\log}$ for the log scheme determined by the log structure associated to the closed point of S . Then it follows immediately from [Hur], § 3.7, § 3.8, that the multiplicative monoid

of sections of O_Y that are invertible on [the open subscheme of Y determined by] Y determines a natural log structure on Y . Denote the resulting log scheme by $Y^{\log}$. Then one verifies immediately that the natural morphism of schemes $Y \rightarrow S$ extends to a proper, log smooth morphism $Y^{\log} \rightarrow S^{\log}$ of fine log schemes. Let y be a geometric point of Y_s . Write $\mathcal{C}_{Y^{\log}, y}^{\text{pf}}$ for the perfection of the stalk of the characteristic $\mathcal{C}_{Y^{\log}}$ at y . Then one verifies immediately that, if the image of y in Y_s is a smooth point that is not a cusp (respectively, a cusp; a node), then $\mathcal{M}_{Y^{\log}, y}^{\text{pf}} \xrightarrow{\sim} \mathbb{Q}_{\geq 0}$ (respectively, $\mathcal{M}_{Y^{\log}, y}^{\text{pf}} \xrightarrow{\sim} \mathbb{Q}_{\geq 0} \times \mathbb{Q}_{\geq 0}$; $\mathcal{M}_{Y^{\log}, y}^{\text{pf}} \xrightarrow{\sim} \mathbb{Q}_{\geq 0} \times \mathbb{Q}_{\geq 0}$).

Fundamental groups: For a connected noetherian scheme S , we shall write Π_S for the étale fundamental group of S , relative to a suitable choice of base-point. Let $\Sigma \subseteq \mathfrak{Primes}$ be a nonempty subset; K a perfect field; X an algebraic variety over K . Then we shall write

$$\Delta_X := \Pi_{X_{\bar{K}}}; \quad \Pi_X^{(\Sigma)} := \Pi_X / \text{Ker}(\Delta_X \twoheadrightarrow \Delta_X^{\Sigma}),$$

where $\Delta_X \rightarrow \Delta_X^{\Sigma}$ denotes the natural surjection. We shall refer to Δ_X^{Σ} (respectively, $\Pi_X^{(\Sigma)}$) as the geometric pro- Σ fundamental group (respectively, geometrically pro- Σ fundamental group) of X .

ちゃんと確認して補正する必要がある。

moduli の記述, Mgr, Mbargr の記述もする

pi1, log scheme, abs Gal group, 分岐理論

1 Preliminaries on Logarithmic Geometry of Hyperbolic Polycurves

In this section, we shall prepare some facts on the geometry of hyperbolic polycurves, which will be used as basic ingredients in the subsequent sections. In particular, we shall verify that the stable compactification of a hyperbolic polycurve admits a natural log structure, which is compatible with a certain stratification. Moreover, we observe that one can associate a semi-graph of anabelioids to each stratum of the stratification and obtain an outer representation on the semi-graph of anabelioids, and that one can obtain specialization morphisms between the semi-graphs of anabelioids associated to the strata, which are compatible with the specialization relation.

Definition 1.1. Let S be a scheme and X a scheme over S .

(i) We shall say that X is a **hyperbolic curve** (of type (g, r)) over S if there exist

- a pair of non-negative integers (g, r) ;
- a scheme X^{cpt} which is smooth, proper, geometrically connected, and of relative dimension 1 over S ;
- a (possibly empty) closed subscheme D of X^{cpt} which is finite étale over S

such that

- $2g - 2 + r > 0$;
- any geometric fiber of X^{cpt} over S is a smooth curve of genus g ;
- the finite étale morphism $D \rightarrow S$ is of degree r ;
- X is isomorphic to $X^{\text{cpt}} \setminus D$ as a scheme over S .

(ii) We shall say that X is a **stable curve** (of type (g, r)) over S if there exist

- a pair of non-negative integers (g, r) ;
- a scheme X^{cpt} which is proper, flat, geometrically connected, and of relative dimension 1 over S ;
- a (possibly empty) closed subscheme D of X^{cpt} which is finite étale over S

such that

- $2g - 2 + r > 0$;
- any geometric fiber of (X^{cpt}, D) over S is a stable curve of genus g , i.e., a curve of genus g with at worst nodal singularities and with r marked points, such that any rational component of the curve contains at least three nodes or marked points;

- the finite étale morphism $D \rightarrow S$ is of degree r ;
- X is isomorphic to $X^{\text{cpt}} \setminus D$ as a scheme over S .

(iii) We shall say that X is a **hyperbolic polycurve** (of relative dimension n) over S (resp. **stable polycurve** (of relative dimension n)) if there exist

- a positive integer n ;
- a factorization of the structure morphism $X \rightarrow S$ (which is not necessarily uniquely determined by X)

$$X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = S$$

such that

- for each $i = 1, \dots, n$, the morphism $X_i \rightarrow X_{i-1}$ is a hyperbolic curve over X_{i-1} (resp. a stable curve over X_{i-1}).

We shall refer to the above morphism $X = X_n \rightarrow X_{n-1}$ as a **parametrizing morphism** for X and refer to the above factorization as a **sequence of parametrizing morphisms** or **parametrization** for X .

Definition 1.2. Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . We shall say that $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ is a **compactified stable model** of X (with respect to its parametrization) if there exist

- open immersion $X_i \hookrightarrow \mathfrak{X}_i$ for each $i = 0, \dots, n$;
- divisors $\mathfrak{D}_i$ on $\mathfrak{X}_i$ for each $i = 1, \dots, n$;

such that

- open immersion $X_0 \hookrightarrow \mathfrak{X}_0$ is a (natural) immersion $\text{Spec } K \hookrightarrow \text{Spec } \mathcal{O}_K$;
- for each $i = 1, \dots, n$, the morphism $(\mathfrak{X}_i, \mathfrak{D}_i) \rightarrow \mathfrak{X}_{i-1}$ is a stable curve over $\mathfrak{X}_{i-1}$;
- for each $i = 1, \dots, n$, the pullback of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \rightarrow \mathfrak{X}_{i-1}$ by the open immersion $X_{i-1} \hookrightarrow \mathfrak{X}_{i-1}$ is a compactification of the hyperbolic curve $X_i \rightarrow X_{i-1}$.

Proposition 1.3 (Log structure on the stable compactification of a hyperbolic polycurve). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. Then there exists an fs log structure on $\mathfrak{X}$ such that the morphism $\mathfrak{X}^{\log} \rightarrow (\text{Spec } \mathcal{O}_K)^{\log}$ is log smooth, where $(\text{Spec } \mathcal{O}_K)^{\log}$ is the log scheme obtained by endowing $\text{Spec } \mathcal{O}_K$ with the log structure associated to the closed point of $\text{Spec } \mathcal{O}_K$ — therefore, the log scheme $\mathfrak{X}^{\log}$ is log regular — and such that the interior of $\mathfrak{X}^{\log}$ (i.e., the locus where the log structure is trivial) coincides with X . Moreover, the log structure on $\mathfrak{X}$ is uniquely determined by the above conditions.

Proof. We shall proceed by induction on the relative dimension n of X over $\text{Spec } K$. The case where $n = 0$ is trivial, so we assume that $n > 0$ and that the proposition holds for any hyperbolic polycurve of relative dimension less than n .

Let $X' = X_{n-1} \rightarrow X_{n-2} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be the hyperbolic polycurve obtained by removing the top level of the parametrization of X . By the induction hypothesis, there exists a log structure on $\mathfrak{X}' = \mathfrak{X}_{n-1} \rightarrow \mathfrak{X}_{n-2} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ such that $(\mathfrak{X}')^{\log} \rightarrow (\text{Spec } \mathcal{O}_K)^{\log}$ is log smooth and such that the interior of $(\mathfrak{X}')^{\log}$ coincides with X' .

Let $\Phi: \mathfrak{X}' \rightarrow \overline{\mathfrak{M}}_{g_n, r_n}$ be the morphism determined by the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \rightarrow \mathfrak{X}_{n-1}$, where g_n and r_n are the genus and the number of marked points of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \rightarrow \mathfrak{X}_{n-1}$. By the induction hypothesis the morphism Φ induces a morphism of fs log schemes $(\mathfrak{X}')^{\log} \rightarrow \overline{\mathfrak{M}}_{g_n, r_n}^{\log}$, where $\overline{\mathfrak{M}}_{g_n, r_n}^{\log}$ is endowed with the natural log structure determined by its boundary, together with the fact that Φ maps the interior of $\mathfrak{X}'^{\log}$ to the interior of $\overline{\mathfrak{M}}_{g_n, r_n}^{\log}$.

By computing the log structures involved, one verifies immediately that the underlying scheme of the fs log fiber product $(\mathfrak{X}')^{\log} \times_{\overline{\mathfrak{M}}_{g_n, r_n}^{\log}} \overline{\mathfrak{C}}_{g_n, r_n}^{\log}$ coincides with the fiber product $\mathfrak{X}' \times_{\overline{\mathfrak{M}}_{g_n, r_n}} \overline{\mathfrak{C}}_{g_n, r_n}$, which is isomorphic to $\mathfrak{X}_n$ by the definition of Φ . Thus, the scheme $\mathfrak{X}_n$ admits a natural fs log structure determined by the natural isomorphism $\mathfrak{X}_n \cong \mathfrak{X}' \times_{\overline{\mathfrak{M}}_{g_n, r_n}} \overline{\mathfrak{C}}_{g_n, r_n}$, and the resulting log scheme $\mathfrak{X}_n^{\log}$ fits into a natural cartesian diagram of fs log schemes.

Since the morphism $\overline{\mathfrak{C}}_{g_n, r_n}^{\log} \rightarrow \overline{\mathfrak{M}}_{g_n, r_n}^{\log}$ is log smooth, it follows that the morphism $\mathfrak{X}_n^{\log} \rightarrow (\mathfrak{X}')^{\log}$ is log smooth. Therefore, the morphism $\mathfrak{X}_n^{\log} \rightarrow (\text{Spec } \mathcal{O}_K)^{\log}$ is also log smooth. In particular, since $(\text{Spec } \mathcal{O}_K)^{\log}$ is log regular, it follows that $\mathfrak{X}_n^{\log}$ is log regular. Moreover, by the computation of the log structures involved, one verifies immediately that the interior of $\mathfrak{X}_n^{\log}$ coincides with X_n . Finally, since the log structure on a log regular fs log scheme is uniquely determined by its interior, it follows that the log structure on $\mathfrak{X}_n$ is uniquely determined by the conditions described in the proposition. This completes the proof of the proposition. $\square$

Lemma 1.4 (Normality of the stable compactification). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. Then $\mathfrak{X}_n$ is integral and normal. Moreover, the complement of X_n in $\mathfrak{X}_n$ is purely of codimension 1.

Proof. In light of the Proposition 1.3, the scheme $\mathfrak{X}_n$ admits a natural log structure such that $\mathfrak{X}_n^{\log}$ is log regular. In particular, the underlying scheme $\mathfrak{X}_n$ of $\mathfrak{X}_n^{\log}$ is normal and the complement of the interior of $\mathfrak{X}_n^{\log}$ (which coincides with X_n [cf. Proposition 1.3]) in $\mathfrak{X}_n$ is purely of codimension 1. Therefore, to verify Lemma 1.4, it suffices to show that $\mathfrak{X}_n$ is connected. However, this follows immediately from the elementary scheme theory and the fact that $\mathfrak{X}_n \rightarrow$

$\mathrm{Spec} \mathcal{O}_K$ is proper and geometrically connected. This completes the proof of the lemma. $\square$

Lemma 1.5 (Uniqueness of the stable compactification of a hyperbolic polycurve). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \mathrm{Spec} K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \mathrm{Spec} \mathcal{O}_K$ with respect to its parametrization. Then any compactified stable model of X with respect to its parametrization is isomorphic to $\mathfrak{X}$ as a sequence of morphisms over $\mathrm{Spec} \mathcal{O}_K$.

Proof. We shall proceed by induction on the relative dimension n of X over $\mathrm{Spec} K$. The case where $n = 0$ is trivial, so we assume that $n > 0$ and that the lemma holds for any hyperbolic polycurve of relative dimension less than n .

Let $\mathfrak{X}' = \mathfrak{X}'_n \rightarrow \mathfrak{X}'_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}'_1 \rightarrow \mathfrak{X}'_0 = \mathrm{Spec} \mathcal{O}_K$ be another compactified stable model of X with respect to its parametrization. Since the moduli stack $\overline{\mathcal{M}}_{g_n, r_n}$ of stable curves of genus g_n with r_n marked points is separated and Deligne-Mumford, it follows that the diagonal morphism $\overline{\mathcal{M}}_{g_n, r_n} \rightarrow \overline{\mathcal{M}}_{g_n, r_n} \times \overline{\mathcal{M}}_{g_n, r_n}$ is proper and unramified; in particular, it is a finite morphism.

Let $\Phi: \mathfrak{X}_{n-1} \rightarrow \overline{\mathcal{M}}_{g_n, r_n}$ (respectively, $\Phi': \mathfrak{X}'_{n-1} \rightarrow \overline{\mathcal{M}}_{g_n, r_n}$) be the morphism determined by the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \rightarrow \mathfrak{X}_{n-1}$ (respectively, the stable curve $(\mathfrak{X}'_n, \mathfrak{D}'_n) \rightarrow \mathfrak{X}'_{n-1}$). Since the restrictions of Φ and Φ' to X_{n-1} coincide, we obtain a commutative diagram

$$\begin{array}{ccc} X_{n-1} & \longrightarrow & \overline{\mathcal{M}}_{g_n, r_n} \\ \downarrow & & \downarrow \Delta \\ \mathfrak{X}_{n-1} & \xrightarrow{(\Phi, \Phi')} & \overline{\mathcal{M}}_{g_n, r_n} \times \overline{\mathcal{M}}_{g_n, r_n} \end{array}$$

Note that Δ is finite (and unramified) morphism since $\overline{\mathcal{M}}_{g_n, r_n}$ is separated and Deligne-Mumford algebraic stack, and the fiber product $\mathfrak{X}_{n-1} \times_{\overline{\mathcal{M}}_{g_n, r_n} \times \overline{\mathcal{M}}_{g_n, r_n}} \overline{\mathcal{M}}_{g_n, r_n}$ is just the Isom-scheme $\mathrm{Isom}_{\mathfrak{X}_{n-1}}(\mathfrak{X}_n, \mathfrak{X}'_n)$, which is representable by a scheme. Since $\mathfrak{X}_{n-1}$ is normal by Lemma 1.4, it follows that the morphism $(\Phi, \Phi'): \mathfrak{X}_{n-1} \rightarrow \overline{\mathcal{M}}_{g_n, r_n} \times \overline{\mathcal{M}}_{g_n, r_n}$ factors through Δ ; in particular, we obtain the equality $\Phi = \Phi'$. Therefore, by the definition involved, we obtain an isomorphism $\mathfrak{X}_n \cong \mathfrak{X}'_n$ over $\mathfrak{X}_{n-1}$, which completes the proof of the lemma. $\square$

Definition 1.6. Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \mathrm{Spec} K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \mathrm{Spec} \mathcal{O}_K$ with respect to its parametrization.

- (i) Let $x \in \mathfrak{X}_n$ be a point. Write x_i for the image of x in X_i for each $i = 0, \dots, n$. We shall say that x is a **stratum point** of $\mathfrak{X}_n$ if the following condition is satisfied:

- for each $i = 1, \dots, n$, the point x_i is either a node of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$, a marked point of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$, or a generic point of an irreducible component of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$.
- (ii) Let x be a stratum point of $\mathfrak{X}_n$. Let x_i be the image of x in $\mathfrak{X}_i$ for each $i = 0, \dots, n$. We shall say that the **VCNGS-signature** of x is the function $\mathbf{sgn}(x): \{1, \dots, n\} \rightarrow \{V, C, N, G, S\}$ defined as follows:
- for each $i = 1, \dots, n$, $\mathbf{sgn}(x)(i) = V$ if x_i is a generic point of an irreducible component of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$;
 - for each $i = 1, \dots, n$, $\mathbf{sgn}(x)(i) = C$ if x_i is a marked point of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$;
 - for each $i = 1, \dots, n$, $\mathbf{sgn}(x)(i) = N$ if x_i is a node of the stable curve $(\mathfrak{X}_i, \mathfrak{D}_i) \times_{\mathfrak{X}_{i-1}} x_{i-1} \rightarrow x_{i-1}$;
 - for $i = 0$, $\mathbf{sgn}(x)(i) = G$ if x_i is the generic point of $\mathrm{Spec} \mathcal{O}_K$;
 - for $i = 0$, $\mathbf{sgn}(x)(i) = S$ if x_i is the closed point of $\mathrm{Spec} \mathcal{O}_K$.
- (iii) Let x be a stratum point of $\mathfrak{X}_n$. We shall say that the **combinatorial dimension** of x is the number of $i \in \{0, \dots, n\}$ such that $\mathbf{sgn}(x)(i) \in \{V, G\}$.

Lemma 1.7 (Stratification of the stable compactification). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \dots \rightarrow X_1 \rightarrow X_0 = \mathrm{Spec} K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \dots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \mathrm{Spec} \mathcal{O}_K$ with respect to its parametrization. Then there exists a stratification of $\mathfrak{X}_n$ such that

- the strata of the stratification are irreducible;
- the set of stratum points of $\mathfrak{X}_n$ coincides with the set of generic points of the strata of the stratification.

The stratification of $\mathfrak{X}_n$ is uniquely determined by the above conditions. Moreover, for each stratum A of the stratification associated to $\mathfrak{X}_n$, the image of A in $\mathfrak{X}_i$ is one of the strata of the stratification associated to $\mathfrak{X}_i$ for each $i = 0, \dots, n-1$.

Proof. First, we shall construct a stratification of $\mathfrak{X}_n$ by induction on the relative dimension n of X over $\mathrm{Spec} K$. The case where $n = 0$ is trivial, so we assume that $n > 0$ and that the lemma holds for any hyperbolic polycurve of relative dimension less than n .

Let π be the morphism $\mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1}$. For each strata A of the stratification of $\mathfrak{X}_{n-1}$ obtained by the induction hypothesis, we shall construct a decomposition of $\pi^{-1}(A)$ by locally closed subsets as follows. Let η_A be the generic point of A . Then the set of stratum points over η_A coincides with the set of nodes, marked points, and generic points of irreducible components of the stable

curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$. For the stratum points s over η_A which are nodes or marked points of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$, we define the subscheme B_s to be the closure of s in $\pi^{-1}(A)$. For the stratum points t over η_A which are generic points of irreducible components of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$, we define the subscheme B_t as follows: let $s_1, \dots, s_m$ be the nodes and marked points of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$ which are contained in the irreducible component corresponding to t , and let $B_{s_1}, \dots, B_{s_m}$ be the strata defined above corresponding to $s_1, \dots, s_m$. Then we define B_t to be the complement of $\bigcup_{i=1}^m B_{s_i}$ in the closure of t in $\pi^{-1}(A)$.

Let $t_1, \dots, t_l$ be the stratum points of $\mathfrak{X}_n$ over η_A which are generic points of irreducible components of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$, and let $s_1, \dots, s_m$ be the stratum points of $\mathfrak{X}_n$ over η_A which are nodes or marked points of the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \times_{\mathfrak{X}_{n-1}} \eta_A \rightarrow \eta_A$. Note that it follows immediately from the construction and induction hypothesis that the strata $B_{t_1}, \dots, B_{t_l}, B_{s_1}, \dots, B_{s_m}$ are irreducible, locally closed in $\mathfrak{X}_n$, and mutually disjoint. Moreover, by the going-down theorem, every point of $\pi^{-1}(A)$ has a generalization which is a stratum point over η_A . Therefore, $B_{t_1}, \dots, B_{t_l}, B_{s_1}, \dots, B_{s_m}$ constitute a decomposition of $\pi^{-1}(A)$ by locally closed subsets. Since every stratum point of $\mathfrak{X}_n$ is over a stratum point of $\mathfrak{X}_{n-1}$, it follows that the decompositions obtained by performing the above construction for all strata of $\mathfrak{X}_{n-1}$ constitute a decomposition of $\mathfrak{X}_n$ such that the set of stratum points of $\mathfrak{X}_n$ coincides with the set of generic points of the strata of the decomposition.

Finally, since the closure of a stratum point x_n of $\mathfrak{X}_n$ is either a finite étale cover over the closure of the image x_{n-1} or an underlying scheme of a stable curve over the closure of the image x_{n-1} of x_n in $\mathfrak{X}_{n-1}$, it follows from the induction hypothesis and the various definitions involved that the above decomposition of $\mathfrak{X}_n$ is indeed a stratification.

The remaining assertions of the lemma follow immediately from the construction and the definitions involved. This completes the proof of the lemma. $\square$

Lemma 1.8 (Dimension of strata of the stable compactification). Let K be a complete discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \dots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \dots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Then for each stratum A of $\mathfrak{A}$, the dimension of the closure of A in $\mathfrak{X}_n$ coincides with the combinatorial dimension of the stratum points corresponding to the stratum A .

Proof. By noting that $\mathcal{O}_K$ is excellent (in particular, universally catenary), it follows immediately from the elementary scheme theory and the various definitions involved. $\square$

Lemma 1.9 (Strata of codimension 0 and 1 of the stable compactification). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \dots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \dots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to

its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n .

- (i) Let x be a stratum point of $\mathfrak{X}_n$, and let d_x be the combinatorial dimension of x . Then the following conditions are equivalent:

- (a) $d_x = n + 1$;
- (b) the stratum point x is a generic point of $\mathfrak{X}_n$.

Moreover, if the above conditions are satisfied, the corresponding stratum coincides with X_n as a subscheme of $\mathfrak{X}_n$.

- (ii) Let x be a stratum point of $\mathfrak{X}_n$, and let d_x be the combinatorial dimension of x . Then the following conditions are equivalent:

- (a) $d_x = n$;
- (b) the stratum point x is a generic point of an irreducible component of the divisor at infinity of $\mathfrak{X}_n$, i.e., a generic point of an irreducible component of the closed subspace $\mathfrak{X}_n \setminus X_n$ of $\mathfrak{X}_n$.

Moreover, for every irreducible component D of the divisor at infinity of $\mathfrak{X}_n$, the generic point of D is a stratum point of $\mathfrak{X}_n$.

Proof. It follows immediately from the definitions involved. $\square$

Definition 1.10. Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n .

- (i) We shall say that an irreducible component D of the divisor at infinity of $\mathfrak{X}_n$, i.e., an irreducible component of the closed subspace $\mathfrak{X}_n \setminus X_n$ of $\mathfrak{X}_n$, is a **divisor of cusp-type** if the following condition is satisfied:

- there exists an element $i \in \{1, \dots, n\}$ such that $\mathfrak{sgn}(\eta_D)(i) = \text{C}$, where η_D is the generic point of D .

Moreover, we shall say that the number of $i \in \{1, \dots, n\}$ such that $\mathfrak{sgn}(\eta_D)(i) = \text{C}$ is the **C-level** of D .

- (ii) We shall say that an irreducible component D of the divisor at infinity of $\mathfrak{X}_n$, i.e., an irreducible component of the closed subspace $\mathfrak{X}_n \setminus X_n$ of $\mathfrak{X}_n$, is a **divisor of vertex-type** if the following condition is satisfied:

- for every $i \in \{1, \dots, n\}$, $\mathfrak{sgn}(\eta_D)(i) = \text{V}$, where η_D is the generic point of D .

Lemma 1.11 (Strata and Divisors of Infinity of the stable compactification). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ with respect to its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Then for each stratum A of $\mathfrak{A}$ and each irreducible component of the divisor at infinity D of $\mathfrak{X}_n$, i.e., each irreducible component of the closed subspace $\mathfrak{X}_n \setminus X_n$ of $\mathfrak{X}_n$, A is either contained in D or disjoint from D .

Proof. By the construction of the stratification of $\mathfrak{X}_n$, which is given in the proof of Lemma 1.7, it follows immediately that the divisor D is a disjoint union of strata of the stratification of $\mathfrak{X}_n$. Therefore, for each stratum A of the stratification of $\mathfrak{X}_n$, the intersection $A \cap D$ is either empty or coincides with A . This completes the proof of the lemma. $\square$

Lemma 1.12 (Strata and Divisors of Infinity of the moduli stack of stable curves). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ with respect to its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_{n-1}$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_{n-1} , and let $\Phi: \mathfrak{X}_{n-1} \rightarrow \overline{\mathfrak{M}}_{g_n, r_n}$ be the morphism determined by the stable curve $(\mathfrak{X}_n, \mathfrak{D}_n) \rightarrow \mathfrak{X}_{n-1}$. Then for each stratum A of $\mathfrak{A}$ and each irreducible component of the divisor at infinity D of the moduli stack $\overline{\mathfrak{M}}_{g_n, r_n}$, i.e., each irreducible component of the boundary $\overline{\mathfrak{M}}_{g_n, r_n} \setminus \mathfrak{M}_{g_n, r_n}$, the intersection $\Phi^{-1}(D) \cap A$ is either empty or coincides with A .

Proof. By noting that $\overline{\mathfrak{M}}_{g_n, r_n}$ is a smooth Deligne-Mumford algebraic stack over $\operatorname{Spec} \mathbb{Z}$, it follows that for each divisor of $\overline{\mathfrak{M}}_{g_n, r_n}$, the pullback of the divisor to $\mathfrak{X}_{n-1}$ is a Cartier divisor. Thus, the assertion follows formally from the Lemma 1.11. This completes the proof of the lemma. $\square$

Definition 1.13. Let $X^{\log}$ be a locally Noetherian fs log scheme.

- (i) We shall say that $X^{\log}$ is **log-constant**, if there exists an fs monoid P and the chart $\underline{P} \rightarrow \mathcal{M}_{X^{\log}}$ of the log structure $\mathcal{M}_{X^{\log}}$ on X such that the induced morphism $\underline{P} \rightarrow \mathcal{C}_{X^{\log}}$ (where $\underline{P}$ is the constant sheaf associated to P) is an isomorphism. We shall refer to such a monoid P as a **type** of $X^{\log}$, and we shall say that $X^{\log}$ is of type P .
- (ii) We shall say that $X^{\log}$ is **étale locally log-constant** at $x \in X$, if there exists an étale neighborhood U of x such that the restriction of $X^{\log}$ to U is log-constant. We shall say that $X^{\log}$ is **étale locally log-constant**, if $X^{\log}$ is étale locally log-constant at every point of X . For each connected component of X , we shall say that a monoid P is a **type** of $X^{\log}$ at the connected component if there exists an étale covering U of the connected component such that the restriction of $X^{\log}$ to U is log-constant of type P . If X is connected, we shall simply say that P is a type of $X^{\log}$.

- (iii) We shall say that $X^{\log}$ is **characteristic-constant**, if the étale sheaf of characteristic $\mathcal{C}_{X^{\log}}$ of $X^{\log}$ is a constant sheaf. Let P be a monoid. Then we shall refer to an isomorphism $\underline{P} \cong \mathcal{C}_{X^{\log}}$ as a **trivialization of characteristic** of $X^{\log}$ (of type P).
- (iv) Let $X^{\log}$ be an étale locally log-constant fs log scheme. Let n be an integer. Then we shall say that $X^{\log}$ is of **log-rank** n if there exists an fs monoid P such that $X^{\log}$ is of type P and the rank of P^{gp} is equal to n .

Remark 1.14. The above definitions are an immediate generalization of the notion of “a morphism of type $\mathbb{N}^{\oplus n}$ ”, which is introduced in [Hoshi1].

Remark 1.15. We can easily generalize the above definitions to the relative situation, i.e., for a morphism of fs log schemes $f: X^{\log} \rightarrow Y^{\log}$ whose underlying morphism of schemes is an isomorphism, we can define the notions of f being log-constant, étale locally log-constant, or characteristic-constant by replacing $\mathcal{C}_{X^{\log}}$ with the étale sheaf of relative characteristic $\mathcal{C}_{X^{\log}/Y^{\log}}$ of f in the above definitions. However, we shall not need such a generalization in this paper, and therefore we do not give a precise definition of these notions.

Lemma 1.16 (Characterization of étale locally log-constant fs log schemes). Let $X^{\log}$ be an fs log scheme. Then the following conditions are equivalent:

- (i) $X^{\log}$ is étale locally log-constant;
- (ii) the étale sheaf of characteristic $\mathcal{C}_{X^{\log}}$ of $X^{\log}$ is étale locally constant.

Proof. It follows immediately from the definitions involved. $\square$

Lemma 1.17 (Correspondence between characteristic-constant log schemes and certain homomorphisms to the Picard group). Let X be a reduced scheme and P be an fs sharp monoid. Write $\mathbf{M}_X$ for the set of isomorphism classes of pairs $(\mathcal{M}, \alpha)$ such that

- $\mathcal{M}$ is an fs log structure on X ;
- Let $X^{\log}$ be the fs log scheme whose underlying scheme is X and whose log structure is $\mathcal{M}$. Then $X^{\log}$ is characteristic-constant and of type P ;
- $\alpha: \underline{P} \rightarrow \mathcal{C}_{(X, \mathcal{M})}$ is an isomorphism.

Write $\iota_X: \mathbf{M}_X \rightarrow \text{Hom}(P, \text{Pic}(X)) = \text{Hom}(P^{\text{gp}}, \text{Pic}(X))$ for the map defined as follows:

- For each element $(\mathcal{M}, \alpha)$ of $\mathbf{M}_X$, write $X^{\log}$ for the fs log scheme whose underlying scheme is X and whose log structure is $\mathcal{M}$. Then we define $\iota_X(\mathcal{M}, \alpha)$ to be the homomorphism $P \rightarrow \text{Pic}(X)$ defined as follows: for each element p of P , we define $\iota_X(\mathcal{M}, \alpha)(p)$ to be the isomorphism class of the invertible sheaf $(\alpha \circ \text{pr})^{-1}(p)$ on X , where $\text{pr}: \mathcal{M}_{X^{\log}} \rightarrow \mathcal{C}_{X^{\log}}$ is the natural projection.

Then the map ι_X is a bijection.

Proof. First, we shall construct the (inverse) map $\kappa_X: \text{Hom}(P, \text{Pic}(X)) \rightarrow \mathbf{M}_X$ of ι_X . Let $f: P \rightarrow \text{Pic}(X)$ be a homomorphism. Write $f^{\text{gp}}: P^{\text{gp}} \rightarrow \text{Pic}(X)$ for the homomorphism induced by f on the groupifications. Temporarily, we take an isomorphism $v: \mathbb{Z}^{\oplus n} \rightarrow P^{\text{gp}}$ of abelian groups, where n is the rank of P^{gp} . For $1 \leq i \leq n$, we write $\mathbf{e}_i$ for the i -th standard basis vector of $\mathbb{Z}^{\oplus n}$. Write $\mathcal{L}_i$ for the invertible sheaf on X corresponding to $f^{\text{gp}}(v(\mathbf{e}_i))$. Then we define an extension $0 \rightarrow \mathcal{O}_X^\times \rightarrow \mathcal{M} \rightarrow \underline{P} \rightarrow 0$ of sheaves on X as follows: we define $\mathcal{M}$ to be the coproduct of $\mathcal{L}_1^{\otimes a_1} \otimes \cdots \otimes \mathcal{L}_n^{\otimes a_n}$ for all $a_1, \dots, a_n \in \mathbb{Z}$ such that $v(a_1 \mathbf{e}_1 + \cdots + a_n \mathbf{e}_n) \in P$ as a sheaf of sets, and we define the monoid structure on $\mathcal{M}$ by the natural (various) isomorphisms. This construction yields an bijective correspondence between the set of extensions of $\underline{P}$ by $\mathcal{O}_X^\times$ and the set of homomorphisms from P to $\text{Pic}(X)$. Next, we shall construct a log structure $\mathcal{M} \rightarrow \mathcal{O}_X$ on X . Let s be a local section of $\mathcal{M}$, and let p be an element of P such that the image of s in $\underline{P}$ coincides with p . Then we define the image of s of the morphism $\mathcal{M} \rightarrow \mathcal{O}_X$ to be s ($\in \mathcal{O}_X^\times$) if $p = 0$ and to be 0 if $p \neq 0$. This construction yields a log structure on X , and the resulting fs log scheme is characteristic-constant and of type P . Thus, we have constructed the map $\kappa_X: \text{Hom}(P, \text{Pic}(X)) \rightarrow \mathbf{M}_X$.

Finally, we verify that ι_X and κ_X are inverse to each other. For each element $(\mathcal{M}, \alpha)$ of $\mathbf{M}_X$, we have an extension $0 \rightarrow \mathcal{O}_X^\times \rightarrow \mathcal{M} \rightarrow \underline{P} \rightarrow 0$ of sheaves on X , and the corresponding homomorphism $P \rightarrow \text{Pic}(X)$. To verify the assertion that ι_X and κ_X are inverse to each other, it suffices to verify that the log structure $\mathcal{M} \rightarrow \mathcal{O}_X$ satisfies the following condition: for each local section s of $\mathcal{M}$ and each element p of P such that the image of s in $\underline{P}$ coincides with p , the image of s by the morphism $\mathcal{M} \rightarrow \mathcal{O}_X$ coincides with s if $p = 0$ and with 0 if $p \neq 0$. The case $p = 0$ is trivial, so we assume that $p \neq 0$. Since the condition is local, we can check it on a smaller open subset.

Let U be an étale open subset of X such that s is defined, and the (local) chart $\varphi: \underline{P}_U \rightarrow \mathcal{M}|_U$ of the log structure $\mathcal{M}$ on X exists. Then s can be written as $\varphi(p) \cdot u$ for some $u \in \mathcal{O}_X^\times(U)$. Since the image of $\varphi(p)$ in $\mathcal{C}$ is nonzero everywhere on U , where $\mathcal{C}$ is the étale sheaf of characteristic of $\mathcal{M}$, it follows that the image of $\varphi(p)$ by the morphism $\mathcal{M} \rightarrow \mathcal{O}_X$ is not invertible at every point of U . Therefore, since X is reduced, the image of $\varphi(p)$ by the morphism $\mathcal{M} \rightarrow \mathcal{O}_X$ is 0. This completes the verification that ι_X and κ_X are inverse to each other, and therefore completes the proof of the lemma. $\square$

Definition 1.18. Let X be a reduced scheme and P be an fs sharp monoid. Let $(X^{\log}, \alpha)$ be a pair consisting of an fs log scheme $X^{\log}$ whose underlying scheme is X and an isomorphism $\alpha: \underline{P} \rightarrow \mathcal{C}_{X^{\log}}$. Let $f: P \rightarrow \text{Pic}(X)$ be the homomorphism corresponding to $(X^{\log}, \alpha)$ via the bijection ι_X in Lemma 1.17. We shall say that the pair $(X^{\log}, \alpha)$ is associated to the homomorphism f .

Lemma 1.19 (Description of the log structure of the stable compactification). Let K be a discrete valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified

stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ with respect to its parametrization. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Let A be a stratum of $\mathfrak{A}$. Then the following assertions hold:

- (i) Write $A^{\log}$ for the fs log scheme whose underlying scheme is A , endowed with the reduced scheme structure, and whose log structure is the pullback of the log structure on $\mathfrak{X}_n^{\log}$. Then $A^{\log}$ is étale locally log-constant.
- (ii) Let η_A be the generic point of A . Let π be the morphism $\mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1}$, and let $\bar{A}$ be the image of A in $\mathfrak{X}_{n-1}$ by π . Let $\eta_{\bar{A}}$ be the generic point of $\bar{A}$. We note that η_A and $\eta_{\bar{A}}$ are stratum points of $\mathfrak{X}_n$ and $\mathfrak{X}_{n-1}$, respectively. Define $A^{\log}$, $\bar{A}^{\log}$ for the fs log schemes as in (i) above. Then the following assertions hold:
 - (a) If $\mathbf{sgn}(\eta_A)(n) = V$, then the morphism $A^{\log} \rightarrow \bar{A}^{\log}$ induced by π is strict, i.e., the type of $A^{\log}$ coincides with the type of $\bar{A}^{\log}$.
 - (b) If $\mathbf{sgn}(\eta_A)(n) = C$, then there exists a natural isomorphism $\mathcal{C}_{A^{\log}} \cong \mathcal{C}_{\bar{A}^{\log}}|_A \oplus \underline{\mathbb{N}}$ of étale sheaves on A , where $\underline{\mathbb{N}}$ is the constant sheaf associated to $\mathbb{N}$.
 - (c) Let P be the type of $A^{\log}$ and $\bar{P}$ be the type of $\bar{A}^{\log}$. If $\mathbf{sgn}(\eta_A)(n) = N$, then there exists a pushout diagram of monoids:

$$\begin{array}{ccc} \mathbb{N} & \longrightarrow & \bar{P} \\ \downarrow \Delta & & \downarrow \\ \mathbb{N}^{\oplus 2} & \longrightarrow & P. \end{array}$$

- (iii) Let η_A be the generic point of A , and let d_A be the combinatorial dimension of η_A . Then the log rank of $A^{\log}$ is equal to $n + 1 - d_A$.

Proof. In light of Lemma 1.12, it follows immediately from the description of the (log) structure of the stable curves (cf. [KatoF]) and the various definitions involved. $\square$

Lemma 1.20 (Finiteness of automorphism groups of fs sharp monoids). Let P be an fs sharp monoid. Then the automorphism group $\operatorname{Aut}(P)$ of P is finite.

Proof. It follows from the well-known fact that the set of irreducible elements of P , i.e., the set of nonzero elements of P which cannot be written as a sum of two nonzero elements of P , is finite and generates P as a monoid, that the automorphism group $\operatorname{Aut}(P)$ of P is finite. This completes the proof of the lemma. $\square$

Lemma 1.21 (Representability of the functor of trivializations of characteristic of étale locally log-constant fs log schemes). Let $X^{\log}$ be a locally Noetherian, connected fs log scheme which is étale locally log-constant. Let P be a monoid

which is a type of $X^{\log}$. Define a functor $F_{X,P}$ from the category of schemes $\mathbf{Sch}$ to the category of sets $\mathbf{Set}$ as follows: for each scheme S , we define $F_{X,P}(S)$ to be the set of pairs (f, α) such that

- the morphism of schemes $f: S \rightarrow X$;
- the isomorphism $\alpha: \underline{P} \rightarrow f^* \mathcal{C}_{X^{\log}}$.

Then the functor $F_{X,P}$ is representable by a scheme, and the (canonical) morphism $F_{X,P} \rightarrow X$ is a finite étale covering.

Proof. Take an étale covering U of X such that the pullback of $\mathcal{C}_{X^{\log}}$ is a constant sheaf. Write F_1 for the scheme

$$\coprod_{V \subset U : \text{conn. comp.}} \coprod_{f \in \text{Hom}(\underline{P}_V, \mathcal{C}_{X^{\log}}|_V)} V.$$

In light of the general theory of descent, and since $\text{Aut}(P)$ is finite by Lemma 1.20, it follows that the scheme F_1 naturally descends to a scheme F_0 which is finite étale over X , and F_0 represents the functor $F_{X,P}$. This completes the proof of the lemma. $\square$

Theorem 1.22 (Exact sequence of log fundamental groups of fs log schemes which are étale locally log-constant). Let $X^{\log}$ be a connected, reduced fs log scheme which is étale locally log-constant. Write X for the underlying scheme of $X^{\log}$, or the log scheme obtained by endowing X with the trivial log structure. Let x be a geometric point of X , and let $\tilde{x}$ be a log geometric point of $X^{\log}$ lying over x . Then there exists an exact sequence of profinite groups

$$\pi_1(X^{\log} \times_X x, \tilde{x}) \rightarrow \pi_1(X^{\log}, \tilde{x}) \rightarrow \pi_1(X, x) \rightarrow 1.$$

Proof. First, we observe that we may assume without loss of generality that X is characteristic-constant. Indeed, by Lemma 1.21, there exists a connected finite étale covering Y of X such that the pullback of $\mathcal{C}_{X^{\log}}$ to Y is a constant sheaf. Write $Y^{\log}$ for the fs log scheme whose underlying scheme is Y and whose log structure is the pullback of the log structure on $X^{\log}$. Then $Y^{\log}$ is characteristic-constant. Now we observe that there exists a pullback diagram of profinite groups:

$$\begin{array}{ccc} \pi_1(Y^{\log} \times_Y y, \tilde{y}) & \longrightarrow & \pi_1(Y^{\log}, \tilde{y}) \\ \downarrow & & \downarrow \\ \pi_1(X^{\log} \times_X x, \tilde{x}) & \longrightarrow & \pi_1(X^{\log}, \tilde{x}), \end{array}$$

for some geometric point y of Y lying over x and some log geometric point $\tilde{y}$ of $Y^{\log}$ lying over $\tilde{x}$. Thus, if we verify the exactness of the sequence in the assertion for $Y^{\log}$, then it follows that the sequence in the assertion is also

exact for $X^{\log}$. Therefore, we may assume without loss of generality that X is characteristic-constant.

Next, we observe that we may assume without loss of generality that $\mathbb{N}^{\oplus n}$ is the type of $X^{\log}$ for some integer n . Indeed, by the elementary monoid theory, there exists an injection $P \rightarrow \mathbb{N}^{\oplus n}$ of monoids for some integer n , which induces an isomorphism $P^{\text{gp}} \cong (\mathbb{N}^{\oplus n})^{\text{gp}}$. By the description of characteristic-constant log schemes, stated in Lemma 1.17, it follows that there exists a morphism $X'^{\log} \rightarrow X^{\log}$ of fs log schemes such that the underlying morphism of schemes is an isomorphism and induced morphism on the characteristic is isomorphic (as a morphism of monoid sheaves) to the injection $\underline{P} \rightarrow \underline{\mathbb{N}^{\oplus n}}$.

We now proceed to show that the (natural) morphism of Galois categories $\text{Fét}(X^{\log}) \rightarrow \text{Fét}(X'^{\log})$ is an equivalence. By the descent theory of log schemes (cf. [Vidal, Proposition 4.4], [Hoshi1, Proposition B.8]), it suffices to verify the case where X is the spectrum of a strictly henselian local ring. However, the assertion follows immediately from the description of the két covering of a strictly henselian local fs log scheme (cf. [Hoshi1, Proposition B.2]).

Finally, we verify the exactness of the sequence in the assertion for $X^{\log}$ such that $\mathbb{N}^{\oplus n}$ is the type of $X^{\log}$ for some integer n . However, it follows immediately from [Hoshi1, Corollary 2]. $\square$

Lemma 1.23 (Slimness of the extension group of slim groups). Let $1 \rightarrow N \rightarrow G \rightarrow Q \rightarrow 1$ be an exact sequence of profinite groups. Assume that N and Q are slim. Then G is also slim.

Proof. Let H be an open subgroup of G . Let $g \in G$ be an element of centralizer of H . Then the image of g in Q is an element of the centralizer of the image of H in Q , and therefore the image of g in Q is trivial. Thus, g is an element of N . Since g is an element of the centralizer of $H \cap N$ in N , together with the assumption that N is slim, it follows that g is trivial. This completes the proof of the lemma. $\square$

Definition 1.24. Let K be a field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be a hyperbolic polycurve over K . Let ℓ be a prime number. We shall say that X is **geometrically pro- ℓ -exact** if for each $0 \leq r < s < t \leq n$, then the sequence of profinite groups

$$1 \rightarrow \Delta_{X_t/X_s}^{\ell} \rightarrow \Delta_{X_t/X_r}^{\ell} \rightarrow \Delta_{X_s/X_r}^{\ell} \rightarrow 1$$

is exact, where Δ_{X_j/X_i}^{ℓ} ($0 \leq i < j \leq n$) is the maximal pro- ℓ quotient of the kernel of the morphism $\Delta_{X_j} \rightarrow \Delta_{X_i}$.

Lemma 1.25 (Slimness of the geometric pro- ℓ fundamental group of a geometrically pro- ℓ -exact hyperbolic polycurve). Let K be a field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be a hyperbolic polycurve over K . Assume that X is geometrically pro- ℓ -exact. Then the geometric pro- ℓ fundamental group Δ_X^{ℓ} of X is slim.

Proof. By Lemma 1.23 and induction on the (relative) dimension of the polycurve, and since the geometric pro- ℓ fundamental group of a hyperbolic curve is slim, the assertion follows immediately. $\square$

Lemma 1.26 (Tameness of the outer representation of a geometrically pro- ℓ -exact hyperbolic polycurve which admits a compactified stable model). Let K be a complete discrete valuation field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be a hyperbolic polycurve over K . Let ℓ be a prime number which is invertible in $\mathcal{O}_K$. Assume X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization, and that X is geometrically pro- ℓ -exact. Then the outer representation $G_K \rightarrow \text{Out}(\Delta_X^\ell)$, which is induced by the exact sequence $1 \rightarrow \Delta_X^\ell \rightarrow \Pi_X^{(\ell)} \rightarrow G_K \rightarrow 1$, factors through the profinite group G_K/P_K .

Proof. Let s be the closed point of $\text{Spec } \mathcal{O}_K$, and let $\bar{s}^{\log}$ be a log geometric point of $\text{Spec } \mathcal{O}_K^{\log}$ lying over s . Consider the commutative diagram of homomorphisms of profinite groups:

$$\begin{array}{ccccccc} 1 & \longrightarrow & \Delta_X & \longrightarrow & \Pi_X & \longrightarrow & \pi_1(\text{Spec } K) \longrightarrow 1 \\ & & \downarrow & & \downarrow & & \downarrow \\ 1 & \longrightarrow & \pi_1(\mathfrak{X}^{\log, \text{geom}}) & \longrightarrow & \pi_1(\mathfrak{X}^{\log}) & \longrightarrow & \pi_1(\text{Spec } \mathcal{O}_K^{\log}) \longrightarrow 1, \end{array}$$

where $\mathfrak{X}^{\log, \text{geom}}$ is the log geometric fiber of $\mathfrak{X}^{\log} \rightarrow \text{Spec } \mathcal{O}_K^{\log}$ at $\bar{s}^{\log}$. We note that the profinite group $\pi_1(\text{Spec } \mathcal{O}_K^{\log})$ is (naturally) isomorphic to the quotient G_K/P_K of G_K by the wild inertia subgroup P_K .

By [Kisin, Theorem 1.4], the morphism $\Delta_X^\ell \rightarrow \pi_1(\mathfrak{X}^{\log, \text{geom}})^\ell$, which is obtained by taking the maximal pro- ℓ quotients of the morphism $\Delta_X \rightarrow \pi_1(\mathfrak{X}^{\log, \text{geom}})$, is an isomorphism. Now, by chasing the diagram above, the assertion in the lemma follows immediately. $\square$

Definition 1.27. Let K be a complete discrete valuation field. We shall refer to the profinite group $(G_K/P_K)/(I_K/P_K)^{(\ell')}$ as the ℓ -**tame absolute Galois group** of K , where $(I_K/P_K)^{(\ell')}$ is the maximal prime-to- ℓ quotient of I_K/P_K . We shall write $G_{K, \ell\text{-tame}}$ for the ℓ -tame absolute Galois group of K .

Definition 1.28. Let K be a field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be a hyperbolic polycurve over K . Let ℓ be a prime number which is invertible in $\mathcal{O}_K$. Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization, and geometrically pro- ℓ -exact.

- (i) Let $G_K \rightarrow \text{Out}(\Delta_X^\ell)$ be the outer representation induced by the exact sequence $1 \rightarrow \Delta_X^\ell \rightarrow \Pi_X^{(\ell)} \rightarrow G_K \rightarrow 1$. By Lemma 1.26, the outer representation $G_K \rightarrow \text{Out}(\Delta_X^\ell)$ factors through the profinite group G_K/P_K ,

where P_K is the wild inertia subgroup of G_K . Since Δ_X^ℓ is slim (cf. Lemma 1.25), we obtain an exact sequence of profinite groups

$$1 \rightarrow \Delta_X^\ell \rightarrow \Pi_{X, \text{tame}}^{(\ell)} \rightarrow G_K/P_K \rightarrow 1,$$

where $\Pi_{X, \text{tame}}^{(\ell)}$ is the pullback of profinite groups $G_K/P_K \times_{\text{Out}(\Delta_X^\ell)} \text{Aut}(\Delta_X^\ell)$.

We shall refer to $\Pi_{X, \text{tame}}^{(\ell)}$ as the **tamely reduced geometrically pro- ℓ fundamental group** of X .

- (ii) We shall refer to the pullback of the profinite group $\Pi_{X, \text{tame}}^{(\ell)} \times_{G_K/P_K} I_K/P_K$ as the **tamely reduced inertia geometrically pro- ℓ fundamental group** of X . We shall write $\Pi_{X, \text{t.iner.}}^{(\ell)}$ for the tamely reduced inertia geometrically pro- ℓ fundamental group of X .

- (iii) We shall say that X is **inertia pro- ℓ -exact** if for each $0 \leq i \leq n$, the outer representation $I_K/P_K \rightarrow \text{Out}(\Delta_{X_i}^\ell)$, which is induced by the exact sequence

$$1 \rightarrow \Delta_{X_i}^\ell \rightarrow \Pi_{X_i, \text{t.iner.}}^{(\ell)} \rightarrow I_K/P_K \rightarrow 1$$

factors through the maximal pro- ℓ quotient $(I_K/P_K)^\ell$ of I_K/P_K . (Here, we recall that the condition inertia pro- ℓ -exactness implies (by definition, as a tautology) the condition geometrically pro- ℓ -exactness.)

- (iv) Assume that X is inertia pro- ℓ -exact. Now we observe that the outer representation $G_K/P_K \rightarrow \text{Out}(\Delta_X^\ell)$, which is induced by the exact sequence in (i) above, factors through the ℓ -tame absolute Galois group $G_{K, \ell\text{-tame}}$ of K . Thus, we obtain an exact sequence of profinite groups

$$1 \rightarrow \Delta_X^\ell \rightarrow \Pi_{X, \ell\text{-tame}}^{(\ell)} \rightarrow G_{K, \ell\text{-tame}} \rightarrow 1,$$

where $\Pi_{X, \ell\text{-tame}}^{(\ell)}$ is the pullback of profinite groups $G_{K, \ell\text{-tame}} \times_{\text{Out}(\Delta_X^\ell)} \text{Aut}(\Delta_X^\ell)$. We shall refer to $\Pi_{X, \ell\text{-tame}}^{(\ell)}$ as the **ℓ -tamely reduced geometrically pro- ℓ fundamental group** of X . We abbreviate the term “ ℓ -tamely reduced geometrically pro- ℓ fundamental group” to “ **ℓ -TR geometrically pro- ℓ fundamental group**”.

- (v) Assume that X is inertia pro- ℓ -exact. We shall refer to the pullback of the profinite group $\Pi_{X, \ell\text{-tame}}^{(\ell)} \times_{G_{K, \ell\text{-tame}}} (I_K/P_K)^\ell$ as the **inertia pro- ℓ fundamental group** of X . We shall write $\Pi_{X, \text{iner.}}^\ell$ for the inertia pro- ℓ fundamental group of X .

Definition 1.29. Let K be a discrete valuation field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ be a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. We shall say that X is **graphically inert** if for each stratum point η of $\mathfrak{X}_i$ ($0 \leq i \leq n-1$), the action of the Galois group $\pi_1(\eta, \bar{\eta})$ on the dual graph of the fiber of $\mathfrak{X}_{i+1} \rightarrow \mathfrak{X}_i$ at η is trivial, where $\bar{\eta}$ is a geometric point of $\mathfrak{X}_i$ lying over η .

Remark 1.30. For a hyperbolic polycurve $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ over a field K , X is graphically inert if and only if for each stratum point η of X_i ($0 \leq i \leq n-1$), the dual graph of the hyperbolic curve $X_{i+1} \times_{X_i} \eta$ over the field $\kappa(\eta)$ is split. Moreover, this condition is equivalent to the condition that for every stratum A of $\mathfrak{X}_i$ ($1 \leq i \leq n$), letting $\bar{A}$ be the image of A in $\mathfrak{X}_{i-1}$, the induced morphism $A \rightarrow \bar{A}$ is geometrically connected. Thus, if X is graphically inert and A is a stratum of $\mathfrak{X}_i$ ($1 \leq i \leq n$) such that $\mathbf{sgn}(\eta_A)(i) = C$ (resp. $\mathbf{sgn}(\eta_A)(i) = N$), where η_A is the generic point of A , then the morphism $A \rightarrow \bar{A}$ is isomorphism (resp. isomorphism).

Definition 1.31. Let K be a complete discrete valuation field, $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . Let K^{ur} be the maximal unramified extension of K , let k be the residue field of K , and let $\bar{k}$ be an algebraic closure of k . Let ℓ be a prime number which is invertible in $\mathcal{O}_K$. Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ with respect to its parametrization, and that X is pro- ℓ -exact and graphically inert. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Let A be a stratum of $\mathfrak{A}$. Let η_A be the generic point of A . Let $a^{\log}$ be a log geometric point of $A^{\log}$ lying over a geometric point a of A .

- (i) Assume that $\mathbf{sgn}(\eta_A)(0) = G$. Then we shall refer to the maximal pro- ℓ quotient of the kernel of the morphism $\pi_1(A^{\log}, a^{\log}) \rightarrow \operatorname{Gal}(K^{\text{ur}}/K)$ as the **inertia pro- ℓ decomposition group** of A , and we shall write D_A^ℓ for the inertia pro- ℓ decomposition group of A .
- (ii) Assume that $\mathbf{sgn}(\eta_A)(0) = S$. Write $\bar{s}^{\log}$ be the log geometric point of $\operatorname{Spec} \mathcal{O}_K^{\log}$ which is image of $a^{\log}$. Then we shall refer to the maximal pro- ℓ quotient of the kernel of the morphism $\pi_1(A^{\log}, a^{\log}) \rightarrow \pi_1(\operatorname{Spec} k^{\log}, \bar{s}^{\log})$ as the **inertia pro- ℓ decomposition group** of A , and we shall write D_A^ℓ for the inertia pro- ℓ decomposition group of A .

Proposition 1.32. Let K be a complete discrete valuation field, and let $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ be a hyperbolic polycurve over K . Let ℓ be a prime number which is invertible in $\mathcal{O}_K$. Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ with respect to its parametrization, and that X is geometrically pro- ℓ -exact and graphically inert. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Let A be a stratum of $\mathfrak{A}$, and let η_A be the generic point of A . Let $a^{\log}$ be a log geometric point of $A^{\log}$ lying over a geometric point a of A . Write $\bar{a}^{\log}$ for the image of $a^{\log}$ in $\bar{A}^{\log}$, and write $\bar{a}$ for the underlying geometric point of $\bar{a}^{\log}$.

Proposition 1.33. かきなおし Then the following assertions hold:

- (i) The (natural) morphism of profinite groups $\pi_1(A^{\log}, a^{\log})^\ell \rightarrow \pi_1(\mathfrak{X}_n^{\log}, a^{\log})^\ell$ is injective.

- (ii) (a) If $\mathbf{sgn}(\eta_A)(n) = V$, then A is smooth over $\bar{A}$. Moreover, there exists a finite étale cover A' of $\bar{A}$ such that, letting Π' denote the corresponding fundamental group, the sequence of profinite groups

$$1 \rightarrow \pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(A^{\log}, a^{\log})^{\ell} \rightarrow \Pi'^{\ell} \rightarrow 1$$

is exact. Moreover, the profinite group $\pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell} \cong \pi_1(A \times_{\bar{A}} \bar{a}, a)^{\ell}$ is isomorphic to the maximal pro- ℓ quotient of the fundamental group of a smooth curve over an algebraically closed field of characteristic 0.

- (b) If $\mathbf{sgn}(\eta_A)(n) = C$, then A is finite étale over $\bar{A}$. Let Π' for the open subgroup of $\pi_1(\bar{A}^{\log}, \bar{a}^{\log})$ corresponding to the strict két covering associated to the morphism $A \rightarrow \bar{A}$. Then the sequence of profinite groups

$$1 \rightarrow \pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(A^{\log}, a^{\log})^{\ell} \rightarrow \Pi'^{\ell} \rightarrow 1$$

is exact. Moreover, the profinite group $\pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell}$ is isomorphic to $\mathbb{Z}_{\ell}(1)$.

- (c) If $\mathbf{sgn}(\eta_A)(n) = N$, then A is finite étale over $\bar{A}$. Let Π' for the open subgroup of $\pi_1(\bar{A}^{\log}, \bar{a}^{\log})$ corresponding to the strict két covering associated to the morphism $A \rightarrow \bar{A}$. Then the sequence of profinite groups

$$1 \rightarrow \pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(A^{\log}, a^{\log})^{\ell} \rightarrow \Pi'^{\ell} \rightarrow 1$$

is exact. Moreover, the profinite group $\pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell}$ is isomorphic to $\mathbb{Z}_{\ell}(1)$.

- (iii) Assume that the polycurve X is graphically inert. Then the natural morphism of profinite groups $\pi_1(A^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(\mathfrak{X}_n^{\log}, a^{\log})^{\ell}$ is injective, and the sequence of profinite groups

$$1 \rightarrow \pi_1(A^{\log} \times_{\bar{A}^{\log}} \bar{a}^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(A^{\log}, a^{\log})^{\ell} \rightarrow \pi_1(\bar{A}^{\log}, \bar{a}^{\log})^{\ell} \rightarrow 1$$

is exact.

Proof. mada kaite nai □

Lemma 1.34. Let K be a complete discrete valuation field, $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \text{Spec } K$ a hyperbolic polycurve over K . Assume that X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \text{Spec } \mathcal{O}_K$ with respect to its parametrization. Let ℓ be a prime number which is invertible in $\mathcal{O}_K$. Assume that X is pro- ℓ -exact and graphically inert. Let $\mathfrak{A}$ be the stratification of $\mathfrak{X}_n$ obtained by applying the construction in Lemma 1.7 to the hyperbolic polycurve X_n . Let A be a stratum of $\mathfrak{A}$. Then the following assertions hold:

- (i) the sequence of profinite groups

$$1 \rightarrow I_A^\ell \rightarrow D_A^\ell \rightarrow \pi_1(A, a)^\ell \rightarrow 1$$

is exact.

- (ii) the pro- ℓ inertia group I_A^ℓ of A is isomorphic to a direct product of r copies of $\mathbb{Z}_\ell(1)$, where r is the log-rank of $A^{\log}$.
- (iii) A has a natural structure of a hyperbolic polycurve over k , where k is the residue field of K as a discrete valuation field, and A is pro- ℓ -exact as a hyperbolic polycurve over k .
- (iv) if the dimension of A is n , then I_A^ℓ is the center of D_A^ℓ . このへんでちょっと調整がほしい

Proof. mada kaite nai

□

チェックポイント: strata vs. stratum 問題, complete うまくいってるか問題, cusp 忘れてないか問題, 概念の名前だいじょうぶか問題, Notation check,

2 Cofinality of l -admissible finite étale coverings

Definition 2.1. Let K be a field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . We shall say that X is **pro- l exact** (with respect to its parametrization) if for each $0 \leq i \leq n$, the (natural) sequence of profinite groups

$$1 \rightarrow \Delta_{X_n/X_i}^l \rightarrow \Pi_{X_n}^l \rightarrow \Pi_{X_i}^l \rightarrow 1$$

is exact. Here, Δ_{X_n/X_i} denotes the kernel of the homomorphism $\Pi_{X_n} \rightarrow \Pi_{X_i}$ induced by the morphism $X_n \rightarrow X_i$.

Definition 2.2. Let K be a valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . Let l be a prime number which is different from the characteristic of the residue field of K . We shall say that X is **l -quasi-fine** (with respect to its parametrization) if X admits a compactified stable model $\mathfrak{X} = \mathfrak{X}_n \rightarrow \mathfrak{X}_{n-1} \rightarrow \cdots \rightarrow \mathfrak{X}_1 \rightarrow \mathfrak{X}_0 = \operatorname{Spec} \mathcal{O}_K$ and X is pro- l exact. We shall say that X is **l -fine** (with respect to its parametrization) if X is l -quasi-fine and for each $i = 1, \dots, n$ and each geometric point $\bar{x}$ of X_{i-1} which lies over the closed point of $\operatorname{Spec} \mathcal{O}_K$, the geometric fiber $X_{i,\bar{x}}$ of the morphism $X_i \rightarrow X_{i-1}$ over $\bar{x}$ is singular.

Theorem 2.3. Let K be a valuation field and $X = X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow X_0 = \operatorname{Spec} K$ a hyperbolic polycurve over K . Let l be a prime number which is different from the characteristic of the residue field of K . Then there exists a finite étale covering $Y \rightarrow X$ such that Y is l -fine (with respect to its parametrization induced by the morphism $Y \rightarrow X$).

Proof. proof here □

3 Dehn-twist Pointed Stable Curves type Outer Representations

4 Reconstruction of Log-full Subgroups

「 l -fine な polycurve が与えられたなら、[HMM] の議論をそのまま援用することで、residually l -cyclotomically full くらいの状況のもとで、log full subgroup を思い出すことができる。ほとんど議論一緒なのでアレだけど、self-contained-ness のこともあって、もう一度かいてみる。ただし、本質的に新しいことはなにもしていないことには注意したい。」

5 Reconstruction of Dimensions via Corank 1 Intersections

6 Reconstruction of the Graph-theoretic Structure of Proper Hyperbolic Polycurves of Dimension 2

References

References

- [KatoF] F. Kato, *Log smooth deformation and moduli of log smooth curves*, Int. J. Math. **11** (2000), no. 2, 215 – 232.
- [MT] S. Mochizuki and A. Tamagawa, *The algebraic and anabelian geometry of configuration spaces*, Hokkaido Math. J. **37** (2008), 75–131.
- [DM] P. Deligne and D. Mumford, *The irreducibility of the space of curves of given genus*, IHES Publ. Math. **36** (1969), 75–109.
- [Hur] S. Mochizuki, *The geometry of the compactification of the Hurwitz scheme*, Publ. Res. Inst. Math. Sci. **31** (1995), 355–441.
- [CmbGC] S. Mochizuki, *A combinatorial version of the Grothendieck conjecture*, Tohoku Math. J. **59** (2007), 455–479.
- [CbGT] Y. Hoshi, A. Minamide, and S. Mochizuki, *Group-theoreticity of numerical invariants and distinguished subgroups of configuration space groups*, Kodai Math. J. **45** (2022), 295–348.
- [NodNon] Y. Hoshi and S. Mochizuki, *On the combinatorial anabelian geometry of nodally nondegenerate outer representations*, Hiroshima Math. J. **41** (2011), 275–342.
- [CbTpI] Y. Hoshi and S. Mochizuki, *Topics surrounding the combinatorial anabelian geometry of hyperbolic curves I: Inertia groups and profinite Dehn twists*, Galois-Teichmüller Theory and Arithmetic Geometry, Adv. Stud. Pure Math., **63**, Math. Soc. Japan (2012), 659–811.
- [Hoshi1] Y. Hoshi, *The exactness of the log homotopy sequence*, Hiroshima Math. J. **39** (2009), 61–121.
- [Vidal] I. Vidal, *Morphismes log étales et descents par homéomorphismes universels*, C. R. Math. Acad. Sci. Paris. **332** (2001), 239–244.
- [Kisin] M. Kisin, *Prime to p fundamental groups and tame Galois actions*, Annals de l’Institut Fourier **50**, no. 4 (2000), 1099–1126.