

Binding of FAD and tryptophan to the tryptophan 6-halogenase Thal is negatively coupled

Ann-Christin Moritzer | Hartmut H. Niemann

Department of Chemistry, Bielefeld University, Bielefeld, Germany

Correspondence

Hartmut Niemann, Department of Chemistry, Bielefeld University, Universitätsstraße 25, 33615 Bielefeld, Germany.
 Email: hartmut.niemann@uni-bielefeld.de

Funding information

Bielefeld University; Deutsche Forschungsgemeinschaft

Abstract

Flavin-dependent halogenases require reduced flavin adenine dinucleotide (FADH₂), O₂, and halide salts to halogenate their substrates. We describe the crystal structures of the tryptophan 6-halogenase Thal in complex with FAD or with both tryptophan and FAD. If tryptophan and FAD were soaked simultaneously, both ligands showed impaired binding and in some cases only the adenosine monophosphate or the adenosine moiety of FAD was resolved, suggesting that tryptophan binding increases the mobility mainly of the flavin mononucleotide moiety. This confirms a negative cooperativity between the binding of substrate and cofactor that was previously described for other tryptophan halogenases. Binding of substrate to tryptophan halogenases reduces the affinity for the oxidized cofactor FAD presumably to facilitate the regeneration of FADH₂ by flavin reductases.

KEY WORDS

cofactor binding, cofactor regeneration, crystal structure, flavin adenine dinucleotide, flavin-dependent halogenase, negative cooperativity, substrate binding, tryptophan halogenase

1 | INTRODUCTION

Flavin-dependent halogenases (FDHs) mediate the regioselective halogenation of a variety of natural products including antibiotics and anticancer compounds.^{1,2} To halogenate their substrates, FDHs require reduced flavin adenine dinucleotide (FADH₂), molecular oxygen, and halide salts (usually Cl⁻ or Br⁻). The halide ion is bound between the protein and the isoalloxazine ring of the cofactor.³ In the first reaction step, FAD(C4a)-OOH is formed.⁴ The flavin hydroperoxide then oxidizes the halide to hypohalous acid, which is assumed to move through the protein toward the ε-amino group of a catalytically essential lysine residue.³ Kinetic analysis established that substrate chlorination occurs after completion of flavin redox reactions and that the substrate does not participate in reactions leading from FADH₂ to FAD(C4a)-oxygenated species to FAD⁴ (Figure S1). In the absence of substrate, a

long-lived chlorinating intermediate is formed, which was proposed to be a lysine chloramine generated by the reaction of HOCl with the active site lysine of the enzyme.⁵ Regioselectivity is achieved by positioning the substrate relative to the ε-amino group of the catalytic lysine.⁶ Depending on whether its C5, C6, or C7 atom is positioned most closely to this lysine, the substrate tryptophan is halogenated in position 5, 6, or 7 by various tryptophan halogenases. So far, crystal structures of one tryptophan 5-halogenase (PyrH⁶), two tryptophan 7-halogenases (PrnA³ and RebH^{5,7}), and several tryptophan 6-halogenases (SttH,⁸ Th-Hal,⁹ Thal,¹⁰ and Tar14¹¹) have been described. Comparison of apo, FAD-bound and L-Trp-bound structures reveals two flexible loops that undergo conformational changes upon binding of substrate or cofactor. When L-Trp is bound in the active site, it is covered by a substrate-binding loop that is disordered in structures without substrate. Binding of FAD usually results in a closed conformation of the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Protein Science* published by Wiley Periodicals, Inc. on behalf of The Protein Society.

FAD loop (e.g., in PrnA, PyrH, SttH, and Tar14), while in apo structures either it adopts an open conformation (e.g., in Thal) or it is not resolved (e.g., in PyrH and BrvH¹²).

To close the reaction cycle, FADH₂ needs to be regenerated. This is achieved by separate NAD(P)H-dependent flavin reductases.¹³ Often a flavin reductase is present in a biosynthetic gene cluster along with an FDH, forming a two-component system, for example, RebH and RebF.^{14,15} Rather little is known about the mechanistic details of FAD reduction. For example, it is not completely clear whether FAD can be reduced while still bound to the FDH or whether it has to dissociate from the halogenase. This may be protein-specific as the FAD affinity appears to vary considerably as inferred from cofactor loading upon purification of overexpressed enzymes.¹⁶

For RebH, a weaker FAD binding was observed in crystals concomitantly soaked with FAD and L-Trp, and a negative coupling between cofactor and substrate binding was proposed.^{5,7} We observed a similar phenomenon in Thal (also known as ThdH), a tryptophan 6-halogenase from *Streptomyces albogriseolus* involved in the biosynthesis of thienodolin.^{17,18} We previously reported the structures of Thal in the apo and Trp-bound forms.¹⁰ During our attempt to obtain the structures of Thal bound to FAD and in complex with both FAD and L-Trp, we found that binding of L-Trp and FAD is negatively coupled as well. Here, we describe these structures of Thal soaked with FAD and with FAD and L-Trp.

2 | RESULTS

2.1 | Structure determination

Because our attempts to cocrystallize Thal with FAD failed, we soaked crystals of apo-Thal with FAD and NaCl or with a combination of L-Trp, FAD, and NaCl resulting in the FAD-Thal and Trp-FAD-Thal structures, respectively (Table 1). As described previously,¹⁰ Thal crystallized as a dimer in the asymmetric unit with the substrate-binding loop located at the dimer interface and the FAD loop on the outside (Figure 1 and Figure S2).

2.2 | Binding of FAD to Thal results in closing of the FAD loop

The FAD-Thal structure contains FAD with an occupancy of 0.9 in both chains (Figure S3). Compared to apo-Thal (PDB ID 6H43) and Trp-Thal (PDB ID 6H44), the conformation of the FAD loop changes to the closed conformation (Figures 1a and 2a,b). Upon switching from the open to closed state, Pro40 undergoes a peptide flip and Glu49, the other flanking residue, flips its side chain by almost 180° as in PyrH.⁶ Conformational switching of the FAD loop probably also occurs


TABLE 1 X-ray data collection and refinement statistics. Values in parentheses are for the highest resolution shell

	FAD-Thal	Trp-FAD-Thal
PDB ID	6SLS	6SLT
Space group	<i>P</i> 6 ₄	<i>P</i> 6 ₄
Unit-cell constants		
<i>a</i> = <i>b</i> , <i>c</i> (Å)	138.6, 142.1	138.3, 142.0
Data collection statistics		
Wavelength (Å)	0.97630	1.000
Resolution range (Å)	50.0–2.32 (2.37–2.32)	50.0–2.70 (2.80–2.70)
No. of reflections	1,391,920/66,881 (measured/unique)	875,709/42,328
Completeness (%)	100 (100)	100 (100)
<i>R</i> _{meas} (%)	6.4 (194.4)	18.1 (178.5)
<i>R</i> _{pim} (%)	1.4 (41.9)	4.0 (39.3)
Redundancy	20.8 (21.4)	20.7 (20.4)
Mean <i>I</i> / <i>σI</i>)	28.5 (1.8)	14.6 (2.1)
CC ½ (%)	100 (81.2)	99.8 (77.5)
Wilson <i>B</i> factor (Å ²)	72	70
Refinement and model statistics		
Resolution range	49.64–2.32 (2.38–2.32)	49.58–2.70 (2.77–2.70)
No. of reflections	63,482 (4673)/ (work/test)	40,099 (2963)/ 3,359 (254) 2,195 (167)
<i>R</i> _{work} (%)	18.0 (29.7)	18.4 (27.8)
<i>R</i> _{free} (%)	22.4 (33.1)	24.1 (35.4)
r.m.s.d. bonds (Å)	0.008	0.009
r.m.s.d. angles (°)	1.507	1.588
No. of atoms	8,585	8,621
Protein	8,308	8,404
Ligands	150	142
Solvent	127	75
Average <i>B</i> factor (Å ²)	77	75
Protein	77	75
Ligands	86	90
Solvent	67	56
Ramachandran		
Favored (%)	95.55	93.70
Allowed (%)	4.36	6.02
Outliers (%)	0.10	0.29

in the other FDHs, although two conformations have only been observed in PyrH.⁶

To address the potential crystal-to-crystal variations of FAD loading and the conformation of flexible loops, we determined the structures of four other Thal crystals soaked

(a) FAD-Thal Dimer



(b) Trp-FAD-Thal Dimer




FIGURE 1 Thal dimer in complex with cofactor or cofactor and substrate. (a) Structure of FAD-Thal. The box and pyramid subdomains are shown in deep teal and green cyan, as well as cyan and pale cyan, respectively. (b) Structure of Trp-FAD-Thal. The box and pyramid subdomains are shown in forest and pale green, as well as split pea green and limon, respectively. The FAD loop that undergoes a conformational change upon FAD binding is highlighted in dark red. The substrate-binding loop is shown in magenta. The substrate L-Trp is shown as a stick model with yellow carbon atoms, FAD and AMP are shown with carbons in pink. An enlarged view of the Trp-FAD-Thal dimer is provided as Figure S2

with FAD and chloride at resolutions of 2.5–2.8 Å. We did not fully refine or deposit these additional structures, but could reliably extract the following features. In all structures both active sites are empty, FAD binds to both monomers with high occupancy and the FAD loop is closed. The substrate-binding loop is defined in chain A of three crystals but disordered in the remaining Thal protomers.

To distinguish between a chloride ion with partial occupancy and water in the canonical halide-binding site formed by the backbone NH groups of Ser360 and Gly361, we collected long-wavelength data sets ($\lambda = 2.1015$ Å) for three crystals with an anomalous redundancy of 9.7 each. Anomalous difference density is visible for sulfur atoms and the phosphates in FAD in all three crystals. Only three of the six

Thal protomers have weak anomalous difference densities in the canonical halide-binding site, but no such density is present in the crystal that we fully refined. Therefore, water A763/B741 occupies the catalytic halide-binding site in the published FAD-Thal structure (PDB ID 6SLS).

There are structural differences between FAD-Thal and Trp-Thal outside the FAD loop and the substrate-binding loop. Gly13 undergoes a peptide flip so that in FAD-Thal, the NH of Gly14 rather than the O of Gly13 points toward the FAD phosphates. Val52 and Pro53 move into the tryptophan-binding site by 0.6 Å–1.1 Å (Figure 2c). These amino acids could be viewed as the continuation of the FAD loop. Thus, subtle structural changes in the tryptophan-binding site might be directly coupled to the binding of FAD. In PyrH, a similar, although




FIGURE 2 Conformational change of the FAD loop and details of substrate-binding site. (a) Overlay of the FAD-binding site of FAD-Thal (pale cyan and dark red) and the FAD loop of chain A of Trp-FAD-Thal (salmon). The cofactor FAD is shown with carbons in yellow and AMP with carbons in grey. The FAD loop is highlighted in dark red in the closed state and in salmon in the open state. In chain A of the Trp-FAD-Thal structure, the phosphate of AMP points out of the FAD-binding site. Several residues are hidden for a better view of the FAD and AMP. (b) Hydrogen bonds between FAD (yellow) and Thal (pale cyan and dark red) or protein-bound waters are shown as dotted yellow lines. (c) Steric conflicts upon placing L-Trp from Trp-Thal in FAD-Thal. L-Trp from Trp-Thal chain A (PDB ID 6H44) was placed in FAD-Thal chain A. Minor clashes are indicated as small green hexagons, more severe clashes are shown as bigger red hexagons. Clashes mainly occur between the indole moiety of L-Trp and the side chains of the conserved residues Pro53, Phe112, Phe465, and Trp466. Conserved residues Tyr454 and Tyr455 are not shown because the substrate-binding loop is disordered. Coloring of the active site residues according to previously published work¹⁰

more pronounced, narrowing of the tryptophan-binding site occurs upon FAD binding. Naismith and coworkers suggested that residues at and following the C-terminal end of the FAD loop act as a conduit for information transfer between the cofactor-binding site and the substrate-binding site.⁶ Further differences between FAD-Thal and Trp-Thal may be functionally important. The side chains of Ser360 and Tyr366 adopt other rotamers. In FAD-Thal, Tyr366 forms a hydrogen bond to the side chain of Asn54, which also interacts with the carboxylate of Glu461. Because Glu461 forms a salt bridge with

the α -amino group of L-Trp, the residues Tyr366, Asn54, and Glu461 might form another path for coupling the FAD-binding site and the substrate-binding site.

2.3 | Simultaneous soaking of FAD and L-Trp impairs the binding of cofactor and substrate

By soaking with FAD, NaCl, and L-Trp, we did not achieve simultaneous high occupancy for FAD and L-Trp in both chains (Figure 1b and Figure S4). Chain A of Trp-FAD-Thal

has a well-defined L-Trp (occupancy of 0.95), but only the adenosine monophosphate (AMP) moiety of FAD is visible in the electron density (occupancy of 0.96). Reduced FAD binding is not due to crystal contacts because Trp-FAD-Thal crystals are isomorphous to FAD-Thal crystals, which show binding of the complete FAD. The FAD loop is not involved in any crystal contacts (Figure S5). In chain B, the complete FAD is defined (occupancy = 0.93), but there is no clear electron density for L-Trp. The substrate-binding loop is ordered in chain A, while it is poorly defined in chain B and, therefore, could not be modeled between residues 454 and 456 (Figure 1b and Figure S2). The FAD loop is closed in chain B, but remains open in chain A despite the clear density for the AMP moiety. This can be rationalized because the adenosine moiety mostly interacts with a nonflexible region (residues 37–40). The phosphate of the AMP moiety does not occupy the same position as in FAD, but points toward the protein surface instead of the binding pocket for the FMN moiety (Figure 2a).

To address potential crystal-to-crystal variation, we determined the structures of additional crystals. We did not fully refine or deposit these additional structures, but could reliably extract the features described below. Four additional Thal crystals cocrystallized or soaked with 5 mM L-Trp (resolution of 2.6 Å–3.0 Å) show a clear difference density for L-Trp in both chains. We also analyzed four additional Thal crystals soaked with L-Trp, FAD, and NaCl at resolutions of 2.6–3.1 Å. One crystal shows essentially the same features as described above, except that in chain A only the adenine instead of the AMP moiety of FAD is well defined. In one crystal, L-Trp and adenosine are visible in both chains. In the last two crystals, L-Trp is present in both chains, there is no convincing density for any part of FAD in chain A and FAD with extremely high B-factors could be placed in chain B. To summarize, when tryptophan is bound, mostly only the AMP or adenosine moiety or a poorly bound FAD with high B-factors was observed.

In the published Trp-FAD-Thal structure, the water molecule B713 occupies the possible chloride position and has a lower B-factor than its surroundings. To distinguish between water and a chloride ion with partial occupancy, we collected long-wavelength data sets ($\lambda = 2.0664 \text{ \AA}$) for three Trp-FAD-Thal crystals, including the crystal in Table 1, with an anomalous redundancy of about 10.2 each. Anomalous difference density is visible for sulfur atoms and some, but not all, phosphates in FAD or AMP. As there is no anomalous difference density in the canonical halide-binding site in all three crystals, we only modeled water in the published structure.

3 | DISCUSSION

Our structures clearly hint to negatively cooperative binding of L-Trp and FAD. Since the structural changes are minor

except for the two flexible loops, it is hard to identify the structural basis of communication between the substrate-binding site and the FAD-binding site. The movement of Val52 and Pro53 into the active site when FAD is bound could lead to clashes of Val52 and Pro53 with L-Trp (Figure 2c). In reverse, binding of L-Trp could push out Val52 and Pro53. As a result, the FAD loop would be more likely to adopt the open conformation when L-Trp is bound, thereby promoting the release of the FAD or at least its FMN moiety. The side chain movement of Tyr366 might represent a second way to couple the FAD and the tryptophan-binding sites via Asn54 and Glu461.

To understand whether negative coupling of substrate and cofactor binding is a more general feature of FDHs, we analyzed all FDHs for which structures with both cofactor and substrate are available. These are the three tryptophan halogenases PrnA,^{3,19,20} RebH,^{5,7} and PyrH,⁶ and the enzymes PltM in complex with phloroglucinol¹⁶ and Mala' bound to premalbrancheamide.²¹

Mala' and PrnA structures provide no hints to negative cooperativity. All structures have one chain per asymmetric unit, and both ligands are bound at occupancies of 1.^{3,19–21} Nevertheless, Zhu et al. mention in their paper about PyrH that “Similar disturbances to the flavin site were observed in [...] PrnA structures when tryptophan was bound.”⁶

PltM structures show negative coupling between FAD and substrate binding. In the presence of the substrate phloroglucinol, FAD is missing from two of four chains in the asymmetric unit and only portions of FAD are resolved in unusual binding poses in the other two chains.¹⁶ Although in Thal the AMP moiety of FAD is better resolved, in phloroglucinol-bound PltM there is no electron density for the ADP part, which is assumed to be flexible. Tsodikov and coworkers suggested that the unusual location of the isoalloxazine moiety may represent a mechanistically important state, in which it can undergo redox chemistry while being sufficiently shielded from solvent.¹⁶

Similarity is highest between Thal and RebH and PyrH. For PyrH, three structures are available with four chains per asymmetric unit each.⁶ In the absence of substrate, all four chains bind FAD and chloride. In the absence of FAD, all four chains bind tryptophan. In the presence of both FAD and L-Trp, FAD was modeled in all chains, but L-Trp was modeled only in chain B. For RebH, seven structures with two chains per asymmetric unit were published, all of them being isomorphous.^{5,7} In FAD-bound RebH without substrate, FAD was modeled with an occupancy of 1 in both chains. The structure of Trp-bound, cofactor-free RebH was obtained by soaking with saturated L-Trp and 5 mM FAD. The FAD is clearly present in this structure, but was not modeled due to disorder.⁵ In the structure with bound FAD and tryptophan, Bitto et al. modeled FAD with an occupancy of 0.8 in chain

A and only the adenosine moiety of FAD with an occupancy of 0.8 in chain B.⁷ Thus, very similar to Thal, binding of tryptophan to RebH destabilizes binding of the FMN moiety in at least one chain of the RebH dimer.

The functional implications of these observations still need to be explored experimentally. Nevertheless, one can put forward a plausible model based on previously published data. Yeh et al. showed for RebH that the formation of hypochlorite, which is coupled to the oxidation of FADH₂ to FAD, can proceed in the absence of the substrate Trp.⁴ It would make sense that the subsequent binding of tryptophan destabilizes binding of the oxidized cofactor FAD to generate an empty cofactor-binding site ready to accept reduced FADH₂ for the next reaction cycle. In the related two-component flavin-dependent monooxygenases, results from many systems indicated that flavin transfer between reductase and monooxygenase can be achieved by free diffusion.²² Alternatively, the structures could suggest that FAD does not exchange completely but stays bound to Thal or RebH via its AMP or adenosine moiety. We interpret the AMP density in our structure as a result of high mobility in the FMN part of FAD rather than hydrolysis of the cofactor. The FMN moiety would thus leave the halogenase and could be reduced by a NAD(P)H-dependent flavin reductase. At least some of the heterologous flavin reductases commonly used to provide FADH₂ for FDHs in *in vitro* halogenation assays like SsuE^{13,16} or Fre²⁰ from *E. coli* accept FMN or even prefer FMN over FAD.^{23,24} The flavin nucleotide-bound crystal structures of SsuE²⁵ and Th-Fre, a flavin reductase from *Bacillus subtilis*,⁹ contain FMN. Thus, it may not be completely unreasonable that the adenosine or the AMP moiety remain bound to the halogenase while the flipped-out FMN moiety can be accessed by a flavin reductase.

4 | MATERIALS AND METHODS

4.1 | Protein expression, purification, and crystallization

Thal was expressed and purified as described.¹⁰ Purified Thal was crystallized using the sitting drop vapor-diffusion method at 20 °C with a drop ratio of 2:1 of protein solution (~15 mg·mL⁻¹) and reservoir solution (0.1 M bicine pH 8.4, 1.6 M K₂HPO₄/KH₂PO₄ or 0.1 M HEPES pH 7.8, 1.3 M K₂HPO₄/NaH₂PO₄). Crystals appeared as hexagonal prisms within 7 days. FAD-Thal crystals were obtained by soaking Thal crystals in reservoir solution additionally containing 12.4 mM FAD and 10 mM NaCl for 45 min. To obtain an L-Trp-FAD bound structure, Thal crystals were soaked in reservoir solution additionally containing 5 mM L-Trp, 5 mM FAD, and 200 mM NaCl for 85 min. For cryoprotection, the crystals

were transferred to reservoir solution supplemented with ~25 % glycerol and ligands before flash cooling in liquid nitrogen.

4.2 | Data collection, structure determination, and refinement

All data were collected at a temperature of 100 K at beamline P13 operated by EMBL Hamburg at the PETRA III storage ring at DESY, Hamburg, Germany.²⁶ The data sets were processed with XDS and scaled with XSCALE.²⁷ Both structures were determined by molecular replacement using the program Phaser²⁸ and the apo-Thal structure (PDB ID 6H43) as search model. The structures were improved by modeling in COOT²⁹ and restrained refinement in Refmac5³⁰ using noncrystallographic symmetry restraints. Structures were built to near completion before placing ligands into difference density. The occupancy of all ligands was refined. The figures of the final structures were generated using PyMOL. The coordinates and structure factors of the FAD-Thal and Trp-FAD-Thal structures were deposited in the Protein Data Bank with accession codes 6SLS and 6SLT, respectively.

ACKNOWLEDGMENTS

The synchrotron MX data were collected at beamline P13 operated by EMBL Hamburg at the PETRA III storage ring (DESY, Hamburg, Germany). We would like to thank Dr. Isabel Bento, Dr. Johanna Hakanpää, and Saravanan Panneerselvam for the assistance in using the beamline. We thank Hannah Minges and Norbert Sewald (Bielefeld University) for the Thal expression plasmid. We acknowledge support for the Article Processing Charge by the Deutsche Forschungsgemeinschaft and the Open Access Publication Fund of Bielefeld University.

CONFLICT OF INTEREST

The authors declare no potential conflict of interest.

ORCID

Ann-Christin Moritzer  <https://orcid.org/0000-0002-7647-9594>
Hartmut H. Niemann  <https://orcid.org/0000-0002-8582-0360>

REFERENCES

1. Weichold V, Milbredt D, Van Pée KH. Specific enzymatic halogenation - from the discovery of halogenated enzymes to their applications *in vitro* and *in vivo*. *Angew Chem Int Ed*. 2016;55:6374–6389.

2. Blasiak LC, Drennan CL. Structural perspective on enzymatic halogenation. *Acc Chem Res.* 2009;42:147–155.
3. Dong C, Flecks S, Unversucht S, Haupt C, van Pee KH, Naismith JH. Tryptophan 7-halogenase (PrnA) structure suggests a mechanism for regioselective chlorination. *Science.* 2005;309: 2216–2219.
4. Yeh E, Cole LJ, Barr EW, Bollinger JM, Ballou DP, Walsh CT. Flavin redox chemistry precedes substrate chlorination during the reaction of the flavin-dependent halogenase RebH. *Biochemistry.* 2006;45:7904–7912.
5. Yeh E, Blasiak LC, Koglin A, Drennan CL, Walsh CT. Chlorination by a long-lived intermediate in the mechanism of flavin-dependent halogenases. *Biochemistry.* 2007;46:1284–1292.
6. Zhu X, De Laurentis W, Leang K, et al. Structural insights into regioselectivity in the enzymatic chlorination of tryptophan. *J Mol Biol.* 2009;391:74–85.
7. Bitto E, Huang Y, Bingman CA, Singh S, Thorson JS, Phillips GN. The structure of flavin-dependent tryptophan 7-halogenase RebH. *Proteins.* 2008;70(1):289–293.
8. Shepherd SA, Menon BRK, Fisk H, et al. A structure-guided switch in the regioselectivity of a tryptophan halogenase. *Chembiochem.* 2016;17:821–824.
9. Menon BRK, Latham J, Dunstan MS, et al. Structure and biocatalytic scope of thermophilic flavin-dependent halogenase and flavin reductase enzymes. *Org Biomol Chem.* 2016;14: 9354–9361.
10. Moritzer AC, Minges H, Prior T, Frese M, Sewald N, Niemann HH. Structure-based switch of regioselectivity in the flavin-dependent tryptophan 6-halogenase Thal. *J Biol Chem.* 2019;294:2529–2542.
11. Luhavaya H, Sigrist R, Chekan JR, McKinnie SMK, Moore BS. Biosynthesis of l-4-chlorokynurenone, an antidepressant prodrug and a non-proteinogenic amino acid found in lipopeptide antibiotics. *Angew Chem Int Ed.* 2019;58:8394–8399.
12. Neubauer PRPR, Widmann C, Wibberg D, et al. A flavin-dependent halogenase from metagenomic analysis prefers bromination over chlorination. *PLoS One.* 2018;13:1–21.
13. Keller S, Wage T, Hohaus K, Hölzer M, Eichhorn E, van Pee KH. Purification and partial characterization of tryptophan 7-halogenase (PrnA) from *Pseudomonas fluorescens*. *Angew Chem Int Ed.* 2000; 39:2300–2302.
14. Yeh E, Garneau S, Walsh CT. Robust in vitro activity of RebF and RebH, a two-component reductase/halogenase, generating 7-chlorotryptophan during rebeccamycin biosynthesis. *Proc Natl Acad Sci U S A.* 2005;102:3960–3965.
15. Sánchez C, Butovich IA, Braña AF, Rohr J, Méndez C, Salas JA. The biosynthetic gene cluster for the antitumor rebeccamycin characterization and generation of indolocarbazole derivatives. *Chem Biol.* 2002;9:519–531.
16. Mori S, Pang AH, Thamban Chandrika N, Garneau-Tsodikova S, Tsodikov OV. Unusual substrate and halide versatility of phenolic halogenase PtM. *Nat Commun.* 2019;10:1–11.
17. Seibold C, Schnerr H, Rumpf J, et al. A flavin-dependent tryptophan 6-halogenase and its use in modification of pyrrolnitrin biosynthesis. *Biocatal Biotransform.* 2006;24:401–408.
18. Milbredt D, Patallo EP, van Pee KH. A tryptophan 6-halogenase and an amidotransferase are involved in thienodolin biosynthesis. *Chembiochem.* 2014;15:1011–1020.
19. Flecks S, Patallo EP, Zhu X, et al. New insights into the mechanism of enzymatic chlorination of tryptophan. *Angew Chem Int Ed.* 2008;47:9533–9536.
20. Shepherd SA, Karthikeyan C, Latham J, et al. Extending the biocatalytic scope of regiocomplementary flavin-dependent halogenase enzymes. *Chem Sci.* 2015;6:3454–3460.
21. Fraley AE, Garcia-Borràs M, Tripathi A, et al. Function and structure of MalA/MalA', iterative halogenases for late-stage C-H functionalization of indole alkaloids. *J Am Chem Soc.* 2017;139: 12060–12068.
22. Sucharitakul J, Tinikul R, Chaiyen P. Mechanisms of reduced flavin transfer in the two-component flavin-dependent monooxygenases. *Arch Biochem Biophys.* 2014;555–556:33–46.
23. Eichhorn E, Van Der Ploeg JR, Leisinger T. Characterization of a two-component alkanesulfonate monooxygenase from *Escherichia coli*. *J Biol Chem.* 1999;274:26639–26646.
24. Fontecave M, Eliasson R, Reichard P. NAD(P)H:Flavin oxidoreductase of *Escherichia coli*. A ferric iron reductase participating in the generation of the free radical of ribonucleotide reductase. *J Biol Chem.* 1987;262:12325–12331.
25. Driggers CM, Dayal PV, Ellis HR, Andrew Karplus PA. Crystal structure of *Escherichia coli* SsuE: Defining a general catalytic cycle for FMN reductases of the flavodoxin-like superfamily. *Biochemistry.* 2014;53:3509–3519.
26. Cianci M, Bourenkov G, Pompidor G, et al. P13, the EMBL macromolecular crystallography beamline at the low-emittance PETRA III ring for high- and low-energy phasing with variable beam focusing. *J Synchrotron Radiat.* 2017;24:323–332.
27. Kabsch W. XDS. *Acta Crystallogr.* 2010;D66:125–132.
28. McCoy AJ, Grosse-Kunstleve RW, Adams PD, Winn MD, Storoni LC, Read RJ. Phaser crystallographic software. *J Appl Cryst.* 2007;40:658–674.
29. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of coot. *Acta Crystallogr.* 2010;D66:486–501.
30. Kovalevskiy O, Nicholls RA, Long F, Carlon A, Murshudov GN. Overview of refinement procedures within REFMAC 5: Utilizing data from different sources. *Acta Crystallogr.* 2018;D74:215–227.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: Moritzer A-C, Niemann HH. Binding of FAD and tryptophan to the tryptophan 6-halogenase Thal is negatively coupled. *Protein Science.* 2019;28:2112–2118. <https://doi.org/10.1002/pro.3739>