

(A)
ITCZ
LLJ
SPA
ATTL
SWW
SACZ
ACC
1
2
3
4
-0.8
0
+0.8
<U500, P_{TT}>
(B)
Elevation (m asl)
200
400
600
800
1000
>1150
School
TT-1
TT-3
TT-2
ATTL
Heusser (1990)
and
Valero-Garcés et al., (2005)
(C)
TT3
TT19-3A
TT19-3A
Depth [cm]
Units Facies
Other Archaeological site
Varela 1976 (TT-1)
Heusser (1990)
Valero-Garcés (2005)
5345 +/- 28
5806 +/- 33
7259 +/- 32
6130 +/- 115
8657 +/- 45
9100 +/- 100
11380 +/- 320
9736 +/- 35
10211 +/- 63
10936 +/- 41
10747 +/- 49
10590 +/- 39
11117 +/- 51
9977 +/- 45
17353 +/- 84
20558 +/- 69
Diatomaceous carbonaceous clays
Grey silts
Diatomaceous clays
Paleosoil
Green clays
2830 +/- 120
6130 +/- 250
9100 +/- 360
9860 +/- 320
11170 +/- 320
12970 +/- 390
14500 +/- 350
21500 +/- 650
6590 +/- 280
9735 +/- 270
11710 +/- 430
19320 +/- 230
20430 +/- 1300
22495 +/- 830
L1: fine dark-gray silty sand with coals and fragments of gastropod shells
L2: coarse gray silt to fine sand,with shell fragments of gastropods and bivalve
L3: coarse grey silt to fine sand with high diatom and carbonate content
L4a: brownish to grayish clay silt (Paleosoil in TT-1)
L4b: coarse silt to fine sand, with shell fragments of gastropods and bivalve
L5: coarse dark grayish green sandy silt, with shell of gastropods and bivalve
L6: brown to greenish clayey fine sandy
L7: fine to medium brown silty -clay with sub-angular and sub-rounded volcanic clasts
L8: Massive silty clays in brown to greenish-brown with sedimentary cracks