Simple Linear Regression

Model conditions

Dr. Maria Tackett



Topics

Topics


List the conditions for simple linear regression

Topics

- List the conditions for simple linear regression
- Use plots of the residuals to check the conditions

Movie ratings data


The data set contains the "Tomatometer" score (critics) and audience score (audience) for 146 movies rated on rottentomatoes.com.



The model

audience =
$$32.316 + 0.519 \times \text{critics}$$

term	estimate	std.error	statistic	p.value
(Intercept)	32.316	2.343	13.795	0
critics	0.519	0.035	15.028	0



1. **Linearity:** There is a linear relationship between the response and predictor variable.


- 1. **Linearity:** There is a linear relationship between the response and predictor variable.
- 2. **Constant Variance:** The variability of the errors is equal for all values of the predictor variable.

- 1. **Linearity:** There is a linear relationship between the response and predictor variable.
- 2. **Constant Variance:** The variability of the errors is equal for all values of the predictor variable.
- 3. **Normality:** The errors follow a normal distribution.

- 1. **Linearity:** There is a linear relationship between the response and predictor variable.
- 2. **Constant Variance:** The variability of the errors is equal for all values of the predictor variable.
- 3. **Normality:** The errors follow a normal distribution.
- 4. **Independence:** The errors are independent from each other.


$$residual = y - \hat{y}$$

Residuals vs. fitted values




Checking linearity

Checking linearity




Checking linearity




There is no distinguishable pattern or structure. The residuals are randomly scattered.

Checking constant variance

Checking constant variance




Checking constant variance




The vertical spread of the residuals is relatively constant across the plot.


Violation: distinguishable pattern



Violation: non-constant variance




Normal quantile plot




Checking normality

Checking normality



Checking normality



Points follow a straight diagonal line on the normal quantile plot.

 We can often check the independence assumption based on the context of the data and how the observations were collected.

- We can often check the independence assumption based on the context of the data and how the observations were collected.
- If the data were collected in a particular order, examine a scatterplot of the residuals versus order in which the data were collected.

- We can often check the independence assumption based on the context of the data and how the observations were collected.
- If the data were collected in a particular order, examine a scatterplot of the residuals versus order in which the data were collected.
- The observations are independent or there is no pattern in the plot of residuals vs. order.

As you check the model conditions, ask if any observed deviation from the model conditions are so great that

As you check the model conditions, ask if any observed deviation from the model conditions are so great that

a different model should be proposed.

As you check the model conditions, ask if any observed deviation from the model conditions are so great that

- a different model should be proposed.
- 2 conclusions drawn from the model should be used with caution.

As you check the model conditions, ask if any observed deviation from the model conditions are so great that

- a different model should be proposed.
- 2 conclusions drawn from the model should be used with caution.
- ✓ If not, the conditions are sufficiently met and we can proceed with the current model.

Recap

Recap

- Used plots of the residuals to check conditions for simple linear regression:
 - Linearity
 - Constant Variance
 - Normality
 - Independence