EHC 26 9–12 December 2026 Berlin (DE) and online
20th European Headache Congress
P550

Scan me, for digital PDF version:

References

COMPARISON OF THE EFFICACY OF NARATRIPTAN AND SUMATRIPTAN FOR ACUTE MIGRAINE TREATMENT: A COMPARATIVE STUDY

Norma Espindola[1], Arthur Soares[1], Daniel Neto[1], Prof. Dr. Wagner Horta[1]
[1] *Universidade de Pernambuco, Faculdade de Ciências Médicas, Recife, Brazil*

Background and Aims

Migraine is a prevalent and disabling neurological disorder that negatively affects quality of life, productivity, and healthcare costs. Diagnosis follows the ICHD-3 criteria, and management includes acute and preventive treatments. Triptans are the drugs of choice for the acute treatment of migraine attacks due to their selective action on the trigeminovascular system. Among the available triptans, sumatriptan (SUM) is the most widely used worldwide, whereas naratriptan (NAR), with its longer half-life and slower onset of action, represents an alternative for specific patient profiles. This study compares the efficacy of SUM and NAR in the acute treatment of migraine.

Objective

To compare the efficacy of sumatriptan and naratriptan in the acute treatment of migraine based on medical record data.

Methods

This prospective study included seven patients diagnosed with migraine who were recruited after signing an informed consent form and followed up, on average, every two months, for a minimum of six months. Patients using SUM or NAR as definitive acute rescue medication were included. The variables analyzed included sex, age, medication used, and time to reduction in migraine attacks. Patient characteristics were described using percentage distributions and frequency distributions. Quantitative variables were summarized using minimum, maximum, mean, median, standard deviation, and 95% confidence intervals. The normality of continuous variables was assessed using the Shapiro–Wilk test when applicable ($n \geq 3$). Response time was compared between groups using the Mann–Whitney test because the SUM group did not follow a normal distribution and the NAR group had an insufficient sample size. Analyses were performed using Epi Info version 7.2.7, with a significance level of 5% and Bonferroni correction for multiple comparisons ($\alpha = 0.025$).

Conclusions

Both triptans demonstrated efficacy in controlling migraine attacks in the cases analyzed, with resolution achieved in 100% of cases in which they were maintained as definitive medication. NAR showed a shorter response time (1.0 ± 0.0 months) than SUM (2.4 ± 2.2 months); however, this difference was not statistically significant after Bonferroni correction ($p = 0.469$; corrected $\alpha = 0.025$). The small sample size was the main limitation of the analysis. The case of treatment failure with NAR followed by a response to SUM illustrates that a lack of response to one triptan does not predict treatment failure with other triptans, a finding consistent with the literature. The exclusively female sample is consistent with the higher prevalence of migraine among women. The choice between triptans should consider their individual pharmacological properties and each patient's clinical response.