

# PREDICTION MODELS BASED ON MAX-STEMS (or harnessing imbalanced data)

Episode Four: Advanced Examinations

Ahmet Furkan EMREHAN

(matahmet@gmail.com)

# PREDICTION MODELS BASED ON MAX-STEMS

- ▶ Episode One: One-Word Based
- ▶ Episode Two: A Combinatorial Approach
- ▶ Episode Three: Effect of Hyperparameters
- ▶ Episode Four: Advanced Examinations

# INTRODUCTION

- ▶ In this chapter, I examine some cases:
- ▶ A) How do Accuracy Rates of Categories change, if the sample is changed ?
- ▶ B) Let's compare all methods for each category
- ▶ C) Can a Trigonometric Approach contribute to hyperparametric model ?

## A) How do Accuracy Rates of Categories change, if the sample is changed ?


(Based on Predict Model 1 (On Word))

Accuracy Rates of Categories					
Sample	DÜNYA	SPOR	SANAT	Teknoloji	
1	0.791	0.167	0.029	0.013	
2	0.798	0.166	0.025	0.012	
3	0.783	0.181	0.023	0.013	
4	0.794	0.169	0.024	0.013	
5	0.792	0.172	0.024	0.012	
6	0.793	0.175	0.024	0.009	
7	0.797	0.172	0.022	0.010	
8	0.809	0.159	0.022	0.010	
9	0.793	0.175	0.024	0.009	
10	0.797	0.172	0.022	0.010	


Result: Stable against samples

Note: Naturally Samples intersect wit each other at %80 level.


## B) Let's compare all methods for each category




## B) Let's compare all methods for each category



## B) Let's compare all methods for each category




## B) Let's compare all methods for each category





## B) Let's compare all methods for each category




## B) Let's compare all methods for each category

- ▶ Conclusion:
- ▶ Whereas NAYN predictor is successful at predicting documents labelled with «DÜNYA», other prediction models (one stems and combinatorial approach) is more successful at predicting documents labelled with other categories.

## C) Can a Trigonometric Approach contribute to hyperparametric model ?


- In this section we use trigonometric approach to hyperparameters such that  $\alpha = \sin(\theta)$  and  $\beta = \cos(\theta)$  where  $\theta \in \left[0, \frac{\pi}{2}\right]$

# General Scheme for Application of Trigonometric Approach to Prediction Models with hyperparameter




## C) Can a Trigonometric Approach contribute to hyperparametric model ?


### ► Results




## C) Can a Trigonometric Approach contribute to hyperparametric model ?



## C) Can a Trigonometric Approach contribute to hyperparametric model ?




## C) Can a Trigonometric Approach contribute to hyperparametric model ?





## C) Can a Trigonometric Approach contribute to hyperparametric model ?



## C) Can a Trigonometric Approach contribute to hyperparametric model ?

- ▶ Conclusion:
- ▶ 1) When  $\theta = \frac{\pi}{2} \rightarrow \alpha = 0, \beta = 1$ , model produces only «DÜNYA» label, then specificity fails. In other saying, predictions of other categories in test set become «DÜNYA»
- ▶ 2) Generally, performance of model based trigonometric functions falls behind that of model based uniform distribution in almost all categories. Only at prediction of documents labelled «Teknoloji», performance of trigonometric approach is a notch better than uniform approach.

# THE END