

Problem statement 1 and 2

For x_1, x_2, \dots, x_n

- (i) Mean = $\frac{\sum_{i=1}^n x_i}{n}$
- (ii) Median if you arrange x_i 's in order for n odd number, median is the $x_{\frac{n}{2}}$ and for even n median is $(x_{\frac{n}{2}} + x_{\frac{n+1}{2}})/2$
- (iii) Mode is the distinct value of x_i with the highest frequency given that the frequency is greater than 1.
- (iv) Standard deviation is the square root of $(\frac{\sum_{i=1}^n x_i^2}{n} - mean^2)$

Using R to calculate the above

```
> #Assignment stats 1
> #
> #Problem Statement 1
> #
> # The marks of the 20 students are stored in a vector marks
> marks = c(6,7,5,7,7,8,7,6,9,7, 4,10,6,8,8,9,5,6,4,8)
>
> # Using the mean function, mean of marks is calculate and stored in the mean_marks
> mean_marks = mean(x = marks)
> mean_marks
[1] 6.85
>
> # Using the median function, median of marks is calculate and stored in the median_marks
> median_marks = median(x = marks)
> median_marks
[1] 7
>
> # creating a function that returns the mode of a vector v called rt_mode.
> mode_fun <- function(w){
+   table_w = table(w)
+   if(which.max(table_w) == 1){
+     return('No mode')
+   }else{
+     as.numeric(names(table_w[which.max(table_w)]))
+   }
+ }
>
> # Mode using the above function
> marks_mode = mode_fun(marks)
> marks_mode
[1] 7
>
> # standard deviation
> std_marks = sqrt(var(marks))
> std_marks
[1] 1.631112
>
>
>
```

```

> #Problem Statement 2
> #Repeating the same steps as on problem statement 1
>
> calls = c(28, 122, 217, 130, 120, 86, 80, 90, 140, 120, 70, 40, 145,
+          113, 90, 68, 174, 194, 170, 100, 75, 104, 97, 75, 123, 100,
+          75, 104, 97, 75, 123, 100, 89, 120, 109)
> #Mean
> mean_calls = mean(x = calls)
> mean_calls
[1] 107.5143
>
> #Median
> median_calls = median(x = calls)
> median_calls
[1] 100
>
> # Mode
> calls_mode = mode_fun(calls)
> calls_mode
[1] 75
>
> # Standard deviation
> std_calls = sqrt(var(calls))
> std_calls
[1] 39.33893

```

answer for Problem 1

- (i) Mean = 6.85
- (ii) Median = 7
- (iii) Mode = 7
- (iv) Standard deviation = 1.6331112

answer for Problem 2

- (i) Mean = 107.5143
- (ii) Median = 100
- (iii) Mode = 75
- (iv) Standard deviation = 39.33893

Problem statement 3

For x_1, x_2, \dots, x_n with probability distribution function $f(x_i)$

- (i) Mean = $\sum_{i=1}^n x_i * f(x_i)$
- (ii) Variance = $\sum_{i=1}^n x_i^2 * f(x_i) - mean^2$

Using R to calculate the above

```

#Problem statement 3
>
> x = c(0,1,2,3,4,5)
> f_x = c(0.09,0.15,0.4,0.25,0.1,0.01)
>
> #Mean
> mean_x = sum(x*f_x)
> mean_x

```

```
[1] 2.15
>
> #variance
> var_x = sum((x^2)*f_x)-mean_x^2
> var_x
[1] 1.2275

>
```

Answer Problem 3

- (i) Mean = 2.15
- (ii) Variance = 1.2275

Problem statement 4

The PDF does not contain the random variable