

Analytic geometry

1. Planes in space $\mathbb{R}^3$

Fix a cartesian frame in space with origin O and triad $\{\vec{i}, \vec{j}, \vec{k}\}$. Every point $M(x, y, z)$ has position vector $\vec{r} = x\vec{i} + y\vec{j} + z\vec{k}$.

Def. For $M_0(x_0, y_0, z_0)$ with $\vec{r}_0 = x_0\vec{i} + y_0\vec{j} + z_0\vec{k}$, the parallelogram rule gives us:

$$\vec{MM}_0 = \vec{r}_0 - \vec{r} = (x_0 - x)\vec{i} + (y_0 - y)\vec{j} + (z_0 - z)\vec{k}$$

1.1. Plane through a point with normal vector:


Def. Let $M_0(x_0, y_0, z_0)$ and $\vec{n} = a\vec{i} + b\vec{j} + c\vec{k} \neq \vec{0}$.

Let P be the plane through M_0 perpendicular to $\vec{n}$. For $M(x, y, z) \in P$:

$$\vec{r} - \vec{r}_0 \perp \vec{n} \Leftrightarrow \vec{n} \cdot (\vec{r} - \vec{r}_0) = 0 \Leftrightarrow a(x - x_0) + b(y - y_0) + c(z - z_0) = 0$$

Obs. Setting $d = -ax_0 - by_0 - cz_0$:

$$ax + by + cz + d = 0$$



This is the general equation of a plane; $\vec{n} = a\vec{i} + b\vec{j} + c\vec{k}$ is normal to it.

ex. The plane xOy passes through O with normal $\vec{k}$: take $a=b=0, c=1$. Its equation is $z=0$.

1.2 Plane through three points:

Def. Let $M_i(x_i, y_i, z_i)$, $i=1,2,3$, be non-collinear and P the plane through them. For $M(x, y, z) \in P$, the four points are coplanar:

$$\begin{vmatrix} 1 & x & y & z \\ 1 & x_1 & y_1 & z_1 \\ 1 & x_2 & y_2 & z_2 \\ 1 & x_3 & y_3 & z_3 \end{vmatrix} = 0$$

1.3. Plane through a point and two dir. vectors:

Def. Let P pass through $M_0(x_0, y_0, z_0)$, parallel to non-collinear $\vec{v}_i = x_i \vec{e}_1 + y_i \vec{e}_2 + z_i \vec{e}_3$, $i=1,2$. For $M \in P$, the vectors $\vec{r} - \vec{r}_0$, $\vec{v}_1$, $\vec{v}_2$ are coplanar $\Leftrightarrow$ their triple product vanishes (only area, no volume):

$$\begin{vmatrix} x-x_0 & y-y_0 & z-z_0 \\ x_1 & y_1 & z_1 \\ x_2 & y_2 & z_2 \end{vmatrix} = 0$$

1.4. The pencil of planes through a line:

Def. Every plane through the intersection line of:

$$(1) a_1x + b_1y + c_1z + d_1 = 0, (2) a_2x + b_2y + c_2z + d_2 = 0$$

has the form:

$$(3) a_1x + b_1y + c_1z + d_1 + \lambda(a_2x + b_2y + c_2z + d_2) = 0$$

2. Straight lines in space:

2.1. Direction ratios and direction cosines:

Def. A direction in space is fixed by $\vec{v} = l\vec{i} + m\vec{j} + n\vec{k} \neq 0$. (l, m, n) are its direction ratios (any proportional triple works as well).

Ob. If $\|\vec{v}\| = 1$, then $l^2 + m^2 + n^2 = 1$ and $l = \vec{v} \cdot \vec{i} = \cos \alpha$, $m = \cos \beta$, $n = \cos \gamma$, where α, β, γ are the angles to the axes - the direction cosines:

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

2.2. Line through a point with a direction vector:

Def. Let d pass through $M_0(x_0, y_0, z_0)$ with direction $\vec{v} = l\vec{i} + m\vec{j} + n\vec{k}$. For $M(x, y, z) \in d$: $\vec{r} - \vec{r}_0 = t\vec{v}$, $t \in \mathbb{R}$:

$$x = x_0 + lt, \quad y = y_0 + mt, \quad z = z_0 + nt$$

$$\Rightarrow \boxed{\frac{x - x_0}{l} = \frac{y - y_0}{m} = \frac{z - z_0}{n} = t}$$

ex. Take $M_0 = O$, $\vec{v} = \vec{i}$: $x_0 = y_0 = z_0 = 0$, $l = 1$, $m = n = 0$:

$$\frac{x - 0}{1} = \frac{y}{0} = \frac{z}{0} \Leftrightarrow \begin{cases} y = 0 \\ z = 0 \end{cases}$$

2.3. Line through two points:


Def. Let $M_0(x_0, y_0, z_0)$, $M_1(x_1, y_1, z_1)$. The line is determined by M_0 and direction $\vec{r}_1 - \vec{r}_0$, giving direction ratios: $(x_1 - x_0, y_1 - y_0, z_1 - z_0)$:

$$\frac{x-x_0}{x_1-x_0} = \frac{y-y_0}{y_1-y_0} = \frac{z-z_0}{z_1-z_0}$$

2.4. Line as the intersection of two planes:

Def. Let $d = P_1 \cap P_2$:

$$\begin{cases} a_1x + b_1y + c_1z + d_1 = 0 \\ a_2x + b_2y + c_2z + d_2 = 0 \end{cases}$$



with normals: $\vec{n}_1 = a_1\vec{i} + b_1\vec{j} + c_1\vec{k}$, $\vec{n}_2 = a_2\vec{i} + b_2\vec{j} + c_2\vec{k}$

Since $\vec{n}_1, \vec{n}_2 \perp d \Rightarrow d \parallel \vec{n}_1 \times \vec{n}_2 \Rightarrow$ the direction
 ration for line d is:


$$\left(e = \begin{vmatrix} b_1 & c_1 \\ b_2 & c_2 \end{vmatrix}, m = \begin{vmatrix} c_1 & a_1 \\ c_2 & a_2 \end{vmatrix}, n = \begin{vmatrix} a_1 & b_1 \\ a_2 & b_2 \end{vmatrix} \right)$$

3. Distance problems:

3.1. Distance from point to line:

Def. Let d pass through M_0 with dir $\vec{v}$ and let A
 be a point. $\text{dist}(A, d)$ is the height of the parallelo-
 gram spanned by $\vec{v}$ and $\vec{M_0A}$.

$$\text{dist}(A, d) = \frac{\|\vec{v} \times \vec{M_0A}\|}{\|\vec{v}\|}$$




3.2. Distance from point to plane:

Def. Let $P: ax + by + cz + d = 0$ and $M_0(x_0, y_0, z_0)$;

Let $M_1(x_1, y_1, z_1)$ be the orthogonal projection of M_0 onto P s.t. $ax_1 + by_1 + cz_1 + d_1 = 0$. The normal $(\vec{n}) = a\vec{i} + b\vec{j} + c\vec{k}$ and $\overrightarrow{M_1 M_0} = (x_0 - x_1)\vec{i} + (y_0 - y_1)\vec{j} + (z_0 - z_1)\vec{k}$.
 $\vec{n} \cdot \overrightarrow{M_1 M_0} = a(x_0 - x_1) + b(y_0 - y_1) + c(z_0 - z_1) = ax_0 + by_0 + cz_0 + d$

$$|\vec{n} \cdot \overrightarrow{M_1 M_0}| = |\vec{n}| \cdot |\overrightarrow{M_1 M_0}| \cdot \cos 0 = \sqrt{a^2 + b^2 + c^2} \cdot \text{dist}(M_0, P) \quad | \Rightarrow$$

$$\boxed{\text{dist}(M_0, P) = \frac{|ax_0 + by_0 + cz_0 + d|}{\sqrt{a^2 + b^2 + c^2}}}$$



Exercises:

$$P: 7(x-1) + 2(y-2) + z = 0$$

$$\vec{n} = 7\vec{i} + 2\vec{j} + \vec{k} \perp P$$

$$M_0(1, 2, 0) \in P$$

$$P_2: x - y + z - 5 = 0, \quad \vec{n} = \vec{i} - \vec{j} + \vec{k} \perp P_2$$

$$M(8, 4, 1) \in P_2$$

$$\text{ex. } P_1: x - y + z - 1 = 0$$

$$d = P_1 \cap P_2$$

$$P_2: x + y + z + 3 = 0$$

$$\begin{cases} x - y + z = 1 \\ x + y + z = -3 \end{cases} \quad \begin{aligned} \vec{n}_1 &= \vec{i} - \vec{j} + \vec{k} \perp P_1 \\ \vec{n}_2 &= \vec{i} + \vec{j} + \vec{k} \perp P_2 \end{aligned}$$

$$\bar{A} = \left[\begin{array}{ccc|c} 1 & -1 & 1 & 1 \\ 1 & 1 & 1 & -3 \end{array} \right] \Rightarrow \text{rang } A = \text{rang } \bar{A}$$

$$\begin{aligned} \left. \begin{array}{l} x, y - \text{f.v.} \\ z = \alpha \in \mathbb{R} \end{array} \right\} & \Rightarrow \begin{aligned} x - y &= 1 - \alpha \\ x + y &= -3 - \alpha \\ 2x &= -2 - 2\alpha \Rightarrow x = -1 - \alpha \\ y &= -2 = -2 + 0 \cdot \alpha \\ z &= \alpha = 0 + 1 \cdot \alpha \end{aligned} \end{aligned}$$

$$M_0(1, -2, 0) \in d$$

$$\vec{v} = -\vec{i} + 0 \cdot \vec{j} + \vec{k} \parallel d$$

$$\vec{n}_1 \times \vec{n}_2 = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -1 & 1 \\ 1 & 1 & 1 \end{vmatrix} = -2\vec{i} + 2\vec{k} = 2\vec{v}$$

$$\frac{x+1}{-1} = \frac{y+2}{0} = \frac{z}{-1}$$