

QFT Lecture 9

Matei Ionita

February 19, 2013

Perturbation theory

$$S = \int d^4x (\mathcal{L}_0 + \mathcal{L}_{int})$$

$$\mathcal{L}_{int} = -\frac{1}{2}(Z_\phi - 1)(\partial\phi)^2 - \frac{1}{2}(Z_m - 1)m^2\phi^2 + \frac{1}{3!}Z_g g\phi^3 + Y\phi + \Lambda$$

$$Z_1[J] = \int D\phi e^{i \int d^4x \mathcal{L}_0 + 1/3! Z_g g\phi^3 + J\phi}$$

$$Z_1[J] = \sum_{v=0}^{\infty} \frac{1}{v!} \left(\frac{i}{3!} Z_g g \int d^3x (\delta/i \delta J_x) \right)^v \sum_{P=0}^{\infty} \frac{1}{P!} \left(\frac{1}{2} \int d^4y d^4z i J_y i J_z \Delta_{yz}/i \right)^P$$

$$Z_1[J] = e^{iW[J]}$$

Where $W[J]$ is the sum over all connected diagrams. For example:

$$\Theta = \frac{1}{12} (iZ_g g)^2 \int d^4x d^4y \left(\frac{\Delta_{xy}}{i} \right)^3$$

$$dumbell = \frac{1}{8} (iZ_g g)^2 \int d^4x d^4y \frac{\Delta_{xy}}{i} \left(\frac{\Delta(0)}{i} \right)^2$$

$$O - * = \frac{1}{2} (iZ_g g) \int d^4x d^4y (iJ_y) \frac{\Delta_{xy}}{i} \frac{\Delta(0)}{i}$$

$$* - * = \frac{1}{2} \int d^4x d^4y (iJ_x) (iJ_y) \frac{\Delta_{xy}}{i}$$

$$* - O - * = \frac{1}{4} (iZ_g g)^2 \int d^4x d^4y d^4a d^4b (iJ_a) (iJ_b) \left(\frac{\Delta_{xy}}{i} \right)^2 \frac{\Delta_{ax}}{i} \frac{\Delta_{yb}}{i}$$

Let's focus for a bit on diagrams with no J. For the free generating function:

$$Z_0[J=0] = 1$$

With ϕ^3 interaction, this no longer holds:

$$Z_1[J=0] = e^{iW[J=0]}$$

i.e. the exponential of the sum of all bubble diagrams. This is not 1 in general. To correct this, we introduce the counterterm Λ . To get 1, we have to choose:

$$-i \int d^4x \Lambda = \Theta + \text{dumbbell} + \dots$$

$$\int d^4x \Lambda = -W(J=0)$$

It's not obvious why this works - why the sum over all bubbles is the volume integral of a constant. To see it, let's look at the Θ diagram:

$$\Theta = \frac{1}{12} (iZ^g g)^2 \int \frac{d^4x d^4y}{i^3} \left(\int \frac{d^4k_1}{2\pi^4} \frac{e^{ik_1(x-y)}}{k_1^2 + m^2 - i\epsilon} \right) (2)(3)$$

$$\Theta = \frac{1}{12} (iZ^g g)^2 \int \frac{d^4x}{i^3} \int \frac{d^4k_2 d^4k_3}{2\pi^8} \frac{1}{(-k_2 - k_3)^2 + m^2 - i\epsilon} \frac{1}{k_1^2 + m^2 - i\epsilon} \frac{1}{k_1^2 + m^2 - i\epsilon}$$

The k integral goes like k^2 - diverges. Thus Λ is infinite. To go around this issue, we introduce the ULTRAVIOLET CUTOFF: integrate only up to $k = k_{UV}$. The infinite Λ problem is very bad because it appears with a - sign in \mathcal{H} . We get a $-\lambda$ energy density for vacuum. Intuitively, the bubble diagrams represent vacuum energy because only J terms are interactions.

Now we'll make the argument that we also want to cancel out $1J$ diagrams. Consider:

$$\langle 0 | \hat{\phi}(x) | 0 \rangle = \left. \frac{\delta Z_1[J]}{i \delta J(x)} \right|_{J=0} = e^{iW[J]} \left. \frac{\delta iW(J)}{\delta iJ_x} \right|_{J=0}$$

The first factor is already 1 by the removal of bubbles. The second factor Gives the derivative of all diagrams with one J . Why do we want this to be 0? This is:

$$\langle 0 | \hat{\phi}(x) | 0 \rangle = \langle 0 | e^{iHt} \hat{\phi}(0, \mathbf{x}) e^{-iHt} | 0 \rangle$$

We can make the exponentials act on the vacuum states, which they leave unchanged. Therefore the 1-point function should be time independent. Then we can take the time to be in the far future or far past, where the field behaves like a free field. Using the expansion of the free field in terms of a and a^\dagger , a annihilates the ket and a^\dagger annihilates the bra. We get 0. An intuitive argument is that one J diagrams mean that a particle interacts with the vacuum; if we want a stable vacuum (one that doesn't randomly poop out particles), we make them 0. For this reason, we introduce the $Y\phi$ term. The vertices will have 1 leg sticking out, so there's only one possible diagram:

$$x - * = (iY) \int d^4x d^4y (iJ_y) \Delta_{xy} / i$$

We want this to cancel out the diagram:

$$O - * = \frac{1}{2}(iZ_g g) \int d^4x d^4y (iJ_y) \frac{\Delta_{xy}}{i} \frac{\Delta(0)}{i}$$

We make the choice:

$$Y = \frac{-Z_g g}{2} \frac{\Delta(0)}{i}$$

This is the $O(g)$ contribution to Y. Choose $O(g^3)$ contributions to Y to cancel out all the g^3 one J diagrams. Then we can do this order by order.