
A
1-year calibration - UK-Erlangen external
N=10 | Filled: overall KM vs mean predicted survival
Open: 3 quantile groups (exploratory_small_sample)
Observed survival probability (KM, 95% CI)
Predicted survival probability
0.00
0.25
0.50
0.75
1.00
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
Observed KM - mean predicted = 0.008 (95% CI: -0.219–0.209)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
Observed KM - mean predicted = -0.04 (95% CI: -0.324–0.223)
B
2-year calibration - UK-Erlangen external
N=10 | Filled: overall KM vs mean predicted survival
Open: 3 quantile groups (exploratory_small_sample)
Observed survival probability (KM, 95% CI)
Predicted survival probability
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
Observed KM - mean predicted = 0.136 (95% CI: -0.079–0.336)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
Observed KM - mean predicted = 0.067 (95% CI: -0.216–0.322)
C
3-year calibration - UK-Erlangen external
N=10 | Filled: overall KM vs mean predicted survival
Open: 3 quantile groups (exploratory_small_sample)
Observed survival probability (KM, 95% CI)
Predicted survival probability
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
Observed KM - mean predicted = -0.026 (95% CI: -0.34–0.286)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
Observed KM - mean predicted = -0.108 (95% CI: -0.493–0.228)