
A
1-year ROC - UK-Erlangen external
N=10 | Deaths by 12 mo=3 | Alive beyond=7 | Censored by 12 mo=0
Sensitivity (TPR)
1 - Specificity (FPR)
0.00
0.25
0.50
0.75
1.00
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
AUC = 0.952 (95% CI: 0.75–1)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
AUC = 0.667 (95% CI: 0.32–1)
B
2-year ROC - UK-Erlangen external
N=10 | Deaths by 24 mo=3 | Alive beyond=5 | Censored by 24 mo=2
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
AUC = 0.933 (95% CI: 0.625–1)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
AUC = 0.533 (95% CI: 0.111–1)
C
3-year ROC - UK-Erlangen external
N=10 | Deaths by 36 mo=5 | Alive beyond=3 | Censored by 36 mo=2
Nomogram (5 vars: Brain metastasis + Age + Grade + Histological subtype + Surgery)
AUC = 0.839 (95% CI: 0.424–1)
AJCC T/N baseline (2 vars: AJCC T stage + AJCC N stage)
AUC = 0.845 (95% CI: 0.446–1)