

Regra da Cadeia

Luis Alberto D'Afonseca

Cálculo de Funções de Várias Variáveis – I



<https://material-didatico.github.io/cfvv1>

Conteúdo

Regra da Cadeia

Caso 1

Caso 2

Exemplos

Lista Mínima

Regra da Cadeia em uma variável

$$h(x) = f(g(x))$$

$$\frac{dh}{dx}(x) = \frac{df}{dg}(g(x)) \frac{dg}{dx}(x)$$

$$h'(x) = f'(g(x))g'(x)$$

Conteúdo

Regra da Cadeia

Caso 1

Caso 2

Exemplos

Lista Mínima

Regra da Cadeia – Caso 1


Uma variável independente e duas variáveis intermediárias

Se $w = f(x, y)$ é diferenciável e se $x = x(t)$ e $y = y(t)$ são deriváveis

a composta $w = f(x(t), y(t))$ será uma função derivável de t e

$$\frac{dw}{dt} = \frac{\partial f}{\partial x}(x(t), y(t)) \frac{dx}{dt}(t) + \frac{\partial f}{\partial y}(x(t), y(t)) \frac{dy}{dt}(t)$$

Regra da Cadeia – Caso 1



Exemplo 1

Calcule a derivada de $w = xy$ com relação a t ao longo do caminho

$$x = \cos(t) \quad y = \sin(t)$$

Qual o valor da derivada em $t = \frac{\pi}{2}$

Exemplo 1 – Solução

Pela regra da cadeia

$$\frac{dw}{dt} = \frac{\partial w}{\partial x} \frac{dx}{dt} + \frac{\partial w}{\partial y} \frac{dy}{dt}$$

Exemplo 1 – Solução

Calculando as derivadas separadamente

$$\frac{\partial w}{\partial x} = \frac{\partial}{\partial x}(xy) = y \frac{\partial x}{\partial x} = y$$

$$\frac{\partial w}{\partial y} = \frac{\partial}{\partial y}(xy) = x \frac{\partial y}{\partial y} = x$$

$$\frac{dx}{dt} = \frac{d}{dt} \cos(t) = -\text{sen}(t)$$

$$\frac{dy}{dt} = \frac{d}{dt} \text{sen}(t) = \cos(t)$$

Exemplo 1 – Solução

$$\begin{aligned}\frac{dw}{dt} &= \frac{\partial w}{\partial x} \frac{dx}{dt} + \frac{\partial w}{\partial y} \frac{dy}{dt} \\&= \frac{\partial}{\partial x}(xy) \frac{d}{dt}(\cos(t)) + \frac{\partial}{\partial y}(xy) \frac{d}{dt}(\sin(t)) \\&= y(-\sin(t)) + x(\cos(t)) \\&= (\sin(t))(-\sin(t)) + (\cos(t))(\cos(t)) \\&= -\sin^2(t) + \cos^2(t) \\&= \cos(2t)\end{aligned}$$

Exemplo 1 – Solução

$$\begin{aligned}\frac{dw}{dt} \left(\frac{\pi}{2} \right) &= \cos(2t) \Big|_{t=\frac{\pi}{2}} \\ &= \cos \left(2 \frac{\pi}{2} \right) \\ &= \cos(\pi) \\ &= -1\end{aligned}$$

Com Três Variáveis


Se $w = f(x, y, z)$ é diferenciável

e se $x = x(t)$, $y = y(t)$ e $z = z(t)$ são deriváveis

a composta $w = f(x(t), y(t), z(t))$ será uma função derivável de t e

$$\begin{aligned}\frac{dw}{dt} &= \frac{\partial f}{\partial x}(x(t), y(t), z(t)) \frac{dx}{dt}(t) \\ &+ \frac{\partial f}{\partial y}(x(t), y(t), z(t)) \frac{dy}{dt}(t) \\ &+ \frac{\partial f}{\partial z}(x(t), y(t), z(t)) \frac{dz}{dt}(t)\end{aligned}$$

Regra da Cadeia – Caso 1



$$\frac{dw}{dt} = \frac{\partial w}{\partial x} \frac{dx}{dt} + \frac{\partial w}{\partial y} \frac{dy}{dt} + \frac{\partial w}{\partial z} \frac{dz}{dt}$$

Conteúdo

Regra da Cadeia

Caso 1

Caso 2

Exemplos

Lista Mínima

Regra da Cadeia – Caso 2

Funções definidas em superfícies

Se $w = f(x, y, z)$ é diferenciável

e se $x = x(r, s)$, $y = y(r, s)$ e $z = z(r, s)$ são deriváveis


a composta $w = f(x(r, s), y(r, s), z(r, s))$ será uma função das deriváveis r e s

e terá duas derivadas parciais

$$\frac{\partial w}{\partial r} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial r}$$

$$\frac{\partial w}{\partial s} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial s}$$

Regra da Cadeia – Caso 2



Exemplo 2

Expresse $\frac{\partial w}{\partial r}$ e $\frac{\partial w}{\partial s}$ em termos de r e s

$$w = x + 2y + z^2 \quad x = \frac{r}{s} \quad y = r^2 + \ln(s) \quad z = 2r$$

Exemplo 2 – Solução

Pela regra da cadeia

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial r}$$

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial s}$$

Exemplo 2 – Solução

Calculando as derivadas separadamente

$$\frac{\partial w}{\partial x} = \frac{\partial}{\partial x} (x + 2y + z^2) = 1$$

$$\frac{\partial w}{\partial y} = \frac{\partial}{\partial y} (x + 2y + z^2) = 2$$

$$\frac{\partial w}{\partial z} = \frac{\partial}{\partial z} (x + 2y + z^2) = 2z$$

Exemplo 2 – Solução

$$\frac{\partial x}{\partial r} = \frac{\partial}{\partial r} \left(\frac{r}{s} \right) = \frac{1}{s}$$

$$\frac{\partial x}{\partial s} = \frac{\partial}{\partial s} \left(\frac{r}{s} \right) = \frac{\partial}{\partial s} (rs^{-1}) = -\frac{r}{s^2}$$

$$\frac{\partial y}{\partial r} = \frac{\partial}{\partial r} (r^2 + \ln(s)) = 2r$$

$$\frac{\partial y}{\partial s} = \frac{\partial}{\partial s} (r^2 + \ln(s)) = \frac{1}{s}$$

$$\frac{\partial z}{\partial r} = \frac{\partial}{\partial r} (2r) = 2$$

$$\frac{\partial z}{\partial s} = \frac{\partial}{\partial s} (2r) = 0$$

Exemplo 2 – Solução

$$\begin{aligned}\frac{\partial w}{\partial r} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial r} \\ &= 1 \frac{\partial x}{\partial r} + 2 \frac{\partial y}{\partial r} + 2z \frac{\partial z}{\partial r} \\ &= 1 \frac{1}{s} + 2(2r) + 2z2 \\ &= \frac{1}{s} + 4r + 4z \\ &= \frac{1}{s} + 4r + 4(2r) \\ &= \frac{1}{s} + 12r\end{aligned}$$

Exemplo 2 – Solução

$$\begin{aligned}\frac{\partial w}{\partial s} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial s} \\&= 1 \frac{\partial x}{\partial s} + 2 \frac{\partial y}{\partial s} + 2z \frac{\partial z}{\partial s} \\&= 1 \left(\frac{-r}{s^2} \right) + 2 \frac{1}{s} + 2z \times 0 \\&= -\frac{r}{s^2} + \frac{2}{s}\end{aligned}$$

Conteúdo

Regra da Cadeia

Caso 1

Caso 2

Exemplos

Lista Mínima

Exemplo 3

Expresse $\frac{\partial w}{\partial r}$ e $\frac{\partial w}{\partial s}$ em termos de r e s

$$w = x^2 + y^2 \quad x = r^3 - s^2 \quad y = r^2 + s^3$$

Exemplo 3 – Solução

Pela regra da cadeia

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r}$$

$$\frac{\partial w}{\partial s} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s}$$

Exemplo 3 – Solução

Calculando as derivadas

$$\frac{\partial w}{\partial x} = \frac{\partial}{\partial x} (x^2 + y^2) = 2x$$

$$\frac{\partial w}{\partial y} = \frac{\partial}{\partial y} (x^2 + y^2) = 2y$$

$$\frac{\partial x}{\partial r} = \frac{\partial}{\partial r} (r^3 - s^2) = 3r^2$$

$$\frac{\partial y}{\partial r} = \frac{\partial}{\partial r} (r^2 + s^3) = 2r$$

$$\frac{\partial x}{\partial s} = \frac{\partial}{\partial s} (r^3 - s^2) = -2s$$

$$\frac{\partial y}{\partial s} = \frac{\partial}{\partial s} (r^2 + s^3) = 3s^2$$

Exemplo 3 – Solução

$$\begin{aligned}\frac{\partial w}{\partial r} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} \\ &= 2x \frac{\partial x}{\partial r} + 2y \frac{\partial y}{\partial r} \\ &= 2x3r^2 + 2y2r \\ &= 6(r^3 - s^2)r^2 + 4(r^2 + s^3)r \\ &= 6r^5 - 6s^2r^2 + 4r^3 + 4s^3r \\ &= 6r^5 + 4r^3 - 6r^2s^2 + 4rs^3\end{aligned}$$

Exemplo 3 – Solução

$$\begin{aligned}\frac{\partial w}{\partial s} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial s} \\ &= 2x \frac{\partial x}{\partial s} + 2y \frac{\partial y}{\partial s} \\ &= 2x(-2s) + 2y3s^2 \\ &= -4(r^3 - s^2)s + 6(r^2 + s^3)s^2 \\ &= -4r^3s + 4s^3 + 6r^2s^2 + 6s^5\end{aligned}$$

Exemplo 4

Considere $z = \sin(x^2 + y^2)$ onde $x = st$ e $y = 2s + t$

Encontre $\frac{\partial z}{\partial s}$

Exemplo 4 – Solução

Pela regra da cadeia

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s}$$

Exemplo 4 – Solução

$$\frac{\partial z}{\partial x} = \frac{\partial}{\partial x} \sin(x^2 + y^2) = \cos(x^2 + y^2) \frac{\partial}{\partial x} (x^2 + y^2) = \cos(x^2 + y^2) 2x$$

$$\frac{\partial z}{\partial y} = \frac{\partial}{\partial y} \sin(x^2 + y^2) = \cos(x^2 + y^2) \frac{\partial}{\partial y} (x^2 + y^2) = \cos(x^2 + y^2) 2y$$

$$\frac{\partial x}{\partial s} = \frac{\partial}{\partial s} (st) = t$$

$$\frac{\partial y}{\partial s} = \frac{\partial}{\partial s} (2s + t) = 2$$

Exemplo 4 – Solução

$$\begin{aligned}\frac{\partial z}{\partial s} &= \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s} \\&= 2x \cos(x^2 + y^2) \frac{\partial x}{\partial s} + 2y \cos(x^2 + y^2) \frac{\partial y}{\partial s} \\&= 2x \cos(x^2 + y^2) t + 2y \cos(x^2 + y^2) 2 \\&= (2xt + 4y) \cos(x^2 + y^2) \\&= [2(st)t + 4(s + t)] \cos((st)^2 + (s + t)^2) \\&= (2st^2 + 4s + 4t) \cos(s^2t^2 + (s + t)^2)\end{aligned}$$

Exemplo 5

Seja $w = e^{xyz}$ onde $x = r^2 + 4s$, $y = 2r - 3s$, $z = rs$

Encontre $\frac{\partial w}{\partial r}$

Exemplo 5 – Solução

Pela regra da cadeia

$$\frac{\partial w}{\partial r} = \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial r}$$

Exemplo 5 – Solução

$$\frac{\partial w}{\partial x} = \frac{\partial}{\partial x} e^{xyz} = yze^{xyz}$$

$$\frac{\partial w}{\partial y} = \frac{\partial}{\partial y} e^{xyz} = xze^{xyz}$$

$$\frac{\partial w}{\partial z} = \frac{\partial}{\partial z} e^{xyz} = xye^{xyz}$$

$$\frac{\partial x}{\partial r} = \frac{\partial}{\partial r} (r^2 + 4s) = 2r$$

$$\frac{\partial y}{\partial r} = \frac{\partial}{\partial r} (2r - 3s) = 2$$

$$\frac{\partial z}{\partial r} = \frac{\partial}{\partial r} (rs) = s$$

Exemplo 5 – Solução

$$\begin{aligned}\frac{\partial w}{\partial r} &= \frac{\partial w}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial w}{\partial y} \frac{\partial y}{\partial r} + \frac{\partial w}{\partial z} \frac{\partial z}{\partial r} \\&= yze^{xyz} \frac{\partial x}{\partial r} + xze^{xyz} \frac{\partial y}{\partial r} + xye^{xyz} \frac{\partial z}{\partial r} \\&= e^{xyz} \left(yz \frac{\partial x}{\partial r} + xz \frac{\partial y}{\partial r} + xy \frac{\partial z}{\partial r} \right) \\&= e^{xyz} (yz2r + xz2 + xys) \\&= e^{xyz} [2(2r - 3s)(rs)r + 2(r^2 + 4s)(rs) + (r^2 + 4s)(2r - 3s)s] \\&= e^{xyz} [2r^2s(2r - 3s) + 2rs(r^2 + 4s) + s(r^2 + 4s)(2r - 3s)]\end{aligned}$$

Exemplo 6

Dada $u = \ln(x^2 + y^2 + z^2)$ com $x = e^t$, $y = e^{-t}$, $z = t^2$

Encontre $\frac{du}{dt}$

Exemplo 6 – Solução

Pela regra da cadeia

$$\frac{du}{dt} = \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial z} \frac{dz}{dt}$$

Exemplo 6 – Solução

$$\frac{\partial u}{\partial x} = \ln' (x^2 + y^2 + z^2) \frac{\partial}{\partial x} (x^2 + y^2 + z^2) = \frac{2x}{x^2 + y^2 + z^2}$$

$$\frac{\partial u}{\partial y} = \ln' (x^2 + y^2 + z^2) \frac{\partial}{\partial y} (x^2 + y^2 + z^2) = \frac{2y}{x^2 + y^2 + z^2}$$

$$\frac{\partial u}{\partial z} = \ln' (x^2 + y^2 + z^2) \frac{\partial}{\partial z} (x^2 + y^2 + z^2) = \frac{2z}{x^2 + y^2 + z^2}$$

Exemplo 6 – Solução

$$\frac{dx}{dt} = \frac{de^t}{dt} = e^t$$

$$\frac{dy}{dt} = \frac{de^{-t}}{dt} = -e^{-t}$$

$$\frac{dz}{dt} = \frac{dt^2}{dt} = 2t$$

Exemplo 6 – Solução

$$\begin{aligned}\frac{du}{dt} &= \frac{\partial u}{\partial x} \frac{dx}{dt} + \frac{\partial u}{\partial y} \frac{dy}{dt} + \frac{\partial u}{\partial z} \frac{dz}{dt} \\&= \frac{2x}{x^2 + y^2 + z^2} \frac{dx}{dt} + \frac{2y}{x^2 + y^2 + z^2} \frac{dy}{dt} + \frac{2z}{x^2 + y^2 + z^2} \frac{dz}{dt} \\&= \frac{2}{x^2 + y^2 + z^2} \left(x \frac{dx}{dt} + y \frac{dy}{dt} + z \frac{dz}{dt} \right) \\&= \frac{2}{x^2 + y^2 + z^2} (xe^t - ye^{-t} + 2zt) \\&= \frac{2e^{2t} - 2e^{-2t} + 4t^3}{x^2 + y^2 + z^2}\end{aligned}$$

Conteúdo

Regra da Cadeia

Caso 1

Caso 2

Exemplos

Lista Mínima

Lista Mínima

Cálculo Vol. 2 do Thomas 12^a ed. – Seção 14.4

1. Estudar o texto da seção
2. Resolver os exercícios: 3, 5, 6, 10, 12, 14, 17

Atenção: A prova é baseada no livro, não nas apresentações