



A High Order Semi-Lagrangian Discontinuous Galerkin Method for the Two-Dimensional Incompressible Euler Equations and the Guiding Center Vlasov Model Without Operator Splitting

Xiaofeng Cai¹ · Wei Guo² · Jing-Mei Qiu¹ 

Received: 23 August 2018 / Revised: 6 November 2018 / Accepted: 27 November 2018 /
Published online: 18 December 2018
© Springer Science+Business Media, LLC, part of Springer Nature 2018

Abstract

In this paper, we generalize a high order semi-Lagrangian (SL) discontinuous Galerkin (DG) method for multi-dimensional linear transport equations without operator splitting developed in Cai et al. (J Sci Comput 73(2–3):514–542, 2017) to the 2D time dependent incompressible Euler equations in the vorticity-stream function formulation and the guiding center Vlasov model. We adopt a local DG method for Poisson’s equation of these models. For tracing the characteristics, we adopt a high order characteristics tracing mechanism based on a prediction-correction technique. The SLDG with large time-stepping size might be subject to extreme distortion of upstream cells. To avoid this problem, we propose a novel adaptive time-stepping strategy by controlling the relative deviation of areas of upstream cells.

Keywords Semi-Lagrangian · Discontinuous Galerkin · Guiding center Vlasov model · Incompressible Euler equations · Non-splitting · Mass conservative · Adaptive time-stepping method

W. Guo: Research is supported by NSF Grant NSF-DMS-1620047. J.-M. Qiu: Research of first and last author is supported by NSF Grant NSF-DMS-1522777 and 1818924, Air Force Office of Scientific Computing FA9550-18-1-0257 and University of Delaware.

✉ Jing-Mei Qiu
jingqiu@udel.edu

Xiaofeng Cai
xfcai@udel.edu

Wei Guo
weimath.guo@ttu.edu

¹ Department of Mathematical Sciences, University of Delaware, Newark, DE 19716, USA

² Department of Mathematics and Statistics, Texas Tech University, Lubbock, TX 79409, USA