

Masum Billal

ADVANCES IN OLYMPIAD INEQUALITIES

Principles and Techniques for Old and New Problems

PREFACE

Inequalities have been extensively studied for at least a couple of centuries. Cauchy was among the first of the mathematicians who had major contributions in this literature. But it was probably not until HARDY et al.¹ we understood that it could be possible to study inequalities in a more systematic way. Since then a good number of books have discussed different aspects of inequalities for example, BECKENBACH and BELLMAN.²

OBJECTIVES

- Dis

¹G. H. HARDY et al., *Inequalities*, Cambridge University Press, 1934, ISBN: 0-521-35880-9.

²EDWIN F. BECKENBACH and RICHARD BELLMAN, *Inequalities*, Springer, 1983.

Contents

1 CLASSICAL INEQUALITIES	1
1.1 INTRODUCTION	1
1.2 WARM UP PROBLEMS	12
1.3 CAUCHY-SCHWARZ	13
1.4 COMPLEX NUMBERS	19
1.5 BERNOULLI AND POWER MEAN INEQUALITY	21
1.6 HOLDER AND MINKOWSKI	23
1.7 REARRANGEMENT AND CHEBYSHEV	25
1.8 JENSEN AND POPOVICIU	26
2 TRADITIONAL PRINCIPLES	29
2.1 ENGEL FORM OF CAUCHY-SCHWARZ	29
2.2 THE BUFFALO WAY	36
2.3 MAJORIZATION AND SYMMETRIC SUMS	40
2.4 KARAMATA	43
2.5 BUNCHING: MUIRHEAD	44
2.6 HOMOGENIZATION AND NORMALIZATION	45
2.7 SUBSTITUTIONS	45
3 ADVANCED TECHNIQUES & INEQUALITIES	47
3.1 TANGENT LINE TRICK	47
3.2 SCHUR-VORNICU-MILDORF INEQUALITY	51
3.3 RADON'S INEQUALITY	54
3.4 SMOOTHING AND ISOLATED FUDGING	56
3.5 THE UVW/PQR/ABC METHOD	56
3.6 MIXING VARIABLES TECHNIQUE	60
3.7 SEPARATION	60
3.8 FLIPPING	62
3.9 NEUTRALIZING DENOMINATORS	66
3.10 DUMBASSING	66
3.11 THE EQUAL VARIABLE METHOD	66

3.12 PROBABILITY IN INEQUALITY	66
4 PRACTICE PROBLEMS	67
5 CONTEST PROBLEMS	69
5.1 ABMO	70
5.2 ALMO	71
5.3 AMC	71
5.4 APC	74
5.5 APMO	76
5.6 AzNO	78
5.7 BkMO	79
5.8 BMO	81
5.9 BNO	81
5.10 BrNO	82
5.11 BuNO	85
5.12 ChMO	87
5.13 IMO	102
5.14 IrMO	105
5.15 MMO	109
5.16 PMO	110
5.17 RNO	113
5.18 SgMO	118
5.19 TNO	119
5.20 USAMO	120
BIBLIOGRAPHY	125
GLOSSARY	129

§1 CLASSICAL INEQUALITIES

§§1.1 INTRODUCTION

Let us start with the most fundamental inequality.

$$x^2 \geq 0 \quad (\ddagger 1.1)$$

The first author calls it the mother of all inequality. Equality occurs if and only if $x = 0$. We can extend it for n variables.

$$x_1^2 + \dots + x_n^2 \geq 0 \quad (\ddagger 1.2)$$

Equality occurs if $x_i = 0$ for all $1 \leq i \leq n$. We immediately get some useful results substituting x with appropriate expressions. Substituting x with $a-b$, we get

$$\begin{aligned} (a-b)^2 &\geq 0 \\ a^2 + b^2 &\geq 2ab \end{aligned}$$

This is true for any real numbers a, b and equality occurs if $a = b$. If a, b are positive, then replacing a and b by \sqrt{a} and \sqrt{b} respectively, we get

$$\begin{aligned} a + b &\geq 2\sqrt{ab} \\ \Leftrightarrow \sqrt{\frac{a}{b}} + \sqrt{\frac{b}{a}} &\geq 2 \\ \Leftrightarrow x + \frac{1}{x} &\geq 2 \end{aligned}$$

where $x = \frac{a}{b}$. This can be generalized to the following result.

THEOREM 1.1 (Arithmetic-Geometric Inequality). *Let a_1, \dots, a_n be positive real numbers. Then*

$$\begin{aligned} \frac{a_1 + \dots + a_n}{n} &\geq \sqrt[n]{a_1 \cdots a_n} \\ \Leftrightarrow a_1 + \dots + a_n &\geq n \sqrt[n]{a_1 \cdots a_n} \\ \Leftrightarrow \left(\frac{a_1 + \dots + a_n}{n} \right)^n &\geq a_1 \cdots a_n \end{aligned}$$

We will show a classical proof of this result here. The proof is due to CAUCHY.¹ Later in § 1.5, § 2.3 we will show more proofs.

Proof.

□

Note the following.

$$\frac{a+b}{2} \geq \sqrt{ab}$$

Now, $\frac{a+b}{2}$ is the *arithmetic mean* of a and b . On the right side, \sqrt{ab} is the *geometric mean* of a and b . So the inequality states that the arithmetic mean of two positive real numbers is greater than or equal to their geometric mean. We can also rewrite it as the following.

$$\begin{aligned} \sqrt{ab} \left(\frac{a+b}{2} \right) &\geq ab \\ \sqrt{ab} &\geq \frac{2ab}{a+b} \\ \sqrt{ab} &\geq \frac{2}{\frac{1}{a} + \frac{1}{b}} \end{aligned}$$

$\frac{2ab}{a+b}$ is the *harmonic mean* of a and b . So, this form of the inequality states that the geometric mean is larger than the harmonic mean. This can be extended for three variables.

$$\frac{a+b+c}{3} \geq \frac{3}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}$$

In fact, this result can be extensively generalized. First, we can consider n variables a_1, \dots, a_n instead of just a and b . Second, we can generalize the fact that $AM \geq GM \geq HM$. We will talk about this generalization in § 1.5.

¹AUG.-LOUIS CAUCHY, *Cours d'Analyse de l'école Royale Polytechnique*, L'Imprimerie Royale, Debure frères, Libraires du Roi et de la Bibliothèque du Roi, 1821.

PROBLEM 1.1. If a is a real number greater than one, prove that $\log a + \log_a e \geq 2$.

Throughout the book, if the base of logarithm is unspecified, then $\log a$ shall mean $\log_e a$. Also, try the next problem in a similar manner.

PROBLEM 1.2. Prove the inequality

$$\frac{x^2}{1+x^4} \leq \frac{1}{2}$$

Recall from the definition of e that

$$\begin{aligned} e^x &= 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots \\ &= \sum_{i \geq 0} \frac{x^i}{i!} \end{aligned}$$

You may also know from elementary differentiation that

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e$$

In other words, if $x_n = \left(1 + \frac{1}{n}\right)^n$, then x_n converges to e . However, we also have the relation that $x_n \leq x_{n+1}$. While we can prove this using induction, we will show a better proof with arithmetic-geometric mean inequality here.

PROBLEM 1.3. Let $x_n = \left(1 + \frac{1}{n}\right)^n$. Prove that $x_n \leq x_{n+1}$.

Solution. Apply the arithmetic-geometric mean inequality for $a_1 = 1, a_2 = 1 + \frac{1}{n}, \dots, a_{n+1} = 1 + \frac{1}{n}$,

$$\begin{aligned} 1 + \left(1 + \frac{1}{n}\right) + \dots + \left(1 + \frac{1}{n}\right) &\geq (n+1) \sqrt[n+1]{1 \cdot \left(1 + \frac{1}{n}\right) \cdots \left(1 + \frac{1}{n}\right)} \\ n+2 &\geq (n+1) \sqrt[n+1]{\left(1 + \frac{1}{n}\right)^n} \\ 1 + \frac{1}{n+1} &\geq \sqrt[n+1]{\left(1 + \frac{1}{n}\right)^n} \\ \left(1 + \frac{1}{n+1}\right)^{n+1} &\geq \left(1 + \frac{1}{n}\right)^n \end{aligned}$$

Note that we can generalize the idea used in this problem.

PROBLEM 1.4. For positive real numbers x, y show that

$$\left(\frac{x+ny}{n+1} \right)^{n+1} \geq xy^n$$

Equality occurs only for $x = y$.

Similarly, we can show the following.

PROBLEM 1.5. Let $y_n = \left(1 + \frac{1}{n}\right)^{n+1}$. Show that $y_n \geq y_{n+1}$.

So y_n is decreasing. Using the two problems above, show that $2 < e < 4$. In fact, we can prove the following using elementary means

$$\lim_{n \rightarrow \infty} y_n = \lim_{n \rightarrow \infty} x_n = e$$

PROBLEM 1.6. For a positive integer n , show that

$$\left(\frac{n+1}{2} \right)^n \geq n!$$

Equality occurs only for $n = 1$.

Solution. We use arithmetic-geometric inequality for $1, 2, \dots, n$.

$$\begin{aligned} 1 + \dots + n &\geq \sqrt[n]{1 \cdots n} \\ \frac{n(n+1)}{2} &\geq \sqrt[n]{n!} \\ \frac{n+1}{2} &\geq \sqrt[n]{n!} \end{aligned}$$

Equality occurs if and only if $1 = \dots = n$ which is possible only when $n = 1$.

Let us again go back to the mother of inequality. Setting $x_1 = a - b, x_2 = b - c, x_3 = c - a$ in § 1.2,

$$\begin{aligned} (a-b)^2 + (b-c)^2 + (c-a)^2 &\geq 0 \\ \Leftrightarrow a^2 + b^2 + c^2 - ab - bc - ca &\geq 0 \\ \Leftrightarrow a^2 + b^2 + c^2 &\geq ab + bc + ca \end{aligned}$$

We could prove this using $a^2 + b^2 \geq 2ab$ repeatedly.

$$\begin{aligned} a^2 + b^2 &\geq 2ab \\ b^2 + c^2 &\geq 2bc \\ c^2 + a^2 &\geq 2ca \\ 2(a^2 + b^2 + c^2) &\geq 2(ab + bc + ca) \\ a^2 + b^2 + c^2 - ab - bc - ca &\geq 0 \end{aligned}$$

Multiplying both sides by $(a + b + c)$ and using the fact $a^3 + b^3 + c^3 - 3abc = (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ca)$,

$$\begin{aligned} a^3 + b^3 + c^3 - 3abc &\geq 0 \\ a^3 + b^3 + c^3 &\geq 3abc \end{aligned}$$

Replacing a^3, b^3, c^3 by u, v, w ,

$$u + v + w \geq 3\sqrt[3]{uvw} \quad (\ddagger 1.3)$$

Before we go into any theory or method of solving problems, here are some basic tactics which are used very often for solving problems. Let $x, y, z > 0$ be real numbers.

(i) If $\frac{1}{x} \geq \frac{1}{y}$, then $\frac{1}{x-a} \geq \frac{1}{y}$ for $a \geq 0$. Similarly, $\frac{1}{x} \geq \frac{1}{y-b}$ for $b \geq 0$. They are true because $x-a \leq x$ implies $\frac{1}{x-a} \geq \frac{1}{x}$ and $y+b \geq y$ implies $\frac{1}{y+b} \leq \frac{1}{y}$.

- (ii) Check if you can assume an ordering on the variables. For example, if the inequality is symmetric or cyclic on x, y, z , you may possibly assume without loss of generality that $x \geq y \geq z$ or $x \leq y \leq z$. We will discuss more on this in § 2.2.
- (iii) If you cannot assume an ordering on the variables e.g. $a \geq b \geq c$, can you assume that it has a *maximal element* e.g. $a = \max(a, b, c)$? Sometimes this helps in unexpected ways. See the example below.
- (iv) See if you can get an *if and only if* way of proving an inequality. We will do this very often. If we can use if and only if (or *iff*), then we are free to prove either the if part or the only if part.
- (v) Check if you can get some familiar expressions with some basic manipulations such as making the numerator or denominator equal. For example, see the transformation in § 1.4. This often helps us get a clue on what to do with the inequality.
- (vi) Does some substitutions such as $x = a + b, y = b + c, z = c + a$ or $x = a - b, y = b - c, z = c - a$ help? One could say we used substitution in our first proof of Nesbitt's inequality (see § 1.5). Also, see the transformation $a - c = a - b + b - c$ used in the example below. We will check more on substitutions in § 2.7.

- (vii) Check if the inequality holds even if you put some restrictions on it. For example, you may be allowed to assume that one of the variables is 1 due to some scaling. We will talk about this in § 2.6.
- (viii) Can you reduce the number of variables without any assumption? See that we have already used it to prove $a^2 + b^2 + c^2 \geq ab + bc + ca$ in a proof above. Also, see a proof of Nesbitt's inequality below where it is enough to prove the inequality for reduced number of variables.
- (ix) Induction works great in many cases. We will use induction on many occasions.

For demonstration purposes, let us prove $a^2 + b^2 + c^2 \geq ab + bc + ca$ again exploiting symmetry. Note that if we let

$$f(a, b, c) = a^2 + b^2 + c^2 - ab - bc - ca$$

then $f(a, b, c)$ is symmetric on a, b, c . We can verify this by the fact that $f(a, b, c) = f(b, a, c) = f(c, a, b)$ and so on. So, without loss of generality, we assume that $a \geq b \geq c$. Then see the following.

$$\begin{aligned} a^2 + b^2 + c^2 &\geq ab + bc + ca \\ a(a-b) + b(b-c) - c(a-c) &\geq 0 \\ a(a-b) + b(b-c) - c(a-b+b-c) &\geq 0 \\ a(a-b) + b(b-c) - c(a-b) - c(b-c) &\geq 0 \\ (a-c)(a-b) + (b-c)^2 &\geq 0 \end{aligned}$$

The last inequality immediately follows from the assumption that $a - c \geq 0, a - b \geq 0, (b - c)^2 \geq 0$. Also, note that we did not actually require the condition $a \geq b \geq c$. Just assuming $a = \max(a, b, c)$ was enough in this case to claim that the inequality holds.

THEOREM 1.2 (*Nesbitt's inequality*). *Let a, b, c be real positive numbers. Then*

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \geq \frac{3}{2}$$

and equality occurs if and only $a = b = c$.

The thing about inequalities is that they can be solved in more than one ways most of the times. We will prove this inequality along with some other classical results such as *Cauchy-Schwarz inequality* in more than one ways. Some proofs will be discussed later when we develop some certain techniques.

For now, we present some proofs using what we have already developed. First, we will try to *familiarize* the expression on the left side.

$$\begin{aligned}
 S &= \frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \\
 &= \frac{a+b+c}{b+c} - 1 + \frac{a+b+c}{c+a} - 1 + \frac{a+b+c}{a+b} - 1 \\
 &= (a+b+c) \left(\frac{1}{b+c} + \frac{1}{c+a} + \frac{1}{a+b} \right) - 3 \\
 &= \frac{1}{2}(a+b+c+a) \left(\frac{1}{b+c} + \frac{1}{c+a} + \frac{1}{a+b} \right) - 3 \quad (\ddagger 1.4)
 \end{aligned}$$

Classical proof. Setting

$$x = a+b, y = b+c, z = c+a \quad (\ddagger 1.5)$$

$$\begin{aligned}
 S &= \frac{1}{2}(x+y+z) \left(\frac{1}{x} + \frac{1}{y} + \frac{1}{z} \right) - 3 \\
 &= \frac{1}{2} \left(1 + \frac{x}{y} + \frac{x}{z} + \frac{y}{x} + 1 + \frac{y}{z} + \frac{z}{x} + \frac{z}{y} + 1 \right) - 3 \\
 &= \frac{3}{2} + \frac{1}{2} \left(\frac{x}{y} + \frac{y}{x} + \frac{y}{z} + \frac{z}{y} + \frac{z}{x} + \frac{x}{z} \right) - 3 \\
 &= \frac{3}{2} + \frac{1}{2} \left(u + \frac{1}{u} + v + \frac{1}{v} + w + \frac{1}{w} \right) - 3
 \end{aligned}$$

where $u = \frac{x}{y}, v = \frac{y}{z}, w = \frac{z}{x}$. Evidently, $u + \frac{1}{u} \geq 2, v + \frac{1}{v} \geq 2, w + \frac{1}{w} \geq 2$ and we have

$$S \geq \frac{3}{2} + \frac{1}{2}(2+2+2) - 3$$

Thus, $S \geq \frac{3}{2}$. Equality occurs if $u = 1, v = 1, z = 1$ or $x = y = z$ or $a = b = c$. \square

Proof by arithmetic-harmonic mean inequality. We write the arithmetic-harmonic mean inequality for u, v, w as below.

$$\begin{aligned}
 \frac{u+v+w}{3} &\geq \frac{3}{\frac{1}{u} + \frac{1}{v} + \frac{1}{w}} \\
 \Leftrightarrow (u+v+w) \left(\frac{1}{u} + \frac{1}{v} + \frac{1}{w} \right) &\geq 9
 \end{aligned}$$

Using this on § 1.4,

$$S \geq \frac{1}{2} \cdot 9 - 2$$

□

Variable reduction proof. Let us clear the denominators in the original inequality.

$$\begin{aligned} \frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} &\geq \frac{3}{2} \\ \Leftrightarrow 2(a^3 + b^3 + c^3) &\geq a^2b + ab^2 + b^2c + bc^2 + c^2a + ca^2 \end{aligned}$$

Notice the cyclic nature in the expression on the right side. There are 6 terms on the right side and if we count each of a^3, b^3, c^3 twice, there are 6 terms on the left side as well. So rearranging the inequality above as below

$$(a^3 + b^3) + (b^3 + c^3) + (c^3 + a^3) \geq (a^2b + ab^2) + (b^2c + bc^2) + (c^2a + ca^2)$$

tells us that if we can prove $x^3 + y^3 \geq x^2y + xy^2$, we will be done if we simply sum them up for $(x, y) = (a, b), (b, c), (c, a)$. Here, we reduced the inequality from 3 variables to 2. And fortunately, this inequality is a lot easier to prove.

$$\begin{aligned} x^3 + y^3 &\geq x^2y + xy^2 \\ \Leftrightarrow x(x^2 - y^2) - y(x^2 - y^2) &\geq 0 \\ \Leftrightarrow (x - y)(x^2 - y^2) &\geq 0 \\ \Leftrightarrow (x - y)^2(x + y) &\geq 0 \end{aligned}$$

The last inequality is evidently true. □

Proof using arithmetic-geometric mean inequality. Using § 1.3,

$$\begin{aligned} (a+b) + (b+c) + (c+a) &\geq 3\sqrt[3]{(a+b)(b+c)(c+a)} \\ \frac{1}{a+b} + \frac{1}{b+c} + \frac{1}{c+a} &\geq 3\sqrt[3]{\frac{1}{(a+b)(b+c)(c+a)}} \end{aligned}$$

We can use this observation on § 1.4 and get the following.

$$\begin{aligned} S &\geq \frac{1}{2} \cdot 3\sqrt[3]{(a+b)(b+c)(c+a)} \cdot 3\sqrt[3]{\frac{1}{(a+b)(b+c)(c+a)}} - 2 \\ &\geq \frac{9}{2} - 2 = \frac{3}{2} \end{aligned}$$

This again proves the inequality. □

ANOVIĆ and PEĆARIĆ² proves the following generalization of Nesbitt's inequality.

THEOREM 1.3 (Generalization of Nesbitt's inequality). *Let x, y, z be positive real numbers and*

$$M = \max \left\{ \frac{x}{y+z} + \frac{2(y+z)}{2x+y+z}, \frac{y}{z+x} + \frac{2(z+x)}{2y+z+x}, \frac{z}{x+y} + \frac{2(x+y)}{2z+x+y} \right\}$$

$$m = \min \left\{ \frac{x}{y+z} + \frac{2(y+z)}{2x+y+z}, \frac{y}{z+x} + \frac{2(z+x)}{2y+z+x}, \frac{z}{x+y} + \frac{2(x+y)}{2z+x+y} \right\}$$

Then

$$\frac{x}{y+z} + \frac{y}{z+x} + \frac{z}{x+y} \geq M \geq m \geq \frac{3}{2}$$

We have already showed that $a+b \geq 2\sqrt{ab}$ which is a special case of the arithmetic-geometric mean inequality. Let us consider the following question. We are given four positive real numbers a, b, c, d such that $S = a+b = c+d$. Which of the products between ab and cd is the smaller one?

$$\begin{aligned} 4ab &= (a+b)^2 - (a-b)^2 \\ &= S^2 - (a-b)^2 \\ 4cd &= (c+d)^2 - (c-d)^2 \\ &= S^2 - (c-d)^2 \end{aligned}$$

As we can see here, the sign in $4ab?4cd$ (?) to be replaced by one of $>, <, \geq, \leq$) will be dictated by which of the differences $a-b, c-d$ is smaller. Since $x^2 \geq 0$, if $(a-b)^2 < (c-d)^2$, we have $4ab > 4cd$. Now, consider the product $a_1 \cdots a_n$. We want to see how the product changes as a_i varies with respect to $\bar{a} = \frac{a_1+\dots+a_n}{n}$.

If all the a_i are equal to each other, then we have nothing to check. Otherwise, there are at least two positive integers i and j such that a_i and a_j are not equal to \bar{a} . Moreover, one of them is greater than \bar{a} and the other is smaller than \bar{a} because all of them cannot be greater (or smaller) than \bar{a} . Without loss of generality, assume that a_1, a_2 are those two numbers and $a_1 = \bar{a} - h, a_2 = \bar{a} + k$. Now, consider two other positive numbers c and d which keeps the sum

²S. IVELIĆ BRA ANOVIĆ and JOSIP PEĆARIĆ, “Generalizations of converse Jensen’s inequality and related results”, *Journal of Mathematical Inequalities*, no. 1 (2011), pp. 43-60, doi: 10.7153/jmi-05-06.

fixed, for example $c = \bar{a}$, $d = \bar{a} + k - h$. We have

$$\begin{aligned} cd &= \bar{a}(\bar{a} + k - h) \\ &= \bar{a}^2 + \bar{a}k - \bar{a}h \\ a_1a_2 &= (\bar{a} - h)(\bar{a} + k) \\ &= \bar{a}^2 + \bar{a}k - \bar{a}h - hk \\ &= cd - hk < cd \\ a_1a_2 \cdots a_n &< cd \cdots a_n \end{aligned}$$

This basically tells us that we can increase the product further if there are other a_i which are not equal to \bar{a} and the product is maximum when all a_i is equal to \bar{a} .

PROBLEM 1.7. Show that for a positive integer n ,

$$n! > \left(\frac{n}{4}\right)^n$$

We can easily prove this with induction. This can be improved to $n! > \left(\frac{n}{3}\right)^n$. In fact, we can prove the following.

$$n! > \left(\frac{n}{e}\right)^n$$

This is another nice result. For example, setting $n = 2019$,

$$2019! > 673^{2019}$$

We can even bound $n!$ from both sides with the next result.

PROBLEM 1.8. For a positive integer n , prove the inequality

$$e \left(\frac{n+1}{e}\right)^{n+1} > n! > \left(\frac{n}{e}\right)^n$$

Equality is not possible because on both sides we have non-integers.

PROBLEM 1.9. Let x, y, z be positive real numbers. Prove that

$$\frac{2}{x+y} + \frac{2}{y+z} + \frac{2}{z+x} \geq \frac{9}{x+y+z}$$

Solution.

A function f is *non-decreasing* (resp. *increasing*) on the interval I if for any $a, b \in I$, $(a - b)(f(a) - f(b)) \geq 0$ (resp. $(a - b)(f(a) - f(b)) > 0$). Similarly, f is *non-increasing* (resp. *decreasing*) on the interval I if for any $a, b \in I$, $(a - b)(f(a) - f(b)) \leq 0$ (resp. $(a - b)(f(a) - f(b)) < 0$). If either of these two conditions apply for f , then f is a *monotone function*. We say that f is *monotonic*. Note that the slope between any two points $(a, f(a))$ and $(b, f(b))$ is $m = \frac{f(a)-f(b)}{a-b}$ and the quantity we have used for the definition is

$$\begin{aligned}(a - b)(f(a) - f(b)) &= (a - b)^2 \frac{f(a) - f(b)}{f(a) - f(b)} \\ &= (a - b)^2 m\end{aligned}$$

So, the sign of this quantity is the same as the sign of the slope m .

A sequence (a_n) is *non-decreasing* if $a_i \leq a_{i+1}$ for all $i \in \mathbb{N}$. (a_n) is *strictly increasing* if $a_i < a_{i+1}$. Similarly, (a_n) is *non-increasing* if $a_i \geq a_{i+1}$ for all i . (a_n) is *strictly decreasing* if $a_i > a_{i+1}$. If (a_n) is either increasing or decreasing, then (a_n) is a *monotone sequence*. We say that (a_n) is *monotonic*.

THEOREM 1.4 (Abel formula). Let (a_n) and (b_n) be two sequences of real numbers. If $c_k = b_1 + \dots + b_k$ for $1 \leq k \leq n$, then

$$a_1 b_1 + \dots + a_n b_n = (a_1 - a_2)c_1 + (a_2 - a_3)c_2 + \dots + (a_{n-1} - a_n)c_n + a_n c_n$$

Proof.

□

THEOREM 1.5 (Abel's Inequality). Let (a_n) and (b_n) be two sequences of real numbers such that $b_1 \geq \dots \geq b_n \geq 0$. For $1 \leq k \leq n$, define

$$\begin{aligned}s_k &= a_1 + \dots + a_k \\ m &= \min_{1 \leq i \leq n} (s_i) \\ M &= \max_{1 \leq i \leq n} (s_i)\end{aligned}$$

Then we have

$$mb_1 \leq a_1 b_1 + \dots + a_n b_n \leq Mb_1$$

Proof. Write the sum $a_1 b_1 + \dots + a_n b_n$ as the following.

$$\begin{aligned}a_1 b_1 + \dots + a_n b_n &= s_1 b_1 + (s_2 - s_1)b_2 + \dots + (s_n - s_{n-1})b_n \\ &= s_1(b_1 - b_2) + s_2(b_2 - b_3) + \dots + s_{n-1}(b_{n-1} - b_n) + s_n b_n\end{aligned}$$

Using $m \leq s_i \leq M$,

$$\begin{aligned} m(b_1 - b_2) &\leq s_1(b_1 - b_2) \leq M(b_1 - b_2) \\ &\vdots \\ m(b_{n-1} - b_n) &\leq s_{n-1}(b_{n-1} - b_n) \leq M(b_{n-1} - b_n) \end{aligned}$$

We additionally have $mb_n \leq s_n b_n \leq Mb_n$. Summing these inequalities together,

$$\begin{aligned} m(b_1 - b_2 + b_2 - b_3 + \dots + b_{n-1} - b_n) + mb_n &\leq s_1(b_1 - b_2) \\ &\quad + \dots + s_{n-1}(b_{n-1} - b_n) + s_n b_n \\ &\leq M(b_1 - b_2 + b_2 - b_3 + \dots \\ &\quad + b_{n-1} - b_n) + Mb_n \\ mb_1 &\leq a_1 b_1 + \dots + a_n b_n \leq Mb_1 \end{aligned}$$

This proves the inequality. \square

§§1.2 WARM UP PROBLEMS

Try the following problems as warm up exercises.

PROBLEM 1.10. For real numbers x, y , prove that

$$\begin{aligned} |x + y| &\leq |x| + |y| \\ ||x| - |y|| &\leq |x + y| \\ (|x| - |y|)^2 &\leq |x^2 - y^2| \end{aligned}$$

When does equality occur in the last inequality?

The first inequality in this problem is also known as the *triangle inequality*.

PROBLEM 1.11. Prove *Jordan's inequality*.

$$\frac{2}{\pi} \leq \frac{\sin \vartheta}{\vartheta} < 1$$

for $0 < |\vartheta| < \frac{\pi}{2}$.

PROBLEM 1.12. Prove Redheffer's inequality.

$$\frac{\sin \vartheta}{\vartheta} \geq \frac{\pi^2 - \vartheta^2}{\pi^2 + \vartheta^2}$$

The two inequalities above are not derived from each other. Also, you may have to put a bit of extra effort to prove them. Use calculus if that gives a faster solution.

PROBLEM 1.13 (IMO 1960). For which real value of x does the inequality

$$\frac{4x^2}{(1 - \sqrt{1 + 2x})^2} < 2x + 9$$

hold?

§§1.3 CAUCHY-SCHWARZ INEQUALITY AND IMPROVEMENTS

Cauchy-Schwarz inequality also known as *Cauchy-Bunyakovsky-Schwarz* inequality is among the most important results for solving problems.

THEOREM 1.6 (Cauchy-Bunyakovsky-Schwarz inequality). Let n be a positive integer, a_1, \dots, a_n and b_1, \dots, b_n be real numbers. Then

$$(a_1^2 + \dots + a_n^2)(b_1^2 + \dots + b_n^2) \geq (a_1 b_1 + \dots + a_n b_n)^2 \quad (\ddagger 1.6)$$

$$\left(\sum_{i=1}^n a_i^2 \right) \left(\sum_{i=1}^n b_i^2 \right) \geq \left(\sum_{i=1}^n a_i b_i \right)^2$$

and equality holds if and only if $\frac{a_1}{b_1} = \dots = \frac{a_n}{b_n}$.

The reason why it is also called Cauchy-Bunyakovsky-Schwarz inequality is that the analog of this inequality for integrals

$$\left(\int_a^b f(x)g(x)dx \right)^2 \leq \left(\int_a^b f^2(x)dx \right) \cdot \left(\int_a^b g^2(x)dx \right) \quad (\ddagger 1.7)$$

appeared in the *Mémoire BUNYAKOVSKY*³ for the first time. This *Mémoire* was published by the Imperial Academy of Sciences of St. Petersburg in BUNYAKOVSKY.⁴ STEELE⁵ states that *Bunyakovsky (1804–1889) had studied in Paris with Cauchy, and he was quite familiar with Cauchy's work on inequalities; so much so that by the time he came to write his Mémoire, Bunyakovsky was content to refer to the classical form of Cauchy' inequality for finite sums simply as well-known.*

This theorem has many proofs and even more applications. We will see later in § 2.1 how powerful just a special case of this inequality can be. Let us start with a proof that is probably the most elegant one. Before we show proofs, we will introduce a notation for shortening the *Cauchy-Bunyakovsky-Schwarz inequality*. Let two vectors (think of a vector as an ordered list of numbers) be $\mathbf{a} = (a_1, \dots, a_n)$ and $\mathbf{b} = (b_1, \dots, b_n)$. The numbers a_1, \dots, a_n inside the vector \mathbf{a} can be called *elements* of \mathbf{a} . So, \mathbf{a} has n elements and so does \mathbf{b} . Denote by $\langle \mathbf{a}, \mathbf{b} \rangle$ the *inner product* of \mathbf{a} and \mathbf{b} defined as

$$\langle \mathbf{a}, \mathbf{b} \rangle = a_1 b_1 + \dots + a_n b_n$$

Then the theorem can be stated as

$$\langle \mathbf{a}, \mathbf{b} \rangle^2 \leq \langle \mathbf{a}, \mathbf{a} \rangle \cdot \langle \mathbf{b}, \mathbf{b} \rangle$$

The inner product has some interesting properties. Note that the mother of inequality in Equation § 1.2 can be stated as

$$\langle \mathbf{x}, \mathbf{x} \rangle \geq 0$$

for any $\mathbf{x} = (x_1, \dots, x_n)$ where x_i is a real number. Similarly, define $\mathbf{y} = (y_1, \dots, y_n)$. Then we have the following.

- (i) $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ if and only if $\mathbf{x} = \mathbf{0}$ i.e. $\mathbf{x} = (0, \dots, 0)$.
- (ii) For any real number α , $\langle \alpha \mathbf{x}, \mathbf{y} \rangle = \alpha \langle \mathbf{x}, \mathbf{y} \rangle$.
- (iii) $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$.

³VIKTOR YAKOVLEVICH BUNYAKOVSKY, *Foundations of the mathematical theory of probabilities*, Imperial Academy of Sciences of St. Petersburg, 1846, p. 495, doi: <http://books.e-heritage.ru/book/10070419>, Page 4.

⁴VIKTOR YAKOVLEVICH BUNYAKOVSKY, “Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies”, *Mémoires de l'Acad. de St.-Pétersbourg*, 7th ser., vol. I, no. 9 (1859), pp. 1-18.

⁵JOHN MICHAEL STEELE, *The Cauchy-Schwarz master class: an introduction to the art of mathematical inequalities*, Cambridge University Press, 2010, Page 10.

(iv) For $\mathbf{z} = (z_1, \dots, z_n)$, $\langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle$.

(v) $\mathbf{x} + \mathbf{y} = (x_1 + y_1, \dots, x_n + y_n)$.

For convenience, we can also define \mathbf{x}^r as

$$\mathbf{x}^r = (x_1^r, \dots, x_n^r)$$

for a real number r . Let us generalize this notation further. Let m be a positive integer. Then for $\mathbf{x} = (x_1, \dots, x_n)$, define

$$\begin{aligned}\|\mathbf{x}\|_m &= \underbrace{\langle \mathbf{x}, \dots, \mathbf{x} \rangle}_{m \text{ times}}^{\frac{1}{m}} \\ &= \sqrt[m]{x_1^m + \dots + x_n^m}\end{aligned}$$

For $m = 2$, we will simply write $\|\mathbf{x}\|$ instead of $\|\mathbf{x}\|_2$ just like we write \sqrt{x} instead of $\sqrt[2]{x}$. So we can write *Cauchy-Bunyakovsky-Schwarz inequality* in a more compact form.

$$\|\mathbf{a}\| \cdot \|\mathbf{b}\| \geq \langle \mathbf{a}, \mathbf{b} \rangle$$

This notion can be generalized even further for arbitrary real numbers. For a real number p , define

$$\|\mathbf{x}\|_p = \sqrt[p]{x_1^p + \dots + x_n^p}$$

This is often called the L_p norm for the vector \mathbf{x} . Furthermore, we can define the addition or subtraction between two vectors as

$$\begin{aligned}\mathbf{a} \pm \mathbf{b} &= (a_1 \pm b_1, \dots, a_n \pm b_n) \\ m\mathbf{a} &= (ma_1, \dots, ma_n)\end{aligned}$$

These notations will help us shorten some long inequalities later. We will see the benefit of these notations shortly in *Hölder's inequality* and *Minkowski's Inequality*. Let us see the proofs now.

Proof by vector. Consider two vectors $\mathbf{a} = (a_1, \dots, a_n)$, $\mathbf{b} = (b_1, \dots, b_n)$ and their modulus

$$\begin{aligned}\|\mathbf{a}\| &= \sqrt{a_1^2 + \dots + a_n^2} \\ \|\mathbf{b}\| &= \sqrt{b_1^2 + \dots + b_n^2}\end{aligned}$$

If ϑ is the minimum angle between them, from the rule of *dot product* we get

$$\begin{aligned} \mathbf{a} \cdot \mathbf{b} &= a_1 b_1 + \dots + a_n b_n \\ \|\mathbf{a}\| \cdot \|\mathbf{b}\| \cdot \cos \vartheta &= a_1 b_1 + \dots + a_n b_n \\ \cos \vartheta &= \frac{a_1 b_1 + \dots + a_n b_n}{\sqrt{a_1^2 + \dots + a_n^2} \sqrt{b_1^2 + \dots + b_n^2}} \end{aligned}$$

Since $-1 \leq \cos \vartheta \leq 1$, after squaring we get

$$\begin{aligned} \cos^2 \vartheta &\leq 1 \\ \left(\frac{a_1 b_1 + \dots + a_n b_n}{\sqrt{a_1^2 + \dots + a_n^2} \sqrt{b_1^2 + \dots + b_n^2}} \right)^2 &\leq 1 \\ \Leftrightarrow (a_1^2 + \dots + a_n^2)(b_1^2 + \dots + b_n^2) &\geq (a_1 b_1 + \dots + a_n b_n)^2 \end{aligned}$$

Equality occurs if and only $\cos \vartheta = 1$ or when \mathbf{a} and \mathbf{b} are parallel. In other words, when we have

$$\frac{a_1}{b_1} = \dots = \frac{a_n}{b_n}$$

□

Classical proof by Schwarz. Consider the quadratic polynomial

$$\begin{aligned} P(x) &= \sum_{i=1}^n (a_i x - b_i)^2 \\ &= x^2 \sum_{i=1}^n a_i^2 - 2x \sum_{i=1}^n a_i b_i + \sum_{i=1}^n b_i^2 \\ &= Ax^2 - Bx + C \end{aligned}$$

where $A = \sum_{i=1}^n a_i^2$, $B = 2 \sum_{i=1}^n a_i b_i$, $C = \sum_{i=1}^n b_i^2$. Setting $x_i \rightarrow a_i x - b_i$ in § 1.2, we see that $P(x) \geq 0$. Then the discriminant of P must be ≤ 0 .

$$\begin{aligned} B^2 - 4AC &\leq 0 \\ 4(a_1 b_1 + \dots + a_n b_n)^2 - 4(a_1^2 + \dots + a_n^2)(b_1^2 + \dots + b_n^2) &\leq 0 \\ \Leftrightarrow (a_1^2 + \dots + a_n^2)(b_1^2 + \dots + b_n^2) &\geq (a_1 b_1 + \dots + a_n b_n)^2 \end{aligned}$$

Equality occurs if and only if $P(x) = 0$ for some x . Then

$$x = \frac{b_1}{a_1} = \dots = \frac{b_n}{a_n}$$

□

Proof by arithmetic-geometric mean. We use the notations from the proof above. If $A = 0$ or $B = 0$, then the inequality is an identity. So, assume that $A, B \neq 0$. Note the following.

$$\begin{aligned} \frac{a_1^2}{A} + \frac{b_1^2}{B} &\geq 2 \frac{a_1 b_1}{\sqrt{AB}} \\ &\vdots \\ \frac{a_n^2}{A} + \frac{b_n^2}{B} &\geq 2 \frac{a_n b_n}{\sqrt{AB}} \end{aligned}$$

Summing them together,

$$\begin{aligned} \frac{a_1^2}{A} + \frac{b_1^2}{B} + \dots + \frac{a_n^2}{A} + \frac{b_n^2}{B} &\geq 2 \frac{a_1 b_1 + \dots + a_n b_n}{\sqrt{AB}} \\ \frac{a_1^2 + \dots + a_n^2}{A} + \frac{b_1^2 + \dots + b_n^2}{B} &\geq 2 \frac{a_1 b_1 + \dots + a_n b_n}{\sqrt{AB}} \\ 1 + 1 &\geq 2 \frac{a_1 b_1 + \dots + a_n b_n}{\sqrt{AB}} \\ AB &\geq (a_1 b_1 + \dots + a_n b_n)^2 \end{aligned}$$

Again, equality holds if and only if $\frac{a_i}{A} = \frac{b_i}{B}$ or $\frac{a_i}{b_i} = \frac{A}{B} = c$, a constant for all i . \square

Proof using the mother of inequality. Let $x_i = \frac{a_i}{A}, y_i = \frac{b_i}{B}$ for $1 \leq i \leq n$. We have

$$\sum_{i=1}^n x_i^2 = \sum_{i=1}^n y_i^2 = 1$$

Rewrite the inequality as

$$\begin{aligned} x_1 y_1 + \dots + x_n y_n &\leq 1 \\ \Leftrightarrow 2(x_1 y_1 + \dots + x_n y_n) &\leq 2 \\ \Leftrightarrow 2(x_1 y_1 + \dots + x_n y_n) &\leq x_1^2 + \dots + x_n^2 + y_1^2 + \dots + y_n^2 \\ \Leftrightarrow (x_1 - y_1)^2 + \dots + (x_n - y_n)^2 &\geq 0 \end{aligned}$$

This is obviously true. \square

Proof using sequences. Define the sequence (S_n) as

$$\begin{aligned} D_n &= a_1 b_1 + \dots + a_n b_n \\ A_n &= a_1^2 + \dots + a_n^2 \\ B_n &= b_1^2 + \dots + b_n^2 \\ S_n &= D_n^2 - A_n B_n \end{aligned}$$

We want to show that $S_n \leq 0$. See the following.

$$\begin{aligned}
 D_{n+1}^2 - D_n^2 &= (D_n + a_{n+1}b_{n+1})^2 - D_n^2 \\
 &= 2a_{n+1}b_{n+1}D_n + a_{n+1}^2b_{n+1}^2 \\
 A_{n+1}B_{n+1} - A_nB_n &= (A_n + a_{n+1}^2)(B_n + b_{n+1}^2) - A_nB_n \\
 &= a_{n+1}^2B_n + b_{n+1}^2A_n + a_{n+1}^2b_{n+1}^2 \\
 S_{n+1} - S_n &= D_{n+1}^2 - A_{n+1}B_{n+1} - D_n^2 + A_nB_n \\
 &= 2a_{n+1}b_{n+1}D_n + a_{n+1}^2b_{n+1}^2 \\
 &\quad - a_{n+1}^2B_n - b_{n+1}^2A_n - a_{n+1}^2b_{n+1}^2 \\
 &= 2a_{n+1}b_{n+1}D_n - a_{n+1}^2B_n - b_{n+1}^2A_n \\
 &= 2a_{n+1}b_{n+1}(D_{n-1} + a_n b_n) - a_{n+1}^2(B_{n-1} \\
 &\quad + b_{n+1}^2) - b_{n+1}^2(A_{n-1} + a_{n+1}^2) \\
 &= 2a_{n+1}b_{n+1}D_{n-1} + 2a_{n+1}b_{n+1}a_n b_n - a_{n+1}^2B_{n-1} \\
 &\quad - a_{n+1}^2b_n^2 - b_{n+1}^2A_{n-1} - a_{n+1}^2b_n^2 \\
 &= 2a_{n+1}b_{n+1}D_{n-1} - a_{n+1}^2B_{n-1} \\
 &\quad - b_{n+1}^2A_{n-1} - (a_{n+1}b_n - a_n b_{n+1})^2 \\
 &\vdots \\
 &= -((a_{n+1}b_n - a_n b_{n+1})^2 + \dots + (a_{n+1}b_1 - a_1 b_{n+1})^2)
 \end{aligned}$$

Clearly, $S_{n+1} - S_n \leq 0$, so

$$S_{n+1} \leq S_n \leq \dots S_1 = 0$$

This proves the inequality. □

The fact about $S_{n+1} - S_n$ also proves the *Lagrange Identity*.

$$\left(\sum_{i=1}^n a_i^2 \right) \left(\sum_{i=1}^n b_i^2 \right) - \left(\sum_{i=1}^n a_i b_i \right)^2 = \sum_{1 \leq i < j \leq n} (a_i b_j - a_j b_i)^2$$

This equation immediately proves the theorem but it is not quite obvious how the identity follows unless someone knows it beforehand. For this reason, some people also call this the Lagrange inequality but we will stick with Cauchy-Schwarz.

Gram's inequality is a very nice generalization of *Cauchy-Bunyakovsky-Schwarz inequality*.

THEOREM 1.7 (Gram's inequality). Let $\mathbf{x}_1 = (x_{11}, \dots, x_{1n}), \dots, \mathbf{x}_n = (x_{n1}, \dots, x_{nn})$ be vectors with n elements. Then

$$\begin{vmatrix} \langle \mathbf{x}_1, \mathbf{x}_1 \rangle & \dots & \langle \mathbf{x}_1, \mathbf{x}_n \rangle \\ & \dots & \\ \langle \mathbf{x}_n, \mathbf{x}_1 \rangle & \dots & \langle \mathbf{x}_n, \mathbf{x}_n \rangle \end{vmatrix} \geq 0$$

Equality occurs if and only if there are real numbers a_1, \dots, a_n such that all of them are not zero at the same time and

$$a_1 \mathbf{x}_1 + \dots + a_n \mathbf{x}_n = \mathbf{0}$$

where $\mathbf{0}$ is the zero vector with every element 0.

This result is very intuitive to say the least. Note that we get *Cauchy-Bunyakovsky-Schwarz inequality* for $n = 2$ since

$$\begin{aligned} & \begin{vmatrix} \langle \mathbf{x}, \mathbf{x} \rangle & \langle \mathbf{x}, \mathbf{y} \rangle \\ \langle \mathbf{y}, \mathbf{x} \rangle & \langle \mathbf{y}, \mathbf{y} \rangle \end{vmatrix} \geq 0 \\ \iff & \langle \mathbf{x}, \mathbf{x} \rangle \cdot \langle \mathbf{y}, \mathbf{y} \rangle - \langle \mathbf{x}, \mathbf{y} \rangle \cdot \langle \mathbf{y}, \mathbf{x} \rangle \geq 0 \end{aligned}$$

implies *Cauchy-Bunyakovsky-Schwarz inequality* due to the fact $\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{y}, \mathbf{x} \rangle$. The condition of equality in Gram's inequality is *linear dependence*. Let $\mathbf{x}_1, \dots, \mathbf{x}_n$ be some vectors with n elements. Then they are *linearly dependent* if and only if there are real numbers a_1, \dots, a_n not all zero such that

$$a_1 \mathbf{x}_1 + \dots + a_n \mathbf{x}_n = \mathbf{0}$$

If such a_1, \dots, a_n do not exist, that is the condition is satisfied only when $a_1 = \dots = a_n = 0$, then $\mathbf{x}_1, \dots, \mathbf{x}_n$ are *linearly independent*.

§§1.4 COMPLEX NUMBERS

We may occasionally need complex numbers. So we provide a brief introduction here. A complex number z is defined as

$$z = x + iy$$

where x, y are real numbers and i is the imaginary unit such that $i^2 = -1$. We call x the real component and y the complex component of z . Complex

numbers follow the same properties as vectors which we established in § 1.3. That is for two complex numbers $z_1 = x_1 + iy_1$ and $z_2 = x_2 + iy_2$,

$$\begin{aligned} z_1 \pm z_2 &= (x_1 \pm x_2) + i(y_1 \pm y_2) \\ \langle z_1, z_2 \rangle &= x_1x_2 + y_1y_2 \end{aligned}$$

Here $\langle z_1, z_2 \rangle$ is the dot product of z_1 and z_2 . We define an additional operation \cdot as

$$z_1 \cdot z_2 = x_1y_2 - x_2y_1$$

The *modulus* of z is similar to L_2 norm.

$$|z| = \sqrt{x^2 + y^2}$$

We have the following property.

$$|z_1z_2| = |z_1| \cdot |z_2|$$

For a complex number $z = x + iy$, the *conjugate* of z is defined as

$$\bar{z} = x - iy$$

So, we have $z\bar{z} = |z|^2 = |\bar{z}|^2$. The *argument* of z is defined as

$$\arg(z) = \tan\left(\frac{y}{x}\right)$$

You can think of it as the angle the point (x, y) creates with the positive X axis and the origin.

PROBLEM 1.14. Prove the triangle inequality for complex numbers.

$$|z_1 + z_2| \leq |z_1| + |z_2|$$

This inequality can be generalized as the following.

$$|z_1 + \dots + z_n| \leq |z_1| + \dots + |z_n|$$

When does equality occur?

§§1.5 BERNOULLI AND POWER MEAN INEQUALITY

The *Bernoulli inequality* is a well known result and it has some nice consequences. We mentioned earlier that we will show more proofs of arithmetic-geometric mean inequality. We will show one such proof here using this inequality.

THEOREM 1.8 (*Bernoulli's inequality*). *Let n be a positive integer and $x > -1$ be a real number. Then*

$$(1 + x)^n \geq 1 + nx$$

This inequality can be generalized as the following result.

THEOREM 1.9 (*Generalized Bernoulli inequality*). *Let $x_1, \dots, x_n > -1$ be real numbers such that either all are positive or all are negative. Then*

$$(1 + x_1) \cdots (1 + x_n) > 1 + x_1 + \dots + x_n$$

Let a_1, \dots, a_n be positive real numbers. Then the *generalized mean* or *power mean of order r* is defined as

$$\mathfrak{M}_r(a_1, \dots, a_n) = \left(\frac{a_1^r + \dots + a_n^r}{n} \right)^{\frac{1}{r}}$$

Note that the arithmetic mean of a_1, \dots, a_n is actually $\mathfrak{M}_1(a_1, \dots, a_n)$. Similarly, the harmonic mean is $\mathfrak{M}_{-1}(a_1, \dots, a_n)$. Moreover, $\mathfrak{M}_0(a_1, \dots, a_n)$ is the geometric mean which we show below.

We may omit the numbers a_1, \dots, a_n and just call it \mathfrak{M}_r instead of $\mathfrak{M}_r(a_1, \dots, a_n)$ if the context is clear. An even better way to denote this would be using $\mathfrak{M}_r(\mathbf{a})$ where $\mathbf{a} = (a_1, \dots, a_n)$.

$$\mathfrak{M}_r(\mathbf{a}) = \frac{\|\mathbf{a}\|_r}{\sqrt[n]{n}}$$

We can also denote the arithmetic, geometric and harmonic means of \mathbf{a} by $\mathfrak{A}(\mathbf{a})$, $\mathfrak{G}(\mathbf{a})$ and $\mathfrak{H}(\mathbf{a})$ respectively. The notations $\mathfrak{M}, \mathfrak{A}, \mathfrak{G}, \mathfrak{H}$ are inspired by HARDY et al.⁶

⁶HARDY et al., *Inequalities* cit.

Next, we show that $\mathfrak{G}(\mathbf{a})$ is the geometric mean.

$$\mathfrak{G}(\mathbf{a}) = \lim_{r \rightarrow 0} \left(\frac{a_1^r + \dots + a_n^r}{n} \right)^{\frac{1}{r}}$$

Then we can write arithmetic-geometric-harmonic mean inequality as $\mathfrak{A}(\mathbf{a}) \geq \mathfrak{G}(\mathbf{a}) \geq \mathfrak{H}(\mathbf{a})$. This is generalized in the next result.

THEOREM 1.10 (*Power mean inequality*). *If r, s are real numbers such that $r \leq s$, then for a vector \mathbf{a} ,*

$$\min\{\mathbf{a}\} \leq \mathfrak{M}_r(\mathbf{a}) \leq \mathfrak{M}_s(\mathbf{a}) \leq \max\{\mathbf{a}\}$$

Equality occurs if and only if $r = s$.

This inequality can be extended further with the notion of what we call *weighted means*. Let $\omega = (w_1, \dots, w_n)$ be a vector of non-negative real numbers such that $w_1 + \dots + w_n = 1$. Then the weighted arithmetic mean of the real numbers a_1, \dots, a_n is

$$\mathfrak{A}(\mathbf{a}, \omega) = w_1 a_1 + \dots + w_n a_n$$

In general, the *weighted power mean of order r* is

$$\mathfrak{M}_r(\mathbf{a}, \omega) = (w_1 a_1^r + \dots + w_n a_n^r)^{\frac{1}{r}}$$

If the context is clear on what the weights are, then we may omit the weight from the notation and simply write $\mathfrak{M}_r(\mathbf{a})$. Let us call ω a *weight vector* if $w_1, \dots, w_n \geq 0$ and $w_1 + \dots + w_n = 1$. We can convert almost any vector of non-negative real numbers into a weight vector. If $\tau = (t_1, \dots, t_n)$ is an arbitrary vector not all elements zero, then

$$\omega = \left(\frac{t_1}{t_1 + \dots + t_n}, \dots, \frac{t_n}{t_1 + \dots + t_n} \right)$$

is a weight vector. The power mean inequality applies to weighted means as well.

THEOREM 1.11 (*Weighted Power Mean Inequality*). *Let \mathbf{a} and ω be vectors with n elements. Then for real numbers r, s such that $r \leq s$,*

$$\min\{\mathbf{a}\} \leq \mathfrak{M}_r(\mathbf{a}, \omega) \leq \mathfrak{M}_s(\mathbf{a}, \omega) \leq \max\{\mathbf{a}\}$$

A special case of this is the *weighted arithmetic-geometric mean inequality*.

$$w_1 a_1 + \dots + w_n a_n \geq a_1^{w_1} \cdots a_n^{w_n} \quad (\dagger 1.8)$$

§§1.6 HÖLDER, MINKOWSKI'S INEQUALITY

THEOREM 1.12 (*Generalized Hölder's Inequality*). *Let x_1, \dots, x_n be vectors of n positive real numbers and ω be a weight vector. Then*

$$\mathfrak{G} \left(\sum_{i=1}^n x_i, \omega \right) \geq \sum_{i=1}^n \mathfrak{G}(x_i, \omega)$$

Equality occurs if any two of the vectors are proportional.

This is not exactly the Hölder's inequality to be precise. This inequality can be specified as Generalized Hölder's inequality which appears in MINKOWSKI.⁷ And HÖLDER⁸ proved Hölder's inequality for which we need to define conjugates. Two real numbers u and v are *conjugate* if

$$\frac{1}{u} + \frac{1}{v} = 1$$

THEOREM 1.13 (*Hölder's inequality*). *Let $u > 1$ be a real number and v be the conjugate of u . If α, β are complex numbers,*

$$\langle \alpha, \beta \rangle \leq \|\alpha\|_u \cdot \|\beta\|_v$$

Equality occurs if α^u and β^v are proportional.

Hölder's inequality can be proved with the help of YOUNG.⁹

THEOREM 1.14 (*Young's inequality*). *Let a, b be positive real numbers and u, v be conjugates. Then*

$$ab \leq \frac{a^u}{u} + \frac{b^v}{v}$$

⁷HERMANN MINKOWSKI, *Geometrie der Zahlen*, Johnson Reprint Corporation, 1968, Page 117.

⁸O. HÖLDER, “Ueber einen Mittelwertsatz”, German, *Gött. Nachr.*, vol. MDCCCLXXXIX (1889), pp. 38-47.

⁹WILLIAM HENRY YOUNG, “On classes of summable functions and their Fourier Series”, *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, vol. LXXXVII, no. 594 (1912), pp. 225-229, doi: 10.1098/rspa.1912.0076.

RAZMINIA¹⁰ presents another proof of *Generalized Hölder's Inequality*. The next result is usually known as *the converse of Hölder's inequality*.

THEOREM 1.15 (*The converse of Hölder's inequality*). *Let \mathbf{a} and \mathbf{b} be vectors of positive real numbers, u and v be conjugates. If B is a positive real number, then a necessary and sufficient condition that*

$$\sum_{i=1}^n a_i^u \leq A$$

is that

$$\langle \mathbf{a}, \mathbf{b} \rangle \leq \sqrt[n]{A} \sqrt[n]{B}$$

holds for all positive real numbers b_1, \dots, b_n such that

$$\sum_{i=1}^n b_i^v \leq B$$

LJAPUNOV¹¹ proves the next theorem.

THEOREM 1.16 (*Ljapunov's inequality*). *Let \mathbf{x}, \mathbf{y} be two vectors of n positive real numbers and ω be a weight vector. If r, s, t are positive real numbers such that $r > s > t$,*

$$\langle \mathbf{x}, \mathbf{y}^s \rangle^{r-t} \leq \langle \mathbf{x}, \mathbf{y}^t \rangle^{r-s} \cdot \langle \mathbf{x}, \mathbf{y}^r \rangle^{s-t}$$

Note that *Generalized Hölder's Inequality* is a generalization of *Cauchy-Bunyakovsky-Schwarz inequality*. We can even prove the *Arithmetic-Geometric Inequality* from *Generalized Hölder's Inequality* in the following way.

$$\begin{aligned} \frac{a_1 + \dots + a_n}{n} &\geq \sqrt[n]{a_1 \cdots a_n} \\ \iff (a_1 + \dots + a_n)^n &\geq n^n a_1 \cdots a_n \end{aligned}$$

This follows from *Generalized Hölder's Inequality* if we set $\mathbf{x}_i = \mathbf{x}$ and $\omega = (\frac{1}{n}, \dots, \frac{1}{n})$. MINKOWSKI¹² proves the following result.

¹⁰K. RAZMINIA, "Hölder's inequality revisited", *The Mathematical Gazette*, vol. ciii, no. 558 (2019), pp. 512–514, doi: 10.1017/mag.2019.117.

¹¹ALEKSANDR MICHAJLOVIĆ LJAPUNOV, "Nouvelle forme du théorème sur la limite de probabilité", "Mémoires de l'Académie impériale des sciences de Saint-Pétersbourg", vol. Série VIII, Tome. 12 (1901), <http://www.sudoc.fr/170960269>.

¹²MINKOWSKI, *Geometrie der Zahlen* cit., Page 115 – 117.

THEOREM 1.17 (Minkowski's Inequality). Let x_1, \dots, x_n be vectors of n positive real numbers, ω be a weight vector and $r \neq 1$ be a real number. Then

$$\mathfrak{M}_r \left(\sum_{i=1}^n x_i, \omega \right) \begin{cases} \geq \sum_{i=1}^n \mathfrak{M}_r(x_i, \omega) & \text{if } r < 1 \\ \leq \sum_{i=1}^n \mathfrak{M}_r(x_i, \omega) & \text{otherwise} \end{cases}$$

§§1.7 REARRANGEMENT AND CHEBYSHEV INEQUALITIES

Two sequences (a_n) and (b_n) are *similarly sorted* if and only for any i and j , we have

$$(a_i - a_j)(b_i - b_j) \geq 0$$

(a_n) and (b_n) are *oppositely sorted* if the inequality is reversed.

THEOREM 1.18 (Chebyshev inequality). Let (a_n) and (b_n) be two similarly sorted sequences. Then

$$\left(\frac{1}{n} \sum_{i=1}^n a_i \right) \cdot \left(\frac{1}{n} \sum_{i=1}^n b_i \right) \leq \frac{1}{n} \sum_{i=1}^n a_i b_i$$

SASSER and SLATER¹³ shows the following result that gives a necessary and sufficient condition for Chebyshev inequality to hold for two sequences (a_n) and (b_n) .

THEOREM 1.19 (Sasser's theorem).

SEITZ¹⁴ proved the following result which combines Cauchy-Schwarz and Chebyshev inequality.

¹³D.W. SASSER and M.L. SLATER, “On the inequality $\sum x_i y_i \geq \frac{1}{n} \sum x_i \sum y_i$ and the van der Waerden permanent conjecture”, *Journal of Combinatorial Theory*, vol. III, no. 1 (1967), pp. 25-33, doi: 10.1016/s0021-9800(67)80012-7.

¹⁴JIRÉ SEITZ, “Une remarque aux inégalités”, *Aktuárské vyd*, vol. VI, no. 4 (1936), pp. 167-171.

THEOREM 1.20 (*Seitz inequality*). Let $(x_n), (y_n), (z_n), (u_n)$ be sequences of real numbers and a_{ij} be a real number for $1 \leq i, j \leq n$. If for all $i < j$ and $r < s$, the conditions

$$\begin{aligned} \begin{vmatrix} x_i & x_j \\ y_i & y_j \end{vmatrix} \cdot \begin{vmatrix} z_r & z_s \\ u_r & u_s \end{vmatrix} &\geq 0 \text{ and} \\ \begin{vmatrix} a_{ri} & a_{rj} \\ a_{si} & a_{sj} \end{vmatrix} &\geq 0 \end{aligned}$$

hold, then we have

$$\frac{\sum_{i,j=1}^n a_{ij}x_iz_j}{\sum_{i,j=1}^n a_{ij}x_iu_j} \geq \frac{\sum_{i,j=1}^n a_{ij}y_iz_j}{\sum_{i,j=1}^n y_iu_j}$$

§§1.8 CONVEXITY, JENSEN AND POPOVICIU'S INEQUALITY

A function $f : I \rightarrow \mathbb{R}$ is called *Jensen convex* on some interval $I = [a, b]$ if for any $x, y \in I$,

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2}$$

f is *Jensen concave* if $-f$ is Jensen convex. Similarly, a sequence of real numbers (a_n) is *Jensen convex* if

$$2a_n \leq a_{n-1} + a_{n+1}$$

holds for $n \in \mathbb{N}$. This concept of convexity was formally introduced first by JENSEN.¹⁵ Jensen also proved the following result.

THEOREM 1.21. Let w_1, \dots, w_n be non-negative rational numbers in the interval I such that $w_1 + \dots + w_n = 1$ and f be a Jensen convex function on I . Then

$$f(w_1x_1 + \dots + w_nx_n) \leq w_1f(x_1) + \dots + w_nf(x_n)$$

¹⁵ JOHAN LUDWIG WILLIAM VALDEMAR JENSEN, “On konvexe funktioner og uligheder mellem middlvaerdier”, *Nyt. Tidsskr. for Math.*, vol. 16B (1905), pp. 49–69; JOHAN LUDWIG WILLIAM VALDEMAR JENSEN, “Sur les fonctions convexes et les inégalités entre les valeurs moyennes”, *Acta Math.*, vol. XXX (1906), pp. 175–193, doi: 10.1007/bf02418571.

But the idea itself was not completely new. HÖLDER¹⁶ proved Theorem 1.21 under the condition that f can be differentiated at least twice and $f''(x) \geq 0$. Ironically, this is exactly the convexity condition from calculus point of view.

Let f be a convex function. For a vector \mathbf{x} , let $f(\mathbf{x})$ be the vector $(f(x_1), \dots, f(x_n))$.

THEOREM 1.22 (Weighted Jensen's inequality). *Let f be a convex function. For a vector \mathbf{x} of real numbers in the domain of f and a weight vector ω ,*

$$f(\mathfrak{A}(\mathbf{x}, \omega)) \leq \mathfrak{A}(f(\mathbf{x}), \omega)$$

THEOREM 1.23 (Petrovic's Inequality). *Let f be a convex function and (x_n) be a sequence of positive real numbers. Then*

$$f(x_1) + \dots + f(x_n) \leq f(x_1 + \dots + x_n) + (n - 1)f(0)$$

THEOREM 1.24 (Huygen's inequality). *Let \mathbf{x} be a vector with n elements and ω be a weight vector. Then*

$$\prod_{i=1}^n (1 + x_i) \geq \mathfrak{G}(\mathbf{x})$$

Note that this is just a special case of Hölder's inequality.

THEOREM 1.25 (Popoviciu's inequality). *Let f be a convex function in the interval $[a, b]$. For any $x, y, z \in [a, b]$,*

$$f\left(\frac{x+y+z}{3}\right) + \frac{f(x)+f(y)+f(z)}{3} \geq \left(f\left(\frac{x+y}{2}\right) + f\left(\frac{y+z}{2}\right) + f\left(\frac{z+x}{2}\right)\right)$$

¹⁶HÖLDER, “Ueber einen Mittelwertsatz” cit.

§2 TRADITIONAL PRINCIPLES

§§2.1 ENGEL FORM OF CAUCHY-SCHWARZ

We mentioned earlier that there is a particular case of Cauchy-Schwarz inequality which is very useful for solving problems. It is known as *Engel form of Cauchy-Schwarz*. Some people also call it *Titu's lemma* or *T2's lemma*. This name became popular among the USA students who attended the IMO training camp after a lecture given by Titu Andreescu at Math Olympiad Summer Program (MOSP) at Georgetown University in June, 2001. Even though it is a direct consequence of Cauchy-Schwarz inequality, it can be proven independently as well. Conversely, Cauchy-Schwarz inequality can be proven using this result.

THEOREM 2.1 (*Engel form of Cauchy-Schwarz*). *Let $a_1, \dots, a_n, b_1, \dots, b_n$ be real numbers. Then*

$$\frac{a_1^2}{x_1} + \dots + \frac{a_n^2}{x_n} \geq \frac{(a_1 + \dots + a_n)^2}{x_1 + \dots + x_n} \quad (\ddagger 2.1)$$

Some writers such as BELLMAN and MITRINOVITCH¹ also call this *Bergström's inequality* due to BERGSTROM.²

¹RICHARD BELLMAN, "Notes on Matrix Theory-IV (An Inequality Due to Bergstrom)", *The American Mathematical Monthly*, vol. LXII, no. 3 (1955), p. 172, doi: 10.2307/2306621; D. S. MITRINOVITCH, "Equivalence of two sets of inequalities", *The Mathematical Gazette*, vol. XLIII, no. 344 (1959), pp. 126-126, doi: 10.2307/3610201.

²H BERGSTROM, 'A triangle-inequality for matrices", *Den Elfte Skandinaviske Matematikerkongress, Trondheim* (1949), pp. 264-267.

Proof using Cauchy-Schwarz. Use Cauchy-Schwarz inequality on

$$\frac{a_1}{\sqrt{x_1}}, \dots, \frac{a_n}{\sqrt{x_n}}; \sqrt{x_1}, \dots, \sqrt{x_n}$$

We get

$$\begin{aligned} \left(\frac{a_1^2}{\sqrt{x_1}^2} + \dots + \frac{a_n^2}{\sqrt{x_n}^2} \right) (\sqrt{x_1}^2 + \dots + \sqrt{x_n}^2) &\geq \left(\frac{a_1}{\sqrt{x_1}} \sqrt{x_1} + \dots + \frac{a_n}{\sqrt{x_n}} \sqrt{x_n} \right)^2 \\ &= (a_1 + \dots + a_n)^2 \end{aligned}$$

This proves the theorem. \square

Proof by induction. We will use induction to prove § 2.1. The inequality is trivial for $n = 1$. For $n = 2$,

$$\begin{aligned} \frac{a^2}{x} + \frac{b^2}{y} &\geq \frac{(a+b)^2}{x+y} \\ \Leftrightarrow \frac{a^2y + b^2x}{xy} &\geq \frac{a^2 + 2ab + b^2}{x+y} \\ \Leftrightarrow a^2xy + b^2x^2 + a^2y^2 + b^2xy &\geq a^2xy + 2abxy + b^2xy \\ \Leftrightarrow (ay - bx)^2 &\geq 0 \end{aligned}$$

This is obviously true. Now, assume that the claim is true for

We get a very useful result as a corollary of this result.

THEOREM 2.2. Let $a_1, \dots, a_n, b_1, \dots, b_n$ be real numbers. Then

$$\frac{a_1^2}{b_1^2} + \dots + \frac{a_n^2}{b_n^2} \geq \left(\frac{a_1 + \dots + a_n}{b_1 + \dots + b_n} \right)^2$$

Proof. By *Engel form of Cauchy-Schwarz*,

$$\begin{aligned} \frac{a_1^2}{b_1^2} + \dots + \frac{a_n^2}{b_n^2} &\geq \frac{(a_1 + \dots + a_n)^2}{b_1^2 + \dots + b_n^2} \\ &\geq \frac{(a_1 + \dots + a_n)^2}{(b_1 + \dots + b_n)^2} \end{aligned}$$

The last inequality follows from the fact that $b_1^2 + \dots + b_n^2 \leq (b_1 + \dots + b_n)^2$. \square

Another result that is similar to *Engel form of Cauchy-Schwarz* is a special case of the *Beckenbach inequality*.

THEOREM 2.3. Let x_1, \dots, x_n and y_1, \dots, y_n be positive real numbers. Then

$$\frac{\sum_{i=1}^n x_i^2}{\sum_{i=1}^n x_i} + \frac{\sum_{i=1}^n y_i^2}{\sum_{i=1}^n y_i} \geq \frac{\sum_{i=1}^n (x_i + y_i)^2}{\sum_{i=1}^n (x_i + y_i)}$$

We will start with another proof of Nesbitt's inequality with the Engel form of Cauchy-Schwarz.

Nesbitt inequality using Engel form. We will use the fact that $a^2 + b^2 + c^2 \geq ab + bc + ca$.

$$\begin{aligned} S &= \frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} \\ &= \frac{a^2}{ab+ca} + \frac{b^2}{bc+ab} + \frac{c^2}{ca+bc} \\ &\geq \frac{(a+b+c)^2}{ab+ca+bc+ab+ca+bc} \\ &= \frac{(a+b+c)^2}{2(ab+bc+ca)} \\ &= \frac{a^2+b^2+c^2+2(ab+bc+ca)}{ab+bc+ca} \\ &= \frac{a^2+b^2+c^2}{2(ab+bc+ca)} + 1 \\ &\geq \frac{1}{2} + 1 = \frac{3}{2} \end{aligned}$$

□

Let us use this inequality to solve some problems.

PROBLEM 2.1. Let n be a positive integer and $a_1, \dots, a_n; b_1, \dots, b_n$ be positive real numbers. Show that

$$(a_1b_1 + \dots + a_nb_n) \left(\frac{a_1}{b_1} + \dots + \frac{a_n}{b_n} \right) \geq (a_1 + \dots + a_n)^2$$

Solution. By Engel form of Cauchy-Schwarz,

$$\begin{aligned} \frac{a_1}{b_1} + \dots + \frac{a_n}{b_n} &= \frac{a_1^2}{a_1b_1} + \dots + \frac{a_n^2}{a_nb_n} \\ &\geq \frac{(a_1 + \dots + a_n)^2}{a_1b_1 + \dots + a_nb_n} \\ \Leftrightarrow (a_1b_1 + \dots + a_nb_n) \left(\frac{a_1}{b_1} + \dots + \frac{a_n}{b_n} \right) &\geq (a_1 + \dots + a_n)^2 \end{aligned}$$

Remark. In order to apply the Engel form, we often need transformations like $a/b = a^2/b^2$. Sometimes we will divide the odd power into an even one and send the remaining one in the denominator, for example,

$$\frac{a^{2n+1}}{b} = \frac{(a^n)^2}{\frac{b}{a}}$$

so that we can then apply it. We will demonstrate these ideas in the following problems.

PROBLEM 2.2. For positive real numbers a, b, c , prove that

$$\frac{a}{b+2c} + \frac{b}{c+2a} + \frac{c}{a+2b} \geq 1$$

Solution. We again resort to the same trick.

$$\begin{aligned} \frac{a}{b+2c} + \frac{b}{c+2a} + \frac{c}{a+2b} &= \frac{a^2}{ab+2ca} + \frac{b^2}{bc+2ab} + \frac{c^2}{ca+2bc} \\ &\geq \frac{(a+b+c)^2}{3(ab+bc+ca)} \\ &= \frac{a^2+b^2+c^2+2(ab+bc+ca)}{3(ab+bc+ca)} \\ &\geq \frac{ab+bc+ca+2(ab+bc+ca)}{3(ab+bc+ca)} \end{aligned}$$

In the last line, we used $a^2 + b^2 + c^2 \geq ab + bc + ca$ which is well known by now.

PROBLEM 2.3. Let x, y, z be positive real numbers. Prove that

$$\frac{2}{x+y} + \frac{2}{y+z} + \frac{2}{z+x} \geq \frac{9}{x+y+z}$$

Solution. This is again a similar problem.

$$\begin{aligned} \frac{2}{x+y} + \frac{2}{y+z} + \frac{2}{z+x} &= \frac{4}{2(x+y)} + \frac{4}{2(y+z)} + \frac{4}{2(z+x)} \\ &\geq \frac{(2+2+2)^2}{2(x+y+y+z+z+x)} \\ &= \frac{36}{4(x+y+z)} \\ &= \frac{9}{x+y+z} \end{aligned}$$

PROBLEM 2.4. Let x, y, z be positive real numbers. Prove that

$$\frac{x^2}{(x+z)(x+y)} + \frac{y^2}{(y+x)(y+z)} + \frac{z^2}{(z+y)(z+x)} \geq \frac{3}{4}$$

Solution. First, see that $(x+z)(x+y) = x^2 + xy + zx + yz$. Next, we apply *Engel form of Cauchy-Schwarz*.

$$\begin{aligned} & \frac{x^2}{(x+z)(x+y)} + \frac{y^2}{(y+x)(y+z)} + \frac{z^2}{(z+y)(z+x)} \\ &= \frac{x^2}{x^2 + xy + zx + yz} + \frac{y^2}{y^2 + yz + xy + zx} + \frac{z^2}{z^2 + zx + yz + xy} \\ &\geq \frac{(x+y+z)^2}{x^2 + y^2 + z^2 + 3(xy + yz + zx)} \\ &= \frac{3}{4} \left(\frac{4(x+y+z)^2}{3(x^2 + y^2 + z^2 + 2(xy + yz + zx) + xy + yz + zx)} \right) \\ &= \frac{3}{4} \left(\frac{3(x+y+z)^2 + (x+y+z)^2}{3((x+y+z)^2 + xy + yz + zx)} \right) \\ &= \frac{3}{4} \left(\frac{3(x+y+z)^2 + x^2 + y^2 + z^2 + 2(xy + yz + zx)}{3(x+y+z)^2 + 3(xy + yz + zx)} \right) \\ &\geq \frac{3}{4} \left(\frac{3(x+y+z)^2 + 3(xy + yz + zx)}{3(x+y+z)^2 + 3(xy + yz + zx)} \right) \\ &= \frac{3}{4} \cdot 1 = \frac{3}{4} \end{aligned}$$

Thus, the inequality is proved.

PROBLEM 2.5 (IMO Shortlist 1993). For positive real numbers a, b, c and d , prove that

$$\frac{a}{b+2c+3d} + \frac{b}{c+2d+3a} + \frac{c}{d+2a+3b} + \frac{d}{a+2b+3c} \geq \frac{2}{3}$$

Solution. Writing

$$\frac{a}{b+2c+3d} = \frac{a^2}{ab+2ca+3ad}$$

and applying *Engel form of Cauchy-Schwarz*, we get

$$\begin{aligned}
 S &= \frac{a}{b+2c+3d} + \frac{b}{c+2d+3a} + \frac{c}{d+2a+3b} + \frac{d}{a+2b+3c} \\
 &= \frac{a^2}{ab+2ca+3ad} + \frac{b^2}{bc+2bd+3ab} + \frac{c^2}{cd+2ca+3bc} + \frac{d^2}{ad+2bd+3cd} \\
 &\geq \frac{(a+b+c+d)^2}{4ab+4bc+4ca+4ad+4bd+4cd} \\
 &= \frac{2}{3} \left(\frac{3(a+b+c+d)^2}{8(ab+bc+ca+ad+bd+cd)} \right) \\
 &= \frac{2}{3} \left(\frac{3(a+b+c+d)^2}{4((a+b+c+d)^2 - (a^2+b^2+c^2+d^2))} \right) \\
 &= \frac{2}{3} \left(\frac{4(a+b+c+d)^2 - (a+b+c+d)^2}{4(a+b+c+d)^2 - 4(a^2+b^2+c^2)} \right)
 \end{aligned}$$

By *Cauchy-Bunyakovsky-Schwarz inequality*,

$$\begin{aligned}
 (a^2 + b^2 + c^2 + d^2)(1^2 + 1^2 + 1^2 + 1^2) &\geq (a+b+c+d)^2 \\
 \Leftrightarrow -4(a^2 + b^2 + c^2 + d^2) &\leq -(a+b+c+d)^2 \\
 \Leftrightarrow 4(a+b+c+d)^2 - 4(a^2 + b^2 + c^2 + d^2) &\leq 4(a+b+c+d)^2 - (a+b+c+d)^2
 \end{aligned}$$

Thus, we have $S \geq \frac{2}{3} \cdot 1 = \frac{2}{3}$.

PROBLEM 2.6 (IMO 1995, problem 2). Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$\frac{1}{a^3(b+c)} + \frac{1}{b^3(c+a)} + \frac{1}{c^3(a+b)} \geq \frac{3}{2}$$

Solution. Write

$$\frac{1}{a^3(b+c)} = \frac{\frac{1}{a^2}}{a(b+c)}$$

and apply *Engel form of Cauchy-Schwarz*.

$$\begin{aligned}
 \frac{1}{a^3(b+c)} + \frac{1}{b^3(c+a)} + \frac{1}{c^3(a+b)} &= \frac{\frac{1}{a^2}}{a(b+c)} + \frac{\frac{1}{b^2}}{b(c+a)} + \frac{\frac{1}{c^2}}{c(a+b)} \\
 &\geq \frac{\left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)^2}{ab + ca + bc + ab + ca + bc} \\
 &= \frac{\left(\frac{ab + bc + ca}{abc}\right)^2}{2(ab + bc + ca)} \\
 &= \frac{ab + bc + ca}{2} \\
 &\geq \frac{3\sqrt[3]{ab \cdot bc \cdot ca}}{2} \\
 &= \frac{3\sqrt[3]{(abc)^2}}{2} \\
 &= \frac{3}{2}
 \end{aligned}$$

Here, we get $ab + bc + ca \geq 3\sqrt[3]{ab \cdot bc \cdot ca}$ by *Arithmetic-Geometric Inequality*.

PROBLEM 2.7 (Tournament of the Towns, 1998). Let a, b, c be positive real numbers. Prove that

$$\frac{a^3}{a^2 + ab + b^2} + \frac{b^3}{b^2 + bc + c^2} + \frac{c^3}{c^2 + ca + a^2} \geq \frac{a + b + c}{3}$$

Solution. We again make the numerator a square using

$$\frac{a^3}{a^2 + ab + b^2} = \frac{a^4}{a^3 + a^2b + ab^2}$$

Then

$$\begin{aligned}
 \frac{a^3}{a^2 + ab + b^2} + \frac{b^3}{b^2 + bc + c^2} + \frac{c^3}{c^2 + ca + a^2} &= \frac{a^4}{a^3 + a^2b + ab^2} + \frac{b^4}{b^3 + b^2c + bc^2} + \frac{c^4}{c^3 + c^2a + ca^2} \\
 &\geq \frac{(a^2 + b^2 + c^2)^2}{a^3 + b^3 + c^3 + ab(a+b) + bc(b+c) + ca(c+a)} \\
 &= \frac{(a^2 + b^2 + c^2)^2}{(a+b+c)(ab+bc+ca)} \\
 &= \frac{a^2 + b^2 + c^2}{a+b+c}
 \end{aligned}$$

The last line follows since $a^2 + b^2 + c^2 \geq ab + bc + ca$. Then the conclusion follows because by *Cauchy-Bunyakovsky-Schwarz inequality*,

$$\begin{aligned} (a^2 + b^2 + c^2)(1^2 + 1^2 + 1^2) &\geq (a + b + c)^2 \\ \Leftrightarrow \frac{a^2 + b^2 + c^2}{a + b + c} &\geq \frac{a + b + c}{3} \end{aligned}$$

§§2.2 THE BUFFALO WAY

The buffalo way is a *bashing* method of solving inequality problems. It usually exploits symmetry or cyclic property and involves tedious calculations, so it is not always elegant. Nonetheless it is a useful tool. The idea is to assume something like $a \leq b \leq c$ without losing any generality, specially when there is symmetry. In general, we want to assume that

$$\begin{aligned} a_1 &\leq a_2 \leq \cdots \leq a_n \\ a_2 &= a_1 + b_1 \\ &\vdots \\ a_n &= a_1 + b_n \end{aligned}$$

We can also assume the following.

$$\begin{aligned} a_2 &= a_1 + c_1 \\ a_3 &= a_2 + c_2 \\ &= a_1 + c_1 + c_2 \\ &\vdots \\ a_n &= a_{n-1} + c_{n-1} \\ &= a_1 + c_1 + \dots + c_{n-1} \end{aligned}$$

where $c_1, \dots, c_{n-1} \geq 0$. Out of these two forms, the latter is actually a consequence of the former one and is usually the more used one. We will see a few problems that implement this technique.

PROBLEM 2.8. Prove that for two non-negative reals x, y

$$x + y \geq 2\sqrt{xy}$$

Solution. Since the inequality is symmetric on x and y , we can assume without loss of generality that $x \leq y$. Let $y = x + k$.

$$\begin{aligned} x + y &\geq 2\sqrt{xy} \\ \Leftrightarrow 2x + k &\geq 2\sqrt{x(x+k)} \\ \Leftrightarrow 4x^2 + 4xk + k^2 &\geq 4x^2 + 4xk \\ \Leftrightarrow k^2 &\geq 0 \end{aligned}$$

which is evident.

Try proving the case $n = 3$ of arithmetic-geometric mean inequality with this technique. You will understand why we said this is often a tedious technique.

PROBLEM 2.9.

$$x + y + z \geq 3\sqrt[3]{xyz}$$

Let us also prove the following using buffalo way.

PROBLEM 2.10. Let $a, b, c > 0$ be real numbers. Prove that

$$a^2 + b^2 + c^2 \geq ab + bc + ca$$

Solution. Assume that $a \leq b \leq c$ and $b = a + x, c = a + x + y$ where $x, y \geq 0$.

$$\begin{aligned} a^2 + (a+x)^2 + (a+x+y)^2 &\geq a(a+x) + (a+x)(a+x+y) + (a+x+y)a \\ \Leftrightarrow 3a^2 + 2x^2 + y^2 + 4ax + 2xy + 2ay &\geq 3a^2 + x^2 + 4ax + 2ay + 2xy \\ \Leftrightarrow x^2 + y^2 &\geq 0 \end{aligned}$$

This is again true.

PROBLEM 2.11. Let $a, b, c \geq 0$ be real numbers such that $a \leq b \leq c$. Prove that

$$(a+b)(c+a)^2 \geq 6abc$$

Solution. Since $a \leq b \leq c$, let $b = a + x, c = a + x + y$.

$$\begin{aligned} (2a+x)(2a+x+y)^2 &\geq 6a(a+x)(a+x+y) \\ \Leftrightarrow (2a+x)(4a^2 + x^2 + y^2 + 4ax + 4ay + 2xy) &\geq 6a(a^2 + 2ax + x^2 + ay + xy) \\ \Leftrightarrow 8a^3 + 12a^2x + 8a^2y + 6ax^2 + 8axy + 2ay^2 + x^3 + 2ax^2 + xy^2 &\geq 6a^3 + 12a^2 + 6ax^2 + 6a^2y + 6axy \\ \Leftrightarrow 2a^3 + 2a^2y + 2axy + 2ay^2 + x^3 + 2x^2y + xy^2 &\geq 0 \end{aligned}$$

This evidently true.

Now, we will prove Nesbitt's inequality using the first form of the buffalo way.

Proof of Nesbitt's inequality by the buffalo way. Let $x \leq y \leq z$ and $y = x + a, z = y + b$ with $a \geq 0, b \geq 0$. We are required to prove that

$$\frac{x}{y+z} + \frac{y}{z+x} + \frac{z}{x+y} \geq \frac{3}{2}$$

Let us express each variable as a parameter of a single variable.

$$\begin{aligned} & \frac{y-a}{2y+b} + \frac{y}{2y+b-a} + \frac{y+b}{2y-a} \geq \frac{3}{2} \\ \Leftrightarrow & \frac{y-a}{2y+b} - \frac{1}{2} + \frac{y}{2y+b-a} - \frac{1}{2} + \frac{y+b}{2y-a} - \frac{1}{2} \geq 0 \\ \Leftrightarrow & \frac{1}{2} \left(\frac{2y-2a-2y-b}{2y+b} + \frac{2y-2y-b+a}{2y+b-a} + \frac{2y+2b-2y+a}{2y-a} \right) \geq 0 \\ \Leftrightarrow & -\frac{2a+b}{2y+b} + \frac{a-b}{2y+b-a} + \frac{2b+a}{2y-a} \geq 0 \end{aligned}$$

The last inequality actually holds because of the following:

$$\begin{aligned} & \frac{2b+a}{2y-a} + \frac{a-b}{2y+b-a} - \frac{2a+b}{2y+b} \geq \frac{2b+a}{2y+b-a} + \frac{a-b}{2y+b-a} - \frac{2a+b}{2y+b} \\ & \geq \frac{2a+2b}{2y+b-a} - \frac{2a+b}{2y+b} \\ & \geq \frac{2a+2b}{2y+b} - \frac{2a+b}{2y+b} \\ & = \frac{a}{2y+b} \end{aligned}$$

which is obviously at least 0. Here, we have used the facts that $\frac{1}{x} \geq \frac{1}{y}$ implies $\frac{1}{x-u} \geq \frac{1}{y}$ and $\frac{1}{x} \geq \frac{1}{y+v}$ for $u, v \geq 0$. \square

The next inequality is a nice application of the buffalo way.

PROBLEM 2.12. Let $x, y, z > 0$ be real numbers such that no two are equal. Show that,

$$\frac{1}{(x-y)^2} + \frac{1}{(y-z)^2} + \frac{1}{(z-x)^2} \geq \frac{4}{xy+yz+zx}$$

Solution. The expression on the left is symmetric on x, y, z . Without loss of generality, assume that $x \leq y \leq z$ and $y = x + a, z = x + a + b$.

$$\begin{aligned} \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{(a+b)^2} &\geq \frac{4}{x(x+a) + (x+a)(x+a+b) + (x+a+b)x} \\ &= \frac{4}{x^2 + xa + x^2 + 2ax + a^2 + bx + ab + x^2 + xa + bx} \\ &\geq \frac{4}{3x^2 + 4xa + 2xb + ab + a^2} \end{aligned}$$

Note that there is no x on the left side of the inequality. So, we can use $x > 0$ to get rid of it.

$$\begin{aligned} 3x^2 + 4xa + 2xb + ab + a^2 &> ab + a^2 \\ \frac{4}{3x^2 + 4xa + 2xb + ab + a^2} &< \frac{4}{ab + a^2} \end{aligned}$$

So, we are done if we can prove the following.

$$\begin{aligned} \frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{(a+b)^2} &\geq \frac{4}{ab + a^2} \\ &= \frac{4}{a(b+a)} \end{aligned}$$

This is not a trivial inequality. We have to show that this actually holds.

$$\begin{aligned} \frac{1}{a^2} + \frac{1}{b^2} - \frac{2}{a(b+a)} + \frac{1}{(a+b)^2} &\geq \frac{2}{a(b+a)} \\ \Leftrightarrow \frac{1}{b^2} + \left(\frac{1}{a} - \frac{1}{a+b}\right)^2 &\geq \frac{2}{a(a+b)} \\ \Leftrightarrow \frac{1}{b^2} + \left(\frac{b}{a(a+b)}\right)^2 &\geq \frac{2}{a(a+b)} \\ \Leftrightarrow \frac{1}{b^2} + \frac{b^2}{a^2(a+b)^2} &\geq \frac{2}{a(a+b)} \end{aligned}$$

The last inequality is true by arithmetic-geometric mean inequality since

$$\begin{aligned} \frac{1}{b^2} + \frac{b^2}{a^2(a+b)^2} &\geq 2\sqrt{\frac{1}{b^2} \cdot \frac{b^2}{a^2(a+b)^2}} \\ &= \frac{2}{a(a+b)} \end{aligned}$$

Solve the next problem as an exercise.

PROBLEM 2.13. For all real $x, y, z \geq 0$, prove that

$$\frac{x^3 + y^3 + z^3}{x^2y + y^2z + z^2x} \geq \sum \frac{x}{y+z} - 1$$

§§2.3 MAJORIZATION AND SYMMETRIC SUMS

Let \mathbf{x} and \mathbf{y} be two vectors of n real numbers. We say that \mathbf{x} *dominates* or *majorizes* \mathbf{y} if

$$\begin{aligned} x_1 &\geq x_2 \geq \dots \geq x_n \\ y_1 &\geq y_2 \geq \dots \geq y_n \\ x_1 + \dots + x_n &= y_1 + \dots + y_n \\ x_1 + \dots + x_k &\geq y_1 + \dots + y_k \end{aligned}$$

for $1 \leq k \leq n-1$. If \mathbf{x} dominates \mathbf{y} (resp. \mathbf{y} is *dominated by* \mathbf{x}), then we write $\mathbf{x} \succ \mathbf{y}$ (resp. $\mathbf{y} \prec \mathbf{x}$). For example, $(4, 0, 0) \succ (3, 1, 0) \succ (2, 2, 0)$. The vectors \mathbf{x} and \mathbf{y} need not be monotonic because we can just sort them into monotonic vectors.

We will also introduce the cyclic and symmetric polynomials and notations juxtaposed with them. The expression $x^2 + y^2 + z^2$ is *symmetric* whereas $x^2y + y^2z + z^2x$ is *cyclic* but not symmetric because $y^2x + z^2y + x^2z \neq x^2y + y^2z + z^2x$. A symmetric polynomial in the variables x_1, \dots, x_n should remain same regardless of the order in which the variables are used. So $f(x_1, \dots, x_n)$ is symmetric iff remains *invariant* for all permutations of x_1, \dots, x_n in the expression unlike the cyclic example we just saw. For example, $xy + yz + zx$ is symmetric and so is xyz . And $\frac{a}{b} + \frac{b}{c} + \frac{c}{a}$ is cyclic but not symmetric. We can use the cyclic and symmetric notations to represent the expressions in a short form. Here are some demonstrations.

$$\begin{aligned} a^2 + b^2 + c^2 &= \sum_{cyc} a^2 \\ a^2b + b^2c + c^2a &= \sum_{cyc} a^2b \\ xy + yz + zx &= \sum_{cyc} xy \end{aligned}$$

Note that even though the expression $xy + yz + zx$ and $a^2 + b^2 + c^2$ are symmetric, we do not consider them symmetric polynomial sums in this notation. A symmetric polynomial sum should have all $n!$ terms in the sum since it is symmetric on all $n!$ permutations of the variables in it. Even if there can be duplicates, the total number of terms should still remain $n!$. For this reason, this sum is often denoted by

$$\sum! \text{ or } \sum_{\sigma}$$

where σ indicates that the sum is taken over all possible permutations. Here are some examples.

$$\begin{aligned}\sum_{sym} a^2 &= 2(a^2 + b^2 + c^2) \\ \sum! x^2 y &= x^2 y + y^2 x + y^2 z + z^2 y + z^2 x + x^2 z \\ \sum_{\sigma} x^3 y &= x^3 y + x^3 z + y^3 z + y^3 x + z^3 x + z^3 y\end{aligned}$$

Let x and a be vectors with n elements. We write

$$\begin{aligned}F(x, a) &= x_1^{a_1} \cdots x_n^{a_n} \\ T(x, a) &= \sum! F(x, a) \\ &= \sum_{sym} F(x, a) \\ &= \sum! x_1^{a_1} \cdots x_n^{a_n}\end{aligned}$$

We will explain it a little bit more. Say, we want to find the symmetric polynomials of the type x^3y for the variables x, y, z . Write it as $x^3y^1z^0$. Now, we fix the powers 3, 1, 0 in their respective places and let x, y, z run through the permutations. Then we get

$$x^3y^1z^0 + x^3z^1y^0 + y^3z^1x^0 + y^3x^1z^0 + z^3x^1y^0 + z^3y^1x^0$$

See that this is exactly what we got in the last line of

$$\sum_{\sigma} x^3 y$$

Some authors write this notation as

$$T[a](x) = T[a_1, \dots, a_n](x_1, \dots, x_n)$$

Simply $T[\mathbf{a}] = T[a_1, \dots, a_n]$ is used if it is clear what \mathbf{x} is.

$$\begin{aligned} T[1, 0, \dots, 0] &= (x_1 + \dots + x_n)(n-1)! \\ T\left[\frac{1}{n}, \dots, \frac{1}{n}\right] &= n! \sqrt[n]{x_1 \cdots x_n} \end{aligned}$$

Then for a vector of positive real numbers \mathbf{x} , we can write the arithmetic-geometric mean inequality as

$$T[1, 0, \dots, 0] \geq T\left[\frac{1}{n}, \dots, \frac{1}{n}\right]$$

We can also define mean values based on the symmetric polynomials. We call

$$\mathfrak{M}[\mathbf{a}](\mathbf{x}) = \frac{T[\mathbf{a}](\mathbf{x})}{n!}$$

the *symmetrical mean*. For example,

$$\begin{aligned} \mathfrak{M}[1, 0 \dots, 0] &= \frac{(n-1)!}{n!}(a_1 + \dots + a_n) \\ &= \mathfrak{A}(\mathbf{a}) \\ \mathfrak{M}\left[\frac{1}{n}, \dots, \frac{1}{n}\right] &= \frac{n!}{n!} a_1^{\frac{1}{n}} \cdots a_n^{\frac{1}{n}} \\ &= \mathfrak{G}(\mathbf{a}) \end{aligned}$$

So, the symmetrical mean is a generalization of \mathfrak{A} and \mathfrak{G} . While we are talking about symmetric sums, we should consider the next identity.

$$(x + a_1) \cdots (x + a_n) = x^n + \binom{n}{1} x^{n-1} D_1 + \binom{n}{2} x^{n-2} D_2 + \dots + \binom{n}{n} D_n$$

where D_k is the sum of products of a_1, \dots, a_n taken k at the same time. If S_k is the coefficient of x^{n-k} in the expansion, then

$$S_k = \binom{n}{k} D_k$$

THEOREM 2.4 (*Newton's inequality*). *For $1 \leq i \leq n-1$, we have*

$$D_i^2 \geq D_{i-1} D_{i+1}$$

THEOREM 2.5 (*Maclaurin's inequality*). *With the same notation as Newton's inequality,*

$$D_1 \geq \sqrt{D_2} \geq \dots \geq \sqrt[n]{D_n}$$

§§2.4 KARAMATA'S INEQUALITY

KARAMATA³ proves the next theorem regarding convex functions when one vector majorizes another.

THEOREM 2.6 (*Karamata's inequality*). *Let f be a convex function and \mathbf{a} and \mathbf{b} be two vectors such that $\mathbf{a} \succ \mathbf{b}$. Then*

$$\sum_{i=1}^n f(a_i) \geq \sum_{i=1}^n f(b_i)$$

The next result first appeared at the forum *Art of Problem Solving*⁴. Daniel S. Liu showed this as a separate result but it also follows from *Karamata's inequality*.

THEOREM 2.7 (*Reverse rearrangement inequality*). *Let σ be a permutation of $\{1, \dots, n\}$ and $(a_n), (b_n)$ be two similarly sorted sequences of positive real numbers. Then*

$$(a_1 + b_1) \cdots (a_n + b_n) \leq (a_1 + b_{\sigma(1)}) \cdots (a_n + b_{\sigma(n)}) \leq (a_1 + b_n) \cdots (a_n + b_1)$$

Later this was generalized to the following result.

THEOREM 2.8 (*Generalized reverse rearrangement inequality*). *Let $(a_1), \dots, (a_m)$ be m similarly sorted sequences of n positive real numbers. Then*

$$\prod_{i=1}^m \sum_{j=1}^n a_{\sigma_j(i), j} \geq \prod_{i=1}^m \sum_{j=1}^n a_{i,j}$$

³JOVAN KARAMATA, “Sur une inégalité relative aux fonctions convexes”, French, *Publ. Math. Univ. Belgrade*, vol. 1 (1932), pp. 145-148.

⁴<https://artofproblemsolving.com/community/q2h617132p3677200>

§§2.5 THE BUNCHING METHOD: MUIRHEAD'S INEQUALITY

MUIRHEAD⁵ proves a very important result regarding symmetric sums and means. It is widely used for solving problems, specially coupled with some other method such as The Buffalo Way. It is also known as *the bunching principle* among the American students.

THEOREM 2.9 (Muirhead's inequality). *Let \mathbf{x} be a vector of positive real numbers and \mathbf{a} be dominated by \mathbf{b} . Then*

$$\mathfrak{M}[\mathbf{a}](\mathbf{x}) \leq \mathfrak{M}[\mathbf{b}](\mathbf{x})$$

PARIS and VENCOVSKÁ⁶ proves a generalization of this inequality but the result is of little practical use for Olympiad purposes.

THEOREM 2.10 (Generalization of Muirhead's inequality). *Let \mathbf{x} and \mathbf{y} be two vectors of positive real numbers such that $\mathbf{x} \succ \mathbf{y}$. If p_1, \dots, p_k are non-negative real numbers, then*

$$\sum_{\substack{S_1 \cup \dots \cup S_r = \{1, \dots, k\} \\ S_i \cap S_j = \emptyset}} \prod_{j=1}^r \left(\sum_{i \in S_j} p_i \right)^{\square x_j} \geq \sum_{\substack{S_1 \cup \dots \cup S_r = \{1, \dots, k\} \\ S_i \cap S_j = \emptyset}} \prod_{j=1}^r \left(\sum_{i \in S_j} p_i \right)^{\square y_j}$$

where \square in $\square x_j$ denotes that in the expansion of this power, we only consider the terms with non-zero power of p_i .

PROBLEM 2.14. Let a, b, c be positive real numbers. Prove that

$$a^7 + b^7 + c^7 \geq a^4b^3 + b^4c^3 + c^4a^3$$

Solution. Since $(7, 0, 0) \succ (4, 3, 0)$, by *Muirhead's inequality* we have

$$\begin{aligned} a^7b^0c^0 + a^7c^0b^0 + b^7c^0a^0 + b^7a^0c^0 + c^7a^0b^0 + c^7b^0a^0 &\geq a^4b^3c^0 + a^4c^3b^0 + b^4c^3a^0 \\ &\quad + b^4a^3c^0 + c^4a^3b^0 + c^4b^3a^0 \\ \iff 2(a^7 + b^7 + c^7) &\geq 2(a^4b^3 + b^4c^3 + c^4a^3) \\ \iff a^7 + b^7 + c^7 &\geq a^4b^3 + b^4c^3 + c^4a^3 \end{aligned}$$

⁵R. F. MUIRHEAD, "Some Methods applicable to Identities and Inequalities of Symmetric Algebraic Functions of n Letters", *Proceedings of the Edinburgh Mathematical Society*, vol. xxi (1902), pp. 144-162, doi: 10.1017/s001309150003460x.

⁶J. B. PARIS and A. VENCOVSKÁ, 'A Generalization of Muirhead's Inequality', *Journal of Mathematical Inequalities*, no. 2 (2009), pp. 181-187, doi: 10.7153/jmi-03-18.

PROBLEM 2.15. Let a, b, c be positive real numbers. Prove that

$$\frac{a^3}{bc} + \frac{b^3}{ca} + \frac{c^3}{ab} \geq a + b + c$$

Solution. We can rearrange the inequality into

$$\begin{aligned} \frac{a^4 + b^4 + c^4}{abc} &\geq a + b + c \\ \Leftrightarrow a^4 + b^4 + c^4 &\geq a^2bc + b^2ca + c^2ab \end{aligned}$$

This follows from *Muirhead's inequality* since $(4, 0, 0) \succ (2, 1, 1)$.

§§2.6 HOMOGENIZATION AND NORMALIZATION

§§2.7 SUBSTITUTIONS

§3 ADVANCED TECHNIQUES & INEQUALITIES

§§3.1 TANGENT LINE TRICK

The *tangent line trick* is a very useful trick which has been around for quite some time. Li¹ demonstrates some problems with this trick. Here, we will discuss this in detail and show how it can tackle problems.

Imagine that we are given an inequality in a_1, \dots, a_n . The inequality can be divided into a sum of expressions which is same for a_1, \dots, a_n . That is it can be written as $f(a_1) + \dots + f(a_n) \geq g(a_1, \dots, a_n)$. Now, we typically try to use Jensen's inequality in cases like this. But in many cases, f is not convex. Even so, we may still be able to prove something like

$$f(x) \geq f(\bar{a}) + (x - \bar{a})f'(\bar{a})$$

where $\bar{a} = \frac{a_1 + \dots + a_n}{n}$. Summing this for $x = a_1, \dots, a_n$, we have

$$f(a_1) + \dots + f(a_n) \geq nf(\bar{a})$$

The motivation sort of comes from basic calculus. By *wishful thinking*, we may be able to prove that the slope between the points $(x, f(x))$ and $(\bar{a}, f(\bar{a}))$ is at least the slope of the tangent line of $f(x)$ at $x = \bar{a}$. This is where the

¹KIN-YIN LI, "Using Tangent Lines to Prove Inequalities", *Mathematical Excalibre*, vol. x, no. 5 (2006), p. 1.

name comes from. This trick is specially useful if you are given the quantity $a_1 + \dots + a_n$. If the expressions involved are homogeneous, then you do not even need this value. You can use homogeneity to impose conditions such as $a + b + c = 1$. For example, see this problem.

PROBLEM 3.1. Let a, b, c be positive real numbers. Prove that

$$\frac{1}{a(b+c)} + \frac{1}{b(c+a)} + \frac{1}{c(a+b)} \geq \frac{27}{2(a+b+c)^2}$$

Solution. The inequality is homogeneous in a, b, c . So without loss of generality, we can normalize the inequality assuming that $a + b + c = 1$. Then $0 < a, b, c < 1$, $\bar{a} = 1/3$ and we get the transformation

$$\frac{1}{a(1-a)} + \frac{1}{b(1-b)} + \frac{1}{c(1-c)} \geq \frac{27}{2}$$

We define

$$f(x) = \frac{1}{x(1-x)}$$

and attempt to prove the inequality that is required by tangent line trick. Since

$$f'(x) = \frac{2x-1}{x^2(1-x)^2}$$

we have $f(1/3) = 9/2$ and $f'(1/3) = -27/4$. Thus, if we can prove the inequality

$$f(x) \geq \frac{9}{2} - \frac{27}{4} \left(x - \frac{1}{3} \right)$$

we are done since we will have

$$\begin{aligned} f(a) + f(b) + f(c) &\geq 3 \cdot f\left(\frac{1}{3}\right) \\ &= \frac{27}{2} \end{aligned}$$

So we should check if the tangent line inequality indeed holds.

$$\begin{aligned} \frac{1}{x(1-x)} &\geq \frac{9}{2} - \frac{27}{4} \left(x - \frac{1}{3} \right) \\ \Leftrightarrow \frac{1}{x(1-x)} &\geq \frac{54 - 81x + 27}{12} \\ &\geq \frac{27(1-x)}{4} \\ \Leftrightarrow x(1-x)^2 &\leq \frac{4}{27} \end{aligned}$$

We can check if this holds. If $g(x) = x(1-x)^2$, then

$$\begin{aligned} g'(x) &= -2x(1-x) + (1-x)^2 \\ &= (1-x)(1-3x) \end{aligned}$$

We have $g'(x) = 0$ if $x \in \{1, 1/3\}$ and

$$\begin{aligned} g''(x) &= -3(1-x) - (1-3x) \\ &= 6x - 4 \end{aligned}$$

Since $0 < x < 1$ and $g''(1) = 2 > 0$, $g''(1/3) = -2 < 0$, we have that $g(x)$ is maximum at $x = 1/3$ in the interval $(0, 1)$. Thus,

$$\begin{aligned} x(1-x)^2 &\leq \frac{1}{3} \left(1 - \frac{1}{3}\right)^2 \\ &= \frac{27}{4} \end{aligned}$$

We are finally done.

Remark. Tangent line is a neat trick but it may always not be pretty. But before you go all in, you can easily check if the desired inequality follows from

$$f(a_1) + \dots + f(a_n) \geq nf(\bar{a})$$

or not. If it does, there is a good chance that $f(x) \geq f(\bar{a}) + (x - \bar{a})f'(\bar{a})$ holds as well. Another indication that this might work is that equality occurs for $a_1 = \dots = a_n = \bar{a}$. In order to avoid calculation with fractional values, we could also assume $a_1 + \dots + a_n = n$ for normalization so $\bar{a} = 1$. Also, we may sometimes have to deal with \leq instead of \geq . Let us see an example of this type below.

PROBLEM 3.2. Given positive real numbers a, b, c such that $a+b+c \geq 3$. Prove that

$$\frac{1}{a^2+b+c} + \frac{1}{a+b^2+c} + \frac{1}{a+b+c^2} \leq 1$$

Solution. We have equality in the case $a = b = c = 1$. So, we may be optimistic that tangent line trick will work here. Using $b+c \geq 3-a$,

$$\frac{1}{a^2+b+c} + \frac{1}{a+b^2+c} + \frac{1}{a+b+c^2} \leq \frac{1}{a^2+3-a} + \frac{1}{b^2+3-b} + \frac{1}{c^2+3-c}$$

So we are done if we can prove that

$$\frac{1}{a^2 + 3 - a} + \frac{1}{b^2 + 3 - b} + \frac{1}{c^2 + 3 - c} \leq 1$$

Let

$$f(x) = \frac{1}{x^2 + 3 - x}$$

We have that

$$f'(x) = \frac{1 - 2x}{(x^2 + 3 - x)^2}$$

Since $f(1) = 1/3$ and $f'(1) = -1/9$, we need to prove the tangent line inequality

$$\begin{aligned} f(x) &\leq \frac{1}{3} - \frac{1}{9}(x - 1) \\ \Leftrightarrow \frac{1}{x^2 + 3 - x} &\leq \frac{4 - x}{9} \\ \Leftrightarrow 5x^2 + 3 - 7x - x^3 &\geq 0 \\ \Leftrightarrow x^3 - 5x^2 + 7x - 3 &\leq 0 \end{aligned}$$

We can see that $(x - 3)$ is a factor of this polynomial. So we can factor it easily.

$$\begin{aligned} \Leftrightarrow x^2(x - 3) - 2x(x - 3) + x - 3 &\leq 0 \\ \Leftrightarrow (x - 3)(x^2 - 2x + 1) &\leq 0 \\ \Leftrightarrow (3 - x)(x - 1)^2 &\geq 0 \end{aligned}$$

This is obviously true. And summing up the tangent line inequality, we get the conclusion.

PROBLEM 3.3 (USAMO 2003). Let a, b, c be positive real numbers. Prove that

$$\frac{(2a + b + c)^2}{2a^2 + (b + c)^2} + \frac{(2b + c + a)^2}{2b^2 + (c + a)^2} + \frac{(2c + a + b)^2}{2c^2 + (a + b)^2} \leq 8$$

Solution. Due to homogeneity, we can assume that $0 < a, b, c < 1$ and $a + b + c = 1$ without loss of generality. Then we are required to prove

$$\frac{(1 + a)^2}{2a^2 + (1 - a)^2} + \frac{(1 + b)^2}{2b^2 + (1 - b)^2} + \frac{(1 + c)^2}{2c^2 + (1 - c)^2} \leq 8$$

We can see that equality occurs for $a = b = c = 1/3$. Define

$$\begin{aligned} f(x) &= \frac{(1+x)^2}{2x^2 + (1-x)^2} \\ &= \frac{(1+x)^2}{3x^2 - 2x + 1} \end{aligned}$$

The tangent line inequality in this case is

$$f(x) \leq f(\bar{a}) + (x - \bar{a})f'(\bar{a})$$

where

$$\begin{aligned} f'(x) &= \frac{2(3x^2 - 2x + 1)(1+x) - 2(1+x)^2(3x - 1)}{(3x^2 - 2x + 1)^2} \\ &= \frac{2(1+x)}{3x^2 - 2x + 1} - \frac{2(1+x)^2(3x - 1)}{(3x^2 - 2x + 1)^2} \end{aligned}$$

We have $f(1/3) = 8/3$ and $f'(1/3) = 4$ so if we can prove that

$$\begin{aligned} f(x) &\leq \frac{8}{3} + 4\left(x - \frac{1}{3}\right) \\ &= \frac{4(3x + 1)}{3} \end{aligned}$$

for $0 < x < 1$, then the inequality follows.

$$\begin{aligned} f(x) &\leq \frac{4(3x + 1)}{3} \\ \Leftrightarrow \frac{(1+x)^2}{3x^2 - 2x + 1} &\leq \frac{12x + 3}{3} \\ \Leftrightarrow 36x^3 - 15x^2 - 2x + 1 &\geq 0 \\ \Leftrightarrow (3x - 1)^2(4x + 1) &\geq 0 \end{aligned}$$

This is obviously true. So, we have the inequality.

§§3.2 SCHUR-VORNICU-MILDORF INEQUALITY

HARDY et al.² states the following as *Schur's inequality*.

²HARDY et al., *Inequalities* cit., Page 64.

THEOREM 3.1 (Schur's inequality). *Let n be a positive integer. If a, b, c are positive real numbers, then*

$$a^n(a-b)(a-c) + b^n(b-c)(b-a) + c^n(c-a)(c-b) \geq 0$$

BARNARD and CHILD³ states this theorem for $n \leq -1$ as well. HARDY et al.⁴ mentions this result in connection with the following result.

THEOREM 3.2. *Consider a vector of positive real numbers \mathbf{x} such that no two elements are equal. If $v \geq 0$ and $\delta > 0$, then*

$$\mathfrak{M}[v + 2\delta, 0, 0, a_4, \dots] - 2\mathfrak{M}[v + \delta, \delta, 0, a_4, \dots] + \mathfrak{M}[v, \delta, \delta, a_4, \dots] \geq 0$$

for a vector of non-negative real numbers \mathbf{a} .

It was Issai Schur who informed HARDY et al.⁵ that Theorem 3.2 does not follow from Muirhead's inequality but from Schur's inequality if we set $n = \frac{v}{\delta}$. So Theorem 3.1 is accredited to Schur following NEVILLE, OPPENHEIM, WATSON, and WRIGHT.⁶ Schur's inequality and its reverse have been generalized in many ways. For example, GUHA⁷ proves the following result.

THEOREM 3.3 (Guha's inequality). *Let a, b, c, u, v, w be positive real numbers such that for a real number p ,*

$$\sqrt[p]{a} + \sqrt[p]{c} \leq \sqrt[p]{b} \quad (\ddagger 3.1)$$

$$\sqrt[p+1]{u} + \sqrt[p+1]{w} \leq \sqrt[p+1]{v} \quad (\ddagger 3.2)$$

If $p > 0$, then

$$abc - vca + wab \geq 0 \quad (\ddagger 3.3)$$

If $-1 < p < 0$, then $\ddagger 3.3$ is reverse. If $p < -1$, then $\ddagger 3.1$ and $\ddagger 3.2$ need to be reversed. Equality occurs if and only if

$$\frac{a^{p+1}}{u^p} = \frac{b^{p+1}}{v^p} = \frac{c^{p+1}}{w^p}$$

³S. BARNARD and J. M. CHILD, *Higher Algebra*, New Academic Science, 2018, Page 217.

⁴HARDY et al., *Inequalities* cit.

⁵Ibid.

⁶E. H. NEVILLE, "Schur's inequality", *The Mathematical Gazette*, vol. XL, no. 333 (1956), pp. 216-216, doi: 10.2307/3608826; A. OPPENHEIM, "2739. Generalisations of Schur's inequality", *The Mathematical Gazette*, vol. XLII, no. 339 (1958), pp. 35-35, doi: 10.2307/3608352; G. N. WATSON, "A trivial inequality", *The Mathematical Gazette*, vol. XL, no. 334 (1956), pp. 288-288, doi: 10.2307/3609631; G.N. WATSON, "Schur's Inequality", *The Mathematical Gazette*, vol. XXXIX, no. 329 (1955), pp. 207-208, doi: 10.2307/3608749; E. M. WRIGHT, "A generalisation of Schur's inequality", *The Mathematical Gazette*, vol. XL, no. 333 (1956), pp. 217-217, doi: 10.2307/3608827.

⁷U. C. GUHA, "Inequalities leading to generalisations of Schur's inequality", *The Mathematical Gazette*, vol. XLVI, no. 357 (1962), pp. 227-229, doi: 10.2307/3614027.

OPPENHEIM and DAVIES⁸ proves the following result which can be thought as the reverse of Schur's inequality.

THEOREM 3.4. *Let $n \geq 3$ be an integer and a_1, \dots, a_n be real numbers. Then*

$$\sum_{i=1}^n (x_i - x_1) \cdots (x_i - x_{i-1}) \cdot (x_i - x_{i+1}) \cdots (x_i - x_n) \geq 0$$

holds for all real numbers x_1, \dots, x_n such that $x_1 \geq \dots \geq x_n$ if

$$\begin{cases} a_1 \geq 0; a_2 \leq \left(a_1^{\frac{1}{2}} + a_3^{\frac{1}{2}}\right)^2; a_3 \geq 0 & \text{if } n = 3 \\ a_2 \leq a_1; (-1)^n(a_{n-1} - a_n) \geq 0; (-1)^{k+1}a_k \geq 0 & \text{if } n \geq 4 \end{cases}$$

where $1 \leq k \leq n-1$ and $k \notin \{2, n-1\}$.

We will first mention a notable generalization of Schur's inequality.

THEOREM 3.5 (Schur-Vornicu-Mildorfinequality). *Let a, b, c be three real numbers and x, y, z be non-negative real numbers. Then*

$$x(a-b)(a-c) + y(b-c)(b-a) + z(c-a)(c-b) \geq 0$$

holds if one of the following conditions hold:

- (1) $a \geq b \geq c$ and $x \geq y$.
- (2) $a \geq b \geq c$ and $z \geq y$.
- (3) $a \geq b \geq c$ and $x+z \geq y$.
- (4) a, b, c are non-negative, $a \geq b \geq c$ and $ax \geq by$.
- (5) a, b, c are non-negative, $a \geq b \geq c$ and $cz \geq by$.
- (6) a, b, c are non-negative, $a \geq b \geq c$ and $ax + cz \geq by$.
- (7) x, y, z are lengths of a triangle.
- (8) x, y, z are square of lengths of a triangle.
- (9) ax, by, cz are the lengths of a triangle.
- (10) There is a convex function f on the interval I such that $x = f(a), y = f(b), z = f(c)$ for non-negative real numbers $a, b, c \in I$.

⁸A. OPPENHEIM and Roy O. DAVIES, "Inequalities of Schur's Type", *The Mathematical Gazette*, vol. XLVIII, no. 363 (1964), pp. 25-27, doi: 10.2307/3614303.

VORNICU⁹ generalized this result to the following.

THEOREM 3.6 (Vornicu-Schur inequality). *Let $a, b, c; x, y, z$ be real numbers such that $a \geq b \geq c$ and either $x \geq y \geq z$ or $z \geq y \geq x$. Let k be a positive integer f be a non-negative real valued convex or monotonic function. Then*

$$f(x)(a-b)^k(a-c)^k + f(y)(b-c)^k(b-a)^k + f(z)(c-a)^k(c-b)^k \geq 0$$

We can easily verify that Schur's inequality follows from Vornicu-Schur inequality by setting $x = a, y = b, z = c, k = 1$ and $f(m) = m^n$.

§§3.3 RADON'S INEQUALITY

RADON¹⁰ proves the following generalization of Engel form of Cauchy-Schwarz inequality.

THEOREM 3.7 (Radon's Inequality). *Let b_1, \dots, b_n be positive real numbers and $a_1, \dots, a_n; m$ be non-negative real numbers. Then*

$$\frac{a_1^{m+1}}{b_1^m} + \dots + \frac{a_n^{m+1}}{b_n^m} \geq \frac{(a_1 + \dots + a_n)^{m+1}}{(b_1 + \dots + b_n)^m}$$

BECKENBACH¹¹ proves a similar result. For a vector of positive real numbers \mathbf{a} and a real number r , let us define

$$\mathfrak{R}_r(\mathbf{a}) = \frac{\sum_{i=1}^n a_i^r}{\sum_{i=1}^n a_i^{r-1}}$$

Note that this is actually a weighted arithmetic mean of \mathbf{a} with the weight vector

$$\omega = \left(\frac{a_1^{r-1}}{\sum_{i=1}^n a_i^{r-1}}, \dots, \frac{a_n^{r-1}}{\sum_{i=1}^n a_i^{r-1}} \right)$$

⁹VALENTIN VORNICU, *Olimpiada de matematica: de la provocare la experienta*, GIL Publishing House, 2003, vol. v.

¹⁰JOHANN RADON, “Über die absolut additiven Mengenfunktionen”, *Wien. Sitz. (IIa)*, vol. CXXII (1913), pp. 1295-1438.

¹¹EDWIN F. BECKENBACH, ‘A Class of Mean Value Functions’, *The American Mathematical Monthly*, vol. LVII, no. 1 (1950), p. 1, doi: 10.2307/2305163.

This is also a generalization of arithmetic, geometric and harmonic means. We get the usual arithmetic and harmonic means if we set $r = 1$ and -1 respectively. $r = \frac{1}{2}$ gives us the geometric mean for $n = 2$.

THEOREM 3.8 (*Beckenbach's inequality*). *Let r be a real number. Then*

$$\mathfrak{R}_r(a+b) \begin{cases} \leq \mathfrak{R}_r(a) + \mathfrak{R}_r(b) & \text{if } 1 \leq r \leq 2 \\ \geq \mathfrak{R}_r(a) + \mathfrak{R}_r(b) & \text{if } 0 \leq r \leq 1 \end{cases}$$

Equality occurs if and only if $r = 1$ or a is proportional to b .

YANG¹² proves the following generalization of *Radon's Inequality*.

THEOREM 3.9 (*Generalized Radon's Inequality*). *Let b_1, \dots, b_n be positive real numbers and*

$$a_1, \dots, a_n; r, s$$

be non-negative real numbers such that $r \geq s + 1$. Then

$$\frac{a_1^r}{b_1^s} + \dots + \frac{a_n^r}{b_n^s} \geq \frac{(a_1 + \dots + a_n)^r}{n^{r-s-1}(b_1 + \dots + b_n)^n}$$

YONGTAO et al.¹³ proves the equivalence between some well known inequalities.

THEOREM 3.10 (*Equivalence of inequalities*). *The following inequalities are mutually equivalent.*

- i. *Bernoulli's inequality*
- ii. *The weighted arithmetic-geometric mean inequality*
- iii. *Hölder's inequality*
- iv. *The weighted power mean inequality*
- v. *Minkowski's inequality*
- vi. *Radon's inequality*

¹²KECHANG YANG, ‘A Note and Generalization of a Fractional Inequality’, *Journal of Yueyang Normal University (Natural Science Edition)*, vol. xv, no. 4 (2002), pp. 9-11, doi: https://caod.oriprobe.com/articles/18339537/A_Note_and_Generalization_of_a_Fractional_Inequality.htm.

¹³LI YONGTAO et al., ‘A note on the proofs of generalized Radon inequality’, *Mathematica Moravica*, vol. xxii, no. 2 (2018), pp. 59-67, doi: 10.5937/matmor18020591.

§§3.4 SMOOTHING AND ISOLATED FUDGING

In many inequalities of the form

$$f(x_1, \dots, x_n) \geq a$$

for some real number a , we may be able to establish the property that f assumes smaller values when the difference between two variables x_i and x_j decreases. In such cases, we can use the fact that f assumes the smallest value when $x_1 = \dots = x_n$. This is known as the *smoothing principle*. Recall that we used a similar argument for arithmetic-geometric mean inequality when we replaced the product $a_1 a_2$ by $\bar{a}(\bar{a} + k - h)$ where $a_1 = \bar{a} - h$ and $a_2 = \bar{a} + k$.

PROBLEM 3.4 (USA 1996). Let a_0, \dots, a_n be real numbers in the interval $(0, \frac{\pi}{2})$. If

$$\tan\left(a_0 - \frac{\pi}{4}\right) + \dots + \tan\left(a_n - \frac{\pi}{4}\right) \geq n - 1$$

prove that

$$\tan a_0 \cdots \tan a_n \geq n^{n+1}$$

Solution. If $x_i = \tan(a_i - \frac{\pi}{4})$ so $-1 < x_i < 1$ and

$$\begin{aligned} x_i &= \frac{\tan a_i - 1}{1 + \tan a_i} \\ &= \frac{y_i - 1}{1 + y_i} \end{aligned}$$

where $y_i = \tan a_i$. Then

$$y_i = \frac{1 + x_i}{1 - x_i}$$

§§3.5 THE UVW/PQR/ABC METHOD

The UVW method was popularized by ROZENBERG.¹⁴ It has also been known as the ABC method or the PQR method although it is unclear what exactly

¹⁴MICHAEL ROZENBERG, “uvw–Method in Proving Inequalities”, *Math. Ed.*, vol. 59-60, no. 3-4 (2011), pp. 6-14, doi: http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=mo&paperid=194&option_lang=eng.

the origin of this method is. Although some people believe it was originated at Vietnam where it was called the ABC (Abstract concreteness) method. Currently, KNUDSEN¹⁵ is the most popular version of this technique but at the core, they all use the same idea. We will try to explain this method as clearly as possible with examples.

Consider an inequality in three variables $a, b, c \in \mathbb{R}$. If the expression is symmetric on a, b, c , then the initial idea of PQR method was to write

$$\begin{aligned} a + b + c &= p \\ ab + bc + ca &= q \\ abc &= r \end{aligned}$$

so that a, b, c are the roots of the equation

$$x^3 - px^2 + qx - r = 0$$

However, nowadays the most popular way to convert these is to write

$$\begin{aligned} a + b + c &= 3u \\ ab + bc + ca &= 3v^2 \\ abc &= w^3 \end{aligned}$$

Hence, the name UVW While working with such transformations, be careful not to assume $v^2 \geq 0$ by default. We have to consider the case where v^2 is negative as well. We do not know for sure that a, b, c are all positive unless it is stated specifically. However, the following result is very nice when they are indeed positive. We will use the notations as stated above throughout this section and the names of the following theorems follow KNUDSEN.¹⁶

THEOREM 3.11 (The Idiot Theorem). *If $a, b, c \geq 0$, then $u \geq v \geq w$.*

Proof. Using $a^2 + b^2 + c^2 \geq ab + bc + ca$ and $a + b + c \geq 3\sqrt[3]{abc}$, we immediately see that

$$(a + b + c)^2 \geq 3(ab + bc + ca)$$

¹⁵MATHIAS TEJS KNUDSEN, “The UVW-method”, *Art of problem solving ()*, doi: <https://artofproblemsolving.com/community/q2h278791p1507763>, <https://artofproblemsolving.com/community/q2h278791p1507763>.

¹⁶Ibid.

which implies $9u^2 \geq 9v^2$ or $u^2 \geq v^2$ and $u \geq v$. For proving $v \geq w$, we can use arithmetic-harmonic mean inequality

$$\begin{aligned} \frac{3}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}} &\leq \sqrt[3]{abc} \\ \Leftrightarrow \frac{3abc}{ab + bc + ca} &\leq \sqrt[3]{abc} \\ \Leftrightarrow ab + bc + ca &\geq 3\sqrt[3]{(abc)^2} \\ \Leftrightarrow 3v^2 &\geq 3w^2 \end{aligned}$$

This proves the theorem. \square

THEOREM 3.12 (The UVW theorem). *Let u, v, w be numbers such that u, v^2, w^3 are real numbers. Then there exists real numbers a, b, c such that*

$$\begin{aligned} a + b + c &= 3u \\ ab + bc + ca &= 3v^2 \\ abc &= w^3 \end{aligned}$$

if and only if $u^2 \geq v^2$ and

$$3uv^2 - 2u^3 - 2\sqrt{(u^2 - v^2)^3} \leq w^3 \leq 3uv^2 - 2u^3 + 2\sqrt{(u^2 - v^2)^3}$$

For proving this theorem, we will need the following result first.

THEOREM 3.13. *a, b, c are real numbers if and only if $(a - b)(b - c)(c - a)$ is a real number.*

Proof. The if part is obvious. So we will focus on the only if part. If $(a - b)(b - c)(c - a)$ is not real, then a, b, c are not real either. Clearly, a, b, c are the roots of $x^3 - 3ux^2 + 3vx - w^3 = 0$ by Vieta's formulas. If one of the roots is complex, say a , then another of b, c is complex, say b . Moreover, we must have $a = z$ and $b = \bar{z}$ for some complex number z since complex roots can appear only in conjugates. Then

$$(a - b)(b - c)(c - a) = (z - \bar{z})(\bar{z} - c)(c - z)$$

Letting $z = u + iv$ for some $u, v \in \mathbb{R}$, we have $\bar{z} = u - iv$ and

$$\begin{aligned} (a - b)(b - c)(c - a) &= -2iv(u - c + iv)(u - c - iv) \\ &= -2iv((u - c)^2 + v^2) \end{aligned}$$

This is obviously a complex number, so the claim is true unless $v = 0$ which cannot hold since $z \notin \mathbb{R}$. \square

Proof of The UVW theorem. For any $x \in \mathbb{R}$, we have $x^2 \geq 0$. By (3.13), $a, b, c \in \mathbb{R}$ implies that $(a - b)(b - c)(c - a) \in \mathbb{R}$ and so $(a - b)^2(b - c)^2(c - a)^2 \in \mathbb{R}$. Then

$$\begin{aligned} & (a - b)^2(b - c)^2(c - a)^2 \geq 0 \\ \iff & 27(-(w^3 - (3uv^2 - 2u^3))^2 + 4(u^2 - v^2)^3) \geq 0 \\ \iff & 4(u^2 - v^2)^3 \geq (w^3 - (3uv^2 - 2u^3))^2 \\ \iff & 2\sqrt{(u^2 - v^2)^3} \geq |w^3 - (3uv^2 - 2u^3)| \end{aligned}$$

This proves the theorem. \square

THEOREM 3.14 (The positivity theorem). *a, b, c are non-negative real numbers if and only if u, v^2, w^3 are non-negative real numbers.*

Proof. The if part is obvious, so we prove the only if part. So we should prove that if any of a, b, c are negative, then at least one of u, v^2, w^3 is negative. It is easy to see that $w^3 = abc$ is negative if one or three of a, b, c are negative. So, we are left with the case where two of a, b, c are negative. Without loss of generality, assume that a and b are negative whereas c is non-negative. Let $a = -x, b = -y$ so that

$$\begin{aligned} a + b + c &= 3u \\ c - x - y &= 3u \\ ab + bc + ca &= 3v^2 \\ &= xy - c(x + y) \end{aligned}$$

From this we can show that at least one of u or v has to be negative. Otherwise, if both u and v are non-negative, then $c - x - y \geq 0$ and $xy - c(x + y) \geq 0$. But this leads to a contradiction since

$$\begin{aligned} xy &\geq c(x + y) \\ &\geq (x + y)^2 \\ &\geq 2xy \end{aligned}$$

by Arithmetic-Geometric Inequality. \square

THEOREM 3.15 (Tej's theorem). *Let a, b, c be non-negative real numbers. Then we have the following:*

1. *If at least one value of w^3 exists for fixed u and v^2 corresponding to a, b, c , then w^3 has a global maximum and minimum (see Global maximum and minimum). This maximum value is achieved when at least two of a, b, c are equal. The minimum value is achieved when two of a, b, c are equal or when one of them is 0.*

2. If at least one value of v^2 exists for fixed u and w^3 corresponding to a, b, c , then v^2 has a global maximum and minimum. This maximum value is achieved when at least two of a, b, c are equal. The minimum value is achieved when two of a, b, c are equal or when one of them is 0.
3. If at least one value of u exists for fixed v^2 and w^3 corresponding to a, b, c , then u has a global maximum and minimum. This maximum value is achieved when at least two of a, b, c are equal. The minimum value is achieved when two of a, b, c are equal or when one of them is 0.

THEOREM 3.16. Every symmetric inequality of degree at most 5 has to be proved only for $a_1 = a_2$ or $a_n = 0$.

Proof. Using Elementary polynomials, □

§§3.6 MIXING VARIABLES TECHNIQUE

HUNG¹⁷

§§3.7 SEPARATION

We prove the arithmetic-geometric inequality yet again. Assume that $x_1 \cdots x_n = 1$ and we want to show that

$$x_1 + \dots + x_n \geq n$$

Assume that the result is valid for n and we want to prove it for $n+1$. Without loss of generality, we can assume that $x_1 \leq 1$ and $x_2 \geq 1$. Otherwise, $x_i \geq 1$ for all i implies that $x_i = 1$ which makes the inequality trivially true.

$$\begin{aligned} (1 - x_1)(x_2 - 1) &\leq 0 \\ \implies x_1 + x_2 &\geq 1 + x_1 x_2 \end{aligned}$$

¹⁷PHAM KIM. HUNG, *Secrets in inequalities*, GIL Publishing House, 2007.

Then we have the following by induction.

$$\begin{aligned} x_1 + \dots + x_{n+1} &\geq 1 + x_1x_2 + x_3 \cdots x_{n+1} \\ &\geq 1 + n \end{aligned}$$

The trick here is to separate the variables in terms of which side of 1 they are on. It is almost the same as being on the same or different sides of a line a point is. See the following problem for a better explanation.

PROBLEM 3.5. Let a, b, c be positive real numbers. Prove that

$$a^2 + b^2 + c^2 + 2abc + 1 \geq 2(ab + bc + ca)$$

Solution. By the *pigeonhole principle*, at least two of a, b, c are on the same side of 1. Without loss of generality, assume that a and b are on the same side of 1. Then

$$\begin{aligned} (a - 1)(b - 1) &\geq 0 \\ \implies ab &\geq a + b - 1 \end{aligned}$$

Then we have

$$\begin{aligned} a^2 + b^2 + c^2 + 2abc + 1 &\geq a^2 + b^2 + c^2 + 2c(a + b - 1) + 1 \\ &= (c - 1)^2 + a^2 + 2ca + 2bc + b^2 \\ &\geq 0 + 2ab + 2bc + 2ca \end{aligned}$$

PROBLEM 3.6 (USAMO 2001, problem 3). Let a, b, c be positive real numbers such that

$$a^2 + b^2 + c^2 + 2abc = 4$$

Prove that,

$$0 \leq ab + bc + ca - abc \leq 2$$

Solution.

PROBLEM 3.7. Let a, b, c be real numbers. Prove that

$$4(1 + a^2)(1 + b^2)(1 + c^2) \geq 3(a + b + c)^2$$

Solution. Again, we can assume without loss of generality that $ab + 1 \geq a + b$.

$$\begin{aligned} (1 + a^2)(1 + b^2) &= 1 + a^2 + b^2 + a^2b^2 \\ &\geq 1 + a^2 + b^2 + (a + b - 1)^2 \\ &= 2(1 + a^2 + b^2) + 2(ab - a - b) \end{aligned}$$

§§3.8 FLIPPING

Sometimes we face inequalities that give us the wrong signs after using some common techniques. We can consider subtracting or dividing some expression to reach the desired form. Let us see how to do that using the following problems.

PROBLEM 3.8. Let x_1, \dots, x_n be real numbers such that $x_1 + \dots + x_n = n$. Prove that,

$$\frac{1}{x_1^2 + 1} + \dots + \frac{1}{x_n^2 + 1} \geq \frac{n}{2}$$

Solution. First, see that $x^2 + 1 \geq 2x$ gives us

$$\frac{1}{x_i^2 + 1} \leq \frac{1}{2x_i}$$

This is the opposite sign of what we want to show. One idea to flip the signs is to send the expressions on the other side like this.

$$\begin{aligned} \frac{1}{x_1^2 + 1} + \dots + \frac{1}{x_n^2 + 1} + n &\geq \frac{n}{2} + n \\ \Leftrightarrow n &\geq \frac{n}{2} + \left(1 - \frac{1}{x_1^2 + 1}\right) + \dots + \left(1 - \frac{1}{x_n^2 + 1}\right) \\ \Leftrightarrow \frac{n}{2} &\geq \frac{x_1^2}{x_1^2 + 1} + \dots + \frac{x_n^2}{x_n^2 + 1} \end{aligned}$$

This is evident since

$$\begin{aligned} \frac{1}{x_i^2 + 1} &\leq \frac{1}{2x_i} \\ \Leftrightarrow \frac{x_i^2}{x_i^2 + 1} &\leq \frac{x_i}{2} \end{aligned}$$

Summing over $1 \leq i \leq n$, we get

$$\begin{aligned} \frac{x_1^2}{x_1^2 + 1} + \dots + \frac{x_n^2}{x_n^2 + 1} &\leq \frac{x_1 + \dots + x_n}{2} \\ &= \frac{n}{2} \end{aligned}$$

PROBLEM 3.9 (PuMaC 2014). Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 = 3$. Prove that

$$\frac{1}{a^3 + 2} + \frac{1}{b^3 + 2} + \frac{1}{c^3 + 2} \geq 1$$

Solution. We again have a similar situation here. Using *Arithmetic-Geometric Inequality*, we get $a^3 + 1 + 1 \geq 3a$ so

$$\frac{1}{a^3 + 2} \leq \frac{1}{3a}$$

Let us use the same strategy as the example right above.

$$\begin{aligned} \frac{1}{a^3 + 2} + \frac{1}{b^3 + 2} + \frac{1}{c^3 + 2} + 3 &\geq 1 + 3 \\ \Leftrightarrow 3 &\geq 1 + \left(1 - \frac{1}{a^3 + 2}\right) + \left(1 - \frac{1}{b^3 + 2}\right) + \left(1 - \frac{1}{c^3 + 2}\right) \\ \Leftrightarrow 2 &\geq \frac{a^3 + 1}{a^3 + 2} + \frac{b^3 + 1}{b^3 + 2} + \frac{c^3 + 1}{c^3 + 2} \end{aligned}$$

We are still facing a problem that the constants in the numerators of each expressions are not vanishing. The reason behind it is the constant 2 which is not fully canceled out. In order to completely cancel it out, we need to subtract the expression from $1/2$ instead of 1.

$$\begin{aligned} \frac{1}{a^3 + 2} + \frac{1}{b^3 + 2} + \frac{1}{c^3 + 2} + \frac{3}{2} &\geq 1 + \frac{3}{2} \\ \Leftrightarrow \frac{3}{2} &\geq 1 + \left(\frac{1}{2} - \frac{1}{a^3 + 2}\right) + \left(\frac{1}{2} - \frac{1}{b^3 + 2}\right) + \left(\frac{1}{2} - \frac{1}{c^3 + 2}\right) \\ \Leftrightarrow \frac{1}{2} &\geq \frac{a^3}{2(a^3 + 2)} + \frac{b^3}{2(b^3 + 2)} + \frac{c^3}{2(c^3 + 2)} \\ \Leftrightarrow 1 &\geq \frac{a^3}{a^3 + 2} + \frac{b^3}{b^3 + 2} + \frac{c^3}{c^3 + 2} \end{aligned}$$

This is again something we can work on.

$$\begin{aligned} \frac{1}{a^3 + 2} &\leq \frac{1}{3a} \\ \Leftrightarrow \frac{a^3}{a^3 + 2} &\leq \frac{a^3}{3a} \end{aligned}$$

Summing over,

$$\frac{a^3}{a^3 + 2} + \frac{b^3}{b^3 + 2} + \frac{c^3}{c^3 + 2} \leq \frac{a^2 + b^2 + c^2}{3}$$

This is exactly what we wanted.

Remark. This can be generalized to the following using the same technique. If a_1, \dots, a_n are positive real numbers such that $a_1^2 + \dots + a_n^2 = n$,

$$\frac{1}{a_1^3 + 2} + \dots + \frac{1}{a_n^3 + 2} \geq \frac{n}{3}$$

PROBLEM 3.10 (IMO 2005, problem 3). Let x, y, z be real numbers such that $xyz \geq 1$. Prove that

$$\frac{x^5 - x^2}{x^5 + y^2 + z^2} + \frac{y^5 - y^2}{x^2 + y^5 + z^2} + \frac{z^5 - z^2}{x^2 + y^2 + z^5} \geq 0$$

Solution. Rearrange the inequality as

$$\begin{aligned} & \frac{x^5 - x^2}{x^5 + y^2 + z^2} + \frac{y^5 - y^2}{x^2 + y^5 + z^2} + \frac{z^5 - z^2}{x^2 + y^2 + z^5} \geq 0 \\ \iff & \frac{x^5 - x^2}{x^5 + y^2 + z^2} + \frac{y^5 - y^2}{x^2 + y^5 + z^2} + \frac{z^5 - z^2}{x^2 + y^2 + z^5} + 3 \geq 3 \\ \iff & \left(1 - \frac{x^5 - x^2}{x^5 + y^2 + z^2}\right) + \left(1 - \frac{y^5 - y^2}{x^2 + y^5 + z^2}\right) + \left(1 - \frac{z^5 - z^2}{x^2 + y^2 + z^5}\right) \leq 3 \\ \iff & \frac{x^2 + y^2 + z^2}{x^5 + y^2 + z^2} + \frac{x^2 + y^2 + z^2}{x^2 + y^5 + z^2} + \frac{x^2 + y^2 + z^2}{x^2 + y^2 + z^5} \leq 3 \end{aligned}$$

Now, there is no obvious way to use something like *Cauchy-Bunyakovsky-Schwarz inequality* or *Engel form of Cauchy-Schwarz* here. And it looks like the denominators are supposed to be in place of the numerators. So, we will try to accomplish that.

$$\begin{aligned} x^5 + y^2 + z^2 &= \frac{x^4}{\frac{1}{x}} + \frac{y^4}{y^2} + \frac{z^4}{z^2} \\ &\geq \frac{(x^2 + y^2 + z^2)^2}{\frac{1}{x} + y^2 + z^2} \\ \iff \frac{x^2 + y^2 + z^2}{x^5 + y^2 + z^2} &\leq \frac{\frac{1}{x} + y^2 + z^2}{x^2 + y^2 + z^2} \end{aligned}$$

So we have

$$\begin{aligned} \frac{\frac{1}{x} + y^2 + z^2}{x^2 + y^2 + z^2} + \frac{x^2 + \frac{1}{y} + z^2}{x^2 + y^2 + z^2} + \frac{x^2 + y^2 + \frac{1}{z}}{x^2 + y^2 + z^2} &\leq \frac{yz + y^2 + z^2}{x^2 + y^2 + z^2} + \frac{x^2 + zx + z^2}{x^2 + y^2 + z^2} + \frac{x^2 + y^2 + xy}{x^2 + y^2 + z^2} \\ &= \frac{2(x^2 + y^2 + z^2) + xy + yz + zx}{x^2 + y^2 + z^2} \\ &\leq \frac{2(x^2 + y^2 + z^2) + x^2 + y^2 + z^2}{x^2 + y^2 + z^2} \end{aligned}$$

This gives us the desired inequality.

PROBLEM 3.11 (APMO 1991, problem 3). Let $a_1, \dots, a_n, b_1, \dots, b_n$ be positive real numbers such that

$$a_1 + \dots + a_n = b_1 + \dots + b_n$$

Show that

$$\frac{a_1^2}{a_1 + b_1} + \dots + \frac{a_n^2}{a_n + b_n} \geq \frac{a_1 + \dots + a_n}{2}$$

Solution. See the following.

$$\begin{aligned} &\frac{a_1^2}{a_1 + b_1} + \dots + \frac{a_n^2}{a_n + b_n} \geq \frac{a_1 + \dots + a_n}{2} \\ \Leftrightarrow &\frac{a_1^2}{a_1 + b_1} + \dots + \frac{a_n^2}{a_n + b_n} + \frac{a_1 + \dots + a_n}{2} \geq a_1 + \dots + a_n \\ \Leftrightarrow &\frac{a_1 + \dots + a_n}{2} \geq \left(a_1 - \frac{a_1^2}{a_1 + b_1}\right) + \dots + \left(a_n - \frac{a_n^2}{a_n + b_n}\right) \\ &= \frac{a_1 b_1}{a_1 + b_1} + \dots + \frac{a_n b_n}{a_n + b_n} \\ \Leftrightarrow &a_1 + \dots + a_n \geq \frac{2a_1 b_1}{a_1 + b_1} + \dots + \frac{2a_n b_n}{a_n + b_n} \\ \Leftrightarrow &\frac{1}{2}(a_1 + b_1 + \dots + a_n + b_n) \geq \frac{2a_1 b_1}{a_1 + b_1} + \dots + \frac{2a_n b_n}{a_n + b_n} \\ \Leftrightarrow &\frac{a_1 + b_1}{2} + \dots + \frac{a_n + b_n}{2} \geq \frac{2a_1 b_1}{a_1 + b_1} + \dots + \frac{2a_n b_n}{a_n + b_n} \end{aligned}$$

The last inequality is evidently true since $\frac{2ab}{a+b}$ is the harmonic mean of a and b . Then we can use the following to verify the inequality.

$$\frac{a+b}{2} \geq \frac{2ab}{a+b}$$

§§3.9 NEUTRALIZING DENOMINATORS

§§3.10 DUMBASSING

§§3.11 THE EQUAL VARIABLE METHOD

CÎRTOAJE¹⁸

§§3.12 PROBABILITY IN INEQUALITY

¹⁸VASILE CÎRTOAJE, “The Equal Variable Method”, *J. Inequal. Pure Appl. Math.*, vol. viii, no. 1 (2007), pp. 1-21, doi: <https://www.emis.de/journals/JIPAM/article828.html#>.

§4 PRACTICE PROBLEMS

We will pose various types of problems here. The problems will not be sorted in any particular order. There will be no categorization and similar problems will not be put together. We did not provide practice problems per section or chapter because in real contest, you will have no way of knowing what technique or theorem to use. This is why we demonstrated some problems while discussing a particular concept but did not add any extra problems. Here, you are free to use any method necessary to solve the problems.

PROBLEM 4.1. Let a, b, c be positive real numbers. Prove that

$$(a^3 + 2)(b^3 + 2)(c^3 + 2) \geq (a + b + c)^3$$

Solution. Rearrange the inequality as

$$(a^3 + 1^3 + 1^3)^{\frac{1}{3}}(1^3 + b^3 + 1^3)^{\frac{1}{3}}(1^3 + 1^3 + c^3)^{\frac{1}{3}} \geq (a^3 1^3 1^3)^{\frac{1}{3}} + (1^3 b^3 1^3)^{\frac{1}{3}} + (1^3 1^3 c^3)^{\frac{1}{3}}$$

which is obviously true by *Generalized Hölder's Inequality*.

PROBLEM 4.2. Let a, b be positive real numbers such that $a + b = 1$. Prove that

$$\frac{1}{a^2} + \frac{1}{b^2} \geq 8$$

Solution. Using *Generalized Hölder's Inequality* on $(a+b), (a+b), (\frac{1}{a^2} + \frac{1}{b^2})$,

$$(a+b)^{\frac{1}{3}}(a+b)^{\frac{1}{3}}\left(\frac{1}{a^2} + \frac{1}{b^2}\right)^{\frac{1}{3}} \geq \left(a \cdot a \cdot \frac{1}{a^2}\right)^{\frac{1}{3}} + \left(b \cdot b \cdot \frac{1}{b^2}\right)^{\frac{1}{3}}$$

PROBLEM 4.3. Let a, b, c be positive real numbers such that $a + b + c = 1$. Prove that

$$4a^3 + 9b^3 + 36c^3 \geq 1$$

Notice that 4, 9, 36 are square numbers while $a + b + c$ is also 1. This along with the terms like $4a^3$ tells us to somehow cancel the factors 4, 9, 36 so that we can use the fact that

$$(a^3)^{\frac{1}{3}} + (b^3)^{\frac{1}{3}} + (c^3)^{\frac{1}{3}} = 1$$

Fortunately, we also have $\frac{1}{2} + \frac{1}{3} + \frac{1}{6} = 1$ so using *Generalized Hölder's Inequality*,

$$\begin{aligned} & \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{6}\right)^{\frac{1}{3}} \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{6}\right)^{\frac{1}{3}} (4a^3 + 9b^3 + 36c^3)^{\frac{1}{3}} \geq \\ & \left(\frac{1}{2} \cdot \frac{1}{2} \cdot 4a^3\right)^{\frac{1}{3}} + \left(\frac{1}{3} \cdot \frac{1}{3} \cdot 9b^3\right)^{\frac{1}{3}} + \left(\frac{1}{6} \cdot \frac{1}{6} \cdot 36c^3\right)^{\frac{1}{3}} \end{aligned}$$

This proves the claim.

§5 CONTEST PROBLEMS

Inequalities have been a part of the mathematical contests for a long time. We have borrowed a lot of problems from these prominent contests. Here is a list of the contests (in no particular order):

- International Mathematical Olympiad (IMO)
- Asian Pacific Mathematical Olympiad (APMO)
- Belarusian National Olympiad (BrNO)
- Bulgarian National Olympiad (BuNO)
- China Mathematical Olympiad (ChMO)
- United States of America Mathematical Olympiad (USAMO)
- Singapore Mathematical Olympiad (SgMO)
- Balkan Mathematical Olympiad (BkMO)
- Abel's Mathematical Contest (Norwegian Mathematical Olympiad) (AMC)
- Austrian-Polish Competition (APC)
- British Mathematical Olympiad (BMO)
- Azerbaijan National Olympiad (AzNO)
- Taiwan National Olympiad (TNO)
- Romania National Olympiad (RNO)

- Albania Mathematical Olympiad (AlMO)
- Poland Mathematical Olympiad (PMO)
- Mediterranean Mathematical Olympiad (MMO)
- Albania-Balkan Mathematical Olympiad (ABMO)
- Brazil National Olympiad (BNO)
- Iranian Mathematical Olympiad (IrMO)

The contests appear in this chapter alphabetically. Problems within a region are ordered based on which year they appeared. In most countries, mathematical Olympiads are followed by team selection tests. However, for better categorization and abbreviation, we have not separated problems based on whether they appeared on a team selection test or national Olympiad. We simply put all the problems under the same region. See DUŠAN et al.¹ for a reference to the IMO problems (at least up to 2009). See DONGPHD and SUUGAKU² for a reference to the APMO problems, although we should warn the reader that the full book is not written in LATEX for some reason.

Recall that for a vector $\mathbf{a} = (a_1, \dots, a_n)$, the arithmetic, geometric and harmonic means are $\mathfrak{A}(\mathbf{a})$, $\mathfrak{G}(\mathbf{a})$, $\mathfrak{H}(\mathbf{a})$ and $\mathfrak{M}_r(\mathbf{a})$ is the power mean of order r . For another vector \mathbf{b} , $\langle \mathbf{a}, \mathbf{b} \rangle$ is the inner product of \mathbf{a} and \mathbf{b} . The L_p norm of \mathbf{a} is $\sqrt[p]{a_1^p + \dots + a_n^p}$.

§§5.1 ABMO

PROBLEM 5.1 (Team Selection Test 2014, problem 1). Prove that for an integer $n > 2$, the following inequality holds:

$$\frac{1}{n+1} \left(1 + \frac{1}{3} + \dots + \frac{1}{2n-1} \right) > \frac{1}{n} \left(\frac{1}{2} + \dots + \frac{1}{2n} \right)$$

Solution.

¹DJUKIĆ DUŠAN et al., *The IMO compendium: a collection of problems suggested for the International Mathematical Olympiads: 1959–2009*, Springer, 2011.

²DONGPHD and SUUGAKU, *APMO 1989–2009 Problems and Solutions*, vnmath.com, 2009.

PROBLEM 5.2 (Team Selection Test 2010, problem 4). Let a, b, c be the sides of a triangle and k be a real number. Prove that

$$a^3 + b^3 + c^3 < k(a + b + c)(ab + bc + ca)$$

holds for $k = 1$. Find the smallest value of k such that the inequality holds.

§§5.2 ALMO

PROBLEM 5.3 (Albania Team Selection Test 2013, problem 2). Let a, b, c, d be positive real numbers. Prove that $x = 3$ is the minimal value for which the following inequality holds:

$$a^x + b^x + c^x + d^x \geq \frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{d}$$

PROBLEM 5.4 (Albania Team Selection Test 2012, problem 1). Find the greatest value of the expression

$$\frac{1}{x^2 - 4x + 9} + \frac{1}{y^2 - 4y + 9} + \frac{1}{z^2 - 4z + 9}$$

where x, y, z are non-negative real numbers such that $x + y + z = 1$.

§§5.3 AMC

PROBLEM 5.5 (2014 Norwegian Mathematical Olympiad, problem 1). Let x, y be non-negative real numbers. Show that

$$x^2 + y^2 + 1 \leq \sqrt{(x^3 + y + 1)(y^3 + x + 1)}$$

PROBLEM 5.6 (2013 Norwegian Mathematical Olympiad, problem 1). Find all real numbers a such that

$$3x^2 + y^2 \geq -ax(x + y)$$

holds for all real numbers x, y .

PROBLEM 5.7 (Norwegian Mathematical Olympiad 2012, problem 4). Let x, y be positive real numbers. Show that

$$\left(1 + \frac{x}{y}\right)^3 + \left(1 + \frac{y}{x}\right)^3 \geq 16$$

PROBLEM 5.8 (Norwegian Mathematical Olympiad 2010, problem 2). Let x be a real number such that $0 < x < 1$. Show that

$$\frac{x^2}{1-x} + \frac{(1-x)^2}{x} \geq 1$$

PROBLEM 5.9 (Norwegian Mathematical Olympiad 2009, problem 4). Show that

$$\left(\frac{2010}{2009}\right)^{2009} > 2$$

PROBLEM 5.10 (Norwegian Mathematical Olympiad 2008, problem 3). (i) Let x, y be positive real numbers such that $x + y = 2$. Show that

$$\frac{1}{x} + \frac{1}{y} \leq \frac{1}{x^2} + \frac{1}{y^2}$$

(ii) Let x, y, z be real numbers such that $x + y + z = 2$. Show that

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} + \frac{4}{9} \leq \frac{1}{x^2} + \frac{1}{y^2} + \frac{1}{z^2}$$

PROBLEM 5.11 (Norwegian Mathematical Olympiad 2006, problem 2). (i) Let a, b be real non-negative numbers. Show that

$$a + b \geq \frac{a^2 + b^2}{2} + \sqrt{ab}$$

(ii) Let a, b be real numbers in the interval $[0, 3]$. Show that

$$\frac{a^2 + b^2}{2} + \sqrt{ab} \geq \frac{(a+b)^2}{2}$$

PROBLEM 5.12 (Norwegian Mathematical Olympiad 2005, problem 4). (a) Let a, b, c be real positive numbers. Show that

$$(a+b)(a+c) \geq 2\sqrt{abc(ab+bc+ca)}$$

(b) Let a, b, c be real numbers such that

$$ab + bc + ca > a + b + c > 0$$

PROBLEM 5.13 (Norwegian Mathematical Olympiad 2003, problem 2). Let a_1, \dots, a_n be n positive integers. Show that

$$\sum_{i=1}^n a_i^3 \geq \left(\sum_{i=1}^n a_i \right)^2$$

PROBLEM 5.14 (Norwegian Mathematical Olympiad 2002, problem 2). Let $n \geq 2$ be a positive integer and $x_1, \dots, x_n, y_1, \dots, y_n$ be positive real numbers such that

$$x_1 + \dots + x_n \geq x_1 y_1 + \dots + x_n y_n$$

Show that

$$x_1 + \dots + x_n \leq \frac{x_1}{y_1} + \dots + \frac{x_n}{y_n}$$

PROBLEM 5.15 (Norwegian Mathematical Olympiad 2000, problem 2). Let a, b, c, d be non-negative real numbers such that $a + b + c + d = 4$. Show that

$$\sqrt{a+b+c} + \sqrt{b+c+d} + \sqrt{c+d+a} + \sqrt{d+a+b} \geq 6$$

PROBLEM 5.16 (Norwegian Mathematical Olympiad 1999, problem 1). If a, b, c, d, e are real numbers, prove that

$$a^2 + b^2 + c^2 + d^2 + e^2 \geq a(b+c+d+e)$$

PROBLEM 5.17 (Norwegian Mathematical Olympiad 1995, problem 4). Let x_i, y_i be positive real numbers for $1 \leq i \leq n$. Prove that

$$\left(\sum_{i=1}^n (x_i + y_i)^2 \right) \left(\frac{1}{\sum_{i=1}^n x_i y_i} \right) \geq 4n^2$$

PROBLEM 5.18 (Norwegian Mathematical Olympiad 1994, problem 3). Let x_1, \dots, x_{1994} be positive real numbers. Prove that

$$\left(\frac{x_1}{x_2} \right)^{\frac{x_1}{x_2}} \cdots \left(\frac{x_{1993}}{x_{1994}} \right)^{\frac{x_{1994}}{x_{1993}}} \geq \left(\frac{x_1}{x_2} \right)^{\frac{x_2}{x_1}} \cdots \left(\frac{x_{1993}}{x_{1994}} \right)^{\frac{x_{1994}}{x_{1993}}}$$

PROBLEM 5.19 (Norwegian Mathematical Olympiad 1993, problem 1). 1. Let a, b, c be sides of a triangle. Prove that

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} < 2$$

PROBLEM 5.20 (Norwegian Mathematical Olympiad 1993, problem 2). Let a, b, c, d are real numbers such that $b < c < d$. Prove that

$$(a+b+c+d)^2 > 8(ac+bd)$$

§§5.4 APC

PROBLEM 5.21 (APC 2006, problem 5). Prove that for all positive integer n and positive real numbers a, b, c , the following inequality holds:

$$\frac{a^{n+1}}{a^n + a^{n-1}b + \dots + b^n} + \frac{b^{n+1}}{b^n + b^{n-1}c + \dots + c^{n-1}} + \frac{c^{n+1}}{c^n + c^{n-1} + \dots + a^n} \geq \frac{a+b+c}{n+1}$$

PROBLEM 5.22 (APC 2005, problem 3). Let a_0, \dots, a_n be real numbers such that

- (a) $0 = a_0 \leq \dots \leq a_n$
- (b) For $0 \leq i < j \leq n$, $a_j - a_i \leq j - i$

Prove that

$$\sum_{i=0}^n a_i^2 \geq \sum_{i=0}^n a_i^3$$

PROBLEM 5.23 (APC 2005, problem 10). Determine all pairs of non-negative integers (k, n) such that the following inequality holds for all positive real numbers x, y :

$$1 + \frac{y^n}{x^k} \geq \frac{(1+y)^n}{(1+x)^k}$$

PROBLEM 5.24 (APC 2003, problem 8). Given real numbers $x_1 \geq \dots \geq x_{2003} \geq 0$. Show that

$$x_1^n - x_2^n + \dots + x_{2003}^n \geq (x_1 - x_2 + \dots + x_{2003})^n$$

for any positive integer n .

PROBLEM 5.25 (APC 2001, problem 3). Let a, b, c be sides of a triangle. Prove that

$$2 \leq \frac{a+b}{c} + \frac{b+c}{a} + \frac{c+a}{b} - \frac{a^3 + b^3 + c^3}{abc} \leq 3$$

PROBLEM 5.26 (APC 2000, problem 9). Let a, b, c be non-negative real numbers such that $a + b + c = 1$. Prove that

$$2 \leq (1 - a^2)^2 + (1 - b^2)^2 + (1 - c^2)^2 \leq (1 + a)(1 + b)(1 + c)$$

PROBLEM 5.27 (APC 1999, problem 2). Find the best possible k, l such that

$$k < \frac{v}{v+w} + \frac{w}{w+x} + \frac{x}{x+y} + \frac{y}{y+z} + \frac{z}{z+x} < l$$

for all positive real numbers v, w, x, y, z .

PROBLEM 5.28 (APC 1998, problem 1). Let x_1, x_2, y_1, y_2 be real numbers such that $x_1^2 + x_2^2 \leq 1$. Prove that

$$(x_1 y_1 + x_2 y_2 - 1)^2 \geq (x_1^2 + x_2^2 - 1)(y_1^2 + y_2^2 - 1)$$

PROBLEM 5.29 (APC 1997, problem 7). Let p, q be arbitrary real numbers.

- (a) Prove that $p^2 + q^2 + 1 > p(q + 1)$.
- (b) Determine the largest possible b such that $p^2 + q^2 + 1 > bp(q + 1)$ for all p, q .
- (c) Determine the largest possible c such that $p^2 + q^2 + 1 > cp(q + 1)$ for all integer p, q .

PROBLEM 5.30 (APC 1995, problem 9). Prove that for all positive integers m, n and all real numbers x, y , the following inequality holds:

$$(n-1)(m-1)(x^{n+m} + y^{n+m}) + (n+m-1)(x^n y^m + x^m y^n) \geq nm(x^{n+m-1}y + y^{n+m-1}x)$$

PROBLEM 5.31 (APC 1993, problem 6). If a, b are non-negative real numbers, prove the inequality

$$\left(\frac{\sqrt{a} + \sqrt{b}}{2} \right)^2 \leq \frac{a + \sqrt[3]{a^2b} + \sqrt[3]{ab^2} + b}{4} \leq \frac{a + \sqrt{ab} + b}{3} \leq \sqrt{\left(\frac{a^{2/3} + b^{2/3}}{2} \right)^3}$$

§§5.5 APMO

PROBLEM 5.32 (2018, problem 2). Let $f(x)$ and $g(x)$ be given by

$$\begin{aligned} f(x) &= \frac{1}{x} + \frac{1}{x-2} + \dots + \frac{1}{x-2018} \\ g(x) &= \frac{1}{x-1} + \frac{1}{x-3} + \dots + \frac{1}{x-2017} \end{aligned}$$

Prove that

$$|f(x) - g(x)| > 2$$

PROBLEM 5.33 (2012, problem 5). Let $n \geq 2$ be an integer and a_1, \dots, a_n be real numbers such that $a_1^2 + \dots + a_n^2 = n$. Prove that

$$\sum_{1 \leq i < j \leq n} \frac{1}{n - a_i a_j} \leq \frac{n}{2}$$

PROBLEM 5.34 (2007, problem 4). Let x, y, z be positive real numbers such that

$$\sqrt{x} + \sqrt{y} + \sqrt{z} = 1$$

Prove that

$$\frac{x^2 + yz}{\sqrt{2x^2(y+z)}} + \frac{y^2 + zx}{\sqrt{2y^2(z+x)}} + \frac{z^2 + xy}{\sqrt{2z^2(x+y)}} \geq 1$$

PROBLEM 5.35 (2005, problem 2). Let a, b, c be positive real numbers such that $abc = 8$. Prove that

$$\frac{a^2}{\sqrt{(1+a^3)(1+b^3)}} + \frac{b^2}{\sqrt{(1+b^3)(1+c^3)}} + \frac{c^2}{\sqrt{(1+c^3)(1+a^3)}} \geq \frac{4}{3}$$

PROBLEM 5.36 (APMO 2004, problem 5). Prove that the inequality

$$(a^2 + 2)(b^2 + 2)(c^2 + 2) \geq 9(ab + bc + ca)$$

holds for all positive real numbers a, b, c .

PROBLEM 5.37 (APMO 2003, problem 4). Let a, b, c be the sides of a triangle with $a + b + c = 1$ and $n \geq 2$ be an integer. Show that

$$\sqrt[n]{a^n + b^n} + \sqrt[n]{b^n + c^n} + \sqrt[n]{c^n + a^n} < 1 + \frac{\sqrt[n]{2}}{2}$$

PROBLEM 5.38 (APMO 2002, problem 1). Let n be a positive integer and a_1, \dots, a_n be a sequence of non-negative integers. Let

$$A_n = \frac{a_1 + \dots + a_n}{n}$$

Show that

$$a_1! \cdots a_n! \geq (\lfloor A_n \rfloor!)^n$$

PROBLEM 5.39 (APMO 2002, problem 4). Let x, y, z be positive numbers such that

$$\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 1$$

Show that

$$\sqrt{x+yz} + \sqrt{y+zx} + \sqrt{z+xy} \geq \sqrt{xyz} + \sqrt{x} + \sqrt{y} + \sqrt{z}$$

PROBLEM 5.40 (APMO 1998, problem 3). Let a, b, c be positive real numbers. Prove that

$$\left(1 + \frac{a}{b}\right) \left(1 + \frac{b}{c}\right) \left(1 + \frac{c}{a}\right) \geq 2 \left(1 + \frac{a+b+c}{\sqrt[3]{abc}}\right)$$

PROBLEM 5.41 (APMO 1996, problem 5). Let a, b, c be the lengths of the sides of a triangle. Prove that

$$\sqrt{a+b-c} + \sqrt{b+c-a} + \sqrt{c+a-b} \leq \sqrt{a} + \sqrt{b} + \sqrt{c}$$

and determine when equality occurs.

PROBLEM 5.42 (APMO 1991, problem 3). Let $a_1, \dots, a_n, b_1, \dots, b_n$ be positive real numbers such that

$$a_1 + \dots + a_n = b_1 + \dots + b_n$$

Show that

$$\frac{a_1^2}{a_1 + b_1} + \dots + \frac{a_n^2}{a_n + b_n} \geq \frac{a_1 + \dots + a_n}{2}$$

Solution. Engel form of Cauchy-Schwarz can be used to solve this easily.

$$\begin{aligned} \frac{a_1^2}{a_1 + b_1} + \dots + \frac{a_n^2}{a_n + b_n} &\geq \frac{(a_1 + \dots + a_n)^2}{a_1 + b_1 + \dots + a_n + b_n} \\ &= \frac{(a_1 + \dots + a_n)^2}{2(a_1 + \dots + a_n)} \\ &= \frac{a_1 + \dots + a_n}{2} \end{aligned}$$

PROBLEM 5.43 (APMO 1990, problem 2). Let a_1, \dots, a_n be positive real numbers and S_k be the sum of the products of a_1, \dots, a_n taken k at a time. Show that

$$S_k S_{n-k} \geq \binom{n}{k}^2 a_1 \cdots a_n$$

PROBLEM 5.44 (APMO 1989, problem 1). Let x_1, \dots, x_n be positive real numbers and

$$S = x_1 + \dots + x_n$$

Prove that

$$(1 + x_1) \cdots (1 + x_n) \leq 1 + S + \frac{S^2}{2!} + \dots + \frac{S^n}{n!}$$

§§5.6 AzNO

PROBLEM 5.45 (2020 National Olympiad, problem 3). a, b, c are positive real numbers such that $a + b + c = 3$. Prove that

$$\sum \frac{a^2 + 6}{2a^2 + 2b^2 + 2c^2 + 2a - 1} \leq 3$$

PROBLEM 5.46 (2015 National Olympiad, problem 1). Let a, b, c be positive real numbers such that $abc = \frac{1}{8}$. Prove that

$$a^2 + b^2 + c^2 + a^2b^2 + b^2c^2 + c^2a^2 \geq \frac{15}{16}$$

PROBLEM 5.47 (2016 Team Selection Test, problem 1, day 3). Let a_1, a_2, \dots be a sequence of positive real numbers such that

$$a_{k+1} \geq \frac{ka_k}{a_k^2 + k - 1}$$

for every positive integer k . Prove that

$$a_1 + \dots + a_n \geq n$$

for every positive integer n .

PROBLEM 5.48 (2016 Balkan Mathematical Olympiad Team Selection Test 4, problem 1). Let a, b, c be non-negative real numbers. Prove that

$$3(a^2 + b^2 + c^2) \geq (a + b + c)(\sqrt{ab} + \sqrt{bc} + \sqrt{ca}) + (a - b)^2 + (b - c)^2 + (c - a)^2 \geq (a + b + c)^2$$

§§5.7 BkMO

PROBLEM 5.49 (2019, problem 2). Let a, b, c be real numbers such that $0 \leq a \leq b \leq c$ and $a + b + c = ab + bc + ca > 0$. Prove that

$$\sqrt{bc}(a + 1) \geq 2$$

Determine when equality occurs.

PROBLEM 5.50 (2015, problem 1). Let a, b, c be positive real numbers. Prove that

$$a^3b^6 + b^3c^6 + c^3a^6 + 3a^3b^3c^3 \geq abc(a^3b^3 + b^3c^3 + c^3a^3) + a^2b^2c^2(a^3 + b^3 + c^3)$$

PROBLEM 5.51 (2012, problem 2). Prove that

$$\sum_{cyc} (x + y) \sqrt{(z + x)(z + y)} \geq 4(xy + yz + zx)$$

for all positive real numbers x, y, z .

PROBLEM 5.52 (2011, problem 2). Given real numbers x, y, z such that $x + y + z = 0$. Prove that

$$\frac{x(x + 1)}{2x^2 + 1} + \frac{y(y + 2)}{2y^2 + 1} + \frac{z(z + 1)}{2z^2 + 1} \geq 0$$

PROBLEM 5.53 (2010, problem 1). Let a, b, c be positive real numbers. Prove that

$$\frac{a^2(b-c)}{a+b} + \frac{b^2(c-a)}{b+c} + \frac{c^2(a-b)}{c+a} \geq 0$$

PROBLEM 5.54 (2008, problem 2). Is there a sequence of positive real numbers a_1, a_2, \dots such that

$$\begin{aligned} a_1 + \dots + a_n &\leq n^2 \\ \frac{1}{a_1} + \dots + \frac{1}{a_n} &\leq 2008 \end{aligned}$$

for all positive integer n .

PROBLEM 5.55 (2006, problem 1). Let a, b, c be positive real numbers. Prove the inequality

$$\frac{1}{a(b+1)} + \frac{1}{b(c+1)} + \frac{1}{c(a+1)} \frac{3}{1+abc}$$

PROBLEM 5.56 (2005, problem 3). Let a, b, c be positive real numbers. Prove the inequality

$$\frac{a^2}{b} + \frac{b^2}{c} + \frac{c^2}{a} \geq a + b + c + \frac{4(a-b)^2}{a+b+c}$$

PROBLEM 5.57 (2001, problem 3). Let a, b, c be positive real numbers such that $a + b + c \leq abc$. Prove that

$$a^2 + b^2 + c^2 \geq \sqrt{3}abc$$

PROBLEM 5.58 (1998, problem 2). Let $n \geq 2$ be an integer and $a_0 < a_1 < \dots < a_{2n+1}$ be real numbers. Prove that

$$\sqrt[n]{a_1} - \sqrt[n]{a_2} + \dots + \sqrt[n]{a_{2n+1}} < \sqrt[n]{a_1 - a_2 + \dots + a_{2n+1}}$$

PROBLEM 5.59 (1993, problem 1). Let a, b, c, d, e, f be real numbers such that $a + b + c + d + e + f = 10$ and

$$(a-1)^2 + (b-1)^2 + (c-1)^2 + (d-1)^2 + (e-1)^2 + (f-1)^2 = 6$$

What is the maximum possible value of f ?

§§5.8 BMO

PROBLEM 5.60 (2005, Round 2, problem 3). Let a, b, c be positive real numbers. Prove that

$$\left(\frac{a}{b} + \frac{b}{c} + \frac{c}{a}\right)^2 \geq (a+b+c) \left(\frac{1}{a} + \frac{1}{b} + \frac{1}{c}\right)$$

PROBLEM 5.61 (2007, Round 1, problem 5). For positive real numbers a, b, c , prove that

$$(a^2 + b^2 + c^2)^2 \geq (a+b+c)(a+b-c)(b+c-a)(c+a-b)$$

PROBLEM 5.62 (2008, Round 1, problem 1). Find the minimum value of $x^2 + y^2 + z^2$ where x, y, z are real numbers such that $x^3 + y^3 + z^3 - 3xyz = 1$.

PROBLEM 5.63 (2010, Round 2, problem 5). For all positive real numbers x, y, z , prove that

$$4(x+y+z)^3 > 27(x^2y + y^2z + z^2x)$$

PROBLEM 5.64 (2011, Round 2, problem 6). Let a, b, c be the sides of a triangle such that $ab + bc + ca = 1$. Prove that

$$(a+1)(b+1)(c+1) < 4$$

PROBLEM 5.65 (2015, Round 1, problem 1). Place the following numbers in order and provide your reasoning to do so:

$$3^{3^4}, 3^{4^3}, 4^{3^3}, 4^{4^3}$$

Here, $a^{b^c} = a^{(b^c)}$ and not $(a^b)^c$ which is a^{bc} .

§§5.9 BNO

PROBLEM 5.66 (2011, problem 6). Let a_1, \dots, a_{2011} be non-negative real numbers such that

$$a_1 + \dots + a_{2011} = \frac{2011}{2}$$

Prove that

$$|(a_1 - a_2) \cdots (a_{2010} - a_{2011})(a_{2011} - a_1)| \leq \frac{3\sqrt{3}}{16}$$

PROBLEM 5.67 (2009, problem 3). Let $n > 3$ be a positive integer and x_1, \dots, x_n be positive real numbers. Find the value of

$$\frac{x_1}{x_n + x_1 + x_2} + \frac{x_2}{x_1 + x_2 + x_3} + \dots + \frac{x_n}{x_{n-1} + x_n + x_1}$$

in terms of n .

PROBLEM 5.68 (2008, problem 3). Let x, y, z be real numbers such that $xy + yz + zx = x + y + z$. Find the minimum value of

$$\frac{x}{x^2 + 1} + \frac{y}{y^2 + 1} + \frac{z}{z^2 + 1}$$

PROBLEM 5.69 (2005, problem 2). Determine the smallest possible C such that the inequality

$$C(x_1^{2005} + x_3^{2005} + x_4^{2005} + x_5^{2005}) \geq x_1 x_2 x_3 x_4 x_5 (x_1^{125} + x_2^{125} + x_3^{125} + x_4^{125} + x_5^{125})^{16}$$

PROBLEM 5.70 (2001, problem 1). Show that for any positive real numbers a, b, c ,

$$(a+b)(a+c) \geq 2\sqrt{abc(a+b+c)}$$

§§5.10 BrNO

PROBLEM 5.71 (2018, problem 9). Let $n \geq 2$ be an integer. Prove the inequality

$$\frac{1}{2!} + \dots + \frac{2^{n-2}}{n!} \leq \frac{3}{2}$$

PROBLEM 5.72 (2015, Day 2, problem 1). Find all real number $x \geq -1$ such that the inequality

$$\frac{a_1 + x}{2} \cdots \frac{a_n + x}{2} \leq \frac{a_1 \cdots a_n + x}{2}$$

holds for all positive integer $n \geq 2$ and for all real numbers $a_1, \dots, a_n \geq 1$.

PROBLEM 5.73 (2014 Final Round). Find all values λ such that the inequality

$$\frac{a+b}{2} \geq \lambda\sqrt{ab} + (1-\lambda)\sqrt{\frac{a^2+b^2}{2}}$$

PROBLEM 5.74 (2014 Final Round). Prove that for all positive real numbers x and y ,

$$\frac{1}{x+y+1} - \frac{1}{(x+1)(y+1)} < \frac{1}{11}$$

PROBLEM 5.75 (2014 Test 2). Let a, b, c be positive real numbers such that

$$ab + bc + ca \geq a + b + c$$

Prove that

$$(a+b+c)(ab+bc+ca) + 3abc \geq 4(ab+bc+ca)$$

PROBLEM 5.76 (2014 Test 4). Let a, b, c be positive real numbers such that $a+b+c=1$. Prove that

$$\frac{a^2}{(b+c)^3} + \frac{b^2}{(c+a)^3} + \frac{c^2}{(a+b)^3} \geq \frac{9}{8}$$

PROBLEM 5.77 (2014 Test 6). Let a, b, c be real numbers in the interval $(0, 2)$ such that $a+b+c = ab+bc+ca$. Prove that

$$\frac{a^2}{a^2-a+1} + \frac{b^2}{b^2-b+1} + \frac{c^2}{c^2-c+1} \leq 3$$

PROBLEM 5.78 (2013 Test 3). Let $n \geq 3$ be an integer and x_1, \dots, x_n be positive real numbers such that $x_1 \cdots x_n = 1$. Prove that

$$\frac{x_1^8}{(x_1^4+x_2^4)x_2} + \frac{x_n^8}{(x_n^4+x_1^4)x_1} \geq \frac{n}{2}$$

PROBLEM 5.79 (2013 Test 7). Given positive real numbers a, b, c such that

$$\frac{1}{ab} + \frac{1}{bc} + \frac{1}{ca} = 1$$

Prove that

$$\frac{a^2+b^2+c^2+ab+bc+ca-3}{5} \geq \frac{a}{b} + \frac{b}{c} + \frac{c}{a}$$

PROBLEM 5.80 (2012 Test 1). Let a, b, c be real numbers such that $0 < a < b < c$. Prove that

$$a^{20}b^{12} + b^{20}c^{12} + c^{20}a^{12} < b^{20}a^{12} + c^{20}b^{12} + b^{20}a^{12}$$

PROBLEM 5.81 (2011 Round 3). Let a, b, x, y be positive real numbers such that

$$ab \geq xa + yb$$

Prove that

$$\sqrt{a+b} \geq \sqrt{x} + \sqrt{y}$$

PROBLEM 5.82 (2011 Round 3). Let a, b, c, k, l and m be positive real numbers such that

$$abc \geq ka + lb + mc$$

Prove that

$$a + b + c \geq \sqrt{3}(\sqrt{k} + \sqrt{l} + \sqrt{m})$$

PROBLEM 5.83 (2011 Final Round). Let a, b, c be positive real numbers such that

$$a^2 + b^2 + c^2 = 3$$

Prove that

$$a + b + c \geq ab + bc + ca$$

PROBLEM 5.84 (2011 Test 5). Let a, b, c be positive real numbers such that

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b} = 1 + \frac{1}{6} \left(\frac{a}{c} + \frac{c}{b} + \frac{b}{a} \right)$$

Prove that

$$\frac{a^3bc}{b+c} + \frac{b^3ca}{c+a} + \frac{c^3ab}{a+b} \geq \frac{1}{6}(ab + bc + ca)^2$$

PROBLEM 5.85 (2010 Final Round). Prove that

$$\frac{a^2}{a+b} + \frac{b^2}{b+c} \geq \frac{3a+2b-c}{4}$$

for all positive real numbers a, b and c .

PROBLEM 5.86 (2010 Test 1). Given non-negative real numbers a, b and c such that $a + b + c = 1$. Prove that

$$(a^2 + b^2 + c^2)^2 + 6abc \geq ab + bc + ca$$

PROBLEM 5.87 (2010 Test 7). Prove that all positive real numbers x, y and z satisfy the inequality

$$x^y + y^z + z^x > 1$$

PROBLEM 5.88 (2010 Test 8). Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$\frac{a}{b(a+b)} + \frac{b}{c(b+c)} + \frac{c}{a(c+a)} \geq \frac{3}{2}$$

PROBLEM 5.89 (2009 Selection and Training Session). Let a, b and c be positive real numbers. Prove that

$$\frac{1}{(a+b)b} + \frac{1}{(b+c)c} + \frac{1}{(c+a)a} \geq \frac{9}{2(ab+bc+ca)}$$

PROBLEM 5.90 (2008, Day 2, problem 1). Let x_1, \dots, x_n be non-negative real numbers. Prove that

$$\frac{x_1(2x_1 - x_2 - x_3)}{x_2 + x_3} + \dots + \frac{x_n(2x_n - x_1 - x_2)}{x_1 + x_2} \geq 0$$

PROBLEM 5.91 (2002). For any positive integers a and b , prove that

$$|a\sqrt{2} - b| > \frac{1}{2(a+b)}$$

PROBLEM 5.92 (2002). Given positive real numbers a, b, c and d . Prove that

$$\sqrt{(a+c)^2 + (b+d)^2} \leq \sqrt{a^2 + b^2} + \sqrt{c^2 + d^2} \leq \sqrt{(a+c)^2 + (b+d)^2} + \frac{2|ad - bc|}{\sqrt{(a+c)^2 + (b+d)^2}}$$

§§5.11 BuNO

PROBLEM 5.93 (2020 Day 1, problem 2). Let b_1, \dots, b_n be non-negative integers and a_0, a_1, \dots, a_n be real numbers such that $b_1 + \dots + b_n = 2$ and $a_0 = a_n = 0$, $|a_i - a_{i-1}| \leq b_i$ for $1 \leq i \leq n$. Prove that

$$\sum_{i=1}^n (a_i + a_{i-1})b_i \leq 2$$

PROBLEM 5.94 (2018, problem 3). Prove that

$$\left(\frac{6}{5}\right)^{\sqrt{3}} > \left(\frac{5}{4}\right)^{\sqrt{2}}$$

PROBLEM 5.95 (2016, problem 3). For positive real numbers a, b, c and d , prove that

$$\frac{a + \sqrt{ab} + \sqrt[3]{abc} + \sqrt[4]{abcd}}{4} \leq \sqrt[4]{a \cdot \frac{a+b}{2} \cdot \frac{a+b+c}{3} \cdot \frac{a+b+c+d}{4}}$$

PROBLEM 5.96 (2009, problem 6). Let $a_1, \dots, a_n, b_1, \dots, b_n$ be arbitrarily taken real numbers and c_1, \dots, c_n be positive real numbers, then

$$\left(\sum_{i,j=1}^n \frac{a_i a_j}{c_i + c_j} \right) \left(\sum_{i,j=1}^n \frac{b_i b_j}{c_i + c_j} \right) \geq \left(\sum_{i,j=1}^n \frac{a_i b_j}{c_i + c_j} \right)^2$$

PROBLEM 5.97 (2008, problem 3). Let n be a natural number and $a_1, \dots, a_n, b_1, \dots, b_n$ be real positive numbers such that $0 \leq a_1 \leq \dots \leq a_n \leq \pi$ and

$$\left| \sum_{i=1}^n b_i \cos k a_i \right| < \frac{1}{k}$$

for all positive integer k . Prove that $b_1 = \dots = b_n = 0$.

PROBLEM 5.98 (2007 Team Selection Test, problem 3). Let $n \geq 2$ be a positive integer. Find the best constant $C(n)$ such that

$$\sum_{i=1}^n x_i \geq C(n) \sum_{1 \leq j < i \leq n} (2x_i x_j + \sqrt{x_i x_j})$$

is true for all $x_i \in (0, 1)$ such that

$$(1 - x_i)(1 - x_j) \geq \frac{1}{4}$$

for $1 \leq i < j \leq n$.

PROBLEM 5.99 (1997, problem 1). Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$\frac{1}{1+b+c} + \frac{1}{1+c+b} + \frac{1}{1+a+b} \leq \frac{1}{2+a} + \frac{1}{2+b} + \frac{1}{2+c}$$

§§5.12 ChMO

PROBLEM 5.100 (2019, problem 1). Let $a, b, c, d, e \geq -1$ be real numbers such that $a + b + c + d + e = 5$. Find the minimum and maximum value of

$$(a+b)(b+c)(c+d)(d+e)(e+a)$$

PROBLEM 5.101 (2018, problem 6). Let n, k be natural numbers such that $n > k$ and a_1, \dots, a_n be real numbers in the interval $(k-1, k)$. Let x_1, \dots, x_n be positive real numbers such that for any subset I of $\{1, 2, \dots, n\}$ with k elements,

$$\sum_{i \in I} x_i \leq \sum_{i \in I} a_i$$

Find the maximum possible value of $x_1 \cdots x_n$.

PROBLEM 5.102 (2017, problem 6). Given an integer $n \geq 2$ and real numbers a, b such that $0 < a < b$. Let x_1, \dots, x_n be real numbers in the interval $[a, b]$. Find the maximum value of

$$\frac{\frac{x_1^2}{x_2} + \dots + \frac{x_{n-1}^2}{x_n} + \frac{x_n^2}{x_1}}{x_1 + \dots + x_n}$$

PROBLEM 5.103 (2015, problem 1). Let z_1, \dots, z_n be complex numbers such that $|z_i - 1| \leq r$ for some real number $r \in (0, 1)$. Show that

$$\left| \sum_{i=1}^n z_i \right| \cdot \left| \sum_{i=1}^n \frac{1}{z_i} \right| \geq n^2(1-r^2)$$

PROBLEM 5.104 (2011, problem 1). Let a_1, \dots, a_n be real numbers. Prove that

$$\sum_{i=1}^n a_i^2 - \sum_{i=1}^n a_i a_{i+1} \leq \left\lfloor \frac{n}{2} \right\rfloor (M - m)$$

where $a_{n+1} = a_1$, $M = \max\{a_1, \dots, a_n\}$, $m = \min\{a_1, \dots, a_n\}$.

PROBLEM 5.105 (2011, problem 5). Let $n \geq 4$ be an integer and $a_1, \dots, a_n, b_1, \dots, b_n$ be non-negative real numbers such that

$$a_1 + \dots + a_n = b_1 + \dots + b_n > 0$$

Find the maximum of

$$\frac{\sum_{i=1}^n a_i(a_i + b_i)}{\sum_{i=1}^n b_i(a_i + b_i)}$$

PROBLEM 5.106 (2009, problem 4). Let $n > 3$ be an integer and a_1, \dots, a_n be real numbers satisfying $\min\{a_i - a_j\} \leq 1$ for $1 \leq i < j \leq n$. Find the minimum value of

$$\sum_{i=1}^n a_i^3$$

PROBLEM 5.107 (2008, problem 3). Let n be a positive integer and $x_1, \dots, x_n, y_1, \dots, y_n$ be real numbers such that

$$\begin{aligned} x_1 &\leq x_2 \dots \leq x_n \\ y_1 &\geq y_2 \dots \geq y_n \\ \sum_{i=1}^n ix_i &= \sum_{i=1}^n iy_i \end{aligned}$$

Show that for any real number α ,

$$\sum_{i=1}^n x_i \lfloor i\alpha \rfloor \geq \sum_{i=1}^n y_i \lfloor i\alpha \rfloor$$

PROBLEM 5.108 (2007, problem 1). Let a, b, c be complex numbers and $|a + b| = m, |a - b| = n$. If $mn \neq 0$, show that

$$\max\{|ac + b|, |a + bc|\} \geq \frac{mn}{\sqrt{m^2 + n^2}}$$

PROBLEM 5.109 (2006, problem 1). Let a_1, \dots, a_k be real numbers such that $a_1 + \dots + a_k = 0$. Prove that

$$\max\{a_1^2, \dots, a_k^2\} \leq \frac{k}{3} ((a_1 - a_2)^2 + \dots + (a_{k-1} - a_k)^2)$$

PROBLEM 5.110 (2006, problem 5). Let (a_n) be a sequence such that $a_1 = \frac{1}{2}, a_{k+1} = -a_k + \frac{1}{2-a_k}$. Prove that

$$\left(\frac{n}{2(a_1 + \dots + a_n)} - 1 \right)^n \leq \left(\frac{a_1 + \dots + a_n}{n} \right)^n \left(\frac{1}{a_1} - 1 \right) \dots \left(\frac{1}{a_n} - 1 \right)$$

PROBLEM 5.111 (2005, problem 1). Let $\vartheta_i \in (-\frac{\pi}{2}, \frac{\pi}{2})$ for $1 \leq i \leq 4$. Prove that, there exists a real number x such that

$$\begin{aligned}\cos^2 \vartheta_1 \cos^2 \vartheta_2 - (\sin \vartheta_1 \sin \vartheta_2 - x)^2 &\geq 0 \\ \cos^2 \vartheta_3 \cos^2 \vartheta_4 - (\sin \vartheta_3 \sin \vartheta_4 - x)^2 &\geq 0\end{aligned}$$

if and only if

$$\sum_{i=1}^4 \sin^2 \vartheta_i \leq 2 \left(1 + \prod_{i=1}^4 \sin \vartheta_i + \prod_{i=1}^4 \cos \vartheta_i \right)$$

PROBLEM 5.112 (2004, problem 2). Let $n \geq 2$ be an integer and a_1, \dots, a_n are positive integers such that $a_1 < \dots < a_n$ and

$$\frac{1}{a_1} + \dots + \frac{1}{a_n} \leq 1$$

Prove that for any real number x ,

$$\left(\sum_{i=1}^n \frac{1}{a_i^2 + x^2} \right)^2 \leq \frac{1}{2} \frac{1}{a_1(a_1 - 1) + x^2}$$

PROBLEM 5.113 (2003, problem 3). Let n be a positive integer. Find the smallest positive real number λ such that for any $x_1, \dots, x_n \in (0, \frac{\pi}{2})$,

$$\prod_{i=1}^n \tan x_i = 2^{\frac{n}{2}}$$

implies

$$\sum_{i=1}^n \cos x_i \leq \lambda$$

PROBLEM 5.114 (2003, problem 3). Let a, b, c, d be positive real numbers such that $ab + cd = 1$ and $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4$ are real numbers such that

$$x_i^2 + y_i^2 = 1$$

for $1 \leq i \leq 4$. Prove that

$$(ax_1 + bx_2 + cx_3 + dx_4)^2 + (ay_4 + by_3 + cy_2 + dy_1)^2 \leq 2 \left(\frac{a^2 + b^2}{ab} + \frac{c^2 + d^2}{cd} \right)$$

PROBLEM 5.115 (2002, problem 1). For every four points P_1, P_2, P_3, P_4 on the plane, find the minimum value of

$$\frac{\sum_{1 \leq i < j \leq 4} P_i P_j}{\min_{1 \leq i < j \leq 4} P_i P_j}$$

PROBLEM 5.116 (2002, problem 3). Let c be a real number such that $c \in (\frac{1}{2}, 1)$. Find the least real number M such that for every integer $n \geq 2$ and real numbers $0 \leq a_1 \leq \dots \leq a_n$, if

$$\frac{1}{n} \sum_{i=1}^n i a_i = c \sum_{i=1}^n a_i$$

then we always have that

$$\sum_{i=1}^n a_i \leq M \sum_{i=1}^m a_i$$

where $m = \lfloor cn \rfloor$.

PROBLEM 5.117 (1999, problem 5). Determine the maximum value of λ such that

$$f(x) \geq \lambda(x - a)^3$$

for all non-negative real number x where

$$f(x) = x^3 + ax^2 + bx + c$$

and has non-negative roots. Find the equality condition.

PROBLEM 5.118 (1998, problem 2). Given a positive integer $n > 1$. Determine with proof if there exists $2n$ distinct positive integers $a_1, \dots, a_n, b_1, \dots, b_n$ such that

$$\begin{aligned} a_1 + \dots + a_n &= b_1 + \dots + b_n \\ n - 1 &> \sum_{i=1}^n \frac{a_i - b_i}{a_i + b_i} > n - 1 - \frac{1}{1998} \end{aligned}$$

PROBLEM 5.119 (1998, problem 6). Let $n \geq 2$ be a positive integer and x_1, \dots, x_n be real numbers such that

$$\sum_{i=1}^n x_i^2 + \prod_{i=1}^{n-1} x_i x_{i+1} = 1$$

For each k , find the maximum value of $|x_k|$.

PROBLEM 5.120 (1997, problem 1). Let x_1, \dots, x_{1997} be real numbers satisfying

$$\begin{aligned} -\frac{1}{\sqrt{3}} &\leq x_i \leq \frac{1}{\sqrt{3}} \\ x_1 + \dots + x_{1997} &= -318\sqrt{3} \end{aligned}$$

Determine the maximum value of

$$x_1^{12} + \dots + x_{1997}^{12}$$

with proof.

PROBLEM 5.121 (1997, problem 6). Let (a_n) be a sequence of real numbers satisfying

$$a_{n+m} \leq a_n + a_m$$

for all non-negative integers m, n . Prove that, if $n \geq m$,

$$a_n \leq ma_1 + \left(\frac{n}{m} - 1\right)a_m$$

PROBLEM 5.122 (1996, problem 2). Let n be a natural number. Suppose that $x_0 = 0$ and $x_i > 0$ for all $i \in \{1, \dots, n\}$. If $\sum_{i=1}^n x_i = 1$, prove that

$$1 \leq \sum_{i=1}^n \frac{x_i}{\sqrt{1+x_0+\dots+x_{i-1}}\sqrt{x_i+\dots+x_n}} < \frac{\pi}{2}$$

PROBLEM 5.123 (1995, problem 1). Let $n \geq 3$ be an integer and $a_1, \dots, a_n, b_1, \dots, b_n$ be real numbers such that

$$a_1 + \dots + a_n = b_1 + \dots + b_n$$

$$0 < a_1 = a_2, a_i + a_{i+1} = a_{i+2}$$

$$0 < b_1 \leq b_2, b_i + b_{i+1} \leq b_{i+2}$$

Prove that $a_{n-1} + a_n \leq b_{n-1} + b_n$.

PROBLEM 5.124 (1995, problem 5). Let a_1, \dots, a_{10} distinct natural numbers such that $a_1 + \dots + a_{10} = 1995$. Find the minimum value of

$$a_1a_2 + a_2a_3 + \dots + a_9a_{10} + a_{10}a_1$$

PROBLEM 5.125 (1993, problem 2). Given a positive integer k and a positive real number a . Find the maximum value of

$$a^{k_1} + \dots + a^{k_r}$$

where $1 \leq r \leq k$ and $k_1 + \dots + k_r = k$.

PROBLEM 5.126 (2019 Team Selection Test, problem 3). Let n be a positive integer and a_1, \dots, a_n be non-negative real numbers such that

$$a_1 + \dots + a_n = 1$$

Find the maximum possible value of

$$\sum_{1 \leq i < j \leq n} \min\{(i-j)^2, (n+i-j)^2 a_i a_j\}$$

PROBLEM 5.127 (2019 Team Selection Test, problem 7). Let x, y, z be complex numbers such that

$$|x|^2 + |y|^2 + |z|^2 = 1$$

Prove that

$$|x^3 + y^3 + z^3 - 3xyz| \leq 1$$

PROBLEM 5.128 (2019 Team Selection Test, problem 5). Find all positive integer n such that for any positive real numbers a, b, c, x, y, z the following conditions hold:

$$\begin{aligned} \max\{a, b, c, x, y, z\} &= a \\ a + b + c &= x + y + z \\ abc &= xyz \\ a^n + b^n + c^n &= x^n + y^n + z^n \end{aligned}$$

PROBLEM 5.129 (2018 Team Selection Test, problem 5). Let n, k be positive integers such that $n > 4k$. Find the minimum value of $\lambda(n, k) = \lambda$ such that for any positive real numbers a_1, \dots, a_n , we have

$$\sum_{i=1}^n \frac{a_i}{\sqrt{a_i^2 + a_{i+1}^2 + \dots + a_{i+k}^2}}$$

where $a_{n+i} = a_i$ for $1 \leq i \leq k$.

PROBLEM 5.130 (2018 Team Selection Test, problem 3). Let $H(n)$ be the harmonic sum

$$1 + \frac{1}{2} + \dots + \frac{1}{n}$$

Prove that there exists a positive constant C such that

$$H(a_1) + \dots + H(a_n) \leq C \sqrt{\sum_{i=1}^n i a_i}$$

for arbitrary positive integer n and positive real numbers a_1, \dots, a_n .

PROBLEM 5.131 (2017 Team Selection Test, problem 2). Let n be positive integers and $x > 1$ be a real number. Prove that

$$\sum_{i=1}^n \frac{\{ix\}}{\lfloor ix \rfloor} < \sum_{i=1}^n \frac{1}{2i-1}$$

where $\{a\}$ and $\lfloor a \rfloor$ is the *decimal* and *integer* portion of a respectively.

PROBLEM 5.132 (2017 Team Selection Test, problem 5). Let $m \geq 2$ be a positive integer and a_1, \dots, a_m be positive real numbers. Prove that

$$(m-1)^{m-1} (x_1^m + \dots + x_m^m) \geq (x_1 + \dots + x_m)^m - m^m x_1 \cdots x_m$$

Find out when equality holds.

PROBLEM 5.133 (2016 Team Selection Test, problem 2). Find the smallest positive real number λ such that for any complex numbers z_1, z_2, z_3 with $|z_i| < 1$ and $z_1 + z_2 + z_3 = 0$,

$$|z_1 z_2 + z_2 z_3 + z_3 z_1|^2 + |z_1 z_2 z_3|^2 < \lambda$$

PROBLEM 5.134 (2016 Team Selection Test, problem 7). Let $n > 1$ be an integer and α be a real number such that $0 < \alpha < 2$ and $a_1, \dots, a_n, c_1, \dots, c_n$ be positive real numbers. For a positive real number y , let

$$f(y) = \left(\sum_{a_i < y} c_i a_i^2 \right)^{\frac{1}{2}} + \left(\sum_{a_i > y} c_i a_i^\alpha \right)^{\frac{1}{\alpha}}$$

If a positive real number x satisfies $x \geq f(y)$ for some positive real number y , prove that $f(x) \leq 8^{\frac{1}{\alpha}} x$.

PROBLEM 5.135 (2015 Team Selection Test, problem 4). Let $n \geq 2$ be an integer and x_1, x_2, \dots , be a non-decreasing monotonous sequence of positive real numbers such that $x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots$ is a non-increasing monotonous sequence. Prove that

$$\frac{\sum_{i=1}^n x_i}{n \sqrt[n]{\sum_{i=1}^n x_i}} \leq \frac{n+1}{2 \sqrt[n]{n!}}$$

PROBLEM 5.136 (2014 Team Selection Test, problem 4). Let (x_n) be a sequence of real numbers and (y_n) be a sequence such that $y_1 = x_1$ and for $n \geq 1$,

$$y_{n+1} = x_{n+1} - \sqrt{\sum_{i=1}^n x_i^2}$$

Find the smallest positive real number λ such that for any (x_n) and positive integer m ,

$$\frac{1}{m} \sum_{i=1}^n x_i^2 \leq \sum_{i=1}^n \lambda^{n-i} y_i^2$$

PROBLEM 5.137 (2014 Team Selection Test, problem 5). Let $n \geq 2$ be a positive integer. Find the greatest constant $\lambda(n)$ such that for any non-zero complex numbers z_1, \dots, z_n ,

$$\sum_{i=1}^n |z_i|^2 \geq \lambda(n) \min_{1 \leq i \leq n} |z_{i+1} - z_i|^2$$

where $z_{n+1} = z_1$.

PROBLEM 5.138 (2013 Team Selection Test, problem 4). Let $n, k > 1$ be integers and a_1, \dots, a_n be non-negative real numbers such that $a_1 \geq \dots \geq a_n$ and

$$\begin{aligned} a_1 + \dots + a_n &= 1 \\ c_1 + \dots + c_m &\leq m^k \end{aligned}$$

for any positive integer $m \leq n$ and non-negative real numbers c_1, \dots, c_m . Find the maximum value of

$$c_1 a_1^k + \dots + c_m a_m^k$$

PROBLEM 5.139 (2013 Team Selection Test, problem 17). Let $n \geq 2$ be an integer and $a_1, \dots, a_n, b_1, \dots, b_n$ be non-negative real numbers. Prove that

$$\left(\frac{n}{n-1} \right)^{n-1} \left(\frac{1}{n} \sum_{i=1}^n a_i^2 \right) + \left(\frac{1}{n} \sum_{i=1}^n b_i^2 \right) \geq \prod_{i=1}^n \sqrt[n]{a_i^2 + b_i^2}$$

PROBLEM 5.140 (2012 Team Selection Test, problem 1). Let n be a positive integer and $x_1, \dots, x_n, y_1, \dots, y_n$ be complex numbers such that $|x_i| = |y_i|$ for $1 \leq i \leq n$. Let

$$\begin{aligned} x &= \frac{1}{n} \sum_{i=1}^n x_i \\ y &= \frac{1}{n} \sum_{i=1}^n y_i \\ z_i &= xy_i - yx_i - x_i y_i \end{aligned}$$

Prove that

$$\sum_{i=1}^n |z_i| \leq n$$

PROBLEM 5.141 (2012 Team Selection Test, problem 4). Let $m, n > 1$ be integers and r, s are real numbers such that $r < s$. Let (a_{ij}) be a $m \times n$ non-zero matrix such that $a_{ij} \geq 0$. Find the maximum value of

$$\frac{\left(\sum_{j=1}^n \left(\sum_{i=1}^m a_{ij}^s\right)^{\frac{r}{s}}\right)^{\frac{1}{r}}}{\left(\sum_{i=1}^m \left(\sum_{j=1}^n a_{ij}^r\right)^{\frac{s}{r}}\right)^{\frac{1}{s}}}$$

PROBLEM 5.142 (2011 Team Selection Test, problem 6). Let n be a positive integer. Find the largest real number λ such that for all positive real numbers x_1, \dots, x_{2n} satisfying

$$\frac{1}{2n} \sum_{i=1}^{2n} (x_i + 2)^n \geq \prod_{i=1}^{2n} x_i$$

the following inequality is also true:

$$\frac{1}{2n} \sum_{i=1}^{2n} (x_i + 1)^n \geq \lambda \prod_{i=1}^{2n} x_i$$

PROBLEM 5.143 (2011 Team Selection Test, problem 7). Let $n \geq 3$ be an integer. Find the largest real number M such that for any positive real numbers x_1, \dots, x_n there is an arrangement y_1, \dots, y_n such that

$$\sum_{i=1}^n \frac{y_i^2}{y_{i+1}^2 - y_{i+1}y_{i+2} + y_{i+2}^2} \geq M$$

where $y_{n+2} = y_2, y_{n+1} = y_1$.

PROBLEM 5.144 (2010 Team Selection Test, problem 1). Let n be a positive integer. The real numbers $a_0, \dots, a_n, b_0, \dots, b_n$ satisfy $a_i + a_{i+1} \geq 0$ for $1 \leq i \leq 2n - 1$ and $a_{2i+1} \leq 0$ for $1 \leq i \leq n - 1$. For any integers p, q such that $0 \leq p \leq q \leq n$, we have

$$\sum_{i=2p}^{2q} b_i > 0$$

Prove that

$$\sum_{i=0}^{2n} (-1)^i a_i b_i \geq 0$$

Determine when equality holds.

PROBLEM 5.145 (2010 Team Selection Test, problem 5). Find all positive real numbers λ such that for all integer $n \geq 2$ and all positive real numbers a_1, \dots, a_n such that $a_1 + \dots + a_n = n$,

$$\sum_{i=1}^n \frac{1}{a_i} - \lambda \prod_{i=1}^n \frac{1}{a_i} \leq n - \lambda$$

PROBLEM 5.146 (2010 Team Selection Test, problem 7). Let $n \geq 2$ be an integer and a be a positive real number. Find the smallest positive real number $M(n, a) = M$ such that

$$\sum_{i=1}^n \frac{1}{a + S - x_i} \geq M$$

where $S = \sum_{i=1}^n x_i$ for any positive real numbers x_1, \dots, x_n .

PROBLEM 5.147 (2010 Team Selection Test 2). Let $n \geq 2$ be an integer and x_1, \dots, x_n be real numbers in the interval $[0, 1]$. Prove that there exists real numbers a_0, \dots, a_n such that

$$\begin{aligned} a_0 + a_n &= 0 \\ |a_i| &\leq 1 \\ |a_i - a_{i-1}| &= x_i \end{aligned}$$

PROBLEM 5.148 (2009 Team Selection Test 2). Let $n \geq 2$ be an integer. Find the maximum constant $\lambda(n)$ so that if a sequence of real numbers a_0, a_1, \dots satisfies $0 = a_0 \leq a_1 \leq \dots \leq a_n$ and for $1 \leq i \leq n-1$, $2a_i \geq a_{i-1} + a_{i+1}$ then

$$\left(\sum_{i=1}^n i a_i \right)^2 \geq \lambda(n) \sum_{i=1}^n a_i^2$$

PROBLEM 5.149 (2009 Team Selection Test, problem 5). Let $m > 1$ be an integer and n be an odd integer such that $3 \leq n < 2m$. Consider a matrix (a_{ij}) such that for any $1 \leq j \leq n$, $a_{1j}, a_{2j}, \dots, a_{mj}$ is a permutation of $1, 2, \dots, m$ and for any $1 \leq i \leq m$ and $1 \leq j \leq n-1$, $|a_{ij} - a_{i(j+1)}| \leq 1$. Find the minimum value of

$$\max_{1 \leq i \leq m} \sum_{j=1}^n a_{ij}$$

PROBLEM 5.150 (2009 Quiz 1, problem 3). Let m, n be positive integers and $x_1, \dots, x_m, y_1, \dots, y_n$ be positive real numbers. Prove that

$$2XY \sum_{i=1}^m \sum_{j=1}^n |x_i - y_j| \geq X^2 \sum_{i=1}^n \sum_{j=1}^n |y_i - y_j| + Y^2 \sum_{i=1}^m \sum_{j=1}^m |x_i - x_j|$$

PROBLEM 5.151 (2009 Quiz 5, problem 3). Let a_1, a_2, a_3, a_4 be non-negative real numbers such that $a_1 + a_2 + a_3 + a_4 = 1$. Prove that

$$\max \left\{ \sum_{i=1}^4 \sqrt{a_i^2 + a_i a_{i-1} + a_{i-1}^2 + a_{i-1} a_{i-2}}, \sum_{i=1}^4 \sqrt{a_i^2 + a_i a_{i+1} + a_{i+1} a_{i+2}} \right\} \geq 2$$

where $a_{i+4} = a_i$.

PROBLEM 5.152 (2008 Team Selection Test, problem 5). Let $m, n > 1$ be integers and (a_{ij}) be a non-zero matrix of non-negative real numbers. Find the minimum and maximum value of

$$\frac{m \sum_{i=1}^m \left(\sum_{j=1}^n a_{ij} \right)^2 + n \sum_{j=1}^n \left(\sum_{i=1}^m a_{ij} \right)^2}{\left(\sum_{i=1}^m \sum_{j=1}^n a_{ij} \right)^2 + mn \sum_{i=1}^m \sum_{j=1}^n a_{ij}^2}$$

PROBLEM 5.153 (2008 Team Selection Test, problem 6). Find the maximum constant M such that for any integer $n \geq 3$, there exists two sequences of positive real numbers a_1, \dots, a_n and b_1, \dots, b_n satisfying

$$\begin{aligned} \sum_{i=1}^n b_i &= 1 \\ 2b_i &\geq b_{i-1} + b_{i+1} \\ a_k^2 &\geq 1 + \sum_{i=1}^k a_i b_i \end{aligned}$$

and $a_n = M$.

PROBLEM 5.154 (2008 Quiz 2 problem 3). Let z_1, z_2, \dots, z_3 be complex numbers such that $|z_i| \leq 1$ and ω_1, ω_2 are the roots of the equation

$$(z - z_1)(z - z_2) + (z - z_2)(z - z_3) + (z - z_3)(z - z_1) = 0$$

Prove that

$$\min\{|z_j - \omega_1|, |z_j - \omega_2|\} \leq 1$$

PROBLEM 5.155 (2008 Quiz 4 problem 2). Let x, y, z be positive real numbers. Prove that

$$\frac{xy}{z} + \frac{yz}{x} + \frac{zx}{y} > 2\sqrt[3]{x^3 + y^3 + z^3}$$

PROBLEM 5.156 (2008 Quiz 5, problem 2). Let $n \geq 2$ be an integer and a_1, \dots, a_n be real numbers not all zero. Determine the necessary and sufficient condition so that there exists a sequence of integers x_1, \dots, x_n which satisfies

$$\begin{aligned} 0 &< x_1 < \dots < x_n \\ a_1 x_1 + \dots + a_n x_n &\geq 0 \end{aligned}$$

PROBLEM 5.157 (2008 Quiz 5, problem 3). Let n be a positive integer and $x_1, \dots, x_n, y_1, \dots, y_n$ be real numbers such that $0 < x_1 \leq \frac{x_2}{2} \leq \dots \leq \frac{x_n}{n}$ and $0 < y_n \leq y_{n-1} \leq \dots \leq y_1$. Prove that

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \left(\sum_{i=1}^n y_i \right) \left(\sum_{i=1}^n \left(x_i^2 - \frac{1}{4} x_i x_{i+1} \right) y_i \right)$$

where $x_0 = 0$.

PROBLEM 5.158 (2007 Team Selection Test, problem 5). Let $n > 1$ be a positive integer and x_1, \dots, x_n be real numbers satisfying $A = |\sum_{i=1}^n x_i| \neq 0$ and $B = \max_{1 \leq i \leq n} |x_i - x_j| \neq 0$. Prove that for any n vectors $\vec{\alpha}_i$ in the plane, there exists a permutation k_1, \dots, k_n of the numbers $1, \dots, n$ such that

$$\left| \sum_{i=1}^n a_i x_{k_i} \vec{\alpha}_i \right| \geq \frac{AB}{2A+B} \max_{1 \leq i \leq n} |\alpha_i|$$

PROBLEM 5.159 (2007 Quiz 2, problem 1). Let u, v, w be positive real numbers such that

$$u + v + w + \sqrt[3]{uvw} = 4$$

Prove that

$$\sqrt{\frac{uv}{w}} + \sqrt{\frac{vw}{u}} + \sqrt{\frac{wu}{v}} \geq u + v + w$$

PROBLEM 5.160 (2007 Quiz 4, problem 1). Let a_1, \dots, a_n be positive real numbers satisfying $a_1 + \dots + a_n = 1$. Prove that

$$(a_1 a_2 + \dots + a_n a_1) \left(\frac{a_1}{a_2^2 + a_2} + \dots + \frac{a_n^2}{a_1^2 + a_1} \right) \geq \frac{n}{n+1}$$

PROBLEM 5.161 (2007 Quiz 5, problem 3). Find the smallest constant k such that

$$\frac{x}{\sqrt{x+y}} + \frac{y}{\sqrt{y+z}} + \frac{z}{\sqrt{z+x}} \leq k \sqrt{x+y+z}$$

PROBLEM 5.162 (2006 Team Selection Test, problem 3). Let n be a positive integer and a_1, \dots, a_n be real numbers. Prove that there exists real numbers b_1, \dots, b_n such that $a_i - b_i$ is a positive integer for $1 \leq i \leq n$ and

$$\sum_{1 \leq i < j \leq n} (b_i - b_j)^2 \leq \frac{n^2 - 1}{12}$$

PROBLEM 5.163 (2006 Team Selection Test, problem 8). Let n be a positive integer and x_1, \dots, x_n be positive real numbers such that $x_1 + \dots + x_n = 1$. Prove that

$$\left(\sum_{i=1}^n \sqrt{x_i} \right) \left(\sum_{i=1}^n \frac{1}{\sqrt{1+x_i}} \right) \leq \frac{n^2}{n+1}$$

PROBLEM 5.164 (2006 Team Selection Test, problem 11). Given positive real numbers x, y, z such that $x + y + z = 1$. Prove that

$$\frac{xy}{\sqrt{xy+yz}} + \frac{yz}{\sqrt{yz+zx}} + \frac{zx}{\sqrt{zx+xy}} \leq \frac{\sqrt{2}}{2}$$

PROBLEM 5.165 (2005 Team Selection Test, problem 4). Let $a_1, \dots, a_6; b_1, \dots, b_6; c_1, \dots, c_6$ be permutations of $1, \dots, 6$. Find the minimum value of $\sum_{i=1}^6 a_i b_i c_i$.

PROBLEM 5.166 (2005 Quiz 1, problem 2). Let a, b, c be non-negative real numbers such that $ab + bc + ca = \frac{1}{3}$. Prove that

$$\frac{1}{a^2 - bc + 1} + \frac{1}{b^2 - ca + 1} + \frac{1}{c^2 - ab + 1} \leq 3$$

PROBLEM 5.167 (2005 Quiz 2, problem 3). Let a, b, c, d be positive real numbers such that $abcd = 1$. Prove that

$$\frac{1}{(1+a)^2} + \frac{1}{(1+b)^2} + \frac{1}{(1+c)^2} + \frac{1}{(1+d)^2} \geq 1$$

PROBLEM 5.168 (2004 Quiz 4, problem 2). Find the greatest positive real number k such that for any positive real numbers a, b, c, d ,

$$(a + b + c)(3^4(a + b + c + d)^5 + 2^4(a + b + c + 2d)^5) \geq kabcd^3$$

PROBLEM 5.169 (2003 Quiz 1, problem 1). x, y, z are positive real numbers such that $x + y + z = xyz$. Find the minimum value of

$$x^7(yz - 1) + y^7(zx - 1) + z^7(xy - 1)$$

PROBLEM 5.170 (2003 Quiz 2, problem 3). Let n be a positive integer and the roots of

$$f(z) = z^n + a_1 z^{n-1} + \dots + a_n$$

are z_1, \dots, z_n where a_1, \dots, a_n are complex numbers. If $\sum_{i=1}^n |a_i|^2 \leq 1$, then prove that $\sum_{i=1}^n |z_i|^2 \leq n$.

PROBLEM 5.171 (2003 Quiz 6, problem 1). Let n be a positive integer and a_1, \dots, a_n, x be real numbers.

$$g(x) = \sum_{i=1}^n a_i \cos ix$$

If $g(x) \geq -1$ for all real number x , then prove that $\sum_{i=1}^n a_i \leq n$.

PROBLEM 5.172 (2003 Quiz 8, problem 3). Let $n \geq 2$ be an integer and a_1, \dots, a_n be positive real numbers not all of which are equal such that

$$\sum_{i=1}^n \frac{1}{a_i^{2n}} = 1$$

Prove that

$$\left(\sum_{i=1}^n a_i^{2n} \right) - n^2 \sum_{1 \leq i < j \leq n} \left(\frac{a_i}{a_j} - \frac{a_j}{a_i} \right)^2 > n^2$$

PROBLEM 5.173 (2002 Team Selection Test, problem 6). Let

$$f(x_1, x_2, x_3) = -2(x_1^3 + x_2^3 + x_3^3) + 3x_1^2(x_2 + x_3) + 3x_2^2(x_3 + x_1) + 3x_3^2(x_1 + x_2) - 12x_1x_2x_3$$

For real numbers r, s, t , define

$$g(r, s, t) = \max_{t \leq x_3 \leq t+2} |f(r, r+2, x_3) + s|$$

Find the minimum value of $g(r, s, t)$.

PROBLEM 5.174 (2001 Team Selection Test, problem 4). Let $n > 3$ be an integer. The real numbers x_1, \dots, x_{n+2} satisfy the condition $0 < x_1 < \dots < x_{n+2}$. Find the minimum possible value of

$$\frac{\left(\sum_{i=1}^n \frac{x_{i+1}}{x_i} \right) \left(\sum_{i=1}^n \frac{x_{i+2}}{x_{i+1}} \right)}{\left(\sum_{i=1}^n \frac{x_{i+1}x_{i+2}}{x_{k+1}^2 + x_kx_{k+2}} \right) \left(\sum_{i=1}^n \frac{x_{i+1}^2 + x_ix_{i+2}}{x_ix_{i+1}} \right)}$$

PROBLEM 5.175 (2001 Team Selection Test, problem 6). Find the maximum value of

$$\max_{1 \leq x \leq 3} |x^3 - ax^2 - bx - c|$$

where a, b, c runs through all real numbers.

PROBLEM 5.176 (1999 Team Selection Test, problem 1). Let n be a positive integer and x_1, \dots, x_n be non-negative real numbers such that $x_1 + \dots + x_n = 1$. Find the largest possible value of $\sum_{i=1}^n (x_i^4 - x_j^5)$.

PROBLEM 5.177 (1998 Team Selection Test, problem 3). For a fixed real number $\vartheta \in [0, \frac{\pi}{2}]$, find the smallest positive real number a for which

$$\frac{\sqrt{a}}{\cos \vartheta} + \frac{\sqrt{a}}{\sin \vartheta} > 1$$

and there exists $x \in \left[1 - \frac{\sqrt{a}}{\sin \vartheta}, \cos \vartheta\right]$ such that

$$\left((1-x)\sin \vartheta - \sqrt{a - x^2 \cos^2 \vartheta}\right)^2 + \left(x \cos \vartheta - \sqrt{a - (1-x)^2 \sin^2 \vartheta}\right)^2 \leq a$$

PROBLEM 5.178 (1996 Team Selection Test, problem 5). Let $n \geq 4$ be an integer and $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_n$ be real numbers such that $\sum_{i=1}^n \alpha_i^2 < 1$ and $\sum_{i=1}^n \beta_i^2 < 1$.

$$\begin{aligned} A^2 &= 1 - \sum_{i=1}^n \alpha_i^2 \\ B^2 &= 1 - \sum_{i=1}^n \beta_i^2 \\ W &= \frac{1}{2} \left(1 - \sum_{i=1}^n \alpha_i \beta_i \right)^2 \end{aligned}$$

Find all real number λ such that

$$x^n + \lambda(x^{n-1} + \dots + x^3 + Wx^2 + ABx + 1) = 0$$

only has real roots.

PROBLEM 5.179 (1995 Team Selection Test, problem 4). Let n be a positive integer and $r_1, \dots, r_n, s_1, \dots, s_n, t_1, \dots, t_n, u_1, \dots, u_n, v_1, \dots, v_n$ be $5n$ real numbers.

$$R = \frac{1}{n} \sum_{i=1}^n r_i$$

$$S = \frac{1}{n} \sum_{i=1}^n s_i$$

$$T = \frac{1}{n} \sum_{i=1}^n t_i$$

$$U = \frac{1}{n} \sum_{i=1}^n u_i$$

$$V = \frac{1}{n} \sum_{i=1}^n v_i$$

Prove that

$$\prod_{i=1}^n \frac{r_i s_i t_i u_i v_i + 1}{r_i s_i t_i u_i v_i - 1} \geq \left(\frac{RSTUV + 1}{RSTUV - 1} \right)^n$$

PROBLEM 5.180 (1993 Team Selection Test, problem 2). Let $n \geq 2$ be an integer and a, b, c, d be positive integers such that $\frac{a}{b} + \frac{c}{d} < 1$ and $a + c \leq n$. Find the maximum value of $\frac{a}{b} + \frac{c}{d}$ for a fixed n .

§§5.13 IMO

PROBLEM 5.181 (IMO 1995, problem 2). Let a, b, c be real numbers such that $abc = 1$. Prove that

$$\frac{1}{a^3(b+c)} + \frac{1}{b^3(c+a)} + \frac{1}{c^3(a+b)} \geq \frac{3}{2}$$

PROBLEM 5.182 (IMO 1999, problem 2). Let $n \geq 2$ be an integer. Find the least constant C such that

$$\sum_{1 \leq i < j \leq n} x_i x_j (x_i^2 + x_j^2) \leq C \left(\sum_{i=1}^n x_i \right)^4$$

holds for all non-negative real numbers x_1, \dots, x_n . When does equality occur?

PROBLEM 5.183 (IMO 2000, problem 2). Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$\left(a - 1 + \frac{1}{b}\right) \left(b - 1 + \frac{1}{c}\right) \left(c - 1 + \frac{1}{a}\right) \leq 1$$

PROBLEM 5.184 (IMO 2001, problem 2). Let a, b, c be positive real numbers. Prove that

$$\frac{a}{\sqrt{a^2 + 8bc}} + \frac{b}{\sqrt{b^2 + 8ca}} + \frac{c}{\sqrt{c^2 + 8ab}} \geq 1$$

PROBLEM 5.185 (IMO 2003, problem 5). Let n be a positive integer and $x_1 \leq \dots \leq x_n$ be real numbers. Prove that

$$\left(\sum_{i,j=1}^n |x_i - x_j| \right)^2 \leq \frac{2(n^2 - 1)}{3} \sum_{i,j=1}^n (x_i - x_j)^2$$

Show that equality holds if and only if x_1, \dots, x_n forms an arithmetic sequence.

PROBLEM 5.186 (IMO 2004, problem 4). Let $n \geq 3$ be an integer. Let t_1, \dots, t_n be positive real numbers such that

$$n^2 + 1 > (t_1 + \dots + t_n) \left(\frac{1}{t_1} + \dots + \frac{1}{t_n} \right)$$

Show that t_i, t_j, t_k are the sides of a triangle for all $1 \leq i < j < k \leq n$.

PROBLEM 5.187 (IMO 2005, problem 3). Let x, y, z be real numbers such that $xyz \geq 1$. Prove that

$$\frac{x^5 - x^2}{x^5 + y^2 + z^2} + \frac{y^5 - y^2}{x^2 + y^5 + z^2} + \frac{z^5 - z^2}{x^2 + y^2 + z^5} \geq 0$$

Solution. We have already solved it in *Problem 3.10*.

PROBLEM 5.188 (IMO 2006, problem 3). Determine the least real number M such that the inequality

$$|ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2)| \leq M(a^2 + b^2 + c^2)^2$$

holds for all real numbers a, b, c .

PROBLEM 5.189 (IMO 2008, problem 2). Let $x, y, z \neq 1$ be real numbers such that $xyz = 1$. Prove that

$$\frac{x^2}{(x-1)^2} + \frac{y^2}{(y-1)^2} + \frac{z^2}{(z-1)^2} \geq 1$$

Also, prove that equality holds for infinitely many rational x, y, z such that $xyz = 1$ and $x, y, z \neq 1$.

PROBLEM 5.190 (IMO 2012, problem 2). Let $n \geq 3$ be a positive integer and a_2, \dots, a_n be positive real numbers such that $a_1 \cdots a_n = 1$. Prove that

$$(1 + a_2)^2 \cdots (1 + a_n)^n \geq n^n$$

PROBLEM 5.191 (IMO 2020, problem 2). Let a, b, c, d be positive real numbers such that $a \geq b \geq c \geq d$ and $a + b + c + d = 1$. Prove that

$$(a + 2b + 3c + 4d)a^a b^b c^c d^d < 1$$

Solution. Since $a + b + c + d = 1$, by *Weighted Power Mean Inequality* on $\omega = (a, b, c, d)$ and $\mathbf{a} = (a, b, c, d)$,

$$a \cdot a + b \cdot a + c \cdot c + d \cdot d \geq a^a b^b c^c d^d$$

So it is enough to prove that

$$(a + 2b + 3c + 4d)(a^2 + b^2 + c^2 + d^2) \leq (a + b + c + d)^3$$

Expanding these we can easily see that the inequality has to follow. But we can prove it in a smarter way.

$$(a + b + c + d)^3 = (a + b + c + d)(a^2 + b^2 + c^2 + d^2 + 2 \sum ab)$$

where the sum runs over all possible $\binom{4}{2}$ pairs. Then

$$\begin{aligned} (a + 2b + 3c + 4d)(a^2 + b^2 + c^2 + d^2) &< (a + b + c + d)^3 \\ \Leftrightarrow (b + 2c + 3d)(a^2 + b^2 + c^2 + d^2) &< 2(a + b + c + d)(\sum ab) \end{aligned}$$

Now, using $a \geq b \geq c$,

$$a^2 + b^2 + c^2 + d^2 \leq a(a + b + c + d)$$

So, it is enough to show that

$$\begin{aligned} a(b + 2c + 3d)(a + b + c + d) &\leq 2(a + b + c + d)(\sum ab) \\ \Leftrightarrow a(b + 2c + 3d) &\leq 2 \sum ab \\ \Leftrightarrow 3da &\leq ab + 2ca \\ \Leftrightarrow da + da + da &\leq ab + ca + ca \end{aligned}$$

This inequality obviously holds.

Remark. We could also use *Weighted Jensen's inequality* after using the fact that $\log(x)$ is concave. The buffalo way works here as well. But the calculation is not going to be pretty if you go that way. This problem was highly criticized within some forums such as the Art of Problem Solving. It was the first inequality problem at the IMO since 2012. A lot of people thought that the days of inequality at the IMO was over. But when this problem appeared at the IMO 2020, many people complained and expressed their disappointment that the *no inequality problem at the IMO* streak was finally broken with such a problem.

PROBLEM 5.192 (IMO Shortlist 2015, A1). Let a, b, c be positive real numbers such that $\min\{ab, bc, ca\} \geq 1$. Prove that

$$\sqrt[3]{(a^2 + 1)(b^2 + 1)(c^2 + 1)} \leq \left(\frac{a + b + c}{3}\right)^2 + 1$$

PROBLEM 5.193 (IMO Shortlist 2015, A8). Determine the largest real number a such that for all $n \geq 1$ and for all real numbers x_0, \dots, x_n satisfying

$$0 = x_0 < x_1 < \dots < x_n$$

we have

$$\frac{1}{x_1 - x_0} + \frac{1}{x_2 - x_1} + \dots + \frac{1}{x_n - x_{n-1}} \geq a \left(\frac{2}{x_1} + \frac{3}{x_2} + \dots + \frac{n+1}{x_n} \right)$$

PROBLEM 5.194 (IMO Shortlist 2018, A7). Find the maximal value of

$$S = \sqrt[3]{\frac{a}{b+7}} + \sqrt[3]{\frac{c}{d+7}} + \sqrt[3]{\frac{d}{a+7}}$$

where a, b, c, d are non-negative real numbers which satisfy $a+b+c+d = 100$.

§§5.14 IrMO

PROBLEM 5.195 (2002 Team Selection Test, problem 6). Let x_1, \dots, x_n be positive real numbers such that $x_1^2 + \dots + x_n^2 = n$, λ be a real number such that $0 \leq \lambda \leq 1$ and s be a positive real number such that $\sum_{i=1}^n x_i \geq s$. Prove that at least

$$\left\lceil \frac{s^2(1-\lambda)^2}{n} \right\rceil$$

of these numbers are larger than $\frac{\lambda s}{n}$.

PROBLEM 5.196 (2003 Round 2, problem 1). Let x, y, z be real numbers such that $xyz = -1$. Prove that

$$x^4 + y^4 + z^4 + 3(x + y + z) \geq \frac{x^2}{y} + \frac{y^2}{z} + \frac{z^2}{x} + \frac{y^2}{x} + \frac{z^2}{x} + \frac{x^2}{z}$$

PROBLEM 5.197 (2005 Team Selection Test, problem 1). Let a_1, \dots, a_n be real numbers and

$$\begin{aligned} \frac{a_1 + \dots + a_n}{n} &= m \\ \frac{a_1^2 + \dots + a_n^2}{n} &= 1 \end{aligned}$$

If there is an i such that $a_i \leq m$, prove that

$$n - i \geq n(m - a_i)^2$$

PROBLEM 5.198 (2006 Team Selection Test, problem 4). Let x_1, \dots, x_n be real numbers. Prove that

$$\sum_{i,j=1}^n |x_i + x_j| \geq n \sum_{i=1}^n |x_i|$$

PROBLEM 5.199 (2008 Team Selection Test, problem 5). Let a, b, c be positive real numbers such that $ab + bc + ca = 1$. Prove that

$$\sqrt{a^3 + a} + \sqrt{b^3 + b} + \sqrt{c^3 + c} \geq 2\sqrt{a + b + c}$$

PROBLEM 5.200 (2009 Team Selection Test, problem 3). Let a, b, c be positive real numbers such that $a + b + c = 3$. Prove that

$$\frac{1}{2 + a^2 + b^2} + \frac{1}{2 + b^2 + c^2} + \frac{1}{2 + c^2 + a^2} \leq \frac{3}{4}$$

PROBLEM 5.201 (2010 Third Round, problem 6). We call a sequence of real numbers a_{10}, \dots, a_{1389} *concave* if $2a_i \geq a_{i-1} + a_{i+1}$ for $0 < i < 1389$. Find the largest real number c such that

$$\sum_{i=0}^{1389} ia_i \geq c \sum_{i=0}^{1389} a_i^2$$

for all concave sequence of non-negative real numbers (a_n) .

PROBLEM 5.202 (2011 Third Round, problem 10). Find the smallest real number k such that

$$\sqrt{(a^2 + 1)(b^2 + 1)(c^2 + 1)} + \sqrt{(b^2 + 1)(c^2 + 1)(d^2 + 1)} + \sqrt{(c^2 + 1)(d^2 + 1)(a^2 + 1)} + \sqrt{(d^2 + 1)(a^2 + 1)} \geq k(a^2 + b^2 + c^2 + d^2)$$

for all real numbers a, b, c, d .

PROBLEM 5.203 (2012 Third Round, problem 10). Let a, b, c be positive real numbers such that $ab + bc + ca = 1$. Show that

$$\sqrt{3}(\sqrt{a} + \sqrt{b} + \sqrt{c}) \leq \frac{a\sqrt{a}}{bc} + \frac{b\sqrt{b}}{ca} + \frac{c\sqrt{c}}{ab}$$

PROBLEM 5.204 (2013 Team Selection Test, problem 11). Let a, b, c be the sides of a triangle such that $a \geq b \geq c$. Prove that

$$\sqrt{a(a+b-\sqrt{ab})} + \sqrt{b(b+c-\sqrt{bc})} + \sqrt{c(c+a-\sqrt{ca})} \geq a+b+c$$

PROBLEM 5.205 (2014 Round 2, problem 3). Let x, y, z be non-negative real numbers such that $x^2 + y^2 + z^2 = 2(xy + yz + zx)$. Prove that

$$\frac{x+y+z}{3} \sqrt[3]{2xyz}$$

PROBLEM 5.206 (2014 Team Selection Test, problem 5). Let n be a positive integer and x_1, \dots, x_{n+1} be positive real numbers such that $x_1 \cdots x_n = 1$. Prove that

$$\sqrt[n]{x_1} + \sqrt[n]{x_2} + \dots + \sqrt[n]{x_n} \geq n^{\sqrt[n]{x_1}} + \dots + n^{\sqrt[n]{x_n}}$$

PROBLEM 5.207 (2014 Team Selection Test 2, problem 5). Let x, y, z be positive real numbers such that

$$x^2 + y^2 + z^2 = x^2y^2 + y^2z^2 + z^2x^2$$

Prove that

$$((x-y)(y-z)(z-x))^2 \leq 2((x^2 - y^2)^2 + (y^2 - z^2)^2 + (z^2 - x^2)^2)$$

PROBLEM 5.208 (2015 Team Selection Test, problem 6). If a, b, c are positive real numbers such that $a + b + c = abc$, prove that

$$\frac{abc}{3\sqrt{2}} \left(\sum_{cyc} \frac{\sqrt{a^3 + b^3}}{ab + 1} \right) \geq \sum_{cyc} \frac{a}{a^2 + 1}$$

PROBLEM 5.209 (2016 Team Selection Test, problem 1). Let a, b, c, d be positive real numbers such that

$$\frac{1}{a+1} + \frac{1}{b+1} + \frac{1}{c+1} + \frac{1}{d+1} = 2$$

Prove that

$$\sqrt{\frac{a^2+1}{2}} + \sqrt{\frac{b^2+1}{2}} + \sqrt{\frac{c^2+1}{2}} + \sqrt{\frac{d^2+1}{2}} \geq 3(\sqrt{a} + \sqrt{b} + \sqrt{c} + \sqrt{d}) - 8$$

PROBLEM 5.210 (2016 Round 2, problem 1). Let a, b, c be positive real numbers such that $c \geq b \geq a$. Prove that

$$\frac{(c-a)^2}{6c} \leq \frac{a+b+c}{3} - \frac{3}{\frac{1}{a} + \frac{1}{b} + \frac{1}{c}}$$

PROBLEM 5.211 (2017 Round 2, problem 4). Let x, y be two positive real numbers such that $x^4 - y^4 = x - y$. Prove that

$$\frac{x-y}{x^6 - y^6} \leq \frac{4}{3}(x+y)$$

PROBLEM 5.212 (2017 Team Selection Test, problem 1). Let a, b, c, d be positive real numbers such that $a + b + c + d = 2$. Prove that

$$\frac{(a+c)^2}{ad+bc} + \frac{(b+d)^2}{ac+bd} + 4 \geq 4 \left(\frac{a+b+1}{c+d+1} + \frac{c+d+1}{a+b+1} \right)$$

PROBLEM 5.213 (2018 Team Selection Test, problem 2). Find the smallest real number k such that

$$\left(\frac{2a}{a-b} \right)^2 + \left(\frac{2b}{b-c} \right)^2 + \left(\frac{2c}{c-a} \right)^2 + k4 \left(\frac{2a}{a-b} + \frac{2b}{b-c} + \frac{2c}{c-a} \right)$$

for all real numbers a, b, c .

PROBLEM 5.214 (2020 Second Round, problem 2). Let x, y, z be positive real numbers such that $x + y + z = 1399$. Find

$$\max ([x]y, [y]z, [z]x)$$

§§5.15 MMO

PROBLEM 5.215 (2018, problem 1). Let $n > 1$ be an integer and a_1, \dots, a_n be real numbers such that $0 \leq a_i \leq \frac{\pi}{2}$. Prove that

$$\left(\frac{1}{n} \sum_{i=1}^n \frac{1}{1 + \sin a_i} \right) \left(1 + \prod_{i=1}^n \sqrt[n]{\sin a_i} \right) \leq 1$$

PROBLEM 5.216 (2017, problem 4). Let x, y, z and a, b, c be positive real numbers such that $a + b + c = 1$. Prove that

$$(x^2 + y^2 + z^2) \left(\frac{a^3}{x^2 + 2y^2} + \frac{b^3}{y^2 + 2z^2} + \frac{c^3}{z^2 + 2x^2} \right) \geq \frac{1}{9}$$

PROBLEM 5.217 (2016, problem 2). Let a, b, c be positive real numbers such that $a + b + c = 3$. Prove that

$$\sqrt{\frac{b}{a^2 + 3}} + \sqrt{\frac{c}{b^2 + 3}} + \sqrt{\frac{a}{c^2 + 3}} \leq \frac{3}{2} \sqrt[4]{\frac{1}{abc}}$$

PROBLEM 5.218 (2014, problem 1). Let n be a positive integer and $a_1, \dots, a_n, b_1, \dots, b_n$ be $2n$ real numbers. Prove that there exists a positive integer $k \leq n$ such that

$$\sum_{i=1}^n |a_i - a_k| \leq \sum_{i=1}^n |b_i - a_k|$$

PROBLEM 5.219 (2013, problem 3). Let x, y, z be positive real numbers such that $x^2y^2 + y^2z^2 + z^2x^2 = 6xyz$. Prove that

$$\sqrt{\frac{x}{x + yz}} + \sqrt{\frac{y}{y + zx}} + \sqrt{\frac{z}{z + xy}} \geq \sqrt{3}$$

PROBLEM 5.220 (2010, problem 2). Let $n > 2$ be an integer and a_1, \dots, a_n be positive real numbers such that $a_1 + \dots + a_n = 1$. Prove that

$$\frac{a_2 \cdots a_n}{a_1 + n - 2} + \frac{a_1 a_3 \cdots a_n}{a_2 + n - 2} + \dots + \frac{a_1 \cdots a_{n-1}}{a_n + n - 2} \leq \frac{1}{(n-1)^2}$$

PROBLEM 5.221 (2009, problem 4). Let x, y, z be positive real numbers. Prove that

$$\sum_{cyc} \frac{xy}{x^2 + xy + y^2} \leq \sum_{cyc} \frac{x}{2x + z}$$

PROBLEM 5.222 (2007, problem 1). Let $x \geq y \geq z$ be real numbers such that $xy + yz + zx = 1$. Prove that $xz < \frac{1}{2}$. Is it possible to improve the constant $\frac{1}{2}$?

PROBLEM 5.223 (2007, problem 4). Let $x > 1$ be a non-integer real number. Prove that

$$\left(\frac{x + \{x\}}{\lfloor x \rfloor} - \frac{\lfloor x \rfloor}{x + \{x\}} \right) + \left(\frac{x + \lfloor x \rfloor}{\{x\}} - \frac{\{x\}}{x + \lfloor x \rfloor} \right) > \frac{9}{2}$$

PROBLEM 5.224 (2007, problem 4). Let $(x_{ij})^{m \times n}$ be a matrix of real numbers such that $0 \leq x_{ij} \leq 1$. Prove that

$$\prod_{j=1}^n \left(1 - \prod_{i=1}^m x_{ij} \right) + \prod_{i=1}^m \left(1 - \prod_{j=1}^n (1 - x_{ij}) \right) \geq 1$$

PROBLEM 5.225 (2004, problem 3). Let a, b, c be positive real numbers such that $ab + bc + ca + 2abc = 1$. Prove that

$$2(a + b + c) + 1 \geq 32abc$$

PROBLEM 5.226 (2003, problem 3). Let a, b, c be non-negative real numbers such that $a + b + c = 3$. Prove that

$$\frac{a}{b^2 + 1} + \frac{b}{c^2 + 1} + \frac{c}{a^2 + 1} \geq \frac{3}{2}$$

PROBLEM 5.227 (2002, problem 4). Let a, b, c be non-negative real numbers such that $a^2 + b^2 + c^2 = 1$. Prove that

$$\frac{a}{b^2 + 1} + \frac{b}{c^2 + 1} + \frac{c}{a^2 + 1} \geq \frac{3}{4} (a\sqrt{a} + b\sqrt{b} + c\sqrt{c})^2$$

PROBLEM 5.228 (1999, problem 3). Let a, b, c be non-zero real numbers and x, y, z be positive real numbers such that $x + y + z = 3$. Prove that

$$\frac{3}{2} \sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}} \geq \frac{x}{1+a^2} + \frac{y}{1+b^2} + \frac{z}{1+c^2}$$

§§5.16 PMO

PROBLEM 5.229 (2019 Finals, problem 5). Let a_0, a_1, \dots be a sequence of positive real numbers such that a_0 is an integer, $a_i \leq a_{i-1} + 1$ for $1 \leq i \leq n$

and

$$\sum_{i=1}^n \frac{1}{a_i} \leq 1$$

Prove that

$$n \leq 4a_0 \sum_{i=1}^n \frac{1}{a_i}$$

PROBLEM 5.230 (2017 Finals, problem 6). Three sequences $a_0, \dots, a_n; b_0, \dots, b_n; c_0, \dots, c_{2n}$ of non-negative real numbers are given such that for all $0 \leq i, j \leq n$, we have $a_i b_j \leq c_{i+j}^2$. Prove that

$$\left(\sum_{i=0}^n a_i \right) \left(\sum_{i=0}^n b_i \right) \leq \left(\sum_{i=0}^{2n} c_i \right)^2$$

PROBLEM 5.231 (2014 Finals, problem 2). Let n be a positive integer, $k \geq 2$ be an integer and $a_1, \dots, a_k; b_1, \dots, b_n$ be integers such that

$$1 < a_1 < \dots < a_k < b_1 < \dots < b_n$$

Prove that if

$$a_1 + \dots + a_k > b_1 + \dots + b_n$$

then

$$a_1 \cdots a_k > b_1 \cdots b_n$$

PROBLEM 5.232 (2013 Finals, problem 5). Let k, m, n be distinct positive integers. Prove that

$$\left(k - \frac{1}{k} \right) \left(m - \frac{1}{m} \right) \left(n - \frac{1}{n} \right) \leq kmn - (k + m + n)$$

PROBLEM 5.233 (2012 Finals, problem 6). Show that for any positive real numbers a, b, c ,

$$\left(\frac{a-b}{c} \right)^2 + \left(\frac{c-a}{b} \right)^2 + \left(\frac{a-b}{c} \right)^2 \geq 2\sqrt{2} \left(\frac{a-b}{c} + \frac{b-c}{a} + \frac{c-a}{b} \right)$$

PROBLEM 5.234 (2008 Finals, problem 1). In each cell of an $n \times n$ matrix, a positive integer at most n^2 is written. In the first row, $1, \dots, n$ are written, in the second $n+1, \dots, 2n$ are written and so on. n numbers are selected such that no two numbers are in the same row or column. If a_i is the number chosen from row i , prove that

$$\frac{1}{a_1} + \frac{2^2}{a_2} + \dots + \frac{n^2}{a_n} \geq \frac{n+2}{2} - \frac{1}{n^2+1}$$

PROBLEM 5.235 (2004 Finals, problem 4). Let a, b, c be real numbers. Prove that

$$\sqrt{2(a^2 + b^2)} + \sqrt{2(b^2 + c^2)} + \sqrt{2(c^2 + a^2)} \geq \sqrt{3(a+b)^2 + 3(b+c)^2 + 3(c+a)^2}$$

PROBLEM 5.236 (2003 Finals, problem 2). Let a be a real number such that $0 < a < 1$. Prove that for all finite strictly increasing sequences k_1, \dots, k_n of non-negative integers,

$$\left(\sum_{i=1}^n a^{k_i} \right)^2 < \left(\frac{1+a}{1-a} \right) \sum_{i=1}^n a^{2k_i}$$

PROBLEM 5.237 (2002 Finals, problem 4). Let $n \geq 3$ be an integer and x_1, \dots, x_n be non-negative real numbers. Prove that at least one of the following inequalities are true.

$$\begin{aligned} \sum_{i=1}^n \frac{x_i}{x_{i+1} + x_{i+2}} &\geq \frac{n}{2} \\ \sum_{i=1}^n \frac{x_i}{x_{i-1} + x_{i-2}} &\geq \frac{n}{2} \end{aligned}$$

where $x_{n+2} = x_2, x_{n+1} = x_1, x_0 = x_n, x_{-1} = x_{n-1}$.

PROBLEM 5.238 (2001 Finals, problem 1). Let n be a positive integer and x_1, \dots, x_n be positive real numbers. Prove that

$$x_1 + \dots + nx_n \leq \frac{n(n-1)}{2} + x_1 + \dots + x_n^n$$

PROBLEM 5.239 (1999 Finals, problem 2). Let n be a positive integer and $a_1, \dots, a_n; b_1, \dots, b_n$ be positive real numbers. Prove that

$$\sum_{1 \leq i < j \leq n} |a_i - a_j| + \sum_{1 \leq i < j \leq n} |b_i - b_j| \leq \sum_{i=1}^n \sum_{j=1}^n |a_i - b_j|$$

PROBLEM 5.240 (1996 Finals, problem 3). Let n be a positive integer and $a_1, \dots, a_n; x_1, \dots, x_n$ be positive real numbers such that

$$a_1 + \dots + a_n = x_1 + \dots + x_n = 1$$

Show that

$$2 \sum_{1 \leq i < j \leq n} x_i x_j \leq \frac{n-2}{n-1} + \sum_{i=1}^n \frac{a_i x_i^2}{1-a_i}$$

When do we have equality?

PROBLEM 5.241 (1995 Finals, problem 1). Let n be a positive integer and $x = (x_1, \dots, x_n)$ be positive real numbers such that $\mathfrak{H}(x) = 1$. Find the smallest possible value of

$$x_1 + \dots + \frac{x_n^n}{n}$$

PROBLEM 5.242 (1994 Finals, problem 3). Let $n > 3$ be an integer and $X = \{x_1, \dots, x_n\}$ be set of distinct real numbers such that $\sum_{i=1}^n x_i = 0$ and $\sum_{i=1}^n x_i^2 = 1$. Show that there are real numbers $a, b, c, d \in X$ such that

$$a + b + c + nabc \leq \sum_{i=1}^n x_i^3 \leq a + b + d + nabd$$

§§5.17 RNO

PROBLEM 5.243 (2019 Team Selection Test, Day 5, problem 1). Determine the largest value of the expression

$$\sum_{1 \leq i < j \leq 4} (x_i + x_j) \sqrt{x_i x_j}$$

where x_1, x_2, x_3, x_4 non-negative real numbers such that $x_1 + x_2 + x_3 + x_4 = 1$. Also, find the specific values where this maximum is achieved.

PROBLEM 5.244 (2018 Team Selection Test, Day 1, problem 1). Let $x_1, \dots, x_n \geq -1$ be real numbers such that $\sum_{i=1}^n x_i^3 = 0$. Find the least constant c such that

$$\sum_{i=1}^n x_i^2 \leq cn$$

PROBLEM 5.245 (2014 Team Selection Test, Day 3, problem 3). Determine the smallest positive constant c such that

$$\sum_{i=1}^n \left(\frac{1}{i} \sum_{j=1}^i x_j \right)^2 \leq c \sum_{i=1}^n x_i^2$$

for all positive integer n and all positive real numbers x_1, \dots, x_n .

PROBLEM 5.246 (2012 Team Selection Test, Day 2, problem 4). Let k be a positive integer. Find the maximum value of

$$a^{3k-1}b + b^{3k-1}c + c^{3k-1}a + k^2 a^k b^k c^k$$

where a, b, c are non-negative real numbers such that $a + b + c = 3k$.

PROBLEM 5.247 (2011 Team Selection Test, Day 3, problem 2). Given real numbers x, y, z such that $x + y + z = 0$. Show that

$$\frac{x(x+2)}{2x^2+1} + \frac{y(y+2)}{2y^2+1} + \frac{z(z+2)}{2z^2+1} \geq 0$$

PROBLEM 5.248 (2011 Team Selection Test, Day 5, problem 2). Let $n \geq 2$ be an integer and x_1, \dots, x_n be positive real numbers such that

$$\sum_{i=1}^n \frac{1}{x_i + 1} = 1$$

and $k > 1$ be a real number. Show that

$$\sum_{i=1}^n \frac{1}{x_i^k + 1} \geq \frac{n}{(n-1)^k + 1}$$

and determine the case of equality.

PROBLEM 5.249 (2010 Team Selection Test 1, problem 2). Let n be a positive integer and a_1, \dots, a_n be positive real numbers. Prove that $f : [0, \infty) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{a_1 + x}{a_2 + x} + \frac{a_2 + x}{a_3 + x} + \dots + \frac{a_n + x}{a_1 + x}$$

is decreasing.

PROBLEM 5.250 (2010 Team Selection 2, problem 1). Let n be a positive integer. Determine the maximum value of

$$\max \left\{ \frac{x_1}{1+x_1}, \dots, \frac{x_n}{1+x_1+\dots+x_n} \right\}$$

as x_1, \dots, x_n runs through all non-negative real numbers such that $x_1 + \dots + x_n = 1$.

PROBLEM 5.251 (2010 Team Selection Test 3, problem 1). Let n be a positive integer and x_1, \dots, x_n be positive real numbers such that $x_1 \cdots x_n = 1$. Prove that

$$\sum_{i=1}^n x_i^n (1+x_i) \geq \frac{n}{2^{n-1}} \prod_{i=1}^n (1+x_i)$$

PROBLEM 5.252 (2009 Team Selection Test, Day 4, problem 1). Let $n \geq 2$ be an integer and x_1, \dots, x_n be positive integers such that $x_1 \leq \dots \leq x_n$ and $x_1 + \dots + x_n = x_1 \cdots x_n$. What is the maximum possible value of $x_1 + \dots + x_n$.

PROBLEM 5.253 (2008 Team Selection Test, Day 1, problem 2). Let $n \geq 2$ be an integer and $a_1, \dots, a_n; b_1, \dots, b_n$ be positive real numbers such that $a_i < b_i$ and

$$b_1 + \dots + b_n < 1 + a_1 + \dots + a_n$$

Prove that there exists a real number c such that

$$(a_i + c + k)(b_i + c + k) > 0$$

for $1 \leq i \leq n$ and any integer k .

PROBLEM 5.254 (2008 Team Selection Test, Day 2, problem 1). Let $n \geq 3$ be an odd integer. Determine the maximum value of

$$\sqrt{|x_1 - x_2|} + \dots + \sqrt{|x_n - x_1|}$$

where x_1, \dots, x_n are real numbers in the interval $[0, 1]$.

PROBLEM 5.255 (2007 Team Selection Test, Day 1, problem 1). Let a_1, \dots, a_n be non-negative real numbers such that $a_1^2 + \dots + a_n^2 = 1$. Find the maximum value of

$$(1 - a_1) \cdots (1 - a_n)$$

PROBLEM 5.256 (2007 Team Selection Test, Day 3, problem 2). Let $n, p \geq 4$ be integers and x_1, \dots, x_n be positive real numbers such that $x_1 + \dots + x_n = n$. Prove that

$$\sum_{i=1}^n \frac{1}{x_i^p} \geq \sum_{i=1}^n x_i^p$$

is false.

PROBLEM 5.257 (2007 Team Selection Test, Day 7, problem 1). Let $n \geq 2$ be an integer and $a_1, \dots, a_n; b_1, \dots, b_n$ be real numbers such that

$$\begin{aligned} \sum_{i=1}^n a_i^2 &= \sum_{i=1}^n b_i^2 = 1 \\ \sum_{i=1}^n a_i b_i &= 0 \end{aligned}$$

Prove that

$$\left(\sum_{i=1}^n a_i \right)^2 + \left(\sum_{i=1}^n b_i \right)^2 \leq n$$

PROBLEM 5.258 (2006 Team Selection Test, Day 1, problem 4). Let n be a positive integer and a_1, \dots, a_n be real numbers such that $|a_i| \leq 1$ for $1 \leq i \leq n$ and $a_1 + \dots + a_n = 0$. Prove that there exists a positive integer $k \leq n$ such that

$$|1 \cdot a_1 + \dots + k a_k| \leq \frac{2k+1}{4}$$

Also show that this is the best bound possible for $n > 2$.

PROBLEM 5.259 (2006 Team Selection Test, Day 2, problem 4). Let n be a positive integer and x_1, \dots, x_n be real numbers. Prove that

$$\sum_{1 \leq i < j \leq n} |x_i + x_j| \geq \frac{n-2}{2} \sum_{i=1}^n |x_i|$$

PROBLEM 5.260 (2005 Team Selection Test, Day 5, problem 2). Let $n \geq 2$ be an integer and x_1, \dots, x_n be positive real numbers such that $x_1 \cdots x_n = 1$. Find the smallest real value $\varrho(n)$ such that for any x_1, \dots, x_n ,

$$\sum_{i=1}^n \frac{1}{x_i} \leq \sum_{i=1}^n x_i^r$$

is true for all $r \geq \varrho(n)$.

PROBLEM 5.261 (2004 Team Selection Test, Day 1, problem 1). Let a_1, \dots, a_4 be the sides of a quadrilateral with perimeter $2s$. Prove that

$$\sum_{i=1}^4 \frac{1}{a_i + s} \leq \frac{2}{9} \sum_{1 \leq i < j \leq 4} \frac{1}{\sqrt{(s - a_i)(s - a_j)}}$$

When does equality hold?

PROBLEM 5.262 (2002 Team Selection Test, Day 3, problem 2). Let $n \geq 4$ be an integer and a_1, \dots, a_n be positive real numbers such that

$$a_1^2 + \dots + a_n^2 = 1$$

Prove that

$$\frac{a_1}{a_1^2 + 1} + \dots + \frac{a_n}{a_n^2 + 1} \geq \frac{4}{5} (a_1 \sqrt{a_1} + \dots + a_n \sqrt{a_n})^2$$

PROBLEM 5.263 (2001 Team Selection Test, Day 1, problem 3). Let a, b, c be the sides of a triangle. Prove that

$$(-a + b + c)(a - b + c) + (a - b + c)(a + b - c) + (a + b - c)(-a + b + c) \leq \sqrt{abc}(\sqrt{a} + \sqrt{b} + \sqrt{c})$$

PROBLEM 5.264 (2000 Team Selection Test, Day 1, problem 2). Let n be a positive integer and x_1, \dots, x_n be real numbers such that $|x_{k+1} - x_k| \leq 1$ for $1 \leq k \leq n - 1$. Prove that

$$\sum_{i=1}^n |x_i| - \left| \sum_{i=1}^n x_i \right| \leq \frac{n^2 - 1}{4}$$

PROBLEM 5.265 (1999 Team Selection Test, Day 2, problem 2). Let n and x_1, \dots, x_n be positive integers. Prove that

$$x_1^2 + \dots + x_n^2 \geq \frac{2n+1}{3}(x_1 + \dots + x_n)$$

PROBLEM 5.266 (1996 Team Selection Test, Day 2, problem 4). If p_1, \dots, p_k are the distinct prime divisors of n , define

$$a_n = \frac{1}{p_1} + \dots + \frac{1}{p_n}$$

Prove that

$$\sum_{i=2}^n a_1 \cdots a_i < 1$$

for $n \geq 2$.

PROBLEM 5.267 (1996 Team Selection Test, Day 3, problem 1). Let $n \geq 3$ be an integer and x_1, \dots, x_{n-1} be non-negative integers such that

$$\begin{aligned}x_1 + \dots + x_{n-1} &= n \\1 \cdot x_1 + \dots + (n-1)x_n &= 2(n-2)\end{aligned}$$

Find the minimum value of

$$\sum_{i=1}^n i(2n-i)x_i$$

PROBLEM 5.268 (1996 Team Selection Test, Day 4, problem 1). Let n be a positive integer and x_1, \dots, x_n be positive real numbers such that

$$x_{n+1} = x_1 + \dots + x_n$$

Prove that

$$\sum_{i=1}^n \sqrt{x_i(x_{n+1}-x_i)} \leq \sqrt{\sum_{i=1}^n x_{n+1}(x_{n+1}-x_i)}$$

PROBLEM 5.269 (1993 Team Selection Test, Day 1, problem 1). Find the maximum possible constant A such that

$$\frac{x}{\sqrt{y^2+z^2}} + \frac{y}{\sqrt{z^2+x^2}} + \frac{z}{\sqrt{x^2+y^2}} \geq A$$

for all positive real numbers x, y, z .

§§5.18 SgMO

PROBLEM 5.270 (Singapore Team Selection Test 2009, problem 2). If a, b, c are three positive real numbers such that $ab + bc + ca = 1$, prove that

$$\sqrt[3]{\frac{1}{a} + 6b} + \sqrt[3]{\frac{1}{b} + 6c} + \sqrt[3]{\frac{1}{c} + 6a} \leq \frac{1}{abc}$$

PROBLEM 5.271 (Singapore Team Selection Test 2008, problem 2). Let x_1, \dots, x_n be positive real numbers such that $x_1 \cdots x_n = 1$. Prove that

$$\sum_{i=1}^n \frac{1}{n-1+x_i} \leq 1$$

PROBLEM 5.272 (Singapore Team Selection Test 2007, problem 2). Prove the inequality

$$\sum_{i < j} \frac{a_i a_j}{a_i + a_j} \leq \frac{n}{2(a_1 + \dots + a_n)} \sum_{i < j} a_i a_j$$

for all positive real numbers a_1, \dots, a_n .

PROBLEM 5.273 (Singapore 2006, problem 2). Let $n > 1$ be an integer and x_1, \dots, x_n be real numbers such that

$$\begin{aligned} |x_1| + \dots + |x_n| &= 1 \quad \text{and} \\ x_1 + \dots + x_n &= 0 \end{aligned}$$

Prove that

$$\left| \frac{x_1}{1} + \dots + \frac{x_n}{n} \right| \leq \frac{1}{2} \left(1 - \frac{1}{n} \right)$$

PROBLEM 5.274 (Singapore 2004, problem 2). Let a, b, c be real numbers such that $ab + bc + ca = 1$ and $0 < a, b, c < 1$. Prove that

$$\frac{a}{1-a^2} + \frac{b}{1-b^2} + \frac{c}{1-c^2} \geq \frac{3\sqrt{3}}{2}$$

§§5.19 TNO

PROBLEM 5.275 (2005 Day 1, problem 1). Let a, b, c be positive real numbers such that $abc = 1$. Prove that

$$1 + \frac{3}{a+b+c} \geq \frac{6}{ab+bc+ca}$$

PROBLEM 5.276 (2005 Day 2, problem 3). Let a_1, \dots, a_{95} be positive real numbers. Prove that

$$\sum_{i=1}^{95} a_i \leq 94 + \prod_{i=1}^n \max\{1, a_i\}$$

PROBLEM 5.277 (2002 Day 1, problem 3). Let $x, y, z, a, b, c, d, e, f$ be real numbers satisfying

$$\begin{aligned}\max\{a, 0\} + \max\{b, 0\} &< x + ay + bz < 1 + \min\{a, 0\} + \min\{b, 0\} \\ \max\{c, 0\} + \max\{d, 0\} &< cx + y + dz < 1 + \min\{c, 0\} + \min\{d, 0\} \\ \max\{e, 0\} + \max\{f, 0\} &< ex + fy + z < 1 + \min\{e, 0\} + \min\{f, 0\}\end{aligned}$$

Prove that $0 < x, y, z < 1$.

PROBLEM 5.278 (2002 Day 2, problem 1). Let x_1, x_2, x_3, x_4 be positive real numbers at most $\frac{1}{2}$. Prove that

$$\frac{x_1 x_2 x_3 x_4}{(1-x_1)(1-x_2)(1-x_3)(1-x_4)} \leq \frac{x_1^4 + x_2^4 + x_3^4 + x_4^4}{(1-x_1)^4 + (1-x_2)^4 + (1-x_3)^4 + (1-x_4)^4}$$

PROBLEM 5.279 (1996 Day 1, problem 2). Let a be a positive real number at most 1 and a_1, \dots, a_{1996} be real numbers such that $a \leq a_i \leq \frac{1}{a_i}$. If k_1, \dots, k_{1996} are any non-negative real numbers such that $k_1 + \dots + k_{1996} = 1$, prove that

$$\left(\sum_{i=1}^{1996} k_i a_i \right) \left(\sum_{i=1}^n \frac{k_i}{a_i} \right) \leq \left(a + \frac{1}{a} \right)^2$$

PROBLEM 5.280 (1996 Day 2, problem 2). Determine integers $a_1, \dots, a_{99} = a_0$ satisfying $|a_k - a_{k+1}| \geq 1996$ for $1 \leq k \leq 1995$ such that

$$\max_{1 \leq i \leq 1995} |a_i - a_{i+1}|$$

is as minimum as possible. What is this minimum value?

PROBLEM 5.281 (1994 Day 1, problem 2). Let a, b, c be positive real numbers and α be a real number.

$$\begin{aligned}f(\alpha) &= abc(a^\alpha + b^\alpha + c^\alpha) \\ g(\alpha) &= a^{2+\alpha}(b+c-a) + b^{2+\alpha}(c+a-b) + c^{2+\alpha}(a+b-c)\end{aligned}$$

Determine $\min |f(\alpha) - g(\alpha)|$ and $\max |f(\alpha) - g(\alpha)|$ if they exist.

§§5.20 USAMO

PROBLEM 5.282 (USAMO 2020, problem 6). Let $n \geq 2$ be an integer and $x_1 \geq \dots \geq x_n, y_1 \geq \dots \geq y_n$ be $2n$ real numbers such that

$$\begin{aligned}0 &= x_1 + \dots + x_n = y_1 + \dots + y_n \\ 1 &= x_1^2 + \dots + x_n^2 = y_1^2 + \dots + y_n^2\end{aligned}$$

Prove that

$$\sum_{i=1}^n (x_i y_i - x_i y_{n+i-1})^2 \geq \frac{2}{\sqrt{n-1}}$$

PROBLEM 5.283 (USAMO 2018, problem 1). Let a, b, c be positive real numbers such that $a + b + c = 4\sqrt[3]{abc}$. Prove that

$$2(ab + bc + ca) + 4 \min\{a^2, b^2, c^2\} \geq a^2 + b^2 + c^2$$

PROBLEM 5.284 (USAMO 2017, problem 6). Find the minimum possible value of

$$\frac{a}{b^3 + 4} + \frac{b}{c^3 + 4} + \frac{c}{d^3 + 4} + \frac{d}{a^3 + 4}$$

given that a, b, c, d are non-negative real numbers such that $a + b + c + d = 4$.

PROBLEM 5.285 (USAMO 2013, problem 4). Find all real numbers $x, y, z \geq 1$ such that

$$\min\{\sqrt{x+xyz}, \sqrt{y+xyz}, \sqrt{z+xyz}\} = \sqrt{x-1} + \sqrt{y-1} + \sqrt{z-1}$$

This is actually a special case of the following.

PROBLEM 5.286. Prove that for real numbers $a, b, c \geq 1$, the following inequality holds:

$$\sqrt{a-1} + \sqrt{b-1} + \sqrt{c-1} \leq \sqrt{a(bc+1)}$$

PROBLEM 5.287 (USAMO 2012, problem 6). Let $n \geq 2$ be an integer and $x_1 \geq \dots \geq x_n, y_1 \geq \dots \geq y_n$ be $2n$ real numbers such that

$$\begin{aligned} 0 &= x_1 + \dots + x_n = y_1 + \dots + y_n \\ 1 &= x_1^2 + \dots + x_n^2 = y_1^2 + \dots + y_n^2 \end{aligned}$$

For each subset A of $\{1, 2, \dots, n\}$, define

$$S_A = \sum_{i \in A} x_i$$

$S_A = 0$ if A is empty. Prove that for any positive number λ , the number of sets A satisfying $S_A > \lambda$ is at most

$$\frac{2^{n-3}}{\lambda^2}$$

For which choices of x_1, \dots, x_n, λ does equality hold?

PROBLEM 5.288 (USAMO 2011, problem 1). Let a, b, c be positive real numbers such that

$$a^2 + b^2 + c^2 + (a + b + c)^2 \leq 4$$

Prove that

$$\frac{ab+1}{(a+b)^2} + \frac{bc+1}{(b+c)^2} + \frac{ca+1}{(c+a)^2} \geq 3$$

PROBLEM 5.289 (USAMO 2009, problem 4). Let $n \geq 2$ be an a_1, \dots, a_n be positive real numbers such that

$$(a_1 + \dots + a_n) \left(\frac{1}{a_1} + \dots + \frac{1}{a_n} \right) \leq \left(n + \frac{1}{2} \right)^2$$

Prove that

$$\max\{a_1, \dots, a_n\} \leq 4 \min\{a_1, \dots, a_n\}$$

PROBLEM 5.290 (USAMO 2004, problem 5). Let a, b, c be positive real numbers. Prove that

$$(a^5 - a^2 + 3)(b^5 - b^2 + 3)(c^5 - c^2 + 3) \geq (a + b + c)^3$$

PROBLEM 5.291 (USAMO 2003, problem 5). Let a, b, c be positive real numbers. Prove that

$$\frac{(2a+b+c)^2}{2a^2+(b+c)^2} + \frac{(2b+c+a)^2}{2b^2+(c+a)^2} + \frac{(2c+a+b)^2}{2c^2+(a+b)^2} \leq 8$$

Solution. We have already solved it in *Problem 3.3*.

PROBLEM 5.292 (USAMO 2001, problem 3). Let a, b, c be positive real numbers such that $a^2 + b^2 + c^2 + abc = 4$. Prove that

$$0 \leq ab + bc + ca - abc \leq 2$$

PROBLEM 5.293 (USAMO 2000, problem 6). Let $a_1, \dots, a_n, b_1, \dots, b_n$ be non-negative real numbers. Prove that

$$\sum_{i,j=1}^n \min\{a_i a_j, b_i b_j\} \leq \sum_{i,j=1}^n \min\{a_i b_j, a_j b_i\}$$

PROBLEM 5.294 (USAMO 1999, problem 4). Let $n > 3$ be an integer and a_1, \dots, a_n be positive real numbers such that

$$\begin{aligned} a_1 + \dots + a_n &\geq n \quad \text{and} \\ a_1^2 + \dots + a_n^2 &\geq n^2 \end{aligned}$$

Prove that $\max\{a_1, \dots, a_n\} \geq 2$.

PROBLEM 5.295 (USAMO 1996, problem 3). Let a_1, \dots, a_n be real numbers in the interval $(0, \pi/2)$ such that

$$\tan\left(a_0 - \frac{\pi}{4}\right) + \dots + \tan\left(a_1 - \frac{\pi}{4}\right) \geq n - 1$$

Prove that

$$\tan a_0 \cdots \tan a_n \geq n^{n+1}$$

Solution. See (3.4).

PROBLEM 5.296 (USAMO 1994, problem 4). Let a_1, \dots, a_n be a sequence of positive real numbers such that

$$\sum_{i=1}^n a_i^2 \geq \sqrt{n}$$

for all n . Prove that

$$\sum_{i=1}^n a_i^2 > \frac{1}{4} \left(1 + \frac{1}{2} + \dots + \frac{1}{n}\right)$$

PROBLEM 5.297 (USAMO 1993, problem 5). Let a_0, \dots, a_n be positive real numbers such that $a_{i-1}a_{i+1} \leq a_i^2$ (such a sequence is called *log concave*). Show that for $n > 1$,

$$\frac{a_0 + \dots + a_n}{n+1} \frac{a_1 + \dots + a_{n-1}}{n-1} \geq \frac{a_0 + \dots + a_{n-1}}{n} \frac{a_1 + \dots + a_n}{n}$$

§5 BIBLIOGRAPHY

- ANOVIĆ, S., IVELIĆ BRA and JOSIP PEĆARIĆ, “Generalizations of converse Jensen’s inequality and related results”, *Journal of Mathematical Inequalities*, no. 1 (2011), pp. 43-60, doi: 10.7153/jmi-05-06.
- BARNARD, S. and J. M. CHILD, *Higher Algebra*, New Academic Science, 2018.
- BECKENBACH, EDWIN F., ‘A Class of Mean Value Functions’, *The American Mathematical Monthly*, vol. LVII, no. 1 (1950), p. 1, doi: 10.2307/2305163.
- BECKENBACH, EDWIN F. and RICHARD BELLMAN, *Inequalities*, Springer, 1983.
- BELLMAN, RICHARD, “Notes on Matrix Theory–IV (An Inequality Due to Bergstrom)”, *The American Mathematical Monthly*, vol. LXII, no. 3 (1955), p. 172, doi: 10.2307/2306621.
- BERGSTRÖM, H, ‘A triangle-inequality for matrices’, *Den Elfte Skandinaviske Matematikerkongress, Trondheim* (1949), pp. 264-267.
- BUNYAKOVSKY, VIKTOR YAKOVLEVICH, *Foundations of the mathematical theory of probabilities*, Imperial Academy of Sciences of St. Petersburg, 1846, p. 495, doi: <http://books.e-heritage.ru/book/10070419>.
- “Sur quelques inégalités concernant les intégrales ordinaires et les intégrales aux différences finies”, *Mémoires de l’Acad. de St.-Pétersbourg*, 7th ser., vol. I, no. 9 (1859), pp. 1-18.
- CAUCHY, AUG.-LOUIS, *Cours d’Analyse de l’école Royale Polytechnique*, L’Imprimerie Royale, Debure frères, Libraires du Roi et de la Bibliothèque du Roi, 1821.
- CIRTOAJE, VASILE, “The Equal Variable Method”, *J. Inequal. Pure Appl. Math.*, vol. viii, no. 1 (2007), pp. 1-21, doi: <https://www.emis.de/journals/JIPAM/article828.html#>.
- DONGPHD and SUUGAKU, *APMO 1989–2009 Problems and Solutions*, vnmath.com, 2009.
- DUŠAN, DJUKIĆ, JANKOVIĆ VLADIMIR, and IVAN MATIC, *The IMO compendium: a collection of problems suggested for the International Mathematical Olympiads: 1959-2009*, Springer, 2011.

- GUHA, U. C., “Inequalities leading to generalisations of Schur’s inequality”, *The Mathematical Gazette*, vol. XLVI, no. 357 (1962), pp. 227-229, doi: 10.2307/3614027.
- HARDY, G. H., J. E. LITTLEWOOD, and PÓLYA G., *Inequalities*, Cambridge University Press, 1934, ISBN: 0-521-35880-9.
- HÖLDER, O., “Ueber einen Mittelwertsatz”, German, *Gött. Nachr.*, vol. MDCCCLXXXIX (1889), pp. 38-47.
- HUNG, PHAM KIM., *Secrets in inequalities*, GIL Publishing House, 2007.
- JENSEN, JOHAN LUDWIG WILLIAM VALDEMAR, “On konvexe funktioner og uligheder mellem middlvaerdier”, *Nyt. Tidsskr. for Math.*, vol. 16B (1905), pp. 49-69.
- “Sur les fonctions convexes et les inégalités entre les valeurs moyennes”, *Acta Math.*, vol. XXX (1906), pp. 175-193, doi: 10.1007/bf02418571.
- KARAMATA, JOVAN, “Sur une inégalité relative aux fonctions convexes”, French, *Publ. Math. Univ. Belgrade*, vol. I (1932), pp. 145-148.
- KNUDSEN, MATHIAS TEJS, “The UVW-method”, *Art of problem solving* (), doi: <https://artofproblemsolving.com/community/q2h278791p1507763>, <https://artofproblemsolving.com/community/q2h278791p1507763>.
- LI, KIN-YIN, “Using Tangent Lines to Prove Inequalities”, *Mathematical Excalibre*, vol. x, no. 5 (2006), p. 1.
- LJAPUNOV, ALEKSANDR MICHAJLOVIČ, “Nouvelle forme du théorème sur la limite de probabilité”, *Mémoires de l’Académie impériale des sciences de Saint-Pétersbourg*, vol. Série VIII, Tome. 12 (1901), <http://www.sudoc.fr/170960269>.
- MINKOWSKI, HERMANN, *Geometrie der Zahlen*, Johnson Reprint Corporation, 1968.
- MITRINOVITCH, D. S., “Equivalence of two sets of inequalities”, *The Mathematical Gazette*, vol. XLIII, no. 344 (1959), pp. 126-126, doi: 10.2307/3610201.
- MUIRHEAD, R. F., “Some Methods applicable to Identities and Inequalities of Symmetric Algebraic Functions of n Letters”, *Proceedings of the Edinburgh Mathematical Society*, vol. XXI (1902), pp. 144-162, doi: 10.1017/s001309150003460x.
- NEVILLE, E. H., “Schur’s inequality”, *The Mathematical Gazette*, vol. XL, no. 333 (1956), pp. 216-216, doi: 10.2307/3608826.
- OPPENHEIM, A., “2739. Generalisations of Schur’s inequality”, *The Mathematical Gazette*, vol. XLII, no. 339 (1958), pp. 35-35, doi: 10.2307/3608352.
- OPPENHEIM, A. and Roy O. DAVIES, “Inequalities of Schur’s Type”, *The Mathematical Gazette*, vol. XLVIII, no. 363 (1964), pp. 25-27, doi: 10.2307/3614303.

- PARIS, J. B. and A. VENCOVSKÁ, “A Generalization of Muirhead’s Inequality”, *Journal of Mathematical Inequalities*, no. 2 (2009), pp. 181-187, doi: 10.7153/jmi-03-18.
- RADON, JOHANN, “Über die absolut additiven Mengenfunktionen”, *Wien. Sitz. (IIa)*, vol. cxxii (1913), pp. 1295-1438.
- RAZMINIA, K., “Hölder’s inequality revisited”, *The Mathematical Gazette*, vol. ciii, no. 558 (2019), pp. 512-514, doi: 10.1017/mag.2019.117.
- ROZENBERG, MICHAEL, “uvw-Method in Proving Inequalities”, *Math. Ed.*, vol. 59-60, no. 3-4 (2011), pp. 6-14, doi: [http://www.mathnet.ru/php/archive.phtml?wshow=paper& jrnid=mo& paperid=194& option_lang=eng](http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=mo&paperid=194&option_lang=eng).
- SASSER, D.W. and M.L. SLATER, “On the inequality $\sum x_i y_i \geq \frac{1}{n} \sum x_i \sum y_i$ and the van der Waerden permanent conjecture”, *Journal of Combinatorial Theory*, vol. iii, no. 1 (1967), pp. 25-33, doi: 10.1016/s0021-9800(67)80012-7.
- SEITZ, JIŘÉ, “Une remarque aux inégalités”, *Aktuárské v dy*, vol. vi, no. 4 (1936), pp. 167-171.
- STEELE, JOHN MICHAEL, *The Cauchy-Schwarz master class: an introduction to the art of mathematical inequalities*, Cambridge University Press, 2010.
- VORNICU, VALENTIN, *Olimpiada de matematica: de la provocare la experienta*, GIL Publishing House, 2003, vol. v.
- WATSON, G. N., “A trivial inequality”, *The Mathematical Gazette*, vol. xl, no. 334 (1956), pp. 288-288, doi: 10.2307/3609631.
- “Schur’s Inequality”, *The Mathematical Gazette*, vol. xxxix, no. 329 (1955), pp. 207-208, doi: 10.2307/3608749.
- WRIGHT, E. M., “A generalisation of Schur’s inequality”, *The Mathematical Gazette*, vol. xl, no. 333 (1956), pp. 217-217, doi: 10.2307/3608827.
- YANG, KECHANG, “A Note and Generalization of a Fractional Inequality”, *Journal of Yueyang Normal University (Natural Science Edition)*, vol. xv, no. 4 (2002), pp. 9-11, doi: https://caod.oriprobe.com/articles/18339537/A_Note_and_Generalization_of_a_Fractional_Inequali.htm.
- YONGTAO, LI, GU XIAN-MING, and JIANCI XIAO, “A note on the proofs of generalized Radon inequality”, *Mathematica Moravica*, vol. xxii, no. 2 (2018), pp. 59-67, doi: 10.5937/matmor18020591.
- YOUNG, WILLIAM HENRY, “On classes of summable functions and their Fourier Series”, *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, vol. lxxxvii, no. 594 (1912), pp. 225-229, doi: 10.1098/rspa.1912.0076.

GLOSSARY

ABMO Albania-Balkan Mathematical Olympiad 70

AlMO Albania Mathematical Olympiad 70

AMC Abel's Mathematical Contest (Norwegian Mathematical Olympiad) 69

APC Austrian-Polish Competition 69

APMO Asian Pacific Mathematical Olympiad 69

AzNO Azerbaijan National Olympiad 69

BkMO Balkan Mathematical Olympiad 69

BMO British Mathematical Olympiad 69

BNO Brazil National Olympiad 70

BrNO Belarusian National Olympiad 69

BuNO Bulgarian National Olympiad 69

ChMO China Mathematical Olympiad 69

Elementary polynomials All symmetric polynomials in x_1, \dots, x_n can be written as a sum of *elementary polynomials*. Elementary polynomials E_k

are defined as

$$\begin{aligned}
 E_0(x_1, \dots, x_n) &= 1 \\
 E_1(x_1, \dots, x_n) &= x_1 + \dots + x_n \\
 E_2(x_1, \dots, x_n) &= x_1x_2 + x_1x_3 + \dots + x_{n-1}x_n \\
 &= \sum_{1 \leq i < j \leq n} x_i x_j \\
 &\vdots \\
 E_k(x_1, \dots, x_n) &= \sum_{1 \leq i_1 < \dots < i_n \leq n} x_{i_1} \cdots x_{i_n}
 \end{aligned}$$

If $k > n$, then $E_k(x_1, \dots, x_n) = 0$. A special case which is very useful for us is

$$\begin{aligned}
 E_1(a, b, c) &= a + b + c \\
 E_2(a, b, c) &= ab + bc + ca \\
 E_3(a, b, c) &= abc
 \end{aligned}$$

We can show that any symmetric polynomial can be written as a sum of elementary polynomials using induction. The main idea behind this is to show that

$$\sigma_k = x_1^k + \dots + x_n^k$$

can be represented in terms of E_0, \dots, E_k . For $k \in \{0, 1\}$, the proof is obvious. 60

Global maximum and minimum A *global minimum (absolute minimum)* and a *global maximum* (resp. *absolute maximum*) is the smallest possible value of a set or a function. 59

IMO International Mathematical Olympiad 69

IrMO Iranian Mathematical Olympiad 70

MMO Mediterranean Mathematical Olympiad 70

PMO Poland Mathematical Olympiad 70

RNO Romania National Olympiad 69

SgMO Singapore Mathematical Olympiad 69

TNO Taiwan National Olympiad 69

USAMO United States of America Mathematical Olympiad 69

INDEX

- arithmetic mean, 2
Arithmetico-Geometric Inequality, 2, 24, 35, 59, 63
Bergström's inequality, 29
Bernoulli inequality, 21
Cauchy-Bunyakovsky-Schwarz, 13
Cauchy-Schwarz, 13
converse of Hölder's inequality, 24
Engel form of Cauchy-Schwarz, 29
Generalized Bernoulli inequality, 21
Generalized Hölder's inequality, 23
geometric mean, 2
Gram's inequality, 18
Hölder's inequality, 23
harmonic mean, 2
inner product, 14
Jensen convex, 26
Jordan's inequality, 12
Karamata's inequality, 43
Lagrange Identity, 18
linearly dependent, 19
linearly independent, 19
Ljapunov's inequality, 24
majorization, 40
Minkowski's inequality, 24
monotone function, 11
mother of all inequality, 1
Muirhead's inequality, 44
Nesbitt's inequality, 6, 9, 31
norm of vector, 15
Power mean, 21
Radon's inequality, 54
Redheffer's inequality, 13
Schur's inequality, 51
symmetrical mean, 42
The buffalo way, 36
the bunching principle, 44
Titu's lemma, 29
triangle inequality, 12
Variable reduction, 8
weighted arithmetic-geometric mean inequality, 22
weighted power mean, 22
Young's inequality, 23