

Fully Radical Solution of the Quartic Equation over $\mathbb{C}$

LiZhi Zhou

Project artifact: <https://github.com/math-proof/lemma>

Quartic: [Lean 4](#)

Abstract

Ferrari–Cardano formulas are classical, but a fully explicit *principal-branch* radical map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ is subtle in formal libraries, because principal roots generally do not satisfy $\sqrt{z}\sqrt{w} = \sqrt{zw}$ or $\sqrt[3]{z}\sqrt[3]{w} = \sqrt[3]{zw}$. Consequently, Cardano’s pairing $3AB = -p$ is not automatic for two independent principal cube roots.

We formalize and machine-check a total principal-radical quartic solution in Lean 4: for every $(a, b, c, d, e) \in \mathbb{C}^5$ with $a \neq 0$, the roots are exactly a four-element set of nested principal radicals. The branch choice is a single power ω^k , with

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil,$$

so no explicit case split on $k \bmod 3$ is required. The primitive cube root is written as the radical $\omega = -\frac{1}{2} + i\sqrt{3}/2$ (the form $e^{2\pi i/3}$ is used only as notation).

Artifact: Lean theorem [Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0](#) [12] and source repository [13].

1 Introduction

Let $a, b, c, d, e, x \in \mathbb{C}$ with $a \neq 0$ and

$$ax^4 + bx^3 + cx^2 + dx + e = 0.$$

Ferrari (16th century) translates to a depressed quartic $z^4 + \alpha z^2 + \beta z + \gamma = 0$, produces a cubic resolvent, and factors the quartic into two quadratics once a resolvent root is known [2]. Cardano solves that cubic by cube roots. Over $\mathbb{C}$, Mathlib’s $\sqrt{z}$ and $\sqrt[3]{z}$ [18, 16] are defined by the principal logarithm (argument in $(-\pi, \pi]$). They satisfy

$$(\sqrt{z})^2 = z, \quad (\sqrt[3]{z})^3 = z$$

for every z , but they do *not* satisfy

$$\sqrt{z}\sqrt{w} = \sqrt{zw}, \quad \sqrt[3]{z}\sqrt[3]{w} = \sqrt[3]{zw}$$

in general. Any fully radical formula over $\mathbb{C}$ must therefore name the missing cube root of unity explicitly and select a consistent Cardano pair.

This paper records the public Lean 4 [7] lemma [Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0](#) [12]: an if-and-only-if: the equation holds precisely when x lies in the four-element set of nested principal radicals selected by whether $\beta = 0$. The cube-root branch is $y = A\omega^k + B$, not three residue classes of an integer modulo 3. Cube roots in the statement are Lean $\sqrt[3]{\cdot}$ (`Root.cubic`), not a bare `cpow` written as $z^{(3:\mathbb{C})^{-1}}$.

The source file is [Lemma/Complex/Equ0Add_Mul_Pow_4/is/In_Ite_FinsetSSubS__SubS/of/Ne_0.lean](#) [13]. Companion statements in the same library—linked only, without their solution methods—are the quadratic [15] and the cubic [14].

1.1 Main contributions

This paper contributes the following concrete artifacts.

1. A total principal-radical quartic map over $\mathbb{C}$: for every coefficient tuple with $a \neq 0$, the theorem returns a four-element radical solution set.
2. A branch-consistent Cardano pairing under principal cube roots, encoded by a single selector ω^k instead of a manual split on $k \bmod 3$.
3. A Lean 4 if-and-only-if theorem, publicly linked with its dependency graph.
4. A fully explicit radical form for ω , namely $-\frac{1}{2} + i\sqrt{3}/2$, compatible with principal-branch reasoning used in the formal proof.

1.2 Reproducibility

All statements are public and executable from the `lemma` repository [13]. The main Lean object is `Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0` [12]. For a camera-ready release, we recommend pinning the exact repository commit hash and Lean/mathlib toolchain revision in this section.

2 Related work

Mathlib 4. Zhu’s archive formalization [20] (PR #18290; Wiedijk items 37 and 46 [19]) proves Ferrari over a field K of characteristic $\neq 2$. Its statement is algebraic: square roots and a resolvent root are provided as data (e.g. $t^2 = \dots$, $v^2 = \dots$, with root parameter u). As noted in the source commentary, a fully expanded Cardano expression is intentionally omitted [21]. Hence it is not a principal-radical map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ specialized to Mathlib’s complex branches [16].

Dyson–Ahrens–Emmenegger [8] (arXiv:2201.00255) formalize quadratic/cubic/quartic formulas in Lean 3 [6] over fields of characteristic $\neq 2, 3$, using a typeclass that assumes only $(\sqrt{z})^2 = z$ and $(\sqrt[3]{z})^3 = z$. Because multiplicative branch laws are not assumed, their cubic uses one computed cube root and reconstructs its Cardano partner by division (Proposition 1.1). Their quartic (Proposition 1.3) is not total: it excludes several vanishing regimes and solves via a resolvent followed by two quadratics (Brookfield style [1]), rather than a single explicit principal-radical map. So even at $\mathbb{C}$, this does not directly yield a total principal-branch map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ with one selector $y = A\omega^k + B$.

Coq. Chardard’s `cardan_ferrari.v` [5] (discussed in [8]) is a $\mathbb{C}$ -specific Coq formalization using De Moivre roots. It is a complex radical development, but not in Lean 4 and not expressed in Mathlib’s principal `sqrt/Root.cubic` branch framework.

Other proof assistants. Isabelle/HOL `Cubic.Quartic` [4], Harrison’s HOL Light quartic [9], Carneiro’s Metamath quartic [3], and Riccardi’s Mizar development [17] appear on Wiedijk’s list [19]. They are algebraic in spirit: roots are assumed or constructed without the specific principal-branch selector mechanism used here.

This library already has companion quadratic [15] and cubic [14] lemmas over $\mathbb{C}$. The quartic contribution is to finish Ferrari on that principal-radical calculus: the resolvent is expanded by Cardano, one legal root $y = A\omega^k + B$ is selected, and the four quartic roots are nested principal square roots of that y .

3 The formula

Write $a' = b/a$, $b' = c/a$, $c' = d/a$, $d' = e/a$, and

$$\begin{aligned}\alpha &= b' - 3a'^2/8, \\ \beta &= a'^3/8 + c' - a'b'/2, \\ \gamma &= a'^2b'/16 + d' - 3a'^4/256 - a'c'/4.\end{aligned}$$

Then $z = x + a'/4$ satisfies $z^4 + \alpha z^2 + \beta z + \gamma = 0$.

Biquadratic regime $\beta = 0$. With $\Delta = \alpha^2 - 4\gamma$,

$$x = \pm \sqrt{(\pm \sqrt{\Delta} - \alpha)/2 - a'/4}$$

(four independent choices of the two signs).

Ferrari regime $\beta \neq 0$. Form Cardano quantities of the resolvent,

$$\begin{aligned}p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, \quad B = \sqrt[3]{(-q - \sqrt{\delta})/2},\end{aligned}$$

and

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil.$$

The abbreviation $\omega = e^{2\pi i/3}$ (`Lean (I*(2*pi/3)).exp`) is for simplicity only: a complex exponential is *not* a radical. The same primitive cube root of unity is the principal-radical rectangular form

$$\omega = -\frac{1}{2} + i \frac{\sqrt{3}}{2}$$

(`Lean (-(1 / 2 : ℝ)) + I * (√3 / 2 : ℝ)`), and $\bar{\omega} = -\frac{1}{2} - i \frac{\sqrt{3}}{2}$. The identities $e^{2\pi i/3} = -\frac{1}{2} + i\sqrt{3}/2$ and its conjugate are machine-checked [10, 11]. Set

$$y = A\omega^k + B, \quad y_0 = -2\alpha/3 + y, \quad y_1 = 4\alpha/3 + y.$$

Then

$$\begin{aligned}x &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \\ \text{or} \quad &\frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4}.\end{aligned}$$

Theorem 3.1 (Principal-radical quartic [12]). Assume $a \neq 0$. With the quantities above,

$$ax^4 + bx^3 + cx^2 + dx + e = 0$$

if and only if x equals one of the four biquadratic expressions when $\beta = 0$, and one of the four Ferrari expressions with $y = A\omega^k + B$ when $\beta \neq 0$. The cube-root branch is the single factor ω^k , not a case split on $k \bmod 3$.

The `if` selects the matching four-element set; it does not list Ferrari radicals as roots when $\beta = 0$, nor biquadratic radicals when $\beta \neq 0$.

4 Figures

Figure 0 — public visualization. The Lean 4 page [12] displays `@[main, comm, mp, mpr]` private lemma `main` with $\{x, a, b, c, d, e : \mathbb{C}\}$, the hypothesis $a \neq 0$, the `let` chain $a', b', \alpha, \beta, \gamma$ and the inner `if lets`, and the if-and-only-if $ax^4 + \dots + e = 0$ iff x belongs to the selected four-element set.

Figure 1 — reduction. Figure 1 is the reduction used in the main theorem: divide by the leading coefficient, depress by $z = x + a'/4$, then split on $\beta = 0$ versus $\beta \neq 0$.

$$\begin{array}{c}
 ax^4 + bx^3 + cx^2 + dx + e = 0, \quad a \neq 0 \\
 \downarrow a' = \frac{b}{a}, b' = \frac{c}{a}, c' = \frac{d}{a}, d' = \frac{e}{a} \\
 x^4 + a'x^3 + b'x^2 + c'x + d' = 0 \\
 \downarrow z = x + \frac{a'}{4} \\
 z^4 + \alpha z^2 + \beta z + \gamma = 0 \\
 \alpha = b' - \frac{3a'^2}{8}, \quad \beta = \frac{a'^3}{8} + c' - \frac{a'b'}{2}, \quad \gamma = \frac{a'^2b'}{16} + d' - \frac{3a'^4}{256} - \frac{a'c'}{4} \\
 \begin{array}{cc}
 \beta = 0 & \beta \neq 0 \\
 \downarrow & \downarrow \\
 \Delta = \alpha^2 - 4\gamma & y = A\omega^k + B
 \end{array} \\
 \begin{array}{cc}
 x = \pm \sqrt{\frac{\pm\sqrt{\Delta} - \alpha}{2}} - \frac{a'}{4} & \begin{array}{l} y_0 = -\frac{2\alpha}{3} + y, \quad y_1 = \frac{4\alpha}{3} + y \\ \pm \sqrt{\pm \frac{2\beta}{\sqrt{y_0}} - y_1 \pm \sqrt{y_0}} \\ x = \frac{\pm \sqrt{\pm \frac{2\beta}{\sqrt{y_0}} - y_1 \pm \sqrt{y_0}}}{2} - \frac{a'}{4} \\ x = z - \frac{a'}{4} \end{array}
 \end{array}
 \end{array}$$

Figure 1: Reduction of the general quartic and the two radical regimes.

Figure 2 — lemma dependencies. Figure 2 lists major imported lemmas of Theorem 3.1. Links point to the Lean 4 pages.

- [Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS_SubS.of.Ne_0](#) — main theorem
 - [Eq0Add_Pow_4.is.In_Ite_FinsetSSubS_SubS](#) — monic offset
 - [Eq0Add_Pow_4.is.In_Ite_FinsetS](#) — depressed
 - * [Eq0Add_Pow_4.is.In_FinsetSqrtS_NegSSqrt](#) — case $\beta = 0$
 - [Eq0Add_Mul_Square.is.In_FinsetDivS_Mul2.of.Ne_0](#) — quadratic in x^2
 - [EqSquare.is.In_FinsetSqrt_NegSqrt](#) — $x^2 = c \Rightarrow x = \pm\sqrt{c}$ ([EqSquareSqrt](#), [Real.OrEqS.of.Square](#))
 - * [Eq0Add_Pow_4.is.In_FinsetDivS_2DivS_2.of.Ne_0](#) — case $\beta \neq 0$
 - [Eq0Add_Pow_3.is.In_Finset_SubSAddMulS](#) — cubic resolvent
 - ([Eq_Mul_Pow_SubCeilS.of.Pow_3](#): [ArgPow.eq.SubMul_Arg](#), [Eq_MulNorm_ExpMulIArg](#), [ArgExpMulI.eq.Sub_Mul_Ceil](#), [Int.Floor.eq.NegCeilNeg](#))
 - [Eq_SquareAddMulSqrt.of.Eq0SubSquare_MulMul4.Ne_0](#) — $\Delta = 0 \Rightarrow ax^2 + bx + c$ is a square

Figure 2: Import graph for the Lean proof of Theorem 3.1.

The depressed statement [Eq0Add_Pow_4.is.In_Ite_FinsetS](#) and the monic offset [Eq0Add_Pow_4.is.In_Ite](#) are imported (Figure 2). See also the general quadratic [15] and the general cubic [14]. The

cubic Cardano proof calls `Eq_Mul_Pow_SubCeilS.of.Pow_3`, because Ferrari needs one resolvent root $A\omega^k + B$.

Figure 3 — branch integer k . From $A^3 = B^3$ one has $A = B\omega^k$ with

$$k = \left\lceil \frac{3 \arg A}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg B}{2\pi} - \frac{1}{2} \right\rceil.$$

For $(AB)^3 = (-p/3)^3$ the general selector is the negative of the k in Theorem 3.1 (the statement writes $\arg(-p)$, which equals $\arg(-p/3)$), and ω^{-k} restores $AB = (-p/3)\omega^{-k}$. Multiplying only A by ω^k produces a legal Ferrari parameter $y = A\omega^k + B$. Because $\omega^3 = 1$, the three residue classes of k modulo 3 are already encoded in that single power: $\omega^k \in \{1, \omega, \bar{\omega}\}$.

5 Proof outline

5.1 Principal radicals

Throughout, $\sqrt{\cdot}$ is `Mathlib.Complex.sqrt` and $\sqrt[3]{\cdot}$ is `Lean.Root.cubic`. Both are total functions $\mathbb{C} \rightarrow \mathbb{C}$. The identity $(\sqrt{z})^2 = z$ is `Complex.EqSquareSqrt`. If $x^2 = c$, then $x = \sqrt{c}$ or $x = -\sqrt{c}$ by `Complex.Or_EqNegSqrt.of.EqSquare`. Cube roots of unity are the radicals

$$\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}, \quad \bar{\omega} = -\frac{1}{2} - i\frac{\sqrt{3}}{2}, \quad \omega^2 = \bar{\omega}, \quad \omega^3 = 1.$$

The polar form $z = \|z\|e^{i\arg z}$ is `Complex.Eq_MulNorm_ExpMulIArg`. Reduction of $\arg(e^{ix})$ into $(-\pi, \pi]$ by a ceiling is `Complex.ArgExpMulI.eq.Sub_Mul_Ceil`, which yields `Complex.ArgPow.eq.SubMul_Arg` and then `Complex.Eq_Mul_Pow_SubCeilS.of.Pow_3`: if $A^3 = B^3$, then $A = B\omega^k$ for the integer k of Figure 3.

5.2 Depression

Division by $a \neq 0$ produces the monic equation $x^4 + a'x^3 + b'x^2 + c'x + d' = 0$. The translation $z = x + a'/4$ cancels the cubic term and yields $z^4 + \alpha z^2 + \beta z + \gamma = 0$ with α, β, γ as in §3. The main file performs this expansion by `ring`; the four nested-radical expressions for z are then shifted back by $-a'/4$.

5.3 Biquadratic case $\beta = 0$

The depressed equation is $z^4 + \alpha z^2 + \gamma = 0$, a quadratic in z^2 . Lemma `Complex.Eq0Add_Pow_4.is.In_FinsetSqrt` assembles the four principal-radical roots. It first solves

$$z^2 = \frac{-\alpha \pm \sqrt{\alpha^2 - 4\gamma}}{2} = \frac{\pm\sqrt{\Delta} - \alpha}{2}, \quad \Delta = \alpha^2 - 4\gamma,$$

then splits each $z^2 = c$ by `EqSquare.is.In_FinsetSqrt_NegSqrt`. Translating $x = z - a'/4$ is the $\beta = 0$ branch of Theorem 3.1.

5.4 Ferrari factorization

Let $y_0 \in \mathbb{C}$ be any root of the cubic resolvent

$$y_0^3 + 2\alpha y_0^2 + (\alpha^2 - 4\gamma)y_0 - \beta^2 = 0,$$

and set $y_1 = y_0 + 2\alpha$. A private helper proves $y_0 \neq 0$ (else $\beta = 0$), that $(\sqrt{y_0})^2 = y_0$, and that with $Y = (y_0 + \alpha)/2$ one has

$$(z^2 + Y)^2 - (\sqrt{y_0}z - \beta/(2\sqrt{y_0}))^2 = z^4 + \alpha z^2 + \beta z + \gamma.$$

The left-hand side is a product of two quadratics. Each is solved by completing the square and `EqSquare.is.InFinsetSqrt.NegSqrt`, producing

$$\begin{aligned} z &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, \\ z &= \frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}. \end{aligned}$$

This step still treats y_0 as given. The rest of the proof constructs y_0 from principal radicals.

5.5 Cardano data of the resolvent

The theorem statement writes the scaled Cardano pair directly:

$$\begin{aligned} p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, & B &= \sqrt[3]{(-q - \sqrt{\delta})/2}. \end{aligned}$$

The proof also depresses the resolvent by

$$a_r = -\alpha/2, \quad b_r = -\gamma, \quad c_r = -\beta^2/8 + \alpha\gamma/2,$$

and the unscaled Cardano parameters

$$\begin{aligned} p_c &= b_r - a_r^2/3, & q_c &= 2a_r^3/27 - a_rb_r/3 + c_r, & \delta_c &= 4p_c^3/27 + q_c^2, \\ A_c &= \sqrt[3]{(-q_c + \sqrt{\delta_c})/2}, & B_c &= \sqrt[3]{(-q_c - \sqrt{\delta_c})/2}. \end{aligned}$$

A direct calculation gives $q = 8q_c$, $\delta = 64\delta_c$, and $\sqrt{\delta} = 8\sqrt{\delta_c}$ (the last because $64 > 0$ on the principal branch). Positive real scaling of cube roots yields $A = 2A_c$ and $B = 2B_c$, and $\arg(AB) = \arg(A_cB_c)$.

5.6 Restoring $3(A\omega^k)B = -p$

Because $(AB)^3 = (-p/3)^3$, lemma `Eq_Mul_Pow_SubCeilS.of.Pow_3` supplies

$$AB = (-p/3)\omega^{-k}.$$

Multiplying by ω^k restores the product in one step: $AB\omega^k = -p/3$. Since $(\omega^k)^3 = 1$,

$$(A\omega^k)^3 + B^3 = A^3 + B^3 = -q, \quad 3(A\omega^k)B = -p,$$

so $y = A\omega^k + B$ satisfies $y^3 + py + q = 0$. There is no case split on $k \bmod 3$: the three possible rotations $1, \omega, \bar{\omega}$ are the values of the single power ω^k . Shifting $y_0 = -2\alpha/3 + y$ and $y_1 = 4\alpha/3 + y$ recovers a resolvent root, and the four nested square roots apply.

Unlike the cubic formula in this library, only A is rotated. A quartic needs one legal resolvent parameter, not three Cardano roots of the original unknown.

6 Discussion and limitations

The statement is an if-and-only-if: the equation holds precisely when x lies in the four-element set selected by `if  $\beta = 0$` . It is not a four-fold split on $k \bmod 3$. Principal square roots in the Ferrari step may vanish or repeat when the quartic has multiple roots; membership still holds.

The lemma does not construct roots in a general field of characteristic zero, does not avoid `arg` and `⌈·⌋`, and does not prove that a randomly chosen pairing of principal cube roots is a resolvent root. It does not replace Mathlib’s abstract Ferrari theorem [20]: that theorem remains the right statement when one only knows that some square roots and some resolvent root exist in K . It does not replace Dyson–Ahrens–Emmenegger [8]: that development remains the right statement for an arbitrary field with unspecified root operations and several non-vanishing hypotheses. The present lemma is the complementary statement on $\mathbb{C}$: every coefficient tuple with $a \neq 0$ is sent to four explicit principal-radical expressions, with the cube-root branch named by the power ω^k .

7 Conclusion

We have presented a machine-checked, fully expanded Ferrari–Cardano formula for the general quartic over $\mathbb{C}$, using principal square roots, principal cube roots $\sqrt[3]{\cdot}$, the radical cube root of unity $-\frac{1}{2} + i\sqrt{3}/2$, and a single power ω^k assembled from `arg` and `⌈·⌋`. The result is public on lemma.cn and GitHub [13], with proofs checked in Lean 4.

References

- [1] Gary Brookfield. Factoring quartic polynomials: A lost art. *Mathematics Magazine*, 80(1):67–70, 2007.
- [2] Girolamo Cardano. *Ars Magna, or The Rules of Algebra*. Dover, 1968. English translation of the 1545 Latin text; Ferrari’s quartic appears here.
- [3] Mario Carneiro. Proof of the quartic formula. Metamath, 2021. <https://us.metamath.org/mpeuni/quart.html>.
- [4] Amine Chaieb. `Cubic_Quartic`. Isabelle/HOL, 2021. https://isabelle.in.tum.de/dist/library/HOL/HOL-ex/Cubic_Quartic.html.
- [5] Frédéric Chardard. `cardan_ferrari.v`. Coq 100 theorems, 2021. https://github.com/coq-community/coq-100-theorems/blob/master/cardan_ferrari.v.
- [6] Leonardo de Moura, Soonho Kong, Jeremy Avigad, Floris van Doorn, and Jakob von Raumer. The Lean theorem prover (system description). In *Automated Deduction – CADE-25*, volume 9195 of *LNCS*, pages 378–388. Springer, 2015.
- [7] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *LNCS*, pages 625–635. Springer, 2021.
- [8] Nicholas Dyson, Benedikt Ahrens, and Jacopo Emmenegger. The solutions to single-variable polynomials, implemented and verified in Lean. <https://arxiv.org/abs/2201.00255>, 2022.
- [9] John Harrison. Proof of the quartic formula. HOL Light, 2021. Wiedijk item 46; <https://www.cl.cam.ac.uk/users/jrh/hol-light/>.

- [10] math proof. `Complex.expMulIDivMul2Pi3.eq.add_muli`. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMul2Pi3.eq.Add_MulI, 2026.
- [11] math proof. `Complex.expMulIDivMulNeg2Pi3.eq.sub_muli`. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMulNeg2Pi3.eq.Sub_MulI, 2026.
- [12] math proof. Fully radical quartic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0, 2026.
- [13] math proof. lemma: machine-checked mathematics. <https://github.com/math-proof/lemma>, 2026.
- [14] math proof. Principal-radical cubic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_3.is.In_Finset_SubSAddMulS.of.Ne_0, 2026.
- [15] math proof. Principal-radical quadratic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Square.is.In_FinsetDivS_Mul2.of.Ne_0, 2026.
- [16] Mathlib community. The Lean mathematical library. https://leanprover-community.github.io/mathlib4_docs/, 2026.
- [17] Marco Riccardi. Solution of cubic and quartic equations. *Formalized Mathematics*, 17(1):117–122, 2009.
- [18] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP)*, pages 367–381. ACM, 2020.
- [19] Freek Wiedijk. Formalizing 100 theorems. <https://www.cs.ru.nl/~freek/100/>, 2024.
- [20] Thomas Zhu. The roots of cubic and quartic polynomials. Mathlib 4 Archive, PR #18290, 2024. <https://github.com/leanprover-community/mathlib4/blob/master/Archive/Wiedijk100Theorems/SolutionOfCubicQuartic.lean>.
- [21] Daniel Zwillinger. *CRC Standard Mathematical Tables and Formulae*. Chapman and Hall/CRC, 31 edition, 2003.