

Fully Radical Solution of the Quartic Equation over $\mathbb{C}$

LiZhi Zhou

Project artifact: <https://github.com/math-proof/lemma>

Quartic: [Lean 4](#), [SymPy](#)

Abstract

Ferrari–Cardano formulas are classical, but a fully explicit *principal-branch* radical map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ is subtle in formal libraries, because principal roots generally do not satisfy $\sqrt{z}\sqrt{w} = \sqrt{zw}$ or $\sqrt[3]{z}\sqrt[3]{w} = \sqrt[3]{zw}$. Consequently, Cardano’s pairing $3AB = -p$ is not automatic for two independent principal cube roots.

We formalize and machine-check a total principal-radical quartic solution in Lean 4: for every $(a, b, c, d, e) \in \mathbb{C}^5$ with $a \neq 0$, the roots are exactly a four-element set of nested principal radicals. The branch choice is a single power ω^k , with

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil,$$

so no explicit case split on $k \bmod 3$ is required. The primitive cube root is written as the radical $\omega = -\frac{1}{2} + i\sqrt{3}/2$ (the form $e^{2\pi i/3}$ is used only as notation).

Artifacts: Lean theorem [Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0](#) [13], SymPy counterpart [Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0](#) [10], and source repository [14].

1 Introduction

Let $a, b, c, d, e, x \in \mathbb{C}$ with $a \neq 0$ and

$$ax^4 + bx^3 + cx^2 + dx + e = 0.$$

Ferrari (16th century) translates to a depressed quartic $z^4 + \alpha z^2 + \beta z + \gamma = 0$, produces a cubic resolvent, and factors the quartic into two quadratics once a resolvent root is known [2]. Cardano solves that cubic by cube roots. Over $\mathbb{C}$, Mathlib’s $\sqrt{z}$ and $\sqrt[3]{z}$ [21, 17] are defined by the principal logarithm (argument in $(-\pi, \pi]$). They satisfy

$$(\sqrt{z})^2 = z, \quad (\sqrt[3]{z})^3 = z$$

for every z , but they do *not* satisfy

$$\sqrt{z}\sqrt{w} = \sqrt{zw}, \quad \sqrt[3]{z}\sqrt[3]{w} = \sqrt[3]{zw}$$

in general. Any fully radical formula over $\mathbb{C}$ must therefore name the missing cube root of unity explicitly and select a consistent Cardano pair.

This paper records the public Lean 4 [7] lemma [Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0](#) [13]: an if-and-only-if: the equation holds precisely when x lies in the four-element set of nested principal radicals selected by whether $\beta = 0$. The cube-root branch is $y = A\omega^k + B$, not three residue classes of an integer modulo 3. Cube roots in the statement are Lean $\sqrt[3]{}$ ([Root.cubic](#)), not a bare `cpow` written as $z^{(3:\mathbb{C})^{-1}}$.

The SymPy [20, 18] counterpart [10] is the one-direction lemma [Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0](#) (the Lean `mp` alias); it has no `is` if-and-only-if. Both

are visualized on lemma.cn. Source files are `Lemma/Complex/Eq0Add_Mul_Pow_4/is/In_Ite_FinsetSSubS__SubS/of/Ne_0.lean` (branch `main`) and `Lemma/Complex/In_Ite_FinsetSSubS__SubS/of/Eq0Add_Mul_Pow_4/Ne_0.py` (branch `master`) [14]. Companion statements in the same library—linked only, without their solution methods—are the quadratic [16] and the cubic [15].

1.1 Main contributions

This paper contributes the following concrete artifacts.

1. A total principal-radical quartic map over $\mathbb{C}$: for every coefficient tuple with $a \neq 0$, the theorem returns a four-element radical solution set.
2. A branch-consistent Cardano pairing under principal cube roots, encoded by a single selector ω^k instead of a manual split on $k \bmod 3$.
3. A Lean 4 if-and-only-if theorem (not only one direction), publicly linked with dependency graph and companion SymPy artifact.
4. A fully explicit radical form for ω , namely $-\frac{1}{2} + i\sqrt{3}/2$, compatible with principal-branch reasoning used in the formal proof.

1.2 Reproducibility

All statements are public and executable from the `lemma` repository [14]. The main Lean object is `Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0` [13], and the SymPy analogue is `Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0` [10]. For a camera-ready release, we recommend pinning the exact repository commit hash and Lean/mathlib toolchain revision in this section.

2 Related work

Mathlib 4. Zhu’s archive formalization [23] (PR #18290; Wiedijk items 37 and 46 [22]) proves Ferrari over a field K of characteristic $\neq 2$. Its statement is algebraic: square roots and a resolvent root are provided as data (e.g. $t^2 = \dots$, $v^2 = \dots$, with root parameter u). As noted in the source commentary, a fully expanded Cardano expression is intentionally omitted [24]. Hence it is not a principal-radical map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ specialized to Mathlib’s complex branches [17].

Dyson–Ahrens–Emmenegger [8] (arXiv:2201.00255) formalize quadratic/cubic/quartic formulas in Lean 3 [6] over fields of characteristic $\neq 2, 3$, using a typeclass that assumes only $(\sqrt{z})^2 = z$ and $(\sqrt[3]{z})^3 = z$. Because multiplicative branch laws are not assumed, their cubic uses one computed cube root and reconstructs its Cardano partner by division (Proposition 1.1). Their quartic (Proposition 1.3) is not total: it excludes several vanishing regimes and solves via a resolvent followed by two quadratics (Brookfield style [1]), rather than a single explicit principal-radical map. So even at $\mathbb{C}$, this does not directly yield a total principal-branch map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ with one selector $y = A\omega^k + B$.

Coq. Chardard’s `cardan_ferrari.v` [5] (discussed in [8]) is a $\mathbb{C}$ -specific Coq formalization using De Moivre roots. It is a complex radical development, but not in Lean 4 and not expressed in Mathlib’s principal $\sqrt{\cdot}/\text{Root.cubic}$ branch framework.

Other proof assistants. Isabelle/HOL `Cubic.Quartic` [4], Harrison’s HOL Light quartic [9], Carneiro’s Metamath quartic [3], and Riccardi’s Mizar development [19] appear on Wiedijk’s list [22]. They are algebraic in spirit: roots are assumed or constructed without the specific principal-branch selector mechanism used here.

This library already has companion quadratic [16] and cubic [15] lemmas over $\mathbb{C}$. The quartic contribution is to finish Ferrari on that principal-radical calculus: the resolvent is expanded by

Cardano, one legal root $y = A\omega^k + B$ is selected, and the four quartic roots are nested principal square roots of that y .

3 The formula

Write $a' = b/a$, $b' = c/a$, $c' = d/a$, $d' = e/a$, and

$$\begin{aligned}\alpha &= b' - 3a'^2/8, \\ \beta &= a'^3/8 + c' - a'b'/2, \\ \gamma &= a'^2b'/16 + d' - 3a'^4/256 - a'c'/4.\end{aligned}$$

Then $z = x + a'/4$ satisfies $z^4 + \alpha z^2 + \beta z + \gamma = 0$.

Biquadratic regime $\beta = 0$. With $\Delta = \alpha^2 - 4\gamma$,

$$x = \pm \sqrt{(\pm \sqrt{\Delta} - \alpha)/2 - a'/4}$$

(four independent choices of the two signs).

Ferrari regime $\beta \neq 0$. Form Cardano quantities of the resolvent,

$$\begin{aligned}p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, \quad B = \sqrt[3]{(-q - \sqrt{\delta})/2},\end{aligned}$$

and

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil.$$

The abbreviation $\omega = e^{2\pi i/3}$ (Lean `(I*(2*pi/3)).exp`) is for simplicity only: a complex exponential is *not* a radical. The same primitive cube root of unity is the principal-radical rectangular form

$$\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$$

(Lean `-(1 / 2 : ℝ) + I * (√3 / 2 : ℝ)`), and $\bar{\omega} = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$. The identities $e^{2\pi i/3} = -\frac{1}{2} + i\sqrt{3}/2$ and its conjugate are machine-checked [11, 12]. Set

$$y = A\omega^k + B, \quad y_0 = -2\alpha/3 + y, \quad y_1 = 4\alpha/3 + y.$$

Then

$$\begin{aligned}x &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \\ \text{or} \quad &\frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4}.\end{aligned}$$

Theorem 3.1 (Principal-radical quartic [13]). Assume $a \neq 0$. With the quantities above,

$$ax^4 + bx^3 + cx^2 + dx + e = 0$$

if and only if x equals one of the four biquadratic expressions when $\beta = 0$, and one of the four Ferrari expressions with $y = A\omega^k + B$ when $\beta \neq 0$. The cube-root branch is the single factor ω^k , not a case split on $k \bmod 3$.

The `if` selects the matching four-element set; it does not list Ferrari radicals as roots when $\beta = 0$, nor biquadratic radicals when $\beta \neq 0$.

4 Figures

Figure 0 — public visualizations. The SymPy page [10] displays the one-direction (mp) statement `In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0` as an interactive theorem document (created 2018-11-29). The Lean 4 page [13] displays `@[main, comm, mp, mpr] private lemma main` with $\{x, a, b, c, d, e : \mathbb{C}\}$, the hypothesis $a \neq 0$, the `let` chain $a', b', \alpha, \beta, \gamma$ and the inner `if` lets, and the if-and-only-if $ax^4 + \dots + e = 0$ iff x belongs to the selected four-element set.

Figure 1 — reduction. Figure 1 is the reduction used in the main theorem: divide by the leading coefficient, depress by $z = x + a'/4$, then split on $\beta = 0$ versus $\beta \neq 0$.

$$\begin{array}{c}
 ax^4 + bx^3 + cx^2 + dx + e = 0, \quad a \neq 0 \\
 \downarrow a' = \frac{b}{a}, b' = \frac{c}{a}, c' = \frac{d}{a}, d' = \frac{e}{a} \\
 x^4 + a'x^3 + b'x^2 + c'x + d' = 0 \\
 \downarrow z = x + \frac{a'}{4} \\
 z^4 + \alpha z^2 + \beta z + \gamma = 0 \\
 \alpha = b' - \frac{3a'^2}{8}, \quad \beta = \frac{a'^3}{8} + c' - \frac{a'b'}{2}, \quad \gamma = \frac{a'^2b'}{16} + d' - \frac{3a'^4}{256} - \frac{a'c'}{4} \\
 \begin{array}{cc}
 \beta = 0 & \beta \neq 0 \\
 \downarrow & \downarrow \\
 \Delta = \alpha^2 - 4\gamma & y = A\omega^k + B
 \end{array} \\
 \begin{array}{cc}
 x = \pm \sqrt{\frac{\pm\sqrt{\Delta} - \alpha}{2}} - \frac{a'}{4} & \begin{array}{l} y_0 = -\frac{2\alpha}{3} + y, \quad y_1 = \frac{4\alpha}{3} + y \\ \pm \sqrt{\frac{2\beta}{\sqrt{y_0}}} - y_1 \pm \sqrt{y_0} \\ x = \frac{\pm \sqrt{\frac{2\beta}{\sqrt{y_0}}} - y_1 \pm \sqrt{y_0}}{2} - \frac{a'}{4} \\ x = z - \frac{a'}{4} \end{array}
 \end{array}
 \end{array}$$

Figure 1: Reduction of the general quartic and the two radical regimes.

Figure 2 — lemma dependencies. Figure 2 lists major imported lemmas of Theorem 3.1. Links point to the Lean 4 pages. SymPy has only the one-direction (of/mp) names, e.g. http://www.lemma.cn/py/?module=Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0, not the Lean `is` if-and-only-if.

The depressed statement `Eq0Add_Pow_4.is.In_Ite_FinsetS` and the monic offset `Eq0Add_Pow_4.is.In_Ite` are imported (Figure 2). See also the general quadratic [16] and the general cubic [15]. The cubic Cardano proof calls `Eq_Mul_Pow_SubCeilS.of.Pow_3`, because Ferrari needs one resolvent root $A\omega^k + B$.

Figure 3 — branch integer k . From $A^3 = B^3$ one has $A = B\omega^k$ with

$$k = \left\lceil \frac{3 \arg A}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg B}{2\pi} - \frac{1}{2} \right\rceil.$$

For $(AB)^3 = (-p/3)^3$ the general selector is the negative of the k in Theorem 3.1 (the statement writes $\arg(-p)$, which equals $\arg(-p/3)$), and ω^{-k} restores $AB = (-p/3)\omega^{-k}$. Multiplying only A by ω^k produces a legal Ferrari parameter $y = A\omega^k + B$. Because $\omega^3 = 1$, the three residue classes of k modulo 3 are already encoded in that single power: $\omega^k \in \{1, \omega, \bar{\omega}\}$.

- `Eq0Add_Mul_Pow_4.is.In.Ite.FinsetSSubS_SubS.of.Ne_0` — main theorem
 - `Eq0Add_Pow_4.is.In.Ite.FinsetSSubS_SubS` — monic offset
 - `Eq0Add_Pow_4.is.In.Ite.FinsetS` — depressed
 - * `Eq0Add_Pow_4.is.In.FinsetSqrtS_NegSSqrt` — case $\beta = 0$
 - `Eq0Add_Mul_Square.is.In.FinsetDivS_Mul2.of.Ne_0` — quadratic in x^2
 - `EqSquare.is.In.FinsetSqrt_NegSqrt` — $x^2 = c \Rightarrow x = \pm\sqrt{c}$ (`EqSquareSqrt`, `Real.OrEqS.of.Square`)
 - * `Eq0Add_Pow_4.is.In.FinsetDivS_2DivS_2.of.Ne_0` — case $\beta \neq 0$
 - `Eq0Add_Pow_3.is.In.Finset_SubSAddMulS` — cubic resolvent
 - (`Eq_Mul_Pow_SubCeilS.of.Pow_3`: `ArgPow.eq.SubMul_Arg`, `Eq_MulNorm_ExpMulIArg`, `ArgExpMulI.eq.Sub_Mul_Ceil`, `Int.Floor.eq.NegCeilNeg`)
 - `Eq_SquareAddMulSqrt.of.Eq0SubSquare_MulMul4.Ne_0` — $\Delta = 0 \Rightarrow ax^2 + bx + c$ is a square

Figure 2: Import graph for the Lean proof of Theorem 3.1.

5 Proof outline

5.1 Principal radicals

Throughout, $\sqrt{\cdot}$ is `Mathlib.Complex.sqrt` and $\sqrt[3]{\cdot}$ is `Lean.Root.cubic`. Both are total functions $\mathbb{C} \rightarrow \mathbb{C}$. The identity $(\sqrt{z})^2 = z$ is `Complex.EqSquareSqrt`. If $x^2 = c$, then $x = \sqrt{c}$ or $x = -\sqrt{c}$ by `Complex.Or_Eq_NegSqrt.of.EqSquare`. Cube roots of unity are the radicals

$$\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}, \quad \bar{\omega} = -\frac{1}{2} - i\frac{\sqrt{3}}{2}, \quad \omega^2 = \bar{\omega}, \quad \omega^3 = 1.$$

The polar form $z = \|z\|e^{i\arg z}$ is `Complex.Eq_MulNorm_ExpMulIArg`. Reduction of $\arg(e^{ix})$ into $(-\pi, \pi]$ by a ceiling is `Complex.ArgExpMulI.eq.Sub_Mul_Ceil`, which yields `Complex.ArgPow.eq.SubMul_Arg` and then `Complex.Eq_Mul_Pow_SubCeilS.of.Pow_3`: if $A^3 = B^3$, then $A = B\omega^k$ for the integer k of Figure 3.

5.2 Depression

Division by $a \neq 0$ produces the monic equation $x^4 + a'x^3 + b'x^2 + c'x + d' = 0$. The translation $z = x + a'/4$ cancels the cubic term and yields $z^4 + \alpha z^2 + \beta z + \gamma = 0$ with α, β, γ as in §3. The main file performs this expansion by `ring`; the four nested-radical expressions for z are then shifted back by $-a'/4$.

5.3 Biquadratic case $\beta = 0$

The depressed equation is $z^4 + \alpha z^2 + \gamma = 0$, a quadratic in z^2 . Lemma `Complex.Eq0Add_Pow_4.is.In.FinsetSqrtS` assembles the four principal-radical roots. It first solves

$$z^2 = \frac{-\alpha \pm \sqrt{\alpha^2 - 4\gamma}}{2} = \frac{\pm\sqrt{\Delta} - \alpha}{2}, \quad \Delta = \alpha^2 - 4\gamma,$$

then splits each $z^2 = c$ by `EqSquare.is.In.FinsetSqrt_NegSqrt`. Translating $x = z - a'/4$ is the $\beta = 0$ branch of Theorem 3.1.

5.4 Ferrari factorization

Let $y_0 \in \mathbb{C}$ be any root of the cubic resolvent

$$y_0^3 + 2\alpha y_0^2 + (\alpha^2 - 4\gamma)y_0 - \beta^2 = 0,$$

and set $y_1 = y_0 + 2\alpha$. A private helper proves $y_0 \neq 0$ (else $\beta = 0$), that $(\sqrt{y_0})^2 = y_0$, and that with $Y = (y_0 + \alpha)/2$ one has

$$(z^2 + Y)^2 - (\sqrt{y_0}z - \beta/(2\sqrt{y_0}))^2 = z^4 + \alpha z^2 + \beta z + \gamma.$$

The left-hand side is a product of two quadratics. Each is solved by completing the square and `EqSquare.is.In.FinsetSqrt.NegSqrt`, producing

$$\begin{aligned} z &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, \\ z &= \frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}. \end{aligned}$$

This step still treats y_0 as given. The rest of the proof constructs y_0 from principal radicals.

5.5 Cardano data of the resolvent

The theorem statement writes the scaled Cardano pair directly:

$$\begin{aligned} p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, & B &= \sqrt[3]{(-q - \sqrt{\delta})/2}. \end{aligned}$$

The proof also depresses the resolvent by

$$a_r = -\alpha/2, \quad b_r = -\gamma, \quad c_r = -\beta^2/8 + \alpha\gamma/2,$$

and the unscaled Cardano parameters

$$\begin{aligned} p_c &= b_r - a_r^2/3, & q_c &= 2a_r^3/27 - a_r b_r/3 + c_r, & \delta_c &= 4p_c^3/27 + q_c^2, \\ A_c &= \sqrt[3]{(-q_c + \sqrt{\delta_c})/2}, & B_c &= \sqrt[3]{(-q_c - \sqrt{\delta_c})/2}. \end{aligned}$$

A direct calculation gives $q = 8q_c$, $\delta = 64\delta_c$, and $\sqrt{\delta} = 8\sqrt{\delta_c}$ (the last because $64 > 0$ on the principal branch). Positive real scaling of cube roots yields $A = 2A_c$ and $B = 2B_c$, and $\arg(AB) = \arg(A_c B_c)$.

5.6 Restoring $3(A\omega^k)B = -p$

Because $(AB)^3 = (-p/3)^3$, lemma `Eq_Mul_Pow_SubCeilS.of.Pow_3` supplies

$$AB = (-p/3)\omega^{-k}.$$

Multiplying by ω^k restores the product in one step: $AB\omega^k = -p/3$. Since $(\omega^k)^3 = 1$,

$$(A\omega^k)^3 + B^3 = A^3 + B^3 = -q, \quad 3(A\omega^k)B = -p,$$

so $y = A\omega^k + B$ satisfies $y^3 + py + q = 0$. There is no case split on $k \bmod 3$: the three possible rotations $1, \omega, \bar{\omega}$ are the values of the single power ω^k . Shifting $y_0 = -2\alpha/3 + y$ and $y_1 = 4\alpha/3 + y$ recovers a resolvent root, and the four nested square roots apply.

Unlike the cubic formula in this library, only A is rotated. A quartic needs one legal resolvent parameter, not three Cardano roots of the original unknown.

6 Discussion and limitations

The statement is an if-and-only-if: the equation holds precisely when x lies in the four-element set selected by `if  $\beta = 0$` . It is not a four-fold split on $k \bmod 3$. Principal square roots in the Ferrari step may vanish or repeat when the quartic has multiple roots; membership still holds.

The lemma does not construct roots in a general field of characteristic zero, does not avoid `arg` and `⌈·⌉`, and does not prove that a randomly chosen pairing of principal cube roots is a resolvent root. It does not replace Mathlib’s abstract Ferrari theorem [23]: that theorem remains the right statement when one only knows that some square roots and some resolvent root exist in K . It does not replace Dyson–Ahrens–Emmenegger [8]: that development remains the right statement for an arbitrary field with unspecified root operations and several non-vanishing hypotheses. The present lemma is the complementary statement on $\mathbb{C}$: every coefficient tuple with $a \neq 0$ is sent to four explicit principal-radical expressions, with the cube-root branch named by the power ω^k .

7 Conclusion

We have presented a machine-checked, fully expanded Ferrari–Cardano formula for the general quartic over $\mathbb{C}$, using principal square roots, principal cube roots $\sqrt[3]{\cdot}$, the radical cube root of unity $-\frac{1}{2} + i\sqrt{3}/2$, and a single power ω^k assembled from `arg` and `⌈·⌉`. The result is public on lemma.cn and GitHub [14], with proofs checked in Lean 4 and an earlier SymPy formulation.

References

- [1] Gary Brookfield. Factoring quartic polynomials: A lost art. *Mathematics Magazine*, 80(1):67–70, 2007.
- [2] Girolamo Cardano. *Ars Magna, or The Rules of Algebra*. Dover, 1968. English translation of the 1545 Latin text; Ferrari’s quartic appears here.
- [3] Mario Carneiro. Proof of the quartic formula. Metamath, 2021. <https://us.metamath.org/mpeuni/quart.html>.
- [4] Amine Chaieb. `Cubic_Quartic`. Isabelle/HOL, 2021. https://isabelle.in.tum.de/dist/library/HOL/HOL-ex/Cubic_Quartic.html.
- [5] Frédéric Chardard. `cardan_ferrari.v`. Coq 100 theorems, 2021. https://github.com/coq-community/coq-100-theorems/blob/master/cardan_ferrari.v.
- [6] Leonardo de Moura, Soonho Kong, Jeremy Avigad, Floris van Doorn, and Jakob von Raumer. The Lean theorem prover (system description). In *Automated Deduction – CADE-25*, volume 9195 of *LNCS*, pages 378–388. Springer, 2015.
- [7] Leonardo de Moura and Sebastian Ullrich. The Lean 4 theorem prover and programming language. In *Automated Deduction – CADE 28*, volume 12699 of *LNCS*, pages 625–635. Springer, 2021.
- [8] Nicholas Dyson, Benedikt Ahrens, and Jacopo Emmenegger. The solutions to single-variable polynomials, implemented and verified in Lean. <https://arxiv.org/abs/2201.00255>, 2022.
- [9] John Harrison. Proof of the quartic formula. HOL Light, 2021. Wiedijk item 46; <https://www.cl.cam.ac.uk/users/jrh/hol-light/>.

- [10] math proof. Fully radical quartic over $\mathbb{C}$ (SymPy). http://www.lemma.cn/py/?module=Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0, 2018.
- [11] math proof. Complex.expmulidivmul2pi3.eq.add_muli. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMul2Pi3.eq.Add_MulI, 2026.
- [12] math proof. Complex.expmulidivmulneg2pi3.eq.sub_muli. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMulNeg2Pi3.eq.Sub_MulI, 2026.
- [13] math proof. Fully radical quartic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0, 2026.
- [14] math proof. lemma: machine-checked mathematics. <https://github.com/math-proof/lemma>, 2026.
- [15] math proof. Principal-radical cubic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_3.is.In_Finset_SubSAddMulS.of.Ne_0, 2026.
- [16] math proof. Principal-radical quadratic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Square.is.In_FinsetDivS_Mul2.of.Ne_0, 2026.
- [17] Mathlib community. The Lean mathematical library. https://leanprover-community.github.io/mathlib4_docs/, 2026.
- [18] Aaron Meurer, Christopher P. Smith, Mateusz Paprocki, Ondřej Čertík, Sergey B. Kirpichev, Matthew Rocklin, Amit Kumar, Sergiu Ivanov, Jason K. Moore, Sartaj Singh, Thilina Rathnayake, Sean Vig, Brian E. Granger, Richard P. Muller, Francesco Bonazzi, Harsh Gupta, Shivam Vats, Fredrik Johansson, Fabian Pedregosa, Matthew J. Curry, Andy R. Terrel, Štěpán Roučka, Ashutosh Saboo, Isuru Fernando, Sumith Kulal, Robert Cimrman, and Anthony Scopatz. SymPy: symbolic computing in Python. *PeerJ Computer Science*, 3:e103, 2017.
- [19] Marco Riccardi. Solution of cubic and quartic equations. *Formalized Mathematics*, 17(1):117–122, 2009.
- [20] SymPy Development Team. SymPy. <https://www.sympy.org/>, 2026.
- [21] The mathlib Community. The Lean mathematical library. In *Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP)*, pages 367–381. ACM, 2020.
- [22] Freek Wiedijk. Formalizing 100 theorems. <https://www.cs.ru.nl/~freek/100/>, 2024.
- [23] Thomas Zhu. The roots of cubic and quartic polynomials. Mathlib 4 Archive, PR #18290, 2024. <https://github.com/leanprover-community/mathlib4/blob/master/Archive/Wiedijk100Theorems/SolutionOfCubicQuartic.lean>.
- [24] Daniel Zwillinger. *CRC Standard Mathematical Tables and Formulae*. Chapman and Hall/CRC, 31 edition, 2003.