

Fully Radical Solution of the Quartic Equation over $\mathbb{C}$

LiZhi Zhou

Project artifact: <https://github.com/math-proof/lemma>

Quartic: [Lean 4](#), [SymPy](#)

Abstract

Ferrari's method reduces a general quartic to a depressed quartic and a cubic resolvent; Cardano then solves that cubic by cube roots. Over an arbitrary field the identities are classical, and Mathlib's archived Ferrari theorem leaves the resolvent root and the square roots as extra data. What is not classical is a *total* map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ in *principal* radicals: Mathlib's $\sqrt{\cdot}$ and $\sqrt[3]{\cdot}$ (Lean `Root.cubic`, notation $\sqrt[3]{\cdot}$) fail the product law, so Cardano's pairing $3AB = -p$ does not hold for two independent principal cube roots.

We state and machine-check such a map in the `lemma` library. Every auxiliary quantity is a principal square or cube root; the missing cube root of unity is the radical $-\frac{1}{2} + i\sqrt{3}/2$ (the exponential $e^{2\pi i/3}$ is an abbreviation only); the branch is the single power ω^k with

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil,$$

not a case split on $k \bmod 3$. The proof is checked in Lean 4 (updated 2026-09-01) and was first stated interactively in SymPy (2018-11-29). The quartic statements are [Lean 4](#) [5] and [SymPy](#) [2]; source is on GitHub [6].

1 Introduction

Let $a, b, c, d, e, x \in \mathbb{C}$ with $a \neq 0$ and

$$ax^4 + bx^3 + cx^2 + dx + e = 0.$$

Ferrari (16th century) translates to a depressed quartic $z^4 + \alpha z^2 + \beta z + \gamma = 0$, produces a cubic resolvent, and factors the quartic into two quadratics once a resolvent root is known. Cardano solves that cubic by cube roots. Over $\mathbb{C}$, Mathlib's $\sqrt{z}$ and $\sqrt[3]{z}$ are defined by the principal logarithm (argument in $(-\pi, \pi]$). They satisfy

$$(\sqrt{z})^2 = z, \quad (\sqrt[3]{z})^3 = z$$

for every z , but they do *not* satisfy

$$\sqrt{z}\sqrt{w} = \sqrt{zw}, \quad \sqrt[3]{z}\sqrt[3]{w} = \sqrt[3]{zw}$$

in general. Any fully radical formula over $\mathbb{C}$ must therefore name the missing cube root of unity explicitly and select a consistent Cardano pair.

This paper records the public Lean 4 lemma `Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS_SubS.of` an if-and-only-if: the equation holds precisely when x lies in the four-element set of nested principal radicals selected by whether $\beta = 0$. The cube-root branch is $y = A\omega^k + B$, not three residue classes of an integer modulo 3. Cube roots in the statement are Lean $\sqrt[3]{\cdot}$ (`Root.cubic`), not a bare `cpow` written as $z^{(3:\mathbb{C})^{-1}}$.

The SymPy counterpart [2] is the one-direction lemma `Complex.In_Ite_FinsetSSubS_SubS.of.Eq0Add_Mul` (the Lean `mp` alias); it has no `is` if-and-only-if. Both are visualized on `lemma.cn`. Source files

are `Lemma/Complex/Eq0Add_Mul_Pow_4/is/In_Ite_FinsetSSubS__SubS/of/Ne_0.lean` (branch `main`) and `Lemma/Complex/In_Ite_FinsetSSubS__SubS/of/Eq0Add_Mul_Pow_4/Ne_0.py` (branch `master`) [6]. Companion statements in the same library—linked only, without their solution methods—are the quadratic [8] and the cubic [7].

2 Related work

Mathlib 4. Zhu’s archive formalization [11] (PR #18290; Wiedijk items 37 and 46 [10]) proves Ferrari over a field K of characteristic not two. Square roots appear as hypotheses $t^2 = \dots$, $v^2 = \dots$, and the resolvent root u is an extra datum. The comment on `quartic_eq_zero_iff` states that an explicit Cardano expression for u “would be too long.” The theorem is not a map $\mathbb{C}^5 \rightarrow \mathbb{C}^4$ in principal radicals, and it is not specific to $\mathbb{C}$ [9].

Dyson–Ahrens–Emmenegger [1] work in Lean 3 in a field equipped with *some* square-root and cube-root operations. The functions are axioms, not Mathlib’s principal branches; the product identity for cube roots is deliberately not assumed. Instantiating the typeclass with $\mathbb{C}$ does not prove the principal-branch formula.

Coq (the development compared in that paper, §7.1) is specific to $\mathbb{C}$ and uses De Moivre n th roots. That is a complex-domain radical solution, but it is not a Lean 4 lemma and does not use Mathlib’s principal $\sqrt{\cdot}$ / `Root.cubic` together with an `arg/[\cdot]` selector.

Isabelle/HOL `Cubic.Quartic` and the HOL Light entry on Wiedijk’s list are algebraic in the same sense as Mathlib: roots are assumed, not constructed by principal functions.

This library already has companion quadratic [8] and cubic [7] lemmas over $\mathbb{C}$. The quartic contribution is to finish Ferrari on that principal-radical calculus: the resolvent is expanded by Cardano, one legal root $y = A\omega^k + B$ is selected, and the four quartic roots are nested principal square roots of that y .

3 The formula

Write $a' = b/a$, $b' = c/a$, $c' = d/a$, $d' = e/a$, and

$$\begin{aligned}\alpha &= b' - 3a'^2/8, \\ \beta &= a'^3/8 + c' - a'b'/2, \\ \gamma &= a'^2b'/16 + d' - 3a'^4/256 - a'c'/4.\end{aligned}$$

Then $z = x + a'/4$ satisfies $z^4 + \alpha z^2 + \beta z + \gamma = 0$.

Biquadratic regime $\beta = 0$. With $\Delta = \alpha^2 - 4\gamma$,

$$x = \pm \sqrt{(\pm \sqrt{\Delta} - \alpha)/2 - a'/4}$$

(four independent choices of the two signs).

Ferrari regime $\beta \neq 0$. Form Cardano quantities of the resolvent,

$$\begin{aligned}p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, \quad B = \sqrt[3]{(-q - \sqrt{\delta})/2},\end{aligned}$$

and

$$k = \left\lceil \frac{3 \arg(-p)}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg(AB)}{2\pi} - \frac{1}{2} \right\rceil.$$

The abbreviation $\omega = e^{2\pi i/3}$ (Lean `(I * (2 * π / 3)).exp`) is for simplicity only: a complex exponential is *not* a radical. The same primitive cube root of unity is the principal-radical rectangular form

$$\omega = -\frac{1}{2} + i \frac{\sqrt{3}}{2}$$

(Lean `(-(1 / 2 : ℝ)) + I * (√3 / 2 : ℝ)`), and $\bar{\omega} = -\frac{1}{2} - i \frac{\sqrt{3}}{2}$. The identities $e^{2\pi i/3} = -\frac{1}{2} + i\sqrt{3}/2$ and its conjugate are machine-checked [3, 4]. Set

$$y = A\omega^k + B, \quad y_0 = -2\alpha/3 + y, \quad y_1 = 4\alpha/3 + y.$$

Then

$$\begin{aligned} x &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2} - \frac{a'}{4} \\ \text{or} \quad &\frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4} \quad \text{or} \quad \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2} - \frac{a'}{4}. \end{aligned}$$

Theorem 3.1 (Principal-radical quartic [5]). *Assume $a \neq 0$. With the quantities above,*

$$ax^4 + bx^3 + cx^2 + dx + e = 0$$

if and only if x equals one of the four biquadratic expressions when $\beta = 0$, and one of the four Ferrari expressions with $y = A\omega^k + B$ when $\beta \neq 0$. The cube-root branch is the single factor ω^k , not a case split on $k \bmod 3$.

The `if` selects the matching four-element set; it does not list Ferrari radicals as roots when $\beta = 0$, nor biquadratic radicals when $\beta \neq 0$.

4 Figures

Figure 0 — public visualizations. The SymPy page [2] displays the one-direction (`mp`) statement `In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0` as an interactive theorem document (created 2018-11-29). The Lean 4 page [5] displays `@[main, comm, mp, mpr] private lemma main` with $\{x, a, b, c, d, e : \mathbb{C}\}$, the hypothesis $a \neq 0$, the `let` chain $a', b', \alpha, \beta, \gamma$ and the inner `if` lets, and the if-and-only-if $ax^4 + \dots + e = 0$ iff x belongs to the selected four-element set.

Figure 1 — reduction. Figure 1 is the reduction used in the main theorem: divide by the leading coefficient, depress by $z = x + a'/4$, then split on $\beta = 0$ versus $\beta \neq 0$.

Figure 2 — lemma dependencies. Figure 2 lists major imported lemmas of Theorem 3.1. Private helpers in the same file are drawn in parentheses. Links point to the Lean 4 pages. SymPy has only the one-direction (`of/mp`) names, e.g. http://www.lemma.cn/py/?module=Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0, not the Lean `is if-and-only-if`.

Related lemmas in the same family, *not* imported by the main theorem: the depressed statement `Imp_OrOrSEqS.Imp_OrOrSEqS.Div_2.of.Eq0Add_Pow_4` and the monic offset `Imp_OrOrSEqS.Sub.Imp_OrOr`. See also the quadratic [8] and the cubic [7]. The quartic proof calls `Eq_Mul_Pow_SubCeilS.of.Pow_3` directly, because Ferrari needs one resolvent root $A\omega^k + B$.

$$\begin{array}{c}
ax^4 + bx^3 + cx^2 + dx + e = 0, \quad a \neq 0 \\
\downarrow a' = \frac{b}{a}, b' = \frac{c}{a}, c' = \frac{d}{a}, d' = \frac{e}{a} \\
x^4 + a'x^3 + b'x^2 + c'x + d' = 0 \\
\downarrow z = x + \frac{a'}{4} \\
z^4 + \alpha z^2 + \beta z + \gamma = 0 \\
\alpha = b' - \frac{3a'^2}{8}, \quad \beta = \frac{a'^3}{8} + c' - \frac{a'b'}{2}, \quad \gamma = \frac{a'^2 b'}{16} + d' - \frac{3a'^4}{256} - \frac{a'c'}{4} \\
\begin{array}{cc}
\beta = 0 & \beta \neq 0 \\
\downarrow & \downarrow \\
\Delta = \alpha^2 - 4\gamma & y = A\omega^k + B
\end{array} \\
\begin{array}{cc}
x = \pm \sqrt{\frac{\pm \sqrt{\Delta} - \alpha}{2}} - \frac{a'}{4} & \begin{array}{l} y_0 = -\frac{2\alpha}{3} + y, \quad y_1 = \frac{4\alpha}{3} + y \\ \pm \sqrt{\frac{2\beta}{\sqrt{y_0}} - y_1 \pm \sqrt{y_0}} \\ x = \frac{\pm \sqrt{\frac{2\beta}{\sqrt{y_0}} - y_1 \pm \sqrt{y_0}}}{2} - \frac{a'}{4} \\ x = z - \frac{a'}{4} \end{array}
\end{array}
\end{array}$$

Figure 1: Reduction of the general quartic and the two radical regimes.

Figure 3 — branch integer k . From $A^3 = B^3$ one has $A = B\omega^k$ with

$$k = \left\lceil \frac{3 \arg A}{2\pi} - \frac{1}{2} \right\rceil - \left\lceil \frac{3 \arg B}{2\pi} - \frac{1}{2} \right\rceil.$$

For $(AB)^3 = (-p/3)^3$ the general selector is the negative of the k in Theorem 3.1 (the statement writes $\arg(-p)$, which equals $\arg(-p/3)$), and ω^{-k} restores $AB = (-p/3)\omega^{-k}$. Multiplying only A by ω^k produces a legal Ferrari parameter $y = A\omega^k + B$. Because $\omega^3 = 1$, the three residue classes of k modulo 3 are already encoded in that single power: $\omega^k \in \{1, \omega, \bar{\omega}\}$.

5 Proof outline

5.1 Principal radicals

Throughout, $\sqrt{\cdot}$ is `Mathlib Complex.sqrt` and $\sqrt[3]{\cdot}$ is `Lean Root.cubic`. Both are total functions $\mathbb{C} \rightarrow \mathbb{C}$. The identity $(\sqrt{z})^2 = z$ is `Complex.EqSquareSqrt`. If $x^2 = c$, then $x = \sqrt{c}$ or $x = -\sqrt{c}$ by `Complex.Or_Eq_NegSqrt.of.EqSquare`. Cube roots of unity are the radicals

$$\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}, \quad \bar{\omega} = -\frac{1}{2} - i\frac{\sqrt{3}}{2}, \quad \omega^2 = \bar{\omega}, \quad \omega^3 = 1.$$

The polar form $z = \|z\|e^{i\arg z}$ is `Complex.Eq_MulNorm_ExpMulIArg`. Reduction of $\arg(e^{ix})$ into $(-\pi, \pi]$ by a ceiling is `Complex.ArgExpMulI.eq.Sub_Mul_Ceil`, which yields `Complex.ArgPow.eq.SubMul_Arg` and then `Complex.Eq_Mul_Pow_SubCeils.of.Pow_3`: if $A^3 = B^3$, then $A = B\omega^k$ for the integer k of Figure 3.

5.2 Depression

Division by $a \neq 0$ produces the monic equation $x^4 + a'x^3 + b'x^2 + c'x + d' = 0$. The translation $z = x + a'/4$ cancels the cubic term and yields $z^4 + \alpha z^2 + \beta z + \gamma = 0$ with α, β, γ as in §3. The main file performs this expansion by `ring`; the four nested-radical expressions for z are then shifted back by $-a'/4$.

- `Eq0Add_Mul_Pow_4.is.In.Ite.FinsetSSubS_SubS.of.Ne_0` — main theorem
 - `OrOrSEqS.of.Eq0Add_Pow_4` — case $\beta = 0$
 - * `OrEqS.Div.of.Eq0Add_Mul_Square.Ne_0` — quadratic in x^2
 - * `Or_Eq_NegSqrt.of.EqSquare` — $x^2 = c \Rightarrow x = \pm\sqrt{c}$
 - `EqSquareSqrt` — $(\sqrt{z})^2 = z$
 - `Real.OrEqS.of.Square`
 - `OrOrSEqS.Div_2.of.Eq0Add_Pow_4.Ne_0` — case $\beta \neq 0$
 - * `(hbranch)` — given a resolvent root y_0 , factor into two quadratics
 - * `Eq_Mul_Pow_SubCeilS.of.Pow_3` — $A^3 = B^3 \Rightarrow A = B \omega^k$
 - `ArgPow.eq.SubMul_Arg`
 - `Eq_MulNorm_ExpMulIArg`
 - `ArgExpMulI.eq.Sub_Mul_Ceil`
 - `Int.Floor.eq.NegCeilNeg`

Figure 2: Import graph for the Lean proof of Theorem 3.1.

5.3 Biquadratic case $\beta = 0$

The depressed equation is $z^4 + \alpha z^2 + \gamma = 0$, a quadratic in z^2 . Lemma `Complex.OrOrSEqS.of.Eq0Add_Pow_4` assembles the four principal-radical roots. It first solves

$$z^2 = \frac{-\alpha \pm \sqrt{\alpha^2 - 4\gamma}}{2} = \frac{\pm\sqrt{\Delta} - \alpha}{2}, \quad \Delta = \alpha^2 - 4\gamma,$$

then splits each $z^2 = c$ by `Or_Eq_NegSqrt`. Translating $x = z - \alpha'/4$ is the $\beta = 0$ branch of Theorem 3.1.

5.4 Ferrari factorization

Let $y_0 \in \mathbb{C}$ be any root of the cubic resolvent

$$y_0^3 + 2\alpha y_0^2 + (\alpha^2 - 4\gamma)y_0 - \beta^2 = 0,$$

and set $y_1 = y_0 + 2\alpha$. A private helper proves $y_0 \neq 0$ (else $\beta = 0$), that $(\sqrt{y_0})^2 = y_0$, and that with $Y = (y_0 + \alpha)/2$ one has

$$(z^2 + Y)^2 - (\sqrt{y_0} z - \beta/(2\sqrt{y_0}))^2 = z^4 + \alpha z^2 + \beta z + \gamma.$$

The left-hand side is a product of two quadratics. Each is solved by completing the square and `Or_Eq_NegSqrt`, producing

$$\begin{aligned} z &= \frac{\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{2\beta/\sqrt{y_0} - y_1} - \sqrt{y_0}}{2}, \\ z &= \frac{\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}, & z &= \frac{-\sqrt{-2\beta/\sqrt{y_0} - y_1} + \sqrt{y_0}}{2}. \end{aligned}$$

This step still treats y_0 as given. The rest of the proof constructs y_0 from principal radicals.

5.5 Cardano data of the resolvent

The theorem statement writes the scaled Cardano pair directly:

$$\begin{aligned} p &= -4\gamma - \alpha^2/3, \\ q &= -2\alpha^3/27 + 8\alpha\gamma/3 - \beta^2, \\ \delta &= 4p^3/27 + q^2, \\ A &= \sqrt[3]{(-q + \sqrt{\delta})/2}, & B &= \sqrt[3]{(-q - \sqrt{\delta})/2}. \end{aligned}$$

The proof also depresses the resolvent by

$$a_r = -\alpha/2, \quad b_r = -\gamma, \quad c_r = -\beta^2/8 + \alpha\gamma/2,$$

and the unscaled Cardano parameters

$$\begin{aligned} p_c &= b_r - a_r^2/3, & q_c &= 2a_r^3/27 - a_rb_r/3 + c_r, & \delta_c &= 4p_c^3/27 + q_c^2, \\ A_c &= \sqrt[3]{(-q_c + \sqrt{\delta_c})/2}, & B_c &= \sqrt[3]{(-q_c - \sqrt{\delta_c})/2}. \end{aligned}$$

A direct calculation gives $q = 8q_c$, $\delta = 64\delta_c$, and $\sqrt{\delta} = 8\sqrt{\delta_c}$ (the last because $64 > 0$ on the principal branch). Positive real scaling of cube roots yields $A = 2A_c$ and $B = 2B_c$, and $\arg(AB) = \arg(A_cB_c)$.

5.6 Restoring $3(A\omega^k)B = -p$

Because $(AB)^3 = (-p/3)^3$, lemma `Eq_Mul_Pow_SubCeilS.of.Pow_3` supplies

$$AB = (-p/3)\omega^{-k}.$$

Multiplying by ω^k restores the product in one step: $AB\omega^k = -p/3$. Since $(\omega^k)^3 = 1$,

$$(A\omega^k)^3 + B^3 = A^3 + B^3 = -q, \quad 3(A\omega^k)B = -p,$$

so $y = A\omega^k + B$ satisfies $y^3 + py + q = 0$. There is no case split on $k \bmod 3$: the three possible rotations $1, \omega, \bar{\omega}$ are the values of the single power ω^k . Shifting $y_0 = -2\alpha/3 + y$ and $y_1 = 4\alpha/3 + y$ recovers a resolvent root, and the four nested square roots apply.

Unlike the cubic formula in this library, only A is rotated. A quartic needs one legal resolvent parameter, not three Cardano roots of the original unknown.

6 Discussion and limitations

The statement is an if-and-only-if: the equation holds precisely when x lies in the four-element set selected by `if` $\beta = 0$. It is not a four-fold split on $k \bmod 3$. Principal square roots in the Ferrari step may vanish or repeat when the quartic has multiple roots; membership still holds.

The lemma does not construct roots in a general field of characteristic zero, does not avoid `arg` and `[·]`, and does not prove that a randomly chosen pairing of principal cube roots is a resolvent root. It does not replace Mathlib's abstract Ferrari theorem [11]: that theorem remains the right statement when one only knows that some square roots and some resolvent root exist in K . The present lemma is the complementary statement on $\mathbb{C}$: every coefficient tuple with $a \neq 0$ is sent to four explicit principal-radical expressions, with the cube-root branch named by the power ω^k .

7 Conclusion

We have presented a machine-checked, fully expanded Ferrari–Cardano formula for the general quartic over $\mathbb{C}$, using principal square roots, principal cube roots $\sqrt[3]{\cdot}$, the radical cube root of unity $-\frac{1}{2} + i\sqrt{3}/2$, and a single power ω^k assembled from `arg` and `[·]`. The result is public on `lemma.cn` and GitHub [6], with proofs checked in Lean 4 and an earlier SymPy formulation.

References

- [1] Nicholas Dyson, Benedikt Ahrens, and Jacopo Emmenegger. The solutions to single-variable polynomials, implemented and verified in Lean, 2022.
- [2] math proof. Fully radical quartic over $\mathbb{C}$ (SymPy). http://www.lemma.cn/py/?module=Complex.In_Ite_FinsetSSubS__SubS.of.Eq0Add_Mul_Pow_4.Ne_0, 2018.
- [3] math proof. Complex.expmulidivmul2pi3.eq.add_muli. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMul2Pi3.eq.Add_MulI, 2026.
- [4] math proof. Complex.expmulidivmulneg2pi3.eq.sub_muli. http://www.lemma.cn/lean/?module=Complex.ExpMulIDivMulNeg2Pi3.eq.Sub_MulI, 2026.
- [5] math proof. Fully radical quartic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_4.is.In_Ite_FinsetSSubS__SubS.of.Ne_0, 2026.
- [6] math proof. lemma: machine-checked mathematics. <https://github.com/math-proof/lemma>, 2026.
- [7] math proof. Principal-radical cubic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Pow_3.is.In_Finset_SubSAddMulS.of.Ne_0, 2026.
- [8] math proof. Principal-radical quadratic over $\mathbb{C}$ (Lean 4). http://www.lemma.cn/lean/?module=Complex.Eq0Add_Mul_Square.is.In_FinsetDivS_Mul2.of.Ne_0, 2026.
- [9] Mathlib community. The Lean mathematical library. https://leanprover-community.github.io/mathlib4_docs/, 2026.
- [10] Freek Wiedijk. Formalizing 100 theorems. <https://www.cs.ru.nl/~freek/100/>, 2024.
- [11] Thomas Zhu. The roots of cubic and quartic polynomials. Mathlib 4 Archive, PR #18290, 2024. <https://github.com/leanprover-community/mathlib4/blob/master/Archive/Wiedijk100Theorems/SolutionOfCubicQuartic.lean>.