

MAT205: Abstract Algebra II

Field Extensions and Roots of Polynomials

Ma, Jia-Jun - Xiamen University Malaysia

Part I

Field Extensions

Definition: Field Extension

Let F and E be fields.

A **field extension** is an inclusion of fields

$$F \subseteq E.$$

We write this as

$$E/F.$$

The notation E/F is **not** quotient notation.

Examples: Field Extensions

Extension	Inclusion
$\mathbb{C}/\mathbb{R}$	$\mathbb{R} \subseteq \mathbb{C}$
$\mathbb{R}/\mathbb{Q}$	$\mathbb{Q} \subseteq \mathbb{R}$
$\mathbb{Q}(\sqrt{2})/\mathbb{Q}$	$\mathbb{Q} \subseteq \mathbb{Q}(\sqrt{2})$
$\mathbb{R}(i)/\mathbb{R}$	$\mathbb{R} \subseteq \mathbb{R}(i) = \mathbb{C}$

Here

$$\mathbb{Q}(\sqrt{2}) = \{a + b\sqrt{2} : a, b \in \mathbb{Q}\}.$$

Non-Examples

$$\mathbb{Q}/\mathbb{Z}$$

is not a field extension.

Reason:

$$\mathbb{Z}$$

is not a field.

Also,

$$\mathbb{R}/\mathbb{C}$$

is not a field extension by inclusion, because

$$\mathbb{C} \not\subseteq \mathbb{R}.$$

Field Extensions as Vector Spaces

If $F \subseteq E$, then E is a vector space over F .

Scalar multiplication is multiplication in E :

$$\lambda \cdot x = \lambda x, \quad \lambda \in F, \quad x \in E.$$

The **degree** of E/F is

$$[E : F] = \dim_F E.$$

Examples: Extension Degree

$$[\mathbb{C} : \mathbb{R}] = 2$$

with basis

$$\{1, i\}.$$

$$[\mathbb{Q}(\sqrt{2}) : \mathbb{Q}] = 2$$

with basis

$$\{1, \sqrt{2}\}.$$

For now, degree is only notation. Its general theory belongs with algebraic extensions.

Part II

Polynomial Equations and Larger Fields

Example: $x^2 - 2$

The equation

$$x^2 - 2 = 0$$

has no solution in $\mathbb{Q}$.

It has solutions in the larger field $\mathbb{Q}(\sqrt{2})$:

$$\sqrt{2}, \quad -\sqrt{2}.$$

Example: $x^2 + 1$

The equation

$$x^2 + 1 = 0$$

has no solution in $\mathbb{R}$.

It has solutions in the larger field

$$\mathbb{C} = \mathbb{R}(i) :$$

$$i, \quad -i.$$

Polynomial Roots in Larger Fields

A polynomial over a field F may have no root in F .

It may have a root after passing to a larger field

$$F \subseteq E.$$

The construction will use quotient rings of the form

$$F[x]/(p(x)).$$

Part III

Irreducible Polynomials and Maximal Ideals

Definition: Irreducible Polynomial

Let F be a field.

A polynomial $p(x) \in F[x]$ is **irreducible over F** if:

1. $p(x) \neq 0$;
2. $p(x)$ is nonconstant;
3. if

$$p(x) = a(x)b(x),$$

then $a(x)$ or $b(x)$ is a unit in $F[x]$.

Units in $F[x]$

Since F is a field, the units in $F[x]$ are exactly the nonzero constant polynomials.

Thus $p(x)$ is irreducible if it has no factorization

$$p(x) = a(x)b(x)$$

with

$$\deg a > 0, \quad \deg b > 0.$$

Examples: Irreducibility

$$x^2 - 2$$

is irreducible over $\mathbb{Q}$.

$$x^2 + 1$$

is irreducible over $\mathbb{R}$.

But $x^2 + 1$ is not irreducible over $\mathbb{C}$, since

$$x^2 + 1 = (x - i)(x + i).$$

$$x^3 - 2$$

is irreducible over $\mathbb{Q}$.

Proposition: Irreducible Implies Maximal

Let F be a field and let $p(x) \in F[x]$ be irreducible.

Then

$$(p(x))$$

is a maximal ideal of $F[x]$.

Therefore

$$F[x]/(p(x))$$

is a field.

Proof: Maximality

Let I be an ideal such that

$$(p(x)) \subseteq I \subseteq F[x].$$

Since $F[x]$ is a PID, write

$$I = (g(x)).$$

The inclusion $(p(x)) \subseteq (g(x))$ means

$$p(x) = g(x)h(x)$$

for some $h(x) \in F[x]$.

Proof: Maximality

Since $p(x)$ is irreducible,

$$p(x) = g(x)h(x)$$

forces $g(x)$ or $h(x)$ to be a unit.

If $g(x)$ is a unit, then

$$I = (g(x)) = F[x].$$

If $h(x)$ is a unit, then

$$I = (g(x)) = (p(x)).$$

So $(p(x))$ is maximal.

Examples: Quotient Fields

Since $x^2 - 2$ is irreducible over $\mathbb{Q}$,

$$\mathbb{Q}[x]/(x^2 - 2)$$

is a field.

Since $x^2 + 1$ is irreducible over $\mathbb{R}$,

$$\mathbb{R}[x]/(x^2 + 1)$$

is a field.

Part IV

Constructing Roots

Notation in $F[x]/(p(x))$

Let $p(x) \in F[x]$ be irreducible and define

$$E = F[x]/(p(x)).$$

Write

$$\alpha = x + (p(x)).$$

Every element of E can be written as

$$a_0 + a_1\alpha + \cdots + a_{n-1}\alpha^{n-1},$$

where $n = \deg p$ and $a_i \in F$.

Constructing a Root

Let $p(x) \in F[x]$ be irreducible and let

$$E = F[x]/(p(x)).$$

The field F embeds into E by sending $a \in F$ to the class of the constant polynomial a .

With this identification,

$$F \subseteq E.$$

So E/F is a field extension.

The Root in the Quotient

Let

$$\alpha = x + (p(x)) \in F[x]/(p(x)).$$

Then

$$p(\alpha) = 0.$$

Indeed, if

$$p(x) = a_0 + a_1x + \cdots + a_nx^n,$$

then

$$p(\alpha) = p(x) + (p(x)) = 0.$$

Theorem: Existence of a Root

Let F be a field and let $f(x) \in F[x]$ be nonconstant.

Then there exists a field extension E/F and an element $\alpha \in E$ such that

$$f(\alpha) = 0.$$

Proof: Existence of a Root

Choose an irreducible factor $p(x)$ of $f(x)$ in $F[x]$.

Write

$$f(x) = p(x)q(x).$$

Let

$$E = F[x]/(p(x)), \quad \alpha = x + (p(x)).$$

Then $p(\alpha) = 0$, so

$$f(\alpha) = p(\alpha)q(\alpha) = 0.$$

Part V

Examples

Example: $x^2 - 2$ over $\mathbb{Q}$

Let

$$E = \mathbb{Q}[x]/(x^2 - 2), \quad \alpha = x + (x^2 - 2).$$

Then

$$\alpha^2 - 2 = 0, \quad \alpha^2 = 2.$$

Every element of E has the form

$$a + b\alpha, \quad a, b \in \mathbb{Q}.$$

Example: $\mathbb{Q}[x]/(x^2 - 2)$

There is an isomorphism

$$\mathbb{Q}[x]/(x^2 - 2) \longrightarrow \mathbb{Q}(\sqrt{2})$$

given by

$$\alpha \longmapsto \sqrt{2}.$$

The quotient construction is an abstract version of adjoining $\sqrt{2}$.

Example: $x^2 + 1$ over $\mathbb{R}$

Let

$$E = \mathbb{R}[x]/(x^2 + 1), \quad \alpha = x + (x^2 + 1).$$

Then

$$\alpha^2 + 1 = 0, \quad \alpha^2 = -1.$$

Every element of E has the form

$$a + b\alpha, \quad a, b \in \mathbb{R}.$$

Example: $\mathbb{R}[x]/(x^2 + 1)$

There is an isomorphism

$$\mathbb{R}[x]/(x^2 + 1) \longrightarrow \mathbb{C}$$

given by

$$\alpha \longmapsto i.$$

So $\mathbb{C}$ can be constructed from $\mathbb{R}$ by adjoining a root of $x^2 + 1$.

Example: $x^3 - 2$ over $\mathbb{Q}$

Let

$$E = \mathbb{Q}[x]/(x^3 - 2), \quad \alpha = x + (x^3 - 2).$$

Then

$$\alpha^3 = 2.$$

Every element of E has the form

$$a + b\alpha + c\alpha^2, \quad a, b, c \in \mathbb{Q}.$$

This construction gives a field containing one root of $x^3 - 2$.

Definition: Adjoining a Root

The notation

$$F(\alpha)$$

means the smallest subfield containing F and α .

In the construction

$$E = F[x]/(p(x)), \quad \alpha = x + (p(x)),$$

one has

$$E = F(\alpha).$$

The element α satisfies

$$p(\alpha) = 0.$$

Exercises

Exercise 1

Let

$$E = \mathbb{Q}[x]/(x^2 - 3), \quad \alpha = x + (x^2 - 3).$$

1. Show that E is a field.
2. Show that $\alpha^2 = 3$.
3. Write every element of E in the form

$$a + b\alpha, \quad a, b \in \mathbb{Q}.$$

4. Construct an isomorphism $E \cong \mathbb{Q}(\sqrt{3})$.

Exercise 2

Let

$$E = \mathbb{Q}[x]/(x^3 - 5), \quad \alpha = x + (x^3 - 5).$$

1. Show that $x^3 - 5$ is irreducible over $\mathbb{Q}$.
2. Show that E is a field.
3. Show that $\alpha^3 = 5$.
4. Write every element of E in the form

$$a + b\alpha + c\alpha^2, \quad a, b, c \in \mathbb{Q}.$$