

Lecture 18: Solvability of Equations

MAT205 Abstract Algebra II

Ma, Jia-Jun

Xiamen University Malaysia

Part I: Solvability by Radicals

From the Quadratic Formula to the Quintic Question

Every root below is built from the coefficients by $+$, $-$, $\times$, $\div$ and extraction of roots.

- **Degree 2.** Over a field of characteristic $\neq 2$, the zeros of $ax^2 + bx + c$ ($a \neq 0$) are $\frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ — one square root.
- **Degrees 3 and 4.** Cardano's formula (1545) solves the cubic and Ferrari's method the quartic, by field operations and cube/square roots.
- **Degree 5.** After years of searching for a “quintic formula”, Abel (1824–1826) proved that a quintic *need not* be solvable by radicals — there is no general formula.

Extensions by Radicals; Solvable by Radicals

Definition. An extension K/F is an *extension by radicals* if there are $\alpha_1, \dots, \alpha_r \in K$ and integers $n_1, \dots, n_r \geq 1$ with

$$K = F(\alpha_1, \dots, \alpha_r), \quad \alpha_1^{n_1} \in F, \quad \alpha_i^{n_i} \in F(\alpha_1, \dots, \alpha_{i-1}) \quad (1 < i \leq r).$$

Equivalently, a tower $F = F_0 \subseteq F_1 \subseteq \dots \subseteq F_r = K$ with $F_i = F_{i-1}(\alpha_i)$ and $\alpha_i^{n_i} \in F_{i-1}$.

Definition. $f \in F[x]$ is *solvable by radicals over F* if its splitting field E is contained in some extension of F by radicals.

Equivalently: every root of f is reached from the coefficients by finitely many $+$, $-$, $\times$, $\div$ and n_i th roots.

Examples: Some Quintics Are Solvable by Radicals

Insolvability does not mean that *no* quintic is solvable, as these examples show.

- $x^5 - 1$ is solvable by radicals over $\mathbb{Q}$: writing $\zeta_5 = e^{2\pi i/5}$ for a primitive 5th root of unity, its splitting field is $\mathbb{Q}(\zeta_5)$ with $\zeta_5^5 = 1 \in \mathbb{Q}$ — a one-step radical extension.
- $x^5 - 2$ is solvable by radicals over $\mathbb{Q}$: its splitting field is $\mathbb{Q}(\sqrt[5]{2}, \zeta_5)$, where $(\sqrt[5]{2})^5 = 2 \in \mathbb{Q}$ and $\zeta_5^5 = 1$ — a two-step radical tower.

So the claim “the quintic is insolvable” means: **there exists** a degree-5 polynomial that is not solvable by radicals. We will exhibit one explicitly.

Galois's Criterion

Recall. A finite group G is *solvable* if it has a subnormal series $\{e\} = G_0 \trianglelefteq G_1 \trianglelefteq \cdots \trianglelefteq G_m = G$ with every quotient G_i/G_{i-1} abelian.

Theorem (Galois). Let F have characteristic 0 and let $f \in F[x]$ have splitting field E . Then

$$f \text{ is solvable by radicals over } F \iff \text{Gal}(E/F) \text{ is a solvable group.}$$

We prove the direction that yields impossibility:

$$f \text{ solvable by radicals} \implies \text{Gal}(E/F) \text{ solvable,}$$

equivalently (contrapositive) a *non-solvable* Galois group forces f *not* solvable by radicals. The converse is also true (stated without proof).

Part II: Radical Towers Have Solvable Galois Groups

The Single-Radical Lemma: $x^n - a$

Lemma. Let F have characteristic 0, let $a \in F$, and let K be the splitting field of $x^n - a$ over F . Then $\text{Gal}(K/F)$ is solvable.

One root extraction contributes only *abelian* layers:

- if F already contains the n th roots of unity, the group is **abelian**;
- in general, adjoin the roots of unity first — two abelian layers, hence solvable.

Here we use that the n th roots of unity $\mu_n = \{1, \zeta_n, \dots, \zeta_n^{n-1}\}$ form a *cyclic* group, being a finite subgroup of $F^\times$.

Proof of the Lemma

Case 1: $\mu_n \subseteq F$. Fix a generator ζ of μ_n . If β is one root of $x^n - a$, all roots are $\beta, \zeta\beta, \dots, \zeta^{n-1}\beta$, so $K = F(\beta)$. Each $\sigma \in \text{Gal}(K/F)$ is determined by $\sigma(\beta) = \zeta^i\beta$; if also $\tau(\beta) = \zeta^j\beta$ then, as $\zeta^i \in F$,

$$(\tau\sigma)(\beta) = \zeta^i\zeta^j\beta = (\sigma\tau)(\beta),$$

so $\text{Gal}(K/F)$ is **abelian**.

Case 2: general F . Let $F' = F(\zeta)$, so $F \leq F' \leq K$. Then F'/F is the splitting field of $x^n - 1$, hence normal, and $\eta \in \text{Gal}(F'/F)$ is determined by $\eta(\zeta) = \zeta^i$; composing shows $\text{Gal}(F'/F)$ is abelian. Over F' we are in Case 1, so $\text{Gal}(K/F')$ is abelian. By the Fundamental Theorem of Galois Theory,

$$\{e\} \trianglelefteq \text{Gal}(K/F') \trianglelefteq \text{Gal}(K/F), \quad \text{Gal}(K/F)/\text{Gal}(K/F') \cong \text{Gal}(F'/F),$$

a subnormal series with abelian quotients. Hence $\text{Gal}(K/F)$ is solvable. ■

Radical Extension $\Rightarrow$ Solvable Galois Group

Theorem. Let F have characteristic 0, let E/F be normal (hence Galois), and suppose $E \subseteq K$ with K/F an extension by radicals. Then $\text{Gal}(E/F)$ is solvable.

Proof. Enlarge K to a *normal* radical extension L/F with $\text{Gal}(L/F)$ solvable. From $K = F(\alpha_1, \dots, \alpha_r)$, $\alpha_i^{n_i} \in F(\alpha_1, \dots, \alpha_{i-1})$, build L in stages: let L_1 be the splitting field of $x^{n_1} - \alpha_1^{n_1}$ over F ; at stage i adjoin the splitting field of

$$\prod_{\sigma \in \text{Gal}(L_{i-1}/F)} (x^{n_i} - \sigma(\alpha_i)^{n_i}) \in F[x]$$

(a product over conjugates, hence over F). Each layer is a tower of $x^n - a$ extensions, so by the Lemma — and since an extension of a solvable group by a solvable group is solvable — $\text{Gal}(L/F)$ stays solvable. The result $L = L_r$ is normal over F , radical, with $K \leq L$.

Since E/F is normal, the Fundamental Theorem gives

$\text{Gal}(E/F) \cong \text{Gal}(L/F)/\text{Gal}(L/E)$, a quotient of a solvable group, hence solvable. ■

Part III: The Insolvability of the Quintic

The General Quintic Has Galois Group S_5

Let $y_1, \dots, y_5$ be independent transcendentals over $\mathbb{Q}$, set $E = \mathbb{Q}(y_1, \dots, y_5)$ and

$$f(x) = \prod_{i=1}^5 (x - y_i).$$

Its coefficients are, up to sign, the elementary symmetric functions $s_1, \dots, s_5$ of the y_i . Put $F = \mathbb{Q}(s_1, \dots, s_5)$; then $f \in F[x]$ and E is the splitting field of f over F .

Each $\sigma \in S_5$ permutes the y_i , inducing $\bar{\sigma} \in \text{Aut}(E)$ that fixes $\mathbb{Q}$; since $\prod (x - y_i) = \prod (x - y_{\sigma(i)})$, it fixes every s_i , hence fixes F . So $S_5 \hookrightarrow \text{Gal}(E/F)$ and $|\text{Gal}(E/F)| \geq 5!$. A degree-5 splitting field has degree $\leq 5!$, so equality holds:

$$\text{Gal}(E/F) \cong S_5.$$

$$\begin{array}{c} E = \mathbb{Q}(y_1, \dots, y_5) \\ \downarrow S_5 \\ F = \mathbb{Q}(s_1, \dots, s_5) \\ \downarrow \\ \mathbb{Q} \end{array}$$

S_5 Is Not Solvable — the General Quintic Is Insolvable

S_5 **is not solvable**. A composition series for S_5 is

$$\{e\} \trianglelefteq A_5 \trianglelefteq S_5,$$

with quotients A_5 and $\mathbb{Z}_2$. Since A_5 is a **non-abelian simple group**, a composition factor of S_5 is non-abelian, so S_5 is not solvable.

Theorem. The general quintic $f(x) = \prod_{i=1}^5 (x - y_i)$ is not solvable by radicals over $F = \mathbb{Q}(s_1, \dots, s_5)$.

Proof. $\text{Gal}(E/F) \cong S_5$ is not solvable; by the contrapositive of the radical theorem, f is not solvable by radicals. ■

The same argument for $n \geq 5$ uses that S_n has the non-abelian simple composition factor A_n : *no general radical formula exists for degree ≥ 5 .*

An Insolvable Quintic over $\mathbb{Q}$: $2x^5 - 5x^4 + 5$


Irreducible. $f(x) = 2x^5 - 5x^4 + 5$ is irreducible over $\mathbb{Q}$ by Eisenstein at $p = 5$: $5 \nmid 2$ (leading), $5 \mid -5, 0, 0, 0, 5$ (lower), and $5^2 = 25 \nmid 5$ (constant).

Exactly three real roots.

$f'(x) = 10x^4 - 20x^3 = 10x^3(x - 2)$, so the critical points are $x = 0$ and $x = 2$, with

$$f(0) = 5 > 0 \text{ (max),} \quad f(2) = -11 < 0 \text{ (min).}$$

With $f \rightarrow -\infty$ as $x \rightarrow -\infty$ and $f \rightarrow +\infty$ as $x \rightarrow +\infty$, the graph crosses the axis exactly **three** times: three real roots and one conjugate pair of non-real roots.



Its Galois Group Is S_5

Let E be the splitting field of $f(x) = 2x^5 - 5x^4 + 5$ over $\mathbb{Q}$.

- **A 5-cycle.** f is irreducible of degree 5, so for a root α , $[\mathbb{Q}(\alpha) : \mathbb{Q}] = 5$ divides $|\text{Gal}(E/\mathbb{Q})|$; by Cauchy's theorem the group has an element of order 5, a 5-cycle on the five roots.
- **A transposition.** Complex conjugation restricts to an automorphism of E ; it fixes the three real roots and swaps the conjugate pair, acting as a transposition.
- **Generation.** A standard fact: a subgroup of S_5 containing a 5-cycle and a transposition is all of S_5 (the transposition and the 5-cycle generate every transposition).

Hence $\text{Gal}(E/\mathbb{Q}) \cong S_5$, which is not solvable, so

$2x^5 - 5x^4 + 5$ is not solvable by radicals over $\mathbb{Q}$. ■

Part IV: Lagrange, Abel, Galois

Lagrange (1770): Resolvents

Resolvent for the cubic. For roots $\alpha_1, \alpha_2, \alpha_3$ and $\omega = e^{2\pi i/3}$, set

$$r = \alpha_1 + \omega\alpha_2 + \omega^2\alpha_3.$$

A 3-cycle of the roots sends $r \mapsto \omega^2 r$, so r^3 is fixed by A_3 and takes 2 **values** under S_3 ; hence r^3 satisfies a quadratic over $\mathbb{Q}(s_1, s_2, s_3)$ — this gives *Cardano's formula*. The analogous quartic resolvent takes 3 values, satisfying a cubic — *Ferrari's resolvent* cubic.

Why degree 5 stalls. For $n \geq 5$ no rational function of the roots takes a small enough number of values under S_n to reduce the problem. Lagrange identified this obstruction but could not prove impossibility.

Modern restatement. Lagrange's action $r \mapsto \omega^i r$ is precisely the Galois action on one radical step: if $\mu_n \subseteq F$ and $\beta^n = a \in F$, then $\text{Gal}(F(\beta)/F)$ is cyclic of order dividing n — Case 1 of Lemma 56.3 (Slide 6). **Each radical step contributes a single cyclic, hence abelian, layer.**

Abel (1824–1826): Insolubility

Theorem (Abel). The general equation of degree $n \geq 5$ is not solvable by radicals over $\mathbb{Q}(s_1, \dots, s_n)$.

Strategy. Suppose a root lies in a radical tower $F = F_0 \subseteq F_1 \subseteq \dots \subseteq F_r$ with $F_i = F_{i-1}(\alpha_i)$, $\alpha_i^{n_i} \in F_{i-1}$. By the Lagrange analysis applied at every step, the symmetry of the roots is built out of r stacked **abelian layers** (Lemma 56.3). Such an action is itself solvable as a group. But the general equation has S_n permuting its roots, and for $n \geq 5$ the group S_n has the non-abelian simple composition factor A_n , hence is *not* solvable — contradiction. In modern terms this is exactly the forward direction *radical* $\Rightarrow$ *solvable Galois group* (Theorem 56.4).

Predecessor. Ruffini (1799) followed the same line, enumerating subgroups of S_5 ; the published proofs had gaps. Abel closed them.

Companion: abelian equations (Abel 1829). If the roots can be ordered so that each $\alpha_{i+1} = \theta_i(\alpha_1)$ for rational maps θ_i that pairwise commute, then f is solvable by radicals. This class is the origin of the term **abelian group**.

Galois (1831): The Criterion

Galois's contribution is the criterion of Slide 4:

f solvable by radicals over $F \iff \text{Gal}(E/F)$ is a solvable group.

Three concepts were created to formulate it.

- **Group of an equation:** permutations of the roots compatible with every rational relation they satisfy — modern $\text{Gal}(E/F)$.
- **Normal subgroup:** $H \trianglelefteq G$ iff $gH = Hg$ for all $g \in G$. Introduced to express recursion: $H \trianglelefteq G$ means the equation factors into an equation with group H and one with group G/H .
- **Solvable group:** a subnormal series with abelian quotients — so the equation reduces step by step to single radical extractions, each a cyclic layer (Slide 7).

Subsumes the earlier results. Abelian Galois group $\Rightarrow$ solvable $\Rightarrow$ solvable by radicals (Abel's abelian equations); the general quintic has group S_5 with composition factor A_5 non-abelian simple, so not solvable, hence not solvable by radicals (Abel's impossibility).