

Last name _____

First name _____

LARSON—MATH 610—CLASSROOM WORKSHEET 36
The Structure of Linear Operators.

Concepts & Notation

- (Sec. 5.3) *permutation*, $\det A$.
- (Sec. 6.2) *characteristic value*, *characteristic vector*, *characteristic polynomial*, *diagonalizable* linear operator.

Review

1. What is the *characteristic polynomial* of a matrix A ? What does it tell us?
2. Does every matrix have characteristic values?
3. What is a *diagonalizable* linear operator T ?

The Structure of a Linear Operator

4. (**Claim:**) Let T be a linear operator on a finite-dimensional space V . Let c_1, \dots, c_k be the distinct characteristic values of T and let W_i be the null space of $(T - c_i I)$. The following are equivalent.
 - (a) T is diagonalizable,
 - (b) The characteristic polynomial for T is:

$$p = (x - c_1)^{d_1} \dots (x - c_k)^{d_k}$$

where $\dim W_i = d_i$ (for $i = 1, \dots, k$),

- (c) $\dim W_1 + \dots + \dim W_k = \dim V$.

5. (**Claim:**) If T is a linear operator from a vector space V over a field \mathbb{F} to itself then $T^2 = T \circ T$ is also in $\mathcal{L}(V)$. And $T^k \in \mathcal{L}(V)$.
6. (**Claim:**) If $T \in \mathcal{L}(V)$ then $cT \in \mathcal{L}(V)$ (for $c \in \mathbb{F}$).
7. (**Claim:**) If $T, T' \in \mathcal{L}(V)$ then $T + T' \in \mathcal{L}(V)$.
8. (**Claim:**) If $T \in \mathcal{L}(V)$ and $p \in \mathbb{F}[x]$ then $p(T) \in \mathcal{L}(V)$.
9. (**Claim:**) If $T \in \mathcal{L}(V)$, $T(\alpha) = c\alpha$ (for $c \in \mathbb{F}$, $\alpha \in V$), and $p \in \mathbb{F}[x]$ then $p(T)(\alpha) = p(c)\alpha$.
10. What does it mean for a polynomial to *annihilate* a linear operator T ?
11. What is the *minimal polynomial* of a linear operator T over a finite-dimensional vector space V ?