

I.9) Principal Components and the Best Low Rank Matrix

The Eckart-Young Theorem states that

$$A_k = \sigma_1 u_1 v_1^T + \dots + \sigma_k u_k v_k^T$$

is the closest rank k matrix to A .

(Here "closest" is with respect to any matrix norm that depends only on the singular values of A , such as the Frobenius norm $\|A\|_F$ or the spectral norm $\|A\|_2$.)

This theorem tells us that k singular vectors explain more of the data than any other set of k vectors: we can choose

$$u_1, \dots, u_k$$

as the basis for the k -dimensional subspace that is closest to the n data points (ie columns of A).

Example:

$$A = \begin{bmatrix} 3 & -4 & 7 & 1 & -4 & -3 \\ 7 & -6 & 8 & -1 & -1 & -7 \end{bmatrix}$$

A is 2×6 :

6 data points in \mathbb{R}^2

Note that the points are centered around


the origin: their

mean is zero. The red line is the span of the singular vector u_1 .

This line minimizes the sum of the squared orthogonal distances from the data points to the line.

(This is different than least squares which minimizes the sum of the squared

vertical distances from the data points



to the line.)

Note that $\{u_1, u_2\}$ are orthonormal.

Thus, column j of A can be written as:

$$a_j = (a_j^T u_1) u_1 + (a_j^T u_2) u_2.$$

$$\text{Then } \|a_j\|_2^2 = |a_j^T u_1|^2 + |a_j^T u_2|^2$$

$$\Rightarrow \sum_{j=1}^n \|a_j\|_2^2 = \sum_{j=1}^n |a_j^T u_1|^2 + \sum_{j=1}^n |a_j^T u_2|^2.$$

The sum of the squared **orthogonal distances** is precisely $\sum_{j=1}^n |a_j^T u_2|^2$ which we can minimize by

maximizing

$$\begin{aligned} \sum_{j=1}^n |a_j^T u_1|^2 &= \sum_{j=1}^n u_1^T a_j a_j^T u_1 = u_1^T \left(\sum_{j=1}^n a_j a_j^T \right) u_1 \\ &= u_1^T (A A^T) u_1. \end{aligned}$$

Which unit vector x maximizes $x^T A A^T x = \|A^T x\|_2^2$? It is the eigenvector of $A A^T$ corresponding to the largest eigenvalue $\lambda_1 = \sigma_1^2$, which is $x = u_1$.

The Statistics Behind PCA

Let $A_0 \in \mathbb{R}^{m \times n}$ be our original data: n points in \mathbb{R}^m . For example, if we measure age, height, and weight of 100 people, then $m=3$ and $n=100$.

	p_1	p_2	...	p_{100}	
age	63	41		21	(years)
height	152	157	...	156	(cm)
weight	48	53		54	(kg)

The mean is $\mu = \frac{1}{n} (A_0 e)$ where $e = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix}$.

sample

Subtracting the mean from each column of A_0 , we get the centered A :

$$A = A_0 - \mu e^T.$$

$$\left[\begin{aligned} \text{Check: } Ae &= A_0 e - \mu(e^T e) \\ &= A_0 e - \frac{1}{n}(A_0 e)n \\ &= A_0 e - A_0 e = 0. \quad \checkmark \end{aligned} \right]$$

The covariance matrix of the data
is ^{sample}

$$S = \frac{1}{n-1} AA^T$$

$$= \frac{1}{n-1} \sum_{j=1}^n a_j a_j^T$$

$$= \frac{1}{n-1} \sum_{j=1}^n (A_{0j} - \mu)(A_{0j} - \mu)^T.$$

The m orthogonal eigenvectors of S are the principal components of A . We compute these principal components by taking the SVD of A .

The total variance is

$$T = \text{trace}(S) = \frac{1}{n-1} (\sigma_1^2 + \dots + \sigma_r^2).$$

The proportion of the total variance explained by the first k principal components is $\frac{\sigma_1^2 + \dots + \sigma_k^2}{\sigma_1^2 + \dots + \sigma_r^2}$ ($k \leq r$).

(numerical demonstration)
