

A Schur-normalization proof for the hyperbolic Cauchy transform

Candidate full solution to Section 3.4(i) of arXiv:math/0204018

Automated Mathematics Run `fa_banach_001`, lane 2

19 July 2026

Status. Candidate full solution to Open Problem (i), subject to human verification. The conclusion applies to the irreducible unitary realizations specified in the question. Reducible parameters are covered sectorwise, with one normalization constant per irreducible sector.

1 Source and exact question

The source is V. V. Kisil, *Spaces of Analytical Functions and Wavelets—Lecture Notes* [1]. It appeared as arXiv:math/0204018 (2002). In Section 3.4, printed page 47 (PDF page 48), item (i) asks:

Demonstrate that Cauchy formula (3.3.23) is an isometry between $F_2(\tilde{T})$ and $H_\sigma(\tilde{D})$ with suitable norms chosen. This almost follows (up to some constant factor) from its property to intertwine two irreducible representations of $SL_2(\mathbb{R})$.

Figure 1 reproduces the source statement.

The Cauchy transform in (3.3.23) is denoted below by

$$W_\sigma : F_2(\tilde{T}) \longrightarrow H_\sigma(\tilde{D}).$$

The source’s Appendix “Boundedness of the Singular Integral Operator W_σ ” treats the four boundary singularities in the principal-value sense and states the resulting L^2 boundedness. The general reduced-wavelet construction proves equivariance, and item (i) explicitly identifies the relevant representations as irreducible. These facts are the only source-specific inputs used below.

2 Definitions and proof intuition

For a unitary representation $\pi : G \rightarrow \mathcal{U}(\mathcal{H})$, its *commutant* consists of bounded operators A satisfying $A\pi(g) = \pi(g)A$ for every $g \in G$. A bounded map $W : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ *intertwines* representations π and ρ if

$$W\pi(g) = \rho(g)W \quad (g \in G).$$

Proof Intuition

Equivariance makes the positive operator W^*W invisible to the group: it belongs to the commutant of the irreducible source representation. Every spectral projection of W^*W is then group invariant. Irreducibility leaves no nontrivial spectral projection, forcing W^*W to be one positive scalar times

where

$$f_p = \int_{\tilde{\mathbb{T}}} tz^{-p} dz f(z).$$

The last integral is similar to the Mellin transform [53, § III.3], [76, Chap. 8, (3.12)], which naturally arises in study of the principal series representations of $SL_2(\mathbb{R})$.

I was pointed by Dr. J. Cnops that for the Cauchy kernel $(-e_1 \mathbf{u} + \mathbf{z})$ there is still a decomposition of the form $(-e_1 \mathbf{u} + \mathbf{z}) = \sum_{j=0}^{\infty} (e_1 \mathbf{u})^j \mathbf{z}^{-j-1}$. In this connection one may note that representations π_1 (3.3.16) and π_σ (3.3.20) for $\sigma = 0$ are unitary equivalent. (this is a meeting point between discrete and principal series). Thus a function theory in $\mathbb{R}^{1,1}$ with the value $\sigma = 0$ could carry many properties known from the complex analysis. $\diamond$

3.4 Open problems

This lecture raises more questions than gives answers. Nevertheless it is useful to state some open problems explicitly.

- (i). Demonstrate that Cauchy formula (3.3.23) is an isometry between $F_2(\tilde{\mathbb{T}})$ and $H_\sigma(\tilde{\mathbb{D}})$ with suitable norms chosen. This almost follows (up to some constant factor) from its property to intertwine two irreducible representations of $SL_2(\mathbb{R})$.
- (ii). Formula (3.3.22) contains Szegő type kernel, which is domain dependent. Integral formula (3.3.23) formulated in terms of analytic kernel. Demonstrate using Stocks theorem that (3.3.23) is true for other suit-

Figure 1: Section 3.4 and the complete text of Open Problem (i) in the source paper [1].

the identity. Thus W already preserves every norm ratio by the same constant. Removing that single constant makes it an isometry. The same identity makes the range closed; equivariance and target irreducibility then force surjectivity.

3 Normalization theorem

Theorem 1 (bounded irreducible intertwiners are scalar unitaries). *Let G be a group, let $\pi : G \rightarrow \mathcal{U}(\mathcal{H}_1)$ and $\rho : G \rightarrow \mathcal{U}(\mathcal{H}_2)$ be irreducible complex unitary representations, and let $0 \neq W \in \mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ satisfy $W\pi(g) = \rho(g)W$ for every $g \in G$. Then there are a unique constant $c > 0$ and a unitary intertwiner $U : \mathcal{H}_1 \rightarrow \mathcal{H}_2$ such that*

$$W = \sqrt{c}U.$$

*Equivalently, $W^*W = cI_{\mathcal{H}_1}$ and $\|Wf\|_{\mathcal{H}_2} = \sqrt{c}\|f\|_{\mathcal{H}_1}$ for all f .*

Proof. Taking adjoints of the intertwining identity, and using unitarity, gives

$$W^*W\pi(g) = W^*\rho(g)W = \pi(g)W^*W \quad (g \in G).$$

Thus the positive operator $A = W^*W$ commutes with $\pi(G)$. By bounded functional calculus, every spectral projection of A also commutes with $\pi(G)$. The range and kernel of such a projection are closed invariant subspaces. Irreducibility therefore makes every spectral projection either 0 or I . Hence the spectrum of A is a singleton and $A = cI$ for some $c \geq 0$. Since $W \neq 0$, we have $c > 0$.

It follows that $U = c^{-1/2}W$ is an isometry, so its range is closed. The range of W (and hence of U) is nonzero and $\rho(G)$ -invariant. Target irreducibility forces this closed range to equal $\mathcal{H}_2$. Therefore U is a surjective isometry, hence unitary. Uniqueness of c follows from $W^*W = cI$. $\square$

Corollary 2 (solution of Section 3.4(i)). *Fix a parameter σ for which the boundary and Hardy realizations in Section 3.3 of [1] are the irreducible unitary representations specified in Open Problem (i), and let W_σ be the bounded nonzero principal-value Cauchy transform (3.3.23). Then there is a constant $c_\sigma > 0$ such that*

$$\|W_\sigma f\|_{H_\sigma(\tilde{D})}^2 = c_\sigma \|f\|_{F_2(\tilde{T})}^2 \quad (f \in F_2(\tilde{T})).$$

Moreover, W_σ maps onto $H_\sigma(\tilde{D})$. With the target norm

$$\|h\|_{H_{\sigma,\text{iso}}} := c_\sigma^{-1/2} \|h\|_{H_\sigma(\tilde{D})},$$

the original Cauchy formula W_σ is an onto isometry.

Proof. The source establishes that W_σ is a bounded intertwiner, while the question specifies irreducibility of the two representations. The Cauchy kernel is not identically zero, so the transform is nonzero. Apply Theorem 1 with $G = SL_2(\mathbb{R})$. $\square$

The normalizing constant can be recovered from any single nonzero test vector:

$$c_\sigma = \frac{\|W_\sigma f\|^2}{\|f\|^2}.$$

The theorem proves that this quotient is independent of the chosen f .

4 Even-Clifford sectors

The source writes parts of the construction using even Clifford-valued functions. If that notation displays the Hilbert realization as an orthogonal sum of irreducible sectors,

$$\mathcal{H}_1 = \bigoplus_{j=1}^m \mathcal{H}_{1,j}, \quad \mathcal{H}_2 = \bigoplus_{j=1}^m \mathcal{H}_{2,j},$$

and $W_\sigma = \bigoplus_j W_{\sigma,j}$ respects the sectors, the same proof gives $W_{\sigma,j}^* W_{\sigma,j} = c_{\sigma,j} I$ on every nonzero irreducible sector. The invariant target norm

$$\|(h_j)_j\|_{\text{iso}}^2 = \sum_j c_{\sigma,j}^{-1} \|h_j\|^2$$

makes the full transform isometric. Thus the Clifford bookkeeping changes only the number of scalar normalizations, not the argument.

5 Verification notes and limitations

1. Proof audit. Boundedness makes W_σ^* available; unitarity is used exactly to move group operators through the adjoint; irreducibility is used twice, first on spectral projections of W^*W and then on the closed range. No density-to-surjectivity shortcut is used.

2. Singular-vacuum issue. The source notes that its vacuum vector is not in L^2 . The proof above does not use that vector as an L^2 element. It starts from the principal-value bounded operator asserted in the source appendix.

3. Reducibility parameters. If a value of σ makes the representation reducible, the single-scalar statement need not hold on the unreduced space. The sectorwise statement above is the correct conclusion. Open Problem (i) itself invokes two irreducible representations, and Corollary 2 matches precisely that scope.

4. What is not computed. The proof determines the normalization intrinsically but does not evaluate c_σ in closed form relative to a preselected geometric normalization of boundary measure. The question asks for suitable norms, so this does not leave a logical gap in the stated isometry problem.

5. Remaining source questions. Section 3.4(ii), concerning domain deformation via Stokes' theorem, and Section 3.4(iii), concerning a principal-series Dirac condition, are not addressed.

6 Bounded novelty check and review recommendation

The run's solution, attempt, and registry indexes were searched for arXiv id 0204018 and the title/keywords. Targeted web/arXiv searches through 19 July 2026 used the exact title and combinations of “Cauchy formula”, “isometry”, “principal series”, “ H_σ ”, and “Kisil”. They found the source paper, the author's earlier paper on principal function theory, and later surveys, but no paper explicitly resolving item (i). Since the argument is the standard commutant proof behind Schur's lemma, the claim of novelty is deliberately modest: this packet records an explicit completion of the source's stated “up to a constant” route.

Human-review recommendation: likely valid. The decisive source check is that the selected σ gives the positive-definite irreducible unitary boundary and Hardy realizations claimed in the question, and that the principal-value transform is nonzero on the selected sector. Once those source hypotheses are confirmed, the normalization proof is complete.

References

- [1] V. V. Kisil, *Spaces of Analytical Functions and Wavelets—Lecture Notes*, arXiv:math/0204018 (2002), especially Sections 2.2, 3.3, 3.4 and the appendix “Boundedness of the Singular Integral Operator W_σ ”.