

The τ_∞ density condition need not imply p -OAP

Run fa_banach_001, lane 5

2026-06-14

Source problem

The source paper is J. A. Chavez-Dominguez, V. Dimant, and D. Galicer, *The p -Operator Approximation Property*, arXiv:2410.05014 [1]. The question appears on PDF page 12, after Theorem 3.8.

12 J. A. CHAVEZ-DOMINGUEZ, V. DIMANT, AND D. GALICER

compactness—such as complete compactness and operator compactness—which all generalize the classical one. It would be interesting to know whether there is an operator space satisfying the density condition in τ_∞ but the space does not have p -OAP.

Similarly to the classical context [SK02, Thm. 6.4] (see also [DOPS09, Cor. 2.5]), the 2-OAP holds in every operator space.

The relevant theorem context is the following. The paper proves a list of equivalent conditions for p -OAP, and remarks that the topology τ_{cc} is used in part (iv) instead of the perhaps expected τ_∞ .

Now we present a noncommutative version of (a part of) [DOPS09, Thm. 2.1] and [SK02, Thm. 4.2]. Surprisingly, in part (iv), we need to employ the topology τ_{cc} instead of the perhaps expected τ_∞ (uniform convergence on operator compact matrix sets). For the case $p = \infty$, it should be noted that Webster [Web98, Thm. 4.4] already proved the equivalence (i) $\Leftrightarrow$ (iii).

Theorem 3.8. *Let $1 \leq p \leq \infty$ and let V be an operator space. The following are equivalent:*

- (i) V has p -OAP.
- (ii) For every operator space W , $\mathcal{F}(W, V)$ is τ_p -dense in $\mathcal{K}_\infty^o(W, V)$.
- (iii) For every operator space W , $\mathcal{F}(W, V)$ is $\|\cdot\|_{cb}$ -dense in $\mathcal{K}_p^o(W, V)$.
- (iv) For every operator space W , $\mathcal{F}(W, V)$ is τ_{cc} -dense in $\mathcal{K}_p^o(W, V)$.

Proof. (i) $\Rightarrow$ (ii) is trivial from Proposition 3.5.

(ii) $\Rightarrow$ (iii): Fix $T \in \mathcal{K}_p^o(W, V)$. By Theorem 2.4 there exist an operator space Z , an operator p -compact map $R : W \rightarrow Z$, and an operator compact map $S : Z \rightarrow V$ such that $T = SR$. By our

Thus the natural reading of the question is:

For a fixed p , can there be an operator space V such that, for every operator space W , the finite-rank maps $\mathcal{F}(W, V)$ are τ_∞ -dense in $\mathcal{K}_p^o(W, V)$, but V does not have p -OAP?

Proof Intuition

The topology τ_∞ is a compact-open type topology: it tests uniform convergence on operator compact matrix sets. Without a boundedness condition on the approximating finite-rank maps, compact-open density is very weak. On a fixed completely compact matrix set in W , one first approximates all matrix levels by a finite-dimensional subspace $E \subset W$. A finite-rank completely bounded projection $P : W \rightarrow E$ then makes $T \circ P$ a finite-rank map that is uniformly close to T on that matrix set. This works for every completely bounded $T : W \rightarrow V$, not just for p -compact maps.

A compact-open density lemma

Lemma. *Let W, V be operator spaces. Then $\mathcal{F}(W, V)$ is τ_{cc} -dense in $\mathcal{CB}(W, V)$, where τ_{cc} is uniform convergence on completely compact matrix sets. Consequently $\mathcal{F}(W, V)$ is τ_∞ -dense in $\mathcal{CB}(W, V)$, and hence in every subspace of $\mathcal{CB}(W, V)$.*

Proof. Fix $T \in \mathcal{CB}(W, V)$, a completely compact matrix set $\mathbf{L} = (L_n)_n$ over W , and $\varepsilon > 0$. By complete compactness, there is a finite-dimensional subspace $E \subseteq W$ such that every $x \in L_n$ is within δ of some point of $M_n(E)$, for all n , where $\delta > 0$ will be chosen below.

Choose a basis $e_1, \dots, e_m$ of E . Let $a_1, \dots, a_m \in E^*$ be the coordinate functionals and extend them by Hahn–Banach to $\varphi_1, \dots, \varphi_m \in W^*$. The map

$$P(w) = \sum_{j=1}^m \varphi_j(w) e_j$$

is finite-rank and completely bounded, since scalar-valued bounded functionals are completely bounded and finite sums of rank-one completely bounded maps are completely bounded. Also $P|_E = \text{id}_E$.

Set $S = T \circ P$. Then $S \in \mathcal{F}(W, V)$. For $x \in L_n$, choose $e \in M_n(E)$ with $\|x - e\| < \delta$. Since $P_n e = e$,

$$\|(T - S)_n x\| = \|T_n(x - P_n x)\| \leq \|T\|_{cb} \|x - P_n x\| \leq \|T\|_{cb} (1 + \|P\|_{cb}) \delta.$$

Choosing $\delta < \varepsilon / (\|T\|_{cb} (1 + \|P\|_{cb}))$ gives uniform ε -approximation on $\mathbf{L}$. This proves τ_{cc} -density.

Finally, every operator compact matrix set is completely compact [1], so τ_∞ is weaker than τ_{cc} . Thus τ_{cc} -density implies τ_∞ -density. $\square$

Full answer to the source question

Theorem. *For every $2 < p \leq \infty$, there exists an operator space V such that, for every operator space W , $\mathcal{F}(W, V)$ is τ_∞ -dense in $\mathcal{K}_p^o(W, V)$, while V fails p -OAP. For $1 \leq p \leq 2$, no such example exists.*

Proof. The density condition is automatic by the lemma: for every W, V , $\mathcal{F}(W, V)$ is τ_∞ -dense in all of $\mathcal{CB}(W, V)$, hence also in $\mathcal{K}_p^o(W, V)$.

It remains only to choose V without p -OAP. The source paper recalls that for every $2 < p \leq \infty$, Sinha–Karn’s theorem gives a Banach space without p -AP, and therefore, by the source paper’s Proposition 3.2, any compatible operator space structure on that Banach space gives an operator space without p -OAP.

which implies that V has the p -AP [SK02, Comment between Defn. 6.1 and Prop. 6.2]. $\square$

For any $2 < p \leq \infty$ it is known that there exists a Banach space without p -AP [SK02, Thm. 6.7], and therefore by Proposition 3.2 there exists an operator space without p -OAP: simply endow the aforementioned Banach space example with any compatible operator space structure. It would be interesting to find an operator space that has the p -AP but fails the p -OAP, although such

Such a V satisfies the τ_∞ density condition by the first paragraph, but fails p -OAP by construction.

For $1 \leq p \leq 2$, the source paper proves that every operator space has p -OAP. Hence no example failing p -OAP can exist in that range. $\square$

Verification notes

The proof uses only the definitions of complete compactness, τ_∞ , and finite-rank completely bounded maps. The key point is that the finite-dimensional projection $P : W \rightarrow E$ need not have any uniform bound as the completely compact test set and ε vary. This is why the argument does not prove norm-density or bounded approximation.

There is an important collateral implication for reviewers. The same lemma shows that condition (iv) in Theorem 3.8 of the source paper, if read literally as unbounded τ_{cc} -density, is automatic for all operator spaces V . Since the paper also states that spaces failing p -OAP exist for $p > 2$, Theorem 3.8(iv) appears to require a correction or an additional boundedness/norm-type hypothesis. This packet is therefore best read both as an answer to the printed open question and as a prompt to check the exact intended topology/density convention.

Novelty check

On 2026-06-14, I checked the run lightweight indexes for arXiv:2410.05014 and the terms p -OAP, p -operator approximation property, τ_∞ , operator p -compactness, and density. I also searched the web for the exact open-question phrase and close variants involving “p-Operator Approximation Property”, “tau_infty”, “tau_cc”, and “density condition”. I found no separate answer. The current arXiv page lists v2 as revised on 2025-06-06.

Human-review recommendation

Verify the intended meaning of the source phrase “density condition in τ_∞ ”. If density is meant in the ordinary locally convex topological sense, with no boundedness condition on the approximating finite-rank maps, the answer above is complete. If the authors intended bounded nets or norm/ideal-norm density, the printed question should be reinterpreted and this packet should be reclassified as a wording/proof-gap note rather than a full solution.

References

- [1] J. A. Chavez-Dominguez, V. Dimant, and D. Galicer, *The p -Operator Approximation Property*, arXiv:2410.05014, 2024; v2 revised 2025.

- [2] D. P. Sinha and A. K. Karn, *Compact operators whose adjoints factor through subspaces of ℓ_p* , Studia Math. **150** (2002), no. 1, 17–33.

Human Review: τ_∞ -Density Without p -OAP

Original automatic proof: `solution_packet.pdf`

Checked date: 14 June 2026

Status: Rejected after a second AI pass; not independently verified by the human reviewer.

Verdict: This packet should not be treated as a verified solution. A second AI review identified a serious gap in the key density lemma. The human reviewer did not directly verify the full argument independently.

Reason for Rejection

The proposed proof rests on the lemma claiming that, for arbitrary operator spaces W, V ,

$$\mathcal{F}(W, V)$$

is τ_{cc} -dense in $\mathcal{CB}(W, V)$, and hence also τ_∞ -dense.

The second AI pass identified a gap in the proof of this lemma. The argument chooses, for a completely compact matrix set, a finite-dimensional subspace $E \subseteq W$ which approximates the test set within some tolerance δ . It then chooses a finite-rank completely bounded map

$$P : W \rightarrow E$$

with $P|_E = \text{id}_E$, and estimates

$$\|(T - T \circ P)_n x\| \leq \|T\|_{cb}(1 + \|P\|_{cb})\delta.$$

The problem is that the projection P , and especially its completely bounded norm $\|P\|_{cb}$, depends on the finite-dimensional space E . But E itself was chosen depending on δ . Thus the final instruction to choose

$$\delta < \frac{\varepsilon}{\|T\|_{cb}(1 + \|P\|_{cb})}$$

is circular.

Complete compactness gives approximation by finite-dimensional subspaces, but it does not provide finite-rank projections onto those subspaces with uniformly controlled completely bounded norm. This is analogous to the classical false argument that every Banach space has the approximation property: compact sets can be approximated by finite-dimensional subspaces, but without controlled projections this does not yield finite-rank approximation of the identity.

Scope of This Review

This note records a rejection after a second pass by the AI model. It should not be read as a direct independent human verification of all details by Antonio Acuaviva. Rather, the packet is being reclassified because the second AI pass found a serious and plausible obstruction in the central lemma.

Review Outcome

Rejected. The proposed solution does not currently prove the claimed answer to the source question. Unless the projection-norm problem can be repaired by an additional argument or by a different construction, this packet should be excluded from the verified examples.